

See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/280727078>

# Plants of the Araceae for malaria and related diseases: a review

RESEARCH · AUGUST 2015

DOI: 10.13140/RG.2.1.4106.2248

---

READS

26

5 AUTHORS, INCLUDING:



**Gina Frausin**

Lancaster University

10 PUBLICATIONS 11 CITATIONS

SEE PROFILE



**Lin Chau Ming**

São Paulo State University

91 PUBLICATIONS 249 CITATIONS

SEE PROFILE



**Adrian Martin Pohlit**

Instituto Nacional de Pesquisas da Amazônia

91 PUBLICATIONS 485 CITATIONS

SEE PROFILE

## Plants of the Araceae for malaria and related diseases: a review

FRAUSIN, G.<sup>1</sup>; LIMA, R.B.S.<sup>1</sup>; HIDALGO, A.F.<sup>1</sup>; MING, L.C.<sup>2</sup>; POHLIT, A.M.<sup>2</sup>

1. Universidade Federal do Amazonas (UFAM), Av. Gal. Rodrigo Octávio, 6200, Coroado I, CEP 69077-000, Manaus-Brazil.

E-mails: ginafrausin@gmail.com, renatabsl@hotmail.com, afreitash@gmail.com;

2. Universidade Estadual de São Paulo (UNESP), R. José Barbosa de Barros, 1780, Campus de Botucatu, CEP 18610-307,

Botucatu-Brazil. E-mail: linming@fca.unesp.br;

3. Instituto Nacional de Pesquisas da Amazonia (INPA), Av. André Araújo, 2936, Petrópolis, CEP 69080-971, Manaus-Brazil.

E-mail: ampohlit@inpa.gov.br.

### **RESUMO: Plantas da família Araceae para a malária e doenças relacionadas: uma**

**revisão.** No presente trabalho realizamos uma revisão das espécies da família Araceae

usadas para tratar malária e seus sintomas. O objetivo foi revelar o grande número de espécies da família usadas no mundo, assim como seu potencial como fontes de produtos

naturais antimaláricos. Foram consultadas as plataformas de busca *SciFinder Scholar*,

*Scielo*, *PubMed*, *ScienceDirect* e *Google books*. Encontramos quarenta e três registros de 36

espécies e 23 generos de Aráceas usadas para malária e seus sintomas. Os generos

neotropicais *Philodendron* Schott e *Anthurium* Schott foram os melhor representados, úteis

para o tratamento da malária, febres, problemas hepáticos e dores de cabeça. Folhas e

tubérculos foram as partes mais utilizadas e decocção foi o método de preparo mais comum.

Os extratos de espécies de Araceae inibem o crescimento *in vitro* do parasito da malária

humana, *Plasmodium falciparum* Welch, e concentrações inibitórias medianas (CI<sub>50</sub>)

significativas foram relatadas para extratos de guaimbê-sulcado (*Rhaphidophora decursiva*

(Roxb.) Schott), aninga (*Montrichardia linifera* (Arruda) Schott), *Culcasia lancifolia* N.E. Br. e

anchomanes do mato (*Anchomanes difformis* (Blume) Engl.), demonstrando o potencial

antimalárico e citotóxico de extratos e subfrações. No único relato sobre os componentes

antimaláricos dessa família, a neolignana polisiforina e o benzoperóxido rafidecurperoxina apresentaram forte inibição das cepas D6 e W2 de *Plasmodium falciparum in vitro* (CI<sub>50</sub> = 368-540 ng/mL). Nenhum estudo sobre a atividade antimalárica *in vivo* em modelo animal foi realizado com espécies da família Araceae. Mais trabalhos biomonitorados pela composição química sobre a atividade antimalárica *in vitro*, assim como estudos *in vivo*, são necessários para aprofundar os conhecimentos sobre potencial antimalárico da família.

**Palavras-chave:** *Anthurium*, *Philodendron*, *Rhaphidophora decursiva*, *Plasmodium falciparum*, planta antimalárica.

**ABSTRACT:** We survey species of the Araceae family traditionally used for malaria and its symptoms. The aim is to reveal the large number of antimalarial Araceae species in use worldwide and their largely unexplored potential as sources of antimalarial natural products. The SciFinder Scholar, Scielo, PubMed, ScienceDirect and Google books search engines were consulted. Forty-three records were found of 36 species and 23 genera of Araceae used for malaria and symptoms. The neotropical genera *Philodendron* Schott and *Anthurium* Schott were the best represented for use in the treatment of malaria, fevers, liver problems and headaches. Leaves and tubers were the parts most used and decoction was the most common preparation method. Extracts of Araceae species inhibit the *in vitro* growth of the human malaria parasite *Plasmodium falciparum* Welch and significant median inhibitory concentrations (IC<sub>50</sub>) for extracts of guaimbê-sulcado (*Rhaphidophora decursiva* (Roxb.) Schott), aninga (*Montrichardia linifera* (Arruda) Schott), *Culcasia lancifolia* N.E. Br. and forest anchomanes (*Anchomanes difformis* (Blume) Engl.) have been reported demonstrating the antimalarial potential and cytotoxicity of extracts and sub-fractions. In the only report on the

antimalarial components of this family, the neolignan polysyphorin and the benzoperoxide raphidecurperoxin exhibited strong *in vitro* inhibition of the D6 and W2 strains of *Plasmodium falciparum* (IC<sub>50</sub> = 368-540 ng/mL). No study on the *in vivo* antimalarial activity in animal models has been conducted on a species of Araceae. More bioguided chemical composition studies on the *in vitro* and also *in vivo* antimalarial activity of the Araceae are needed to further the knowledge of the antimalarial potential of this family.

**Keywords:** *Anthurium*, *Philodendron*, *Rhaphidophora decursiva*, *Plasmodium falciparum*, antimalarial plant.

## INTRODUCTION

Malaria is caused by protozoans of the genus *Plasmodium* that are transmitted to humans by the bite of infected, female *Anopheles* mosquitos. Malaria symptoms include fevers, headaches, vomiting and chills that appear 10-15 days after infection (WHO 2014). Despite advances in its control and prevention in the past decade, malaria continues to be one of the world's major transmissible diseases. It is responsible for high mortality in tropical and sub-tropical regions of the planet (Muñoz et al. 2000; Adebayo & Krettli 2011; Tsabang et al. 2011). Several factors contribute to the unacceptably high malaria morbidity and mortality rates. The major human malaria parasites *Plasmodium falciparum* Welch and *P. vivax* Grassi & Feletti are acquiring resistance to the most important drugs, the quinoline antimalarials and artemisinin derivatives. Also, many ecosystems favor the ready reproduction and propagation of malaria vectors: mosquitos of the genus *Anopheles* Meigen.

In tropical regions where malaria is endemic, alternative therapies based on traditionally used antimalarial plants are used (Milliken 1997a; Willcox et al. 2004; Blair et al.

2005). New drugs introduced into the therapeutic arsenal are mostly derived from natural products (Newman & Cragg 2012). Plants provide secondary metabolites that are useful for the treatment of protozoan diseases such as malaria (Pohlit et al. 2013), leishmaniasis, and African and American trypanosomiasis (Schmidt et al. 2012a, 2012b). In fact, traditionally-used antimalarial plants are the origin of the alkaloid quinine (isolated from species of *Cinchona*) and the sesquiterpene artemisinin (isolated from *Artemisia annua* L.) that gave rise last century to the synthetic quinoline antimalarials (chloroquine, *etc.*) and semi-synthetic artemisinin derivatives (sodium artesunate, *etc.*). These two classes are the basis of artemisinin-combined therapies (ACTs) now used worldwide. Plants also help combat malaria by providing mosquito repellent and insecticidal oils (citronella, neem, *etc.*), solvent extracts and isolated chemicals (chrysanthamic acid, nicotine, *etc.*) that have given rise to the pyrethroid, neonicotinoid and other insecticides and repellents (Pohlit et al. 2011a; 2011b).

The Araceae is one of the most botanically diverse families within the monocotyledons. It is comprised of *ca.* 105 genera and 3,300 species worldwide (Croat 1983; Mayo et al. 1997; Croat 1998; Ribeiro et al. 1999; Coelho 2000; Vargas 2002). The greatest diversity of its species is in tropical America (Croat 1998). Brazil is among the countries having the largest variety of Araceae, numbering 30 genera and 700 species (Mayo et al. 1997).

Over 800 species of Araceae are of economic importance (ornamental, edible, medicinal). For example, about 10% of the world population consumes taro corms (rhizomes of *Colocasia esculenta* (L.) Schott), that represent the most cultivated species of Araceae. It is a foodstuff and foodstock for animals (Pedralli 2002). Most species of Araceae are ornamental plants the most important of which are from the genera *Anthurium* Schott, *Philodendron* Schott, *Dieffenbachia* Schott, *Monstera* Adams and *Zantedeschia* Spreng. (Pedralli 2002). In the Amazon region, many species of Araceae are used for medicinal

purposes, including the treatment of malaria and associated fevers and the most important species are from the genus *Philodendron* Schott (Milliken et al. 1997a, 1997b; Kvist et al. 2006; López et al. 2006). Species of Araceae traditionally used for malaria, fevers, headaches and liver disorders are reviewed. Relevant literature that reveals the antimalarial potential of extracts and isolated compounds, including median inhibitory concentrations (IC<sub>50</sub>) against *Plasmodium falciparum*, are also reviewed.

## **SURVEY METHODS**

This survey was performed mainly on the specialized literature published from July, 2010 until January, 2014, the first record on this topic was in 1977. Searches were performed in the Scifinder Scholar, Scielo, PubMed, Science Direct databases. In general, searches were performed using the terms *Araceae antimalarial plants*, *Araceae fever*, *Araceae malaria* and *Araceae medicinal plants* and library book collections and documents were used at the following institutions: HUAZ Herbarium of the University of Amazonia (Colombia) and the National Institute for Amazon Research (Brazil). The data were compiled and organized in an Excel (Microsoft) spreadsheet containing information on species name, regions where the plant is used, preparation methods, parts of the plants used and where available, information on chemical composition and pharmacological properties.

## **SPECIES OF ARACEAE USED IN THE TREATMENT OF MALARIA**

Forty-three bibliographic sources were found on plants used to treat malaria and symptoms of malaria in the Araceae family. These works describe 38 species in 22 genera (Table 1). *Philodendron* Schott was the most cited genus with seven uses registered by communities in South American countries, including Brazil, Colombia, Ecuador, Peru and

French Guyana (Figure 1). Antimalarial plants of the Araceae family are also found in Central America, Africa, Western Europe, Asia and Southeast Asia.

Twenty-one species of Araceae are specifically used to treat malaria and many are from South America. Thus, the aerial part of folha cheirosa, yeuri cumare (*Anthurium oxycarpum* Poepp.) is used by the Tacana Amerindians of Bolivia (Deharo et al. 2001) and macerates and decoctions of cipo de tara or tracua (*Philodendron cf. linnaei* Kunth) are indicated by the Tirios Amerindians of Suriname (Lopéz et al. 2006). Also, taioba (*Xanthosoma sagittifolium* (L.) Schott) is a widely cultivated tuber that is a foodstuff and foodstock for animals (Aiyeloja et al. 2006) and is used as an antimalarial along the Manso River in Minas Gerais, Brazil (Reskalla 2001). In northern Brazil, an antimalarial aerial root infusion of a species of *Philodendron* is prepared by the native Watorik Yanomami (Milliken 1997). In Togo, Africa, whole plant decoctions of tonflo (*Pistia stratiotes* L.) are used as antimalarials (Kyei et al. 2012).

## **ARACEAE SPECIES FOR FEVERS, HEADACHES AND OTHER SYMPTOMS OF MALARIA**

Several plants have uses both in the treatment of malaria and also in the the treatment of the symptoms of malaria. Thus, whole plant decoctions of *P. stratiotes* are used in Togo to treat fevers and malaria infections (Lahitte et al. 1998; Kyei et al. 2012).

Some species of Araceae are not used specifically to treat malaria, but are used to treat symptoms often associated with malaria such as fevers and headaches. So, the indigenous Wayãpi in French Guyana treat fevers with a decoction of Tapi`Ykũ (*Philodendron linnaei* Kunth) (Grenand et al. 1987). Similarly, the Tukano of southeast Colombia treat headaches by placing fresh inflorescences of madona lily (*Spathiphyllum floribundum* (Linden

& André) N.E. Br.) on the forehead (Croat 1994). *Culcasia lancifolia* N.E. Br. has been used by traditional healers to treat headaches, fevers and vomiting (Lekana-Douki et al. 2011).

## **TRADITIONAL METHODS OF PREPARATION OF REMEDIES**

Decoction was the method of preparation for antimalarial remedies based on species of Araceae. Leaves and the tubercles were the parts most often cited. Few details on the method of preparation are included in many reports (Table 1). However, some methods of preparation and treatments are especially interesting. Three slices of *hoa pouk*, also called *habarala* (*Alocasia macrorrhizos* (L.) G. Don) is boiled with 3 pieces of sugar cane and 7 seeds of unpolished rice in the Province of Khammouane, Laos. A non-malarious person who drinks the decoction is said to become itchy whereas someone with malaria is said to feel good and is then expected to continue treatment with the decoction (Shirayama et al. 2006). On the Pacific Coast of Colombia, the leaves of *mano*, also known as *guaco de mata* or *tres dedos* (*Anthurium cf. tridigitatum* Engl.) are macerated and rubbed on the body as a rheumatic treatment for malaria (Blair et al. 2005). Similarly, the Achuar Jívaro of Peru apply hot leaves of a species of *Monstera* on the skin to relieve liver pain (Lewis et al. 1977, cited by Croat 1994).


Therapeutic preparations often involve species from Araceae and other families. *Corazón de Jesús* or *Jesus's heart* (*Caladium bicolor* Vent.) leaves are boiled in 1 L of water until half the volume evaporates and then lemon is added (*Citrus × limon* (L.) Osbeck). The resulting liquid is allowed to sit to catch the morning dew and consumed on an empty stomach for 9 consecutive mornings in the treatment of inflamed liver (Blair et al. 2005).

## **DISTRIBUTION OF ARACEAE SPECIES USED FOR MALARIA AND RELATED SYMPTOMS**



Species of Araceae are used for the treatment of malaria and its symptoms throughout the tropical regions of the world (Figure 1). Poverty and lack of access to health services in many countries make these plants an important alternative for the treatment of malaria. Reports from the African nations of Ivory Coast, Kenya, Benin, Gabon and Togo have revealed eight antimalarial species belonging to the genera *Amorphophallus* Blume ex Decne, *Anchomanes* Schott, *Culcasia* P. Beauv., *Homalomena* Schott, *Pistia* L. and *Pothos* L. In the Amazon region, the largest numbers of Araceae species are used as antimalarials. The neotropical genera *Philodendron* Schott and *Anthurium* Schott are used by these Amerindian ethnic groups: Yanomami (Brazil), Tirios (Suriname), Wayãpi (French Guyana), Makuna and Mirañá (Colombia) and Secoya and Tacana, and by African descendants (Pacific Coast, Colombia). Interestingly, no genus of antimalarial Araceae was reportedly used on both the African and American Continents thus providing evidence for the local distribution of the species used.

**FIGURE 1.** World locations where species of Araceae are used to treat malaria and its symptoms.



Note: An orange colored point indicates where an Araceae species has been reported as being used to treat malaria or its symptoms. This map was generated based on the countries represented in Table 1.

### **ANTIPLASMODIAL ACTIVITY AND TOXICITY OF SPECIES OF ARACEAE**

Extracts of plants of the Araceae exhibit antimalarial potential and also cytotoxicity. The hexane and ethanol extracts of the stems of the aquatic plant aninga (*Montrichardia linifera* (Arruda) Schott) were inactive *in vitro* against *P. falciparum* Dd2 strain. However, the dichloromethane fraction of the ethanol extract exhibited high antiplasmodial activity ( $IC_{50} < 10 \mu\text{g/mL}$ ) and toxicity to *Artemia salina* (Amarante et al. 2011). *M. linifera* from Belém, Brazil exhibited leaf ethanol extracts with no toxicity to *A. salina* ( $DL_{50} > 500 \mu\text{g/mL}$ ) and moderate inhibition of *P. falciparum* W2 strain ( $IC_{50} = 11.7 \mu\text{g/mL}$ ) (Costa et al. 2009).


In other work, dichloromethane root extracts of *Culcasia lancifolia* inhibited the growth of FCB and W2 strains of *P. falciparum* ( $IC_{50} = 8.9$  and  $10.0 \mu\text{g/mL}$ , respectively) and exhibited toxicity to MRC-5 cells. The root methanol extract exhibited moderate antiplasmodial activity ( $IC_{50} = 25.0$  and  $16.0 \mu\text{g/mL}$  against FCB and W2 strains of *P. falciparum*, respectively) (Lekana-Douki et al. 2011). Root extracts of forest anchomanes (*Anchomanes difformis* (Blume) Engl.) failed to inhibit ( $IC_{50} > 100 \mu\text{g/mL}$ ) the 3D7 strain of *P. falciparum* and were cytotoxic (Bero et al. 2009).

Water and ethanol extracts of the leaves of water lettuce (*Pistia stratiotes* L.) were apparently well-tolerated and exhibited antiarthritic and antipyretic effects in formalin-induced arthritis and LPS-induced fever in Sprague-Dawley rats (Kyei et al. 2012). On the other hand, the sap of *Caladium bicolor* is toxic and can even produce asphyxia (Flores et al. 2001). Isolation of the antiplasmodial and cytotoxic components from the extracts of plants of the Araceae is necessary for a better understanding of the medicinal potential of this family.

## CHEMISTRY OF ANTIMALARIAL SPECIES OF ARACEAE

Oxalic acid is frequently deposited as crystals of calcium oxalate (Figure 2) in plants of the Araceae and is responsible for the toxicity of some genera (*e.g. Dieffenbachia* Schott). A large variety of anthocyanines have been identified in the flowers, fruit, leaves and leaf stems of 59 species of Araceae (Williams et al. 1981). The most commonly occurring pigment in these species was cyanidin 3-*O*-rutinoside, however, pelargonidin 3-rutinoside is also regularly found in the Araceae.

Leaves and branches of guaimbê-sulcado (*Rhaphidophora decursiva* (Roxb.) Schott) were extracted with methanol. The dry extract was defatted with hexanes. The chloroform soluble fraction of this extract was evaporated and chromatographed on silica gel using an acetone-chloroform gradient. Further normal phase, flash or reverse-phase chromatographies on the resulting fractions provided 14 compounds. Six of these compounds exhibited *in vitro* antiplasmodial activity (Zhang et al. 2001). The most active against *P. falciparum* D6 and W2 strains were the neolignan *threo*-polysyphorin (IC<sub>50</sub> = 404 and 368 ng/mL, respectively; Figure 2) and the benzoperoxide raphidecurperoxin (IC<sub>50</sub> = 540 and 420 ng/mL, respectively; Figure 2). The neolignans raphidecursinol A and B and lignans grandisin and epigrandisin were less active. To our knowledge, the above is the only report on antimalarial components from the Araceae.

Figure 2. Antiplasmodial chemical constituents of *Rhaphidophora decursiva*.

## FINAL REMARKS

A number of South American Araceae species are traditionally used in the treatment of malaria and its symptoms. Despite the *in vitro* inhibitory activity of extracts, fractions and isolated constituents against the human malaria parasite *Plasmodium falciparum* we found no *in vivo* antimalarial study in animal models for this family. Studies on the *in vitro* and *in vivo* antimalarial activity are needed to further explore the antimalarial potential of the Araceae.

## ACKNOWLEDGEMENTS

The authors thank the Herbarium at the Universidade de La Amazonia in Colombia for the use of bibliographic materials. Funding was provided by grants from the Brazilian National

Council for Scientific and Technological Development (CNPQ, National Malaria Network and Bionorth Program), the Amazonas State Research Support Foundation (FAPEAM, PRONEX).

G. F., R. B. S. L. and A. M. P. would like to recognize the following scholarships received from CNPq: DTI, GD (383557/2010-0), (554317/2010-9) and PQ (311.649/2011-4), respectively.

## REFERENCES

ADAMS, M. *et al.* Malaria in the renaissance: Remedies from European herbals from the 16<sup>th</sup> and 17th century. **Journal of Ethnopharmacology**, v.133 (2), p. 278-288, 2011.

ADEBAYO, J.O.; KRETTLI, A.U. 2011. Potential antimalarials from Nigerian plants: a review. **Journal of Ethnopharmacology**, v.133, n.2, 27, p. 289-302, 2011.

AIYELOJA A. A; BELLO O.A. Ethnobotanical potentials of common herbs in Nigeria: a case study of Enugustate. **Educational Research and Review**, v.1, n.1, p.16-22, 2006.

ALMEIDA, R.N. *et al.* Plants with central analgesic activity. **Phytomedicine**, v.8, n.4, p.310-322, 2001.

AMARANTE *et al.* Estudo fitoquímico biomonitorado pelos ensaios de toxicidade frente à *Artemia salina* e de atividade antiplasmodica do caule de aninga (*Montrichardia linifera*). **Acta Amazonica**, v.41, n.3, p.431-43, 2011.

ATINDEHOU, K., SCHMID, C., BRUN, R., KONÉ, M.W. TRAORE, D. Antitrypanosomal and antiplasmodial activity of medicinal plants from C<sup>o</sup>te d'Ivoire. **Journal of Ethnopharmacology**, v.90, p. 221–227. 2004.

BENSKY, D.; GAMBLE, A. **El chino medica medicina herbal materia**, Seattle (USA): Eastland Press Inc., 1993. Cited by WILLCOX, M. *et al.* **Traditional herbal medicines for modern times-traditional medicinal plants and malaria**. Boca Raton (USA): CRC press, 2004. 508p.

BERO, J. *et al.* *In vitro* antiplasmodial activity of plants used in Benin in traditional medicine to treat malaria. **Journal of Ethnopharmacology**, v.122, n.3, 21, p.439 – 444, 2009.

BLAIR, S.; MADRIGAL, B. **Plantas antimaláricas de Tumaco: Costa Pacífica colombiana**. 1 ed., Medellín: University of Antioquia, 2005. 347p.

BRANDÃO, M. *et al.* Antimalarial experimental chemotherapy using natural products. **Ciência e Cultura**, v.37, p.1152-1163, 1985.

CHINCHILLA-CARMONA, M. *et al.* Evaluación *in vivo* de la actividad antimalarica de 25 plantas provenientes de una reserva de conservación biológica de Costa Rica. **Revista Chilena de Historia Natural**, v.84, p.115-123, 2011.

COELHO, N. M. A. *Philodendron* Schott (Araceae): Morfología e taxonomía das especies da Reserva da Macae da Cima – Nova Friburgo, Rio de Janeiro, Brasil. **Rodriguesia**, v. 51, n.78/79, p.21-68, 2000.

COSTA, E.S.S. *et al.* Estudos farmacognósticos, fitoquímicos, atividade antiplasmódica e toxicidade em *Artemia salina* de extrato etanólico de folhas de *Montrichardia linifera* (Arruda) Schott, Araceae. **Revista Brasileira de Farmacognosia**, v.19, n.4, p.834-838, 2009.

CROAT, T. B. A revision of the genus *Anthurium* (Araceae) of Mexico and Central America. Part. I: Mexico and middle America. **Annals of Missouri Botanical Garden**, v.70, n.2, p.211-416, 1983.

CROAT, T.B. The use of the new world Araceae as drug plants. **Journal of Japanese Botany**, v.69, n.4, p.185-203, 1994.

CROAT, T.B. History and current status of systematic research with Araceae. **Aroideana**, v.21, p. 26-145, 1998.

- DEHARO, E. *et al.* A search for natural bioactive compounds in Bolivia through a multidisciplinary approach. Part V. Evaluation of the antimalarial activity of plants used by the Tacana Indians. **Journal of Ethnopharmacology**, v.77, p.91-98, 2001.
- EDIRISINGHE, S. **Plant Based Antimalarials of the Indigenous Herbal Traditions of Sri Lanka**. Sri Lanka: National Science Foundation, 1999. 23p.
- FLORES, J.S. *et al.* Plantas de la flora yucatanense que provocan alguna toxicidad en el humano. **Revista Biomédica**, v.12, p.86-96, 2001.
- GOSLING, D. **Religion and ecology in India and Southeast Asia**. London: Routledge, 2001. 215p.
- GRENAND, P. *et al.* **Pharmacopées traditionnelles en Guyane: Créoles, Wayãpi, Palikur**. Paris: Orstom, 1987. 569p.
- KOUDOUVO, K. *et al.* An ethnobotanical study of antimalarial plants in Togo Maritime Region. **Journal of Ethnopharmacology**, v.134, p.183-190, 2011.
- KULIP, J. An ethnobotanical survey of medicinal and other useful plants of Muruts in Sabah, Malaysia. **Telopea**, v.10, n.1, p.81-98, 2003.
- KVIST, L.P. *et al.* Identification and evaluation of Peruvian plants used to treat malaria and leishmaniasis. **Journal of Ethnopharmacology**, v.106, p.390-402, 2006.
- KYEI, S. *et al.* The efficacy of aqueous and ethanolic leaf extracts of *Pistia stratiotes* Linn. in the management of arthritis and fever. **Journal of Medical and Biomedical Sciences**, v.1 n.2, p.29-37, 2012.
- LAHITTE, H.J. *et al.* **Plantas medicinales Rioplatenses**. Buenos Aires: Editorial L.O.L.A., 1998, 240p.

LEAMAN, D. A. *et al.* Malaria remedies of the Kenyah of the Apo Kayan, East Kalimantan, Indonesian Borneo: a quantitative assessment of local consensus as an indicator of biological efficacy. **Journal of Ethnopharmacology**, v.49, p.1-16, 1995.

LEKANA-DOUKI, J. B. *et al.* *In vitro* antiplasmodial activity and cytotoxicity of nine plants traditionally used in Gabon. **Journal of Ethnopharmacology**, v.133, p.1103-1108, 2011.

LENSE, O. Biological screening of selected traditional medicinal plants species utilized by local people of Manokwari, West Papua Province. **Musantara Biociencia**, v.3, n.3, p.145-150, 2011.

LÓPEZ, R. *et al.* **Manual de identificación de especies no maderables del corregimiento de Tarapacá**. Bogotá: Instituto Amazónico de Investigaciones Científicas (SINCHI) – GTZ (Germany), 2006. 120p.

MARLES, R.J. *et al.* A contribution to the ethnopharmacology of the lowland quichua people of amazonian Ecuador. **Revista da Academia Colombiana de Ciencias Exactas, Físicas e Naturales**, v.16, n.63, p.111-120, 1988.

MAYO, S. J. *et al.* **The genera of Araceae**. Kew (UK): Royal Botanic Garden, 1997. 370p.

MILLIKEN, W. Traditional antimalarial medicine in Roraima, Brazil. **Economic Botany**, v.51, n.3, p.212-237, 1997a.

MILLIKEN, W. **Plants for malaria plants for fever: medicinal species in Latin America - a bibliographic survey**. Kew (UK): Royal Botanic Garden, 1997b. 116p.

MUÑOZ, V. *et al.* A search for natural bioactive compounds in Bolivia through a multidisciplinary approach Part I. Evaluation of the antimalarial activity of plants used by the ChacoboIndians. **Journal of Ethnopharmacology**, v.69, p.127-137, 2000.

NEWMAN, D.J.; CRAGG, G.M. Natural products as sources of new drugs over the 30 years from 1981 to 2010. **Journal of Natural Products**, v.75, p.311-335, 2012.



PEDRALLI. Dioscoreaceae e Araceae: aspectos taxonômicos, etnobotânicos e espécies nativas com potencial para melhoramento genético. In: II Simpósio Nacional sobre as Culturas do Inhame e do Taro. João Pessoa: EMEPA-PB, v.2, 2002. 234 p.

POHLIT, A.M. *et al.* Patent literature on mosquito repellent inventions which contain plant essential oils-a review. **Planta Medica**, v.77, p.598-617, 2011a.

POHLIT, A.M. *et al.* Plant extracts, isolated phytochemicals, and plant-derived agents which are lethal to arthropod vectors of human tropical diseases-A review. **Planta Medica**, v.77, p.618-630, 2011b.

POHLIT, A.M. *et al.* Amazonian plant natural products: perspectives for discovery of new antimalarial drug leads. **Molecules**, v.18, p.9219-9240, 2013.

PRAKASH, K. C. Ethnomedicinal botany of the Apatani in the Eastern Himalayan region of India. **Journal of Ethnobiology and Ethnomedicine**, v.1, p.1-8, 2005.

PRAVAKAR-PADHIAL. **Arum Family (Araceae): Taro**, in:

<http://agriorissa.blogspot.co.uk/2011/10/arum-family-araceae-taro.html>

Access on 9/10/2011.

RAHMATULLAH, M. *et al.* A survey of medicinal plants in two areas of Dinajpur district, Bangladesh including plants which can be used as functional foods. **American-Eurasian Journal of Sustainable Agriculture**, v.3, n.4, p.862-876, 2009.

RESKALLA, A. **Inhame, saúde como alimento e lucro como produto**. Belo Horizonte: Estado de Minas/Agropecuário, p. 6-7, 2001. Cited by: PEDRALLI, G. Dioscoreaceae e Araceae: aspectos taxonômicos, etnobotânicos e espécies nativas com potencial para melhoramento genético. In: II Simpósio Nacional sobre as Culturas do Inhame e do Taro. João Pessoa: EMEPA-PB, v.2, 2002. 234 p.

RIBEIRO, J. E. *et al.* **Flora da Reserva Ducke**. Guia de identificação das plantas vasculares de uma floresta de terra-firme na Amazônia Central. Manaus: INPA/DFID, 1999. 816p.

SCHMIDT, T.J. *et al.* The potential of secondary metabolites from plants as drugs or leads against protozoan neglected diseases-part I. **Current Medicinal Chemistry**, v.19, p.2128-2175, 2012a.

SCHMIDT, T.J. *et al.* The potential of secondary metabolites from plants as drugs or leads against protozoan neglected diseases-part II. **Current Medicinal Chemistry**, v.19, p.2176-2228, 2012b.

SCHULTES, R. E.; RAFFAUF, R.F. **The healing forest: medicinal and toxic plants of the northwest Amazonia**. Portland (USA): Dioscorides Press, 1990. 484p.

SCHULTES, R. E.; RAFFAUF, R.F. **De plantis toxicariis e mundo novo tropicale commentationes XXXIX febrifuges of northwest Amazonia**, 1994. 68p.

SHIRAYAMA, Y. *et al.* Modern medicine and indigenous health beliefs: malaria control alongside “sadsana-phae”(animist belief system) in LAO PDR. **Southeast Asian Journal of Tropical Medicine and Public Health** v.37, p.622-629, 2006.

TRAN, Q. *et al.* *In vitro* antiparasitic activity of antimalarial medicinal plants used in Vietnamese traditional medicine. **Journal of Ethnopharmacology**, v.86, n.2-3, p.249-252, 2003.

TSABANG, N. *et al.* Ethnopharmacological survey of Annonaceae medicinal plants used to treat malaria in four areas of Cameroon. **Journal of Ethnopharmacology**, v.139, p.171-180, 2011.

TSHIBANGU, N.J. *et al.* Screening of African medicinal plants for antimicrobial and enzyme inhibitory activity. **Journal of Ethnopharmacology**, v.80, p.25-35, 2002.

VARGAS, W.G. **Guía ilustrada de las plantas de las montañas del Quindío y los andes centrales**. Manizales (Colombia): Centro Editorial Universidad de Caldas, 2002. 813p.

World Health Organization (WHO). **Tratamiento del paludismo grave**. Manual practico. 3ed. Geneva (Switzerland): WHO, 2014. 83p.

WILLCOX, M. *et al.* **Traditional herbal medicines for modern times-traditional medicinal plants and malaria**. Boca Raton (USA): CRC press, 2004. 508p.

WILLIAMS, C.A. *et al.* Anthocyanin pigments and leaf flavonoids in the family Araceae. **Phytochemistry**, v.20, n.2, p.217-234, 1981.

ZHANG, H. *et al.* Antimalarial compounds from *Rhaphidophora decursiva*. **Journal of Natural Products**, v.64, p.772-777, 2001.

**TABLE 1.** Species of the Araceae family used in the treatment of malaria and malaria symptoms.

Species	Local name	People, Region where used	Use	Part used	Preparation	Source
<i>Alocasia macrorrhizos</i> (L.) G. Don (APGIII 2009) [Name cited: <i>Alocasia macrorrhiza</i> (L.) Schott]	Habarala; Hoa Pouk	Sri Lanka; Bourapar District, Khammouane Province LAO	Malaria	Tuber	Boil 3 slices with 3 parts sugar cane and 7 whole grains of rice, drink.	Edirisinghe, 1999; Shirayama et al. 2006
<i>Amorphophallus bequaertii</i> De Wilde	Mbandakabiri, Ikomalyakabiri	Africa	Malaria, fever	Tuber	Place fresh tuber in hot water, filter, use filtrate for rectal injection	Tshibangu et al. 2002
<i>Anchomanes difformis</i> (Blume) Engl.	forest anchomanes	Benin, Côte d'Ivoire	Malaria, analgesic	Leaves, Rhizomes/Roots	-	Adjanohoun et al. 1989 cited in Bero et al. 2009; Almeida et al. 2001; Atindehou et al. 2004
<i>Anthurium</i> sp.		Ingano Macoa Amerindians, Mocoa, Putumayo, Colombia	Fever sores	Sap	Caustic sap is applied directly on fever sores	Schultes & Raffauf 1990
<i>Anthurium uleanum</i> Engl.		Secoya Amerindians, Amazonia	Headaches	Roots	Decoction	Schultes & Raffauf, 1990 (based on Vickers 220); Croat, 1994
<i>Anthurium oxycarpum</i> Poeppig	folha cheirosa; yeuri cumare	Indigenous Tacana, Bolivia	Malaria	Aerial parts	-	Deharo et al. 2001
<i>Anthurium</i> cf. <i>tridigitatum</i>	Mano, guaco de mata, tres dedos, corrientoso, deshinchador	Costa pacífica, Colombia	Rheumatic pain caused by malaria	Leaves	Macerated leaves rubbed over whole body	Blair et al. 2005
<i>Arisaema triphyllum</i> (L.) Schott	-	Menomini and Iroquois tribes	Headaches	-	-	Croat, 1994
<i>Arum maculatum</i> L.	Wild arum	Europe	Malaria	-	-	Zwinger, 1696, cited by Adams 2011

<i>Caladium bicolor</i> Vent.	Pana, corazón de Jesús	Costa pacífica, Colombia	Swollen liver, liver tonic	Leaves	Boil 5-6 leaves in 1 L of water until ½ volume evaporated. Add lemon, allow to sit in morning dew. Drink 9 mornings before eating	Blair et al. 2005
<i>Colocasia affinis</i> Schott		Eastern Himalayan India	Febrifuge	Leaves		Prakash Kala, 2005
<i>Colocasia esculenta</i> (L.) Schott	Taro (saru)	Brazil	Malaria		-	Pravakar Padhial, 2011
<i>Culcasia lancifolia</i> N.E. Br.	-	Kele, Gabon	Fever		-	Lekana-Douki, et al. 2011
<i>Cyrtosperma</i> <i>johnstonii</i> N.E. Br.	-	India	Headaches	Leaves, flowers, roots/rhizome or whole plant	-	Gosling, 2001
<i>Dieffenbachia</i> sp.	Patiquina	Loreto, Perú	Malaria	Leaves	-	Kvist et al. 2006
<i>Homalomena</i> <i>propingna</i> [presumably <i>Homalomena</i> <i>propinqua</i> Schott]	Nyato	Muruts, Sabah, Malaysia	Chills	-	Heated, placed on forehead	Kulip, 2003
<i>Homalomena rubra</i> Hassk.	Lung bala	Kenya	Splenomegaly, malaria, rubefacient	Roots	Decoction	Leaman et al. 1995
<i>Lasia spinosa</i> (L.) Thwaites	-	Vietnam	Malaria	Whole plant	-	Tran et al. 2003
<i>Monstera</i> sp.	-	Achuar Jívaro, Peru	Liver pain	Leaves	Hot leaves applied to skin over the liver to alleviate pain	Lewis et al.1977. cited by Croat 1994
<i>Monstera deliciosa</i> Liebm.	-	Brazil	Malaria	-	-	Brandão et al. 1985
<i>Monstera spruceana</i> (Schott) Engl.	-	Achuar Jívaro, Peru	Liver	Leaves	Hot leaves applied to skin over the liver to alleviate pain	Lewis et al. 1977 cited by Croat 1994

<i>Philodendron</i> sp.	-	Indigenous Watorik Yanomami, Brazil	Malaria	Roots	Infusion of aromatic, aerial roots, oral	Milliken, 1997a
<i>Philodendron</i> sp.		Quichua, Ecuador	Hepatitis			Marles et al.1988
<i>Philodendron</i> cf. <i>deflexum</i> Poepp. ex Schott	Taracua; bejuco agraz, meonomisi (Makuna), pjimacu (Yukuna); huncageme (Huaorani), u'cu (Secoya), ya'i (Siona); chuhudaek, itininga, pantorrilla	Brazil, Colombia (Indigenous Makuna), Ecuador, Peru	Fever	Roots	Softer pendant vines are macerated and drunk in cold water for fever	López et al. 2006
<i>Philodendron</i> <i>fragantissimum</i> (Hook.) Kunth	-	Tropical Americas	Liver pain relief	Leaves	Cold poultice of the plant placed over the liver, for pain. Leaves may also be heated	Lewis et al. 1977, cited by Croat 1994
<i>Philodendron</i> cf. <i>linnaei</i> Kunth	Cipo de tara, tracua; bejuco quemador; orejita de venado, ellapapar	Brazil, Colombia, Peru, Suriname (indigenous Tirios)	Malaria	Whole plant	Decoction of macerated plant for washings	López et al. 2006
	Tapi`Ykū	Wayapi, French Guiana	Fever associated with tabu	Leaves	Decoction for external washings	Grenand et al. 1987
<i>Philodendron</i> cf. <i>uleanum</i> Engl.	-	Mirafña, Colombia	Malaria, fever	-	Infusion	Schultes & Raffauf, 1994
<i>Pinellia ternata</i> (Thunb.) Ten. ex Breitenb.	-	-	Intense fever	Rhizome	In mixture with <i>Artemisia annua</i>	Bensky et al. 1993
<i>Pistia stratiotes</i> L.	Tonflo	Maritime Region, Togo	Malaria	Whole plant	Decoction, taken orally	Koudouvo et al. 2011
<i>Pothos ovatifolius</i> Engl.	Aka malung	Kenya	Splenomegaly, malaria	Leaves	Poultice	Leaman et al. 1995
<i>Rhaphidophora</i> <i>decursiva</i> (Roxb.) Schott	- Guaimbê-sulcado	Brazil	Malaria	-	-	Zhang et al. 2001

<i>Rhaphidophora pertusa</i> (Roxb.) Schott		Manokwari, (West Papua Province), Merdey, Wasior, Indonesia	Liver diseases	Leaves	-	Lense, 2011
<i>Scindapsus officinalis</i> (Roxb.) Schott	Takbir gach	Bangladesh	Fever, pain	Leaves	Powdered leaves are drunk for 21 days	Rahmatullah et al. 2009
<i>Spathiphyllum floribundum</i> (Linden & André) N.E. Br.	Madonna Lily	Indigenous Tukano and Gwanano (near the Papurf River), Southeast Colombia	Headaches	Fresh inflorescences	Placed directly on forehead	Croat, 1994
<i>Symplocarpus foetidus</i> (L.) W. Salisb.	-	Micmac Indians	Headaches	Leaves	Inhalation of volatile oils from ground leaves	Lewis and Elvin 1977, cited by Croat 1994
<i>Xanthosoma atrovirens</i> K. Koch & C.D. Bouché	-	-	For swollen liver after bouts of swamp or malaria fever	Leaves	Fresh leaves applied to body	Peckolt 1893 and Uphof 1959, cited by Croat 1994.
<i>Xanthosoma sagittifolium</i> (L.) Schott	Taioba, taiá	Rio Manso-MG, Brazil	Malaria	-	-	Reskalla, 2001, cited by Pedralli, 2002
<i>Xanthosoma undipes</i> (K. Koch & C.D. Bouché) K. Koch	Tall elephant's ear	Costa Rica	Malaria	-	-	Chinchilla-Carmona et al. 2011

---