

# PROCEEDINGS B

## On potential ocular artifacts in infant EEG: A reply to comments by Köster

Journal:	<i>Proceedings B</i>
Manuscript ID	RSPB-2016-1285
Article Type:	Invited Reply
Date Submitted by the Author:	08-Jun-2016
Complete List of Authors:	Kampis, Dora; Central European University, Parise, Eugenio; Lancaster University, Department of Psychology; Lancaster University, Flyde College Csibra, Gergely; Central European University, Department of Cognitive Science; University of London, Department of Cognitive Science Kovács, Ágnes; Central European University,
Subject:	Cognition < BIOLOGY, Neuroscience < BIOLOGY
Keywords:	EEG, infants, gamma-band oscillations, microsaccades
Proceedings B category:	Neuroscience

SCHOLARONE™  
Manuscripts

1           On potential ocular artifacts in infant EEG: A reply to comments by Köster  
2   Kampis, D., Parise, E., Csibra, G., & Kovács, Á.M.

3

4

5

6           Köster's comment on Kampis, et al. [2] adopts an objection that was put  
7 forward previously regarding the interpretation of scalp-recorded gamma-band  
8 EEG activity in adults as a correlate of object processing. Gamma-band (over 25  
9 Hz) oscillatory activity has been consistently found to signal object processing in  
10 various populations, such as non-human primates, human adults, and human  
11 infants. However, Yuval-Greenberg and colleagues [3] reported that in human  
12 adults saccadic spike potentials (SP), co-occurring with micro-saccades (MS),  
13 contribute to this signal, and questioned the neural origins of the oscillatory  
14 activation found in earlier studies. In response to this, specific tools have been  
15 developed to remove possible MS-related artifacts from adult EEG data (e.g.,  
16 Hassler et al. [4]).

17

18           Köster [1] points out that analogous attempts have not been implemented in  
19 infancy research. We argue that while this is indeed the case, there are several  
20 theoretical and methodological considerations that cast doubt on whether it is  
21 necessary or possible to apply these tools to infant EEG recordings.

22

23           First, the algorithm applied on adult EEG to remove MS-related artifacts  
24 would not be applicable to infant recordings as it is. Hassler et al. [4] propose a  
25 two-step method, which consists of detecting and then removing SPs that

26 accompany MSs. The first step of this method detects SPs based on their  
27 characteristics in adult EEG. However, Csibra et al. [5] found no saccade-related  
28 SPs in infants younger than 12 months, and even at this age SPs differed greatly  
29 in amplitude and in morphology from those reported in adults. Because of this,  
30 the algorithms used with adults to detect SPs would simply not be applicable to  
31 infant EEG. The second step of Hassler et al [4], using independent component  
32 analysis (ICA) to remove MS-related SPs from the signal, also seems unfeasible to  
33 apply directly on infant data given the nature of infant EEG recordings. As Köster  
34 [1] rightly points out, performing ICA requires a vast amount of data to produce  
35 valid results. As an estimate, finding  $N$  stable components in  $N$ -channel data  
36 requires more than  $3 \cdot N^2$  sample points at each channel [6]. In EEG recordings  
37 at 128 channels and 500 Hz sampling rate (like in our study) this requirement  
38 demands more than 90 seconds of perfectly clean EEG on *all* channels. In most  
39 infant EEG studies (especially ones with relatively longer trials and dynamic  
40 stimuli), recordings are regularly contaminated by movement artifacts, and the  
41 cleaned data are much sparser than what might be required by ICA.

42

43 Furthermore, to our knowledge no one has managed to identify and measure  
44 MSs in infants so far, and therefore it is not known in what form they occur at  
45 this early age. While the appropriate tools are available (eye-trackers with a high  
46 enough sampling rate), it would be a separate methodological challenge to keep  
47 young infants' head sufficiently stable for accurately measuring MSs. Therefore  
48 even in case of successful co-recording of EEG and eye-movements, it is unclear  
49 how MSs (and/or SPs) should be detected. Because of this, at the moment it is  
50 not possible to remove any potential MS-related artifacts from infant EEG, and

51 we agree with Köster [1] that we cannot decisively exclude the possibility that  
52 microsaccades contaminate gamma-band responses in infants.

53

54 To estimate the likelihood of eye-movement contamination of our measures  
55 in Kampis et al. [2], we performed an additional analysis on our time-frequency  
56 data from Study 1. To approximate a measure of eye-movement-related activity,  
57 we estimated the bipolar horizontal EOG signal in our recordings by subtracting  
58 the activation at the two electrodes closest to the outer canthi of the eyes  
59 (channels 1 and 32) from each other. We then subjected this signal to the same  
60 time-frequency analysis as our original data and correlated the resulted gamma  
61 activation in this EOG signal with the activation we obtained in our original  
62 analyses. If eye-movements induced the gamma-band activation found in our  
63 study, then activations at the temporal channels would likely be correlated with  
64 the EOG signal. However, this correlation was not significant either in Segment 1  
65 ( $r = .347$ ,  $p = .205$  in Occlusion condition - for activations in the Occlusion  
66 condition during Segment 1, see Figure 1; and  $r = .239$ ,  $p = .390$  in Control  
67 condition), or in Segment 2 ( $r = -.059$ ,  $p = .835$  in Occlusion condition, and  $r = -.099$ ,  
68  $p = .725$  in Control condition). Based on this analysis it seems unlikely that our  
69 findings originate from eye-movements.

70




Figure 1. Gamma-band activation in the eye channel (channel 32 subtracted from channel 1), and temporal channels (channels 40, 41, 46, 47, 51, 97, 98, 102, 103, 109). The red rectangle marks the frequency and time window used in the analyses in Kampis et al. (2015)

71

72        Additionally, beyond the methodological challenge to detect MS-related  
 73 artifacts in infant EEG, several findings (including some mentioned by Köster  
 74 [1]) of scalp recorded gamma-band activity during object processing in infants  
 75 would not be easily explained by MS patterns. First, in many cases, there were no  
 76 visual differences during the measurement periods between the experimental  
 77 and control conditions, and therefore it is not clear why MSs would show a  
 78 different pattern [e.g. 7,8]. Second, many of the studies reported gamma-band  
 79 activity over temporal areas [e.g. 2,7], whereas MS-related SPs were found  
 80 mostly around the midline in adults [3]. Third, while MS-related SPs were shown  
 81 to manifest themselves in a time window of approximately 200-350 ms after

82 stimulus onset, many studies have used different time windows for analyses [e.g.  
83 2,9], and in some cases it is not obvious what should count as stimulus onset, as  
84 activation was measured after a longer sequence of events [2,7]. Finally, as  
85 Melloni et al. [10] pointed out in their response to the paper demonstrating MS-  
86 related gamma activity, MS-related EEG effects should show a broadband  
87 response, whereas many studies report effects in narrower gamma ranges, and  
88 this observation applies to infant recordings as well.

89

90 In sum, on one hand the tools developed for MS-related artifact removal  
91 from adult EEG are not used currently in infant EEG because they are not  
92 straightforwardly applicable to infant data. Once our understanding of the  
93 characteristics of infant EEG and (oculo-)motor development reaches the  
94 necessary level, it will be possible to return to these concerns and address them.

95

96 On the other hand it is not clear whether this issue has to be addressed in  
97 infants, as the factors that were found to induce possible artifacts in adult studies  
98 are not simply hard to measure but might not be present (or might have radically  
99 different characteristics) in young infants. With regard to our own data [2], it  
100 seems unlikely that the gamma-band activation in temporal areas was due to  
101 infants' eye-movements during the observation of the events (Figure 1). Finally,  
102 some recent results, discussed in Köster [1] as well, suggest that gamma-band  
103 oscillations, even in the adult literature, provide us with a valid tool to  
104 investigate object representations [11].

105

106

107

## References

108

109

1. Köster M. 2016 What about microsaccades in infants?. *Proc. R. Soc. B*

110

2. Kamps D., Parise E., Csibra G. and Kovács Á.M. 2015 Neural signatures for

111

sustaining object representations attributed to others in preverbal human

112

infants. *In Proc. R. Soc. B.* 282: 20151683.

113

<http://dx.doi.org/10.1098/rspb.2015.1683>

114

3. Yuval-Greenberg S., Tomer O., Keren A. S., Nelken I., Deouell, L. Y. 2008

115

Transient induced gamma-band response in EEG as a manifestation of

116

miniature saccades. *Neuron* 58, 429-441.

117

[doi:10.1016/j.neuron.2008.03.027](https://doi.org/10.1016/j.neuron.2008.03.027)

118

4. Hassler U., Barreto N. T., Gruber T. 2011 Induced gamma band responses

119

in human EEG after the control of miniature saccadic artifacts.

120

*Neuroimage* 57, 1411-1421. [doi:10.1016/j.neuroimage.2011.05.062](https://doi.org/10.1016/j.neuroimage.2011.05.062)

121

5. Csibra G., Tucker L.A., Volein Á., Johnson M.H. 2000 Cortical development

122

and saccade planning: the ontogeny of the spike potential. *Neuroreport*,

123

11(5), 1069-1073.

124

6. EEGLAB Tutorial. 2009 Retrieved June 7, 2016, from

125

[http://sccn.ucsd.edu/wiki/EEGLAB\\_Wiki#EEGLAB\\_Tutorial](http://sccn.ucsd.edu/wiki/EEGLAB_Wiki#EEGLAB_Tutorial)

126

7. Kaufman, J., Csibra, G., Johnson, M.H. 2003 Representing occluded objects

127

in the human infant brain. *Proc. R. Soc. B Biol. Sci.*, 270, S140-S143

128

8. Leung S., Mareschal D., Rowsell R., Simpson D., Iaria L., Grbic A., Kaufman,

129

J. 2016 Oscillatory activity in the infant brain and the representation of

130

small numbers. *Front. Sys. Neurosci.* 10. [doi:10.3389/fnsys.2016.00004](https://doi.org/10.3389/fnsys.2016.00004)

- 131 9. Kaufman, J., Csibra, G., Johnson, M.H. 2005 Oscillatory activity in the infant  
132 brain reflects object maintenance. *Proc. Natl. Acad. Sci. USA* 102, 15271-  
133 15274
- 134 10. Melloni L, Schwiedrzik CM, Wibral M, Rodriguez E, Singer W. 2008  
135 Response to: Yuval-Greenberg et al, "Transient Induced Gamma-Band  
136 Response in EEG as a Manifestation of Miniature Saccades." *Neuron* 58,  
137 429-441.
- 138 11. Köster M., Frieze U., Schöne B., Trujillo-Barreto N., Gruber, T. 2014 Theta-  
139 gamma coupling during episodic retrieval in the human EEG. *Brain Res.*  
140 1577, 57-68.

141

142

143 Data availability

144 The EEG dataset used in the analyses reported in this article are available at:

145 doi:10.5061/dryad.k1r85.

146