



Bracher, Michael and Corner, Jessica and Wagland, Richard (2016) Exploring experiences of cancer care in Wales - a thematic analysis of free-text responses to the 2013 Wales Cancer Patient Experience Survey (WCPES). *BMJ Open* . ISSN 2044-6055 (In Press)

**Access from the University of Nottingham repository:**

<http://eprints.nottingham.ac.uk/35525/1/BMJopen-2016-011830%20R1%20%28WCPES%29%20%28Mike%20Bracher%20Jessica%20Corner%20Richard%20Wagland%29%20UoS%2008%2006%2016.pdf>

**Copyright and reuse:**

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: [http://eprints.nottingham.ac.uk/end\\_user\\_agreement.pdf](http://eprints.nottingham.ac.uk/end_user_agreement.pdf)

**A note on versions:**

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact [eprints@nottingham.ac.uk](mailto:eprints@nottingham.ac.uk)

# BMJ Open

## Exploring experiences of cancer care in Wales – a thematic analysis of free-text responses to the 2013 Wales Cancer Patient Experience Survey (WCPES).

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2016-011830.R1
Article Type:	Research
Date Submitted by the Author:	08-Jun-2016
Complete List of Authors:	Bracher, Mike; University of Southampton, Corner, Jessica; University of Nottingham, Research and Knowledge Exchange Wagland, Richard; University of Southampton, Faculty of Health Sciences
<b>Primary Subject Heading</b>:	Oncology
Secondary Subject Heading:	Qualitative research, Research methods, Patient-centred medicine, Communication, Health services research
Keywords:	cancer, patient experience, QUALITATIVE RESEARCH, free text, wales, outcomes

SCHOLARONE™  
Manuscripts

1 Exploring experiences of cancer care in Wales – a thematic analysis of free-text responses to the  
2 2013 Wales Cancer Patient Experience Survey (WCPES).  
3  
4  
5  
6  
7

8 Michael Bracher PhD Research Fellow<sup>1</sup>  
9

10 Dame Jessica Corner PhD Dean of Health Sciences<sup>1</sup>  
11

12 Richard Wagland PhD Senior Research Fellow<sup>1</sup>  
13  
14  
15  
16  
17  
18

19 1. Faculty of Health Sciences, University of Southampton, Highfield, Southampton SO17 1BJ  
20  
21  
22

23 Corresponding Author: Michael Bracher: [mb5v07@soton.ac.uk](mailto:mb5v07@soton.ac.uk)  
24

25 Faculty of Health Sciences, University of Southampton, Highfield, Southampton SO17 1BJ  
26  
27  
28

29 **Word count:** Abstract: 300; Manuscript 5734 words  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

#### 43 **Acknowledgement** 44

45 The authors would also like to acknowledge Dr. Carol Rivas (University of Southampton), for her  
46 helpful comments on earlier drafts of this paper.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

1 **Abstract: (300 words max)**

2

3 **Objectives**

4 To provide the first systematic analysis of a national (Wales) sample of free-text comments from  
5 cancer patients, to determine emerging themes and insights regarding experiences of cancer care in  
6 Wales.  
7

8

9

10 **Design:**

11 Thematic analysis of free-text data from a population-based survey.

12

13

14 **Setting and Participants:**

15 Adult patients with a confirmed cancer diagnosis treated within a three month period during 2012 in  
16 the seven Health Boards and one trust providing cancer care in Wales.

17

18

19 **Main outcome measures:**

20 Free-text categorised by theme, coded as positive or negative, with ratios. Overarching themes are  
21 identified incorporating comment categories.  
22

23

24

25 **Methods:**

26 4,672 respondents (of n=7352 survey respondents) provided free-text comments. Data was coded  
27 using a multistage approach; (1) coding of comments into general categories (e.g. Nursing, Surgery  
28 etc.), (2) coding of sub-categories within main categories (e.g. Nursing Care, Nursing Communication  
29 etc.), (3) cross-sectional analysis to identify themes cutting across categories (4) mapping of  
30 categories/sub-categories to corresponding closed questions in the WCPES data for comparison.  
31

32

33

34 **Results:**

35 Most free-text respondents (82%, n 3818) provided positive comments about their cancer care, with  
36 49% (n=2313) giving a negative comment (ratio: 0.6:1, negative-to-positive). 3172 respondents (67.9%  
37 of free-text respondents) provided a comment mapping to one of four overarching themes:  
38 communication (n=1673, 35.8% free-text respondents, a ratio of 1.0:1); waiting during the treatment  
39 and/or post-treatment phase (n=923, 19.8%, ratio: 1.5:1); staffing and resource levels (n=671, 14.4%  
40 ratio: 5.3:1); speed and quality of diagnostic care (n=374, 8.0%, ratio: 1.5 : 1). Within these areas,  
41 constituent sub-themes are discussed.  
42

43

44

45 **Conclusions:**

46 This study presents specific areas of concern for cancer patients, and reveals a number of themes  
47 present across the cancer journey. While the majority of comments were positive, analysis reveals  
48 concerns shared by significant numbers of respondents. Timely communication can help to manage  
49 these anxieties, even where delays or difficulties in treatment may be encountered.  
50

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Strengths and limitations of the study**

- Provides further detail on closed measures in population-based survey.
- Indicates area of concern not addressed by closed measures.
- Volume of comments and ratios of negative to positive comments in specific areas indicate areas of particular concern.

**Key words:** cancer, patient experience, PROM, PREM, PES, qualitative, free-text, outcomes, Wales.

For peer review only

## INTRODUCTION

The global burden of cancer disease is growing worldwide [1]. Increasing numbers of people in the UK are affected by cancer diagnosis and treatment, with lifetime risk being projected at 1 in 2 for those born from the early 1960s onward [2]. However, research has indicated that survival rates for all cancers combined has increased substantially since 1971, and that more people are living longer with, and beyond, their cancer [3]. Patient experiences of cancer treatment and support, through diagnostic, treatment, and post-treatment phases, are therefore areas of significant and growing public concern.

There is increasing recognition in Europe and North America that the quality and effectiveness of services are best evaluated from the patient's perspective [4]. Patient-reported outcome measures (PROMs) and experience measures (PREMs) are commonly used to explore patients' views on their care and treatment [5,6], and frequently include open-ended questions. Open questions can enhance understanding of responses to closed questions by providing greater detail on experiences, and allowing respondents to offer information that may not be elicited through closed measures. However, these data often remain unexplored, due to the time and resource-heavy process of analysing large free-text sets [7].

In the UK, the NHS (National Health Service) Cancer Reform Strategy and Outcomes Strategy for Cancer documents [8] highlight the important role of patient experiences in measuring and improving clinical quality, and national surveys have been undertaken to determine the quality of experience of cancer patients and survivors [9–11]. In England, the Cancer Patient Experience Survey (CPES) has been conducted annually since 2010, and continues to provide useful insights into patient experiences of cancer treatment and care [12]. Data from responses to closed questions in this survey has been used in previous work by Bone et al. (2014) to explore variations in overall satisfaction with care by socio-demographic, patient, clinical and trust-related factors [14].

1 Elsewhere, analysis of free-text responses to the CPES from patients identified with London Trusts  
2  
3 has been undertaken by Wiseman et al. (2015). In 2012, the Cancer Delivery Plan for NHS Wales has  
4  
5 recognised the important of patient experience and established a commitment to produce a national  
6  
7 survey [15]. The first Wales Cancer Patient Experience Survey (WCPES) was conducted in 2013  
8  
9 through a partnership between the Welsh Government and Macmillan Cancer Support, and was  
10  
11 administered by Quality Health. In common with the England CPES, closed questions in the Wales  
12  
13 survey related to a number of topic areas, for example: seeing your GP; diagnostic tests; clinical  
14  
15 nurse specialist; support for people with cancer [13]. The majority of respondents related positive  
16  
17 experiences of their care; however, there also exist groups of patients who report less positive  
18  
19 experience in a variety of areas.  
20  
21  
22  
23  
24

25 The present study was commissioned by Macmillan Cancer Support to analyse the content of the  
26  
27 free-text responses, provide more information on specific experiences of care and treatment,  
28  
29 identify any areas that had not been covered by quantitative measures, and thereby facilitate mixed  
30  
31 methods descriptive analysis of the data.  
32  
33  
34  
35

## 36 **METHODS**

### 37 **Study design**

38  
39 This investigation involves analysis of secondary data from a population-based postal survey  
40  
41 undertaken in Wales in 2013 of all individuals aged  $\geq 16$  years with a primary diagnosis of cancer<sup>1</sup>,  
42  
43 who were admitted to an NHS hospital as an inpatient or as a day case patient, and were discharged  
44  
45 from hospital between 01/09/2012 and 30/03/2013 (n=10,945)[12]. Survey results were published  
46  
47 in January 2014[16] with a response rate of 69% (n=7352 patients). Results from the closed  
48  
49  
50  
51  
52  
53

---

54 <sup>1</sup> 'Data definitions identifying patients qualifying for inclusion were based on the ICD10 codes of C00-C99, and  
55 D05 were used. Patients with an ICD10 code of C44 (other malignant neoplasms of the skin), and C84 (some  
56 Haematology codes) were excluded from the sample by agreement. There are only very small numbers of such  
57 patients with these codes.' [16]  
58  
59  
60

1 questions demonstrate a positive experience of cancer care in Wales, with 89% of patients rating  
2 their care as excellent (58%) or very good (31%).  
3  
4  
5

### 6 7 8 **Cohort identification**

9  
10 The seven health boards and one trust treating adult patients with cancer in Wales were included.  
11 Patients were identified from data provided by health boards/trusts, selected from local patient  
12 administration systems [16].  
13  
14

### 15 16 17 18 19 20 **Questionnaire and design content**

21  
22 Questionnaires included questions on socio-demographics, quality of treatment and care, disease  
23 status and long term conditions (LTCs) [16]. Three free-text comment boxes were placed at the end  
24 of the questionnaire, following the closed questions: "Was there anything particularly good about  
25 your NHS cancer care?"; "Was there anything that could have been improved?"; "Any other  
26 comments?" (these questions are identical to those used in the 2013 CPES for England).  
27  
28  
29  
30  
31  
32

### 33 34 35 36 **Survey process**

37  
38 The survey was distributed by post, with two reminders sent out to non-responders only [16].  
39 Covering letters were sent out on Health Board/Trust headed paper and signed by a member of the  
40 Health Board/Trust staff [16]. Survey and covering letters were sent out in both English and Welsh  
41 language versions. An enclosed language leaflet offered translation services and a pre-paid return  
42 envelope was included so patients could respond without financial cost [16]. 4576 free-text  
43 respondents used English language (63.6% of total English language respondents to the full survey<sup>2</sup>),  
44  
45  
46  
47  
48  
49  
50

51  
52  
53  
54  
55  
56  
57 <sup>2</sup> n=7190

58 6  
59  
60



1 while 96 (59.3% of Welsh Language respondents<sup>3</sup> to the full survey) provided free-text responses in  
2  
3 Welsh language, which were translated into English for analysis.  
4  
5  
6

### 7 **Ethics and governance**

8  
9 Approval was given by the National Information Governance Board (NIGB) and the Confidentiality  
10 Advisory Group CAG 3-04(PR2)/2013. The survey was performed as a service evaluation. Survey  
11  
12 respondents had access to a telephone support line to discuss issues raised by the survey.  
13  
14  
15  
16

### 17 **ANALYSIS**

18  
19 Data were subjected to a thematic analysis, informed by multi-stage coding (see figure 1) of free-text  
20  
21 data [17]. The coding taxonomy was developed inductively from the data using the NVivo  
22  
23 Qualitative data analysis software package.  
24  
25  
26

27  
28 **Stage 1:** Stage one involved analysis of semantic content of the entire free-text data set (i.e. whether  
29  
30 comments contained references to nursing care, surgery, chemotherapy etc.) and whether  
31  
32 comments<sup>4</sup> were of a positive or negative nature. A coding taxonomy was produced for sorting  
33  
34 qualitative data into categories of patient experience, developed by one researcher (MB) in  
35  
36 collaboration with two further researchers (RW, JC) (Table 2). Once the main taxonomy had been  
37  
38 established (i.e. it accommodated the majority of comments without need for additional categories),  
39  
40 a sample of 200 randomly selected comments was double-coded by two researchers (MB, RW).  
41  
42 Coding agreement between the researchers was 80% (Cohen's Kappa), and conflicts were resolved  
43  
44 through discussion.  
45  
46  
47  
48  
49  
50  
51  
52

---

53  
54 <sup>3</sup> n=162

55 <sup>4</sup> Respondent/response counts in this article refer to incidence of individual respondents within a given code or  
56 theme (i.e. a respondent would only be counted once towards the total for 'Surgery Positive' even if the  
57 comment referred to two different incidents of surgical care).  
58

1 **Stage 2:** Individual categories were subjected to a second stage of more detailed sorting into sub-  
2 categories. For example, at stage one, comments relating to nursing care were coded to the  
3 categories 'Nursing' and then 'Nursing Positive' or 'Nursing Negative'. At stage two, comments  
4 within these categories were sorted further according to what specifically was 'positive' or 'negative'  
5 about the care (e.g. patient perceptions of information provided by nurses or the manner in which  
6 they were dealt with by staff)<sup>5</sup>.  
7  
8  
9  
10  
11  
12

13  
14  
15  
16 **Stage 3:** Categories/sub-categories were subjected to cross-sectional analysis to highlight common  
17 themes present across different categories (e.g. communication)(Table 3).  
18  
19

20  
21  
22 **Stage 4:** Comparisons between results of closed question and free-text responses were conducted  
23 where there was appropriate correspondence.  
24  
25  
26

#### 27 28 29 **FINDINGS:**

30  
31 4672 patients provided free-text comments in the survey, representing 64% of those who returned  
32 questionnaires. Self-reported demographic data on age, sex, long-standing condition, employment  
33 status, ethnicity and sexual orientation, were collected (data on tumour type was provided through  
34 local patient administration systems). The profile of participants who provided free-text comments  
35 was representative of all WCPES survey respondents (n=7352) (see Appendix, Table 1). Most free-  
36 text respondents (82%, n=3818) provided a positive comment about their cancer care, with 49%  
37 (n=2313) providing a negative comment, giving an overall positive ratio of 0.61:1 (see Appendix,  
38 Table 2).  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 Stage 3 analysis identified four major overarching themes, incorporating the majority of stage 1 and  
49 2 text categories: *communication; waiting; staffing and resource levels; speed and quality of*  
50 *diagnostic services*. 3172 respondents (63.9% of total free-text respondents) provided a comment  
51  
52  
53  
54

55  
56  
57 <sup>5</sup> A full breakdown of the stage 2 analysis is included as an additional file with this submission, due to size.  
58

1 relating to one of the four themes identified, of which 1948 (41.7% of total free-text respondents)  
2  
3 were negative, and 1276 (27.3% of total free-text respondents) were positive (overall negative ratio  
4  
5 of 1.53:1) (see Appendix, Table 3).  
6  
7  
8  
9  
10

### 11 **1. Communication**

12  
13  
14 The largest single theme was *communication* (1673 respondents) with a balanced ratio of 1.01:1,  
15  
16 representing 35.8% of free-text respondents and 22.8% of survey respondents. Comments relating  
17  
18 to communication fell into two sub-themes; *communication between patients and staff*; and  
19  
20 *communication between staff and/or institutions* (i.e. sharing information).  
21  
22  
23  
24

#### 25 ***Communication between patients and staff***

26  
27 A third (31.5%) of free-text respondents (n=1472) provided a comment relating to communication  
28  
29 with healthcare staff, of which 854 were negative and 846 positive (a balanced ratio: 1.01:1) (see  
30  
31 Appendix, Table 3). Responses in this sub-theme were of two broad types: those commenting on  
32  
33 the *quality and/or availability of information provided by staff*; and those referring to the *manner in*  
34  
35 *which patients perceived that they had been dealt with by staff*.  
36  
37  
38  
39

#### 40 ***Quality and/or availability of information provided by staff***

41  
42 Comments on information provision cut across a wide range of treatment types and staff  
43  
44 areas/specialties, with the majority of negative responses related primarily to the availability of  
45  
46 adequate information on treatment/care, (269 respondents, 5.7% of free-text respondents).  
47  
48  
49  
50

51 *Sometimes it feels like you have to tease information out of doctors - it doesn't seem to be*  
52  
53 *given readily, you just have to ask the right questions. (female, aged 35-44 years)*  
54  
55  
56  
57

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Comments such as these indicate the additional communicative work described by many respondents as necessary to obtain sufficient information from clinical staff. Conversely in the positive comments, (143 respondents, 3.1% of free-text respondents), we observe statements indicating satisfaction when information provision and access to specialist staff was perceived to be adequate.

*The doctors / surgeons at [hospital removed] were excellent and caring, explaining all that was happen[ing] or about to happen (male, aged 65-74 years)*

Having access to sufficient information in a timely manner during treatment represents a significant area of concern for many free-text respondents, can be seen as limiting the extra communicative work that some patients perceived in needing to 'tease information' from staff, potentially lessening the 'burden of treatment' [18].

#### *Perceptions of staff manner in interactions with patients*

227 respondents (4.9% of free-text respondents) gave a negative comment about the manner in which they felt that they had been dealt with by healthcare staff. For some, this related to presentation of their diagnosis (54 respondents).

*The original time I was told I had terminal cancer and nothing could be done for me was handled very badly. There was no support at all and the doctor was in and out of the room in about 6 minutes. It was as if my life counted for nothing, as if I was being thrown away. (male, aged 55-64 years)*

*I took great exception to the manner in which I was told, no privacy, no family member present, and people each side of me could hear. (female, aged 75-84 years)*

1  
2  
3 Complaints were observed relating to inadvertent disclosure of cancer diagnosis (of which  
4 participants had previously been unaware), as well as not having family or loved ones present when  
5 told. Other respondents perceived negative attitudes amongst consultants and specialists (22  
6 respondents), hospital doctors (35 respondents) or nursing staff (47 respondents).  
7  
8  
9  
10

11  
12  
13  
14 *As a former employee of NHS, I have the greatest respect for the ward staff who work*  
15 *exceedingly hard, but the attitude of some of the medics and other disciplines need to be*  
16 *visited. Sometimes I felt I was treated like a piece of meat or idiot as medics discussed me*  
17 *with colleagues, without ever talking to me directly (female, aged 55-64 years)*  
18  
19  
20  
21

22  
23  
24  
25 *Some good nurses – but in equal amount, some very lazy, gossiping and bad tempered*  
26 *nurses too. (female, aged 55-64 years)*  
27  
28  
29

30  
31 However, greater numbers of respondents (n=544) praised the manner of staff during their  
32 treatment journey, including: *nursing staff* (202 respondents); *hospital doctors* (136 respondents);  
33 *consultants and specialist medics* (101 respondents); and *'surgical staff'* (47 respondents).  
34  
35  
36  
37

38  
39  
40 *The consultant and registrar are most informative and to the point. They always discuss...the*  
41 *way forward with my treatment. I have every confidence in them. (male, aged 65-74 years)*  
42  
43

44 *The nurses that administer the area and in my case carried out tests were very caring and*  
45 *efficient but very obviously overloaded with work. (male, aged 65-74 years)*  
46  
47  
48

49  
50  
51 Respondent comments highlighted personal qualities (e.g. kindness, empathy, sympathy, respect,  
52 compassion) in interactions with staff as positive experiences of their treatment, as well as the  
53  
54  
55

1 professionalism of staff involved in their care (e.g. that staff were helpful, efficient, competent etc. ),  
2  
3 despite challenging workloads.  
4  
5  
6

### 7 ***Communication between staff and/or institutions***

8  
9 252 respondents commented on inter-/intra-agency communication between staff and/or  
10 institutions (e.g. information sharing between specialists and GPs where the latter were not  
11 identified as the source of the problem, sharing of notes and/or test results between hospitals/sites  
12 etc.). Of these, 208 described negative experiences, while 44 gave positive responses (a ratio of  
13 3.8:1). Negative respondents gave general comments relating to this area of communication.  
14  
15  
16  
17  
18  
19

20  
21  
22 *At times, a lack of clear communication between different departments/clinics made the*  
23 *situation more and more difficult... (female, aged 55-64 years)*  
24  
25  
26

27 Given that this theme references examples of communication in which patients were not involved  
28 directly, the generality of the comments is perhaps unsurprising. Nonetheless, they indicate a sizable  
29 number of respondents that perceived poor communication between staff and/or institutions  
30 involved in their care. 44 respondents reported positive experiences, highlighting the beneficial  
31 impact that this had on their care.  
32  
33  
34  
35  
36  
37  
38  
39

40 *I appreciate the communication between hospital, GP, out of hours etc. It means I don't have*  
41 *to repeat myself so often. It also means I have instant treatment. (female, aged 55-64 years)*  
42  
43  
44  
45

46 Positive respondents often associated perceptions of good communication between staff and/or  
47 institutions with speediness of treatment.  
48  
49  
50  
51  
52

## 53 **2. *Waiting during the treatment and post-treatment phase***

54  
55  
56  
57  
58 12  
59  
60

1 923 free-text respondents (19.8%) provided a comment about waiting times during the treatment  
2  
3 and/or post-treatment phases, either the interval before consultations/treatment (738 respondents,  
4  
5 15.8%) or the time spent waiting in hospitals on the day of appointments (n=194, 4.2%) (see  
6  
7 Appendix, Table 3).  
8  
9

### 10 ***Waiting for appointments***

11  
12 In the closed questions (question 2), 78% of respondents (n=5520) reported they had been seen 'as  
13  
14 soon as necessary' by an oncologist, with 12% (n = 839) feeling that they 'should have been seen a  
15  
16 bit sooner' and 10% (n = 685) indicating that they 'should have been seen a lot sooner'  
17  
18 (QualityHealth, 2014). In the free-text portion of the survey, 397 respondents (8.5% of free-text  
19  
20 respondents) gave negative comments, while 342 (7.3% of free-text respondents) provided a  
21  
22 positive response. The majority of responses in this sub-theme did not map to a specific area of  
23  
24 treatment or care, but instead referred to consultations or 'treatment' related appointments in  
25  
26 general terms.  
27  
28  
29  
30  
31  
32

33 *The wait to start treatment is too long. I was initially told I should start treatment by August.*

34  
35 *I have an appointment for [date removed]. The long delay is disappointing. I was diagnosed*  
36  
37 *in February. (male, aged 65-74 years)*  
38  
39  
40  
41

42 Positive comments often appeared in the context of broader comments relating to the entire  
43  
44 journey.  
45  
46  
47

48  
49 *I went to my GP on the Thursday and I was seen by the following week. The consultant in the*  
50  
51 *hospital which I had biopsies taken and told that same day I had cancer, and it was dealt*  
52  
53 *with very quickly and I was very happy with the care I was given and how quickly it was*  
54  
55 *treated. (female, aged 55-64 years)*  
56  
57

58 13  
59  
60

1  
2  
3 Comments praising the swiftness of appointments during treatment were often attended by  
4 expressions of confidence in and satisfaction with the overall package of care given to respondents.  
5  
6  
7

8  
9  
10 ***Waiting on the day.***

11 194 respondents (4.1%) provided a comment about waiting times on the day of their appointments  
12 during the treatment phase, of which 163 responses (3.5%) were negative and 31 were positive  
13 (0.7%). Once again, the majority of these responses were of a general nature, referring only to  
14 events such as 'treatment(s)' or 'appointment(s)'. The vast majority of negative comments  
15 concerned (unspecified) clinic appointments, and most commonly referred to delays of around 1.5-2  
16 hours beyond the appointed time.  
17  
18  
19  
20  
21  
22  
23

24  
25 *Sometimes as an outpatient with an appointment, the wait is too long! Eg 1 1/2 to 2 hours,*  
26 *even when you arrive well before time. (female, aged 18- 24 years)*  
27  
28

29  
30  
31 It is also important to note that many respondents providing such comments added qualifications  
32 indicating their perception of services being under pressure, as a reason for these delays (e.g.  
33 'nurses [were] literally running from one patient to the next').  
34  
35  
36  
37  
38

39  
40 A smaller sub-set of respondents (n=31) reported good or acceptable waiting times on the day of  
41 their appointments.  
42  
43  
44  
45

46  
47 *Appointments have been kept on time and in my view within reasonable waiting time. (male,*  
48 *aged 65-74 years)*  
49  
50

51  
52  
53 While for some waiting times on the day of appointments were not an issue, a greater number of  
54 respondents reported difficulties in this area. For some, protracted waiting times were a source of  
55  
56  
57



1 additional problems and discomfort relating to their condition (e.g. bladder problems), social and  
2  
3 employment commitments and car park charges.  
4  
5

### 6 7 8 **3. Staffing and resource levels** 9

10 Concerns about staffing and resources were expressed by a significant number of free-text  
11 respondents (n=671, 14.4%), of which 568 responses (12.2%) were negative, and 107 (2.3%) were  
12 positive (see Appendix, Table 3). These responses cut across a number of areas of the cancer  
13 journey, the largest of which was *availability of aftercare* (312 respondents).  
14  
15  
16  
17  
18  
19

#### 20 ***Availability of aftercare*** 21

22 This sub-theme was comprised of 217 negative and 98 positive responses (ratio: 2.21:1). Negative  
23 comments identified a lack of general aftercare provision following the completion of treatment,  
24 whether chemotherapy, radiotherapy, surgery or other treatment programmes, and this was also  
25 true for respondents giving otherwise positive responses.  
26  
27  
28  
29  
30  
31  
32

33 *When discharged from completing radiotherapy I felt quite alone as there had been so much*  
34  
35 *support before (female, aged 45-54 years)*  
36  
37  
38  
39

40 The generality of negative comments appears indicative of a profound gap in services after  
41 treatment has finished. Support from specialist medical and nursing staff, as well as emotional,  
42 social and psychological support while recovering from cancer treatment, were unmet needs  
43 reported by many respondents. Concerns included fear of recurrence linked to a lack of clear plans  
44 for determining success of treatment, or for long term monitoring. Several respondents described  
45 actual recurrence of cancer, and reported that its discovery was delayed due to failure to conduct  
46 what they considered to be appropriate follow up investigations.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Conversely, comments from the smaller group who provided positive responses reflected experiences of security from regular monitoring, following completion of treatment.

*I am being monitored regularly and feel looked after. The specialists are very professional and I felt confident in their care (female, aged 55-64 years)*

Aftercare was one of the few areas of treatment where negative responses greatly outnumbered positives, and in some cases the former accompanied otherwise positive responses praising many or all other aspects of their cancer journey.

#### **General comments about staffing levels (nursing and medical staff)**

267 free-text respondents (5.7%) gave comments relating to staffing levels in hospital settings, of which all but one were negative. While approximately half of respondents (n=141) in this sub-theme gave comments of a general nature (e.g. referring to 'staff' but not specifying a particular speciality), 131 respondents referred to inadequate provision of nursing staff (in general terms), while 17 made (similarly general) comments about hospital doctors.

*The nursing staff on the wards work very hard but are very overworked. Staffing levels need to be improved. (female, aged 55-64 years)*

These comments mirror responses to the closed section of the WCPES survey, in which 29% (n=1229) of respondents indicated that 'there were sometimes enough [nurses] on duty'; 11% (n=478) indicated that there were rarely or never enough on duty; while 60% (n=2580) agreed that 'there were always or nearly always enough on duty' (QualityHealth, 2014).

### ***Availability and quality of staff on hospital wards at evening and weekends***

62 respondents provided negative comments regarding out of hours and weekend care with respect to the quality and availability of staff, while eight provided positive responses (ratio: 7.5:1). All positive comments were of a general nature, (e.g. 'good care at the weekend'). Negative comments presented concerns about staffing levels at weekends and during the night in hospital wards, as well as examples of poor care (again, particularly during the night). Noise levels during the night, and difficulties obtaining out of hours advice and/or treatment for problems arising during treatment were also significant concerns. Some of these comments were general, reflecting concerns around quality of care.

*Night time on the ward was awful due to it being short staffed. (female, aged 65-74 years)*

The night staff could have been more respectful it was difficult to sleep because of noise from them, and my bell wasn't answered. *(female, aged 65-74 years)*

These comments were not matched by a significant number of positive comments regarding out of hours/weekend care. While the responses can only reflect the experiences of 60 respondents, the specificity of some of the comments (e.g. noise on hospital wards) suggests that there may be specific areas of concern.

### ***Availability of specialist nursing staff***

Almost all negative comments (n=63, 1.35%) related to availability of specialist nursing staff to answer questions and provide information about treatment.

1 *I felt I needed specialist nurse support (phone or personal contact) following my 3 operations,*  
2  
3 *especially I experienced difficulty with chemotherapy. Needed emotional support, although I*  
4  
5 *did not contact anyone. I live alone and did ask if there was any advice on home support, no*  
6  
7 *action (female, aged 55-64 years)*  
8  
9

10  
11 Many respondents perceived specialist nursing staff to be highly pressurised, and linked this  
12  
13 perception to lack of availability. However, the majority of comments relating to communication  
14  
15 with specialist nursing staff, both in hospital settings (e.g. during chemotherapy or radiotherapy  
16  
17 treatment) and away from hospitals (e.g. district nurse / keyworker visits, access to CNS) were  
18  
19 positive (n=106). Indeed, there was a high ratio of positive over negative (1:0.33) comments relating  
20  
21 to specialist nurses amongst free-text respondents.  
22  
23

24  
25 *I have found my nurse specialist to be very helpful and always has time to listen to my*  
26  
27 *concerns. She will always do her best to answer my questions. She always returns calls*  
28  
29 *(female, aged 69-74 years)*  
30  
31

32  
33 Comments reflect experiences of high standards in both information provision and the manner in  
34  
35 which the information was provided by specialist nursing staff, often in spite of significant pressures  
36  
37 on their time and resources as perceived by patients.  
38  
39

#### 40 41 42 **4. Speed and quality of diagnostic care**

43  
44 411 free-text respondents (8.8%) gave comments relating to the speed and quality of care during the  
45  
46 diagnostic phase of the cancer journey, with further comments relating to the *GP role in diagnosis*  
47  
48 (n=211) and *investigations and diagnostic procedures* (n=193)(see Appendix, Table 3).  
49  
50

#### 51 52 53 ***General Practitioner (GP) role in diagnosis***

1 General practitioners (GPs) were the only specialist staff category in which negative free-text  
2 responses outnumbered positives (ratio: 1.53:1). Amongst closed, tick-box questions, 78% (n=5520)  
3 of respondents reported they had been seen by a hospital doctor as soon as necessary following  
4 referral, 12% (n=839) felt that they 'should have been seen a bit sooner', and 10% (n=685) 'a lot  
5 sooner'[19]. Amongst free-text comments, n=80 (%) respondents described delays in referral by  
6 their GP for further investigation of their symptoms, 16 of whom for what they considered an  
7 unwillingness to refer.  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19

*I had to bypass my GP to get an endoscope test, after numerous requests explaining how ill I  
20 felt. The endoscopy dept. discovered the cancer. (male, aged 75-84 years)*  
21  
22  
23  
24

25 Of particular concern was a subset of respondents (n=35) who described inaccurate diagnosis of  
26 their cancer prior to correct diagnosis. This was seen to delay treatment often by months, and in  
27 some cases a year or more.  
28  
29  
30  
31  
32

*The GP got my condition completely wrong. He had it fixed in his mind that I had  
33 haemorrhoids. Finally my daughter took me to A&E, where they discovered an obstruction.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60*  
*The following day I had an endoscopy, which revealed a tumour. That day I had the  
colostomy. (male, aged 75-84 years)*

In contrast, positive comments (n=52) on GP care tended to be more general, but almost all referred  
to the speed with which presenting symptoms had been investigated, including referral for further  
investigations.

1                    *The speed at which my GP referred me to a specialist was phenomenal! It gave me a feeling*  
2  
3                    *of confidence in the NHS service at a time when I was very frightened. (male, aged 55-64*  
4  
5                    *years)*  
6  
7  
8  
9

10                    Positive responses were often allied with more general comments expressing feelings of satisfaction  
11  
12                    and reassurance in terms of the overall care and treatment received during the cancer journey.  
13

### 14                    ***Investigations and diagnostic services***

15  
16                    173 (3.70%) respondents gave negative comments regarding delays relating to initial diagnostic  
17  
18                    procedures. Of these, 94 were general or miscellaneous comments regarding delays and/or access  
19  
20                    to diagnostic services in the initial stages of the cancer journey.  
21  
22  
23

24  
25  
26  
27                    *A simple colonoscopy at the early stages would save a lot of pain and suffering and a much*  
28  
29                    *cheaper option. (male, age unavailable)*  
30  
31

32  
33                    36 negative responses included reference to perceived inaccurate or mistaken diagnosis.  
34  
35

36  
37  
38                    *I believe the cancer was missed in earlier mammograms. (female, aged 55-64 years)*  
39  
40

41  
42                    While delays and accuracy in diagnostic services were of concern to some, 57 other respondents  
43  
44                    praised the speed of investigative services.  
45  
46

47  
48                    *The speed with which my diagnostic test, scans and surgery were organised. All the doctors*  
49  
50                    *exuded a sense of urgency which I found reassuring. (male, aged 75-84 years)*  
51  
52

1 Delays (whether attributed to waiting times or inaccuracies in diagnosis) were linked to concerns  
2  
3 around cancer progression, implications for treatment response, risk of poorer outcomes and  
4  
5 additional suffering. Conversely, swiftness of diagnosis was associated with expressions of  
6  
7 satisfaction and confidence in the process.  
8  
9

## 11 **DISCUSSION:**

14 Analysis of free-text comments within the WCPES survey complements the formal closed questions  
15  
16 by allowing patients to indicate the issues most important to them and provides important insights  
17  
18 of the experience of patients. The high response rate to the free-text question (64% of those who  
19  
20 returned questionnaires) indicates that patients actively engage with the opportunity to provide  
21  
22 comments relating to their experiences. They also reflect the findings of the closed questions, that  
23  
24 most respondents had a positive overall experience of cancer care. In terms of potential  
25  
26 improvement, the themes indicate the impact that uncertainty can have on patient experiences,  
27  
28 particularly around perceptions of delays in diagnosis and treatment, or of poor communication  
29  
30 during treatment.  
31  
32

36 For patients who suspect they might have cancer, delay also causes additional psychological distress,  
37  
38 which has been shown to correlate positively with the length of that delay [20]. Previous research  
39  
40 has highlighted the presence of free-text comments relating to delays in referral within the CPES  
41  
42 England (for London trusts) [21]. Elsewhere it has been indicated that patients are often not  
43  
44 satisfied with the time it took for the GP to identify their problem and for a diagnosis to be reached  
45  
46 [22]. Delays for investigations and referral can be caused through 'misdiagnosis' with GPs either  
47  
48 treating patients symptomatically or relating symptoms to a health problem other than cancer, while  
49  
50 for some cancers this could be linked to inadequate patient examination, use of inappropriate tests  
51  
52 or failing to follow-up negative or inconclusive test results [23]. A recent international survey-based  
53  
54 study of primary care physicians (PCPs) demonstrated a correlation suggesting a relationship  
55  
56

1 between PCP willingness to act on presentation of symptoms, and cancer survival [24]. Percentages  
2 of PCP respondents in Wales<sup>6</sup> that indicated willingness to act on clinical scenarios given in the  
3 survey were the lowest for all but two of these scenarios (in which they were second lowest) [24].  
4 These percentages were correlated with survival rates (both 1 year and conditional five year survival)  
5 that were either lowest or second/joint-second lowest for all of the cancer types [24]. These  
6 findings support patient concerns about a lack of willingness to refer for further investigations at the  
7 GP level, which may be indicative of systemic problems at the GP level requiring further investigation.  
8 One factor may be communication between and access to support from secondary care, as the  
9 authors also reported that PCPs in the UK were the only groups in their study in which most PCPs did  
10 not report ready access to secondary care advice about investigation or referral for suspected cancer  
11 [24].  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

27 Uncertainty can be understood as a common feature of cancer patient experience, and one that can  
28 likely be reduced but not eradicated completely [25,26]. Our findings indicate consequences of  
29 uncertainties for patient experiences in treatment and post-treatment, and areas to which attention  
30 may be paid in reducing them. Patients in this study often communicated perceptions of mitigating  
31 factors in the issues that they experienced, for example, in highlighting the dedication of staff in  
32 circumstances of perceived understaffing. Such comments indicate that where delays and/or  
33 uncertainties relating to treatment were present, perceptions of being informed and having a point  
34 of contact to ask questions were linked to a greater tolerance for difficulties faced. Recent evidence  
35 suggests that patients want more information concerning effects of treatment, and also that cancer  
36 patients continue to receive what they perceive as sub-optimal levels of information and  
37 preparation [21,27,28]. A wider range of unmet needs have been identified for those post  
38 treatment or in survivorship relating to emotional and social support, quality of life, long term  
39 functioning and finance [29–31]. A lack of clarity regarding the process of care has also been  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

---

55  
56  
57 <sup>6</sup> Approximately n=217 participants (11.7% crude response rate, 1861 invited) [18]



1 identified as an issue for survivors post treatment, in part associated with less contact with services  
2  
3 [32]. Such support and guidance have been indicated as important factors in patients' satisfaction  
4  
5 with their care [22], but this requires sufficient and accessible specialist staff. Inadequate staffing  
6  
7 levels were perceived as a problem in this study (echoing observations from Wiseman et al.'s (2015)  
8  
9 analysis of CPES England free-text data from London trusts) [21]. In the present study , this was  
10  
11 particularly true of accessibility of specialist nurses, and recent evidence shows that care  
12  
13 coordination and emotional support and support for the control of side-effects are better in Trusts /  
14  
15 Hospitals with more specialist nurses [33]. It is probable that inadequate levels of staffing will also  
16  
17 contribute to other problems experienced by patients, such as instances of uncoordinated care, lack  
18  
19 of individualised care and waiting for treatment and pain control [19].  
20  
21  
22  
23  
24

25 In the post-treatment phase, previous research has indicated that patients can often feel 'cut adrift'  
26  
27 by the health system after the period of hospital treatment and are left feeling vulnerable and  
28  
29 isolated [34], a finding echoed by many patients in this study. Evidence indicates that approximately  
30  
31 30% to 50% of cancer survivors have unmet needs, mainly for psychological support and coping with  
32  
33 fear of recurrence [35–37]. While unmet needs reduce for some patients in the months following  
34  
35 treatment, one study found that for 60% of these patients the situation did not improve over a six  
36  
37 month period [34]. Finally, patients' comments within the WCPES often did not describe specific  
38  
39 issues related to aftercare, other than to describe its lack, which reinforces findings from previous  
40  
41 studies [11,21].  
42  
43  
44  
45  
46

47 Our analysis of the free-text data has been used by Macmillan Cancer Support to gain further insight  
48  
49 into the extent and quality of person-centred care in Wales, and to support the organisation's key  
50  
51 policy calls for provision of a Cancer Nurse Specialist as the key worker for every patient diagnosed  
52  
53 with cancer, as well as a holistic needs assessment and written care plan. It has also formed part of  
54  
55 evidence submissions from Macmillan in response to Welsh Government consultations, and the  
56  
57

1 National Assembly's Health and Social Care Committee inquiry, focussing on understanding progress  
2  
3 in implementing Welsh Government's Cancer Delivery Plan.  
4  
5  
6

### 7 **STUDY LIMITATIONS**

8  
9 Data were volunteered by individuals and not reported against a pre-determined structured list of  
10 topics, and therefore are not necessarily representative. Recall and response bias may also be  
11 present. The detail provided in the comments is constrained by the brevity of the response format  
12 (i.e. a hand-written box) and so may not be as empirically rich as other forms of qualitative data (e.g.  
13 semi-structured interview data or longitudinal diaries). Positive comments tended to be of a more  
14 general quality and scope than negative comments, and that a far greater proportion of positive  
15 responses were not identified with a specific area (3% of negative respondents vs. 22 % of positive  
16 respondents). Therefore in more specific categories/themes, numbers of negative respondents may  
17 be close to or outnumber positive ones, despite positive responses outnumbering negatives overall.  
18 Counts relating to comments refer to numbers of respondents providing comments in specific  
19 categories/themes, and as such negative and positive comments in a given area may not equal the  
20 total amount of respondents (i.e. because individual respondents may have provided both negative  
21 and positive comments). Counts do not account for the strength of comments or their seriousness  
22 (e.g. a negative comment concerning quality of meals counts towards a total in the same way as a  
23 more serious complaint relating to poor care or treatment).  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

### 44 **FUTURE RESEARCH**

45  
46 Manual coding of free-text affords the most thorough means to analyse these data thematically;  
47 however, working with a large corpus is a labour intensive process, and larger projects may require  
48 additional methods for manipulation and sorting of free- text, in order to produce thematic analyses  
49 at the level of detail in the present study. Our previous work with colleagues on survey data from  
50 colorectal cancer patients has used text mining techniques to automate sorting of responses into  
51  
52  
53  
54  
55  
56  
57

1 broad categories for manual coding [38]. In addition, the PRESENT (Patient Reported Experience  
2 Survey Engineering of Natural Text) project currently underway will explore and develop methods  
3 for working with these data using text engineering[39].  
4  
5

6  
7 In their analysis of England CPES free-text data from cancer patients within London NHS trusts,  
8 Wiseman et al. noted that a number of patients described care outside of their assigned trusts,  
9 and/or sought to identify closed-question responses with areas outside of London [21]. Both types  
10 of response were observed in the present study, and therefore future research might seek to  
11 examine associations between specific treatment sites and responses. Such work would be of  
12 benefit in assessing and developing the ability of surveys such as CPES to reflect the complexities of  
13 cancer care pathways[21].  
14  
15  
16  
17  
18  
19  
20  
21

## 22 23 24 25 **CONCLUSION:**

26 This study has illustrated the value of free-text analysis for exploring patient experiences of cancer  
27 care, and for complementing and extending findings from closed questions. As the first systematic  
28 analysis of free-text data from a national sample of cancer patient experiences, it has presented  
29 specific areas of concern for cancer patients, as well as areas of good practice, and revealed themes  
30 present across the cancer journey. The volume of comments within specific themes, as well as ratios  
31 of negative-to-positive comments, indicate areas of potential concern. Our work on the WCPES has  
32 also highlighted an area of potential significance with regard to the reliability of survey data at  
33 greater levels of specificity (i.e. the site level). These findings have been discussed in the context of  
34 existing issues in cancer care, and in so doing have presented areas of specific attention for policy  
35 makers and further research.  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

## 55 **Contributorship statement**

56  
57  
58 25  
59  
60

1 All three authors have made substantial original contributions to this work, participated in drafting  
2 and revision, approved the final submission, and have agreed to be accountable for all aspects of the  
3 work.  
4

5 Mike Bracher lead on analysis of the data, writing of the paper, and establishment of the analytical  
6 taxonomy. Jessica Corner contributed to the development of the thematic taxonomy, write up of  
7 the findings, and contributed throughout the drafting process. Richard Wagland conducted analysis  
8 of the data, contributed to development of the thematic taxonomy, write-up of findings and  
9 discussion, and participated throughout the drafting process.  
10  
11

### 12 **Competing interests**

13 The authors have no competing interests.  
14

### 15 **Funding**

16 This study was funded by Macmillan Cancer Support.  
17

### 18 **Data sharing statement**

19 Source data for the study are free-text responses to the Welsh Cancer Patient Experience Survey  
20 (WCPES) for 2013. These data are available from the survey provider (Quality Health, UK)  
21

22 <https://www.quality-health.co.uk/> - email: [info@quality-health.co.uk](mailto:info@quality-health.co.uk)  
23  
24

25 No further data specific to this study are available.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## REFERENCES


- 1 International Agency for Research on Cancer (IARC). All Cancers (excluding non-melanoma skin cancer) Estimated Incidence, Mortality and Prevalence Worldwide in 2012. Cancer Fact Sheets. 2016.[http://globocan.iarc.fr/Pages/fact\\_sheets\\_cancer.aspx](http://globocan.iarc.fr/Pages/fact_sheets_cancer.aspx) (accessed 2 May2016).
- 2 Ahmad AS, Ormiston-Smith N, Sasieni PD. Trends in the lifetime risk of developing cancer in Great Britain: comparison of risk for those born from 1930 to 1960. *Br J Cancer* 2015;**112**:943–7. doi:10.1038/bjc.2014.606
- 3 Quaresma M, Coleman MP, Rachet B. 40-year trends in an index of survival for all cancers combined and survival adjusted for age and sex for each cancer in England and Wales, 1971-2011: A population-based study. *Lancet* 2015;**385**:1206–18. doi:10.1016/S0140-6736(14)61396-9
- 4 Darzi A. High Quality care for all: NHS Next Stage Review final report. London, UK: : Department of Health 2008. [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/228836/7432.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/228836/7432.pdf)
- 5 Black N. *Patient reported outcome measures could help transform healthcare*. 2013. doi:10.1136/bmj.f167
- 6 Lipscomb J, Gotay CC, Snyder CF. Patient-reported outcomes in cancer: a review of recent research and policy initiatives. *CA Cancer J Clin* 2007;**57**:278–300. doi:10.3322/ca.57.5.278
- 7 Boynton PM, Greenhalgh T. Selecting, designing, and developing your questionnaire. *Bmj* 2004;**328**:1312–5.
- 8 DoH. Improving Outcomes: A Strategy for Cancer. Department of Health 2011. <http://www.cancerscreening.nhs.uk/breastscreen/improving-outcomes-strategy-for-cancer.pdf>
- 9 Glaser A, Fraser L, Corner J. Patient-reported outcomes of cancer survivors in England 1-5 years after diagnosis: a cross-sectional study. . *BMJ Open* 2013.
- 10 DoH. Cancer Patient Experience Survey 2011/12. . London, UK: : Department of Health. 2012. [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/212860/Cancer-Patient-Experience-Survey-National-Report-2011-12](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/212860/Cancer-Patient-Experience-Survey-National-Report-2011-12)
- 11 Downing A, Morris EJ a, Richards M, *et al*. Health-related quality of life after colorectal cancer in England: a patient-reported outcomes study of individuals 12 to 36 months after diagnosis. *J Clin Oncol* 2015;**33**:616–24. doi:10.1200/JCO.2014.56.6539
- 12 Quality Health. National Cancer Patient Experience Survey. 2014.
- 13 Quality Health. 2014 National Cancer Patient Experience Survey Questionnaire (PDF). 2014.<https://www.quality-health.co.uk/resources/surveys/national-cancer-experience-survey/2014-national-cancer-patient-experience-survey/2014-national-cancer-patient-experience-survey-materials/689-2013-national-cancer-patient-experience-survey-questionnaire-pdf/file> (accessed 25 May2016).

27

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 14 Bone A, Mc Grath-Lone L, Day S, *et al.* Inequalities in the care experiences of patients with cancer: analysis of data from the National Cancer Patient Experience Survey 2011-2012. *BMJ Open* 2014;**4**:e004567. doi:10.1136/bmjopen-2013-004567
- 15 NHS WALES. Together for Health – Delivering End of Life Care A Delivery Plan up to 2016 for NHS Wales. Cardiff, Wales: 2016. <http://wales.gov.uk/docs/dhss/consultation/120913publicen.pdf>
- 16 Quality Health. Wales Cancer Patient Experience Survey. Wales: : NHS Wales 2014. <http://wales.gov.uk/topics/health/publications/health/reports/report13/?lang=en>
- 17 Mason J. *Qualitative Researching*. SAGE Publications 2002. <http://books.google.co.uk/books?id=WDT7AwAAQBAJ>
- 18 May C, Montori VM, Mair FS. We need minimally disruptive medicine. *BMJ* 2009;**339**.<http://www.bmj.com/content/339/bmj.b2803.abstract>
- 19 Bracher M, Wagland R, Corner J. Exploration and analysis of free-text comments from the 2013 Wales Cancer Patient Experience Survey (WCPES). Southampton, UK: 2014. <http://www.macmillan.org.uk/Documents/Fundraising/InYourLocalArea/Wales/WCPESReportFINALpdf.pdf> (accessed 27 Apr2015).
- 20 Risberg T, Sørbye S, Norum J, *et al.* Diagnostic delay causes more psychological distress in female than in male cancer patients. *Anticancer Res* Published Online First: 1995.<http://europepmc.org/abstract/med/8687166> (accessed 29 Apr2015).
- 21 Wiseman T, Lucas G, Sangha A, *et al.* Insights into the experiences of patients with cancer in London: framework analysis of free-text data from the National Cancer Patient Experience Survey 2012/2013 from the two London Integrated Cancer Systems. *BMJ Open* 2015;**5**:e007792. doi:10.1136/bmjopen-2015-007792
- 22 Davidson R, Mills ME. Cancer patients' satisfaction with communication, information and quality of care in a UK region. *Eur J Cancer Care (Engl)* 2005;**14**:83–90. doi:10.1111/j.1365-2354.2005.00530.x
- 23 Macleod U, Mitchell ED, Burgess C, *et al.* Risk factors for delayed presentation and referral of symptomatic cancer: evidence for common cancers. *Br J Cancer* 2009;**101**:S92–101. doi:10.1038/sj.bjc.6605398
- 24 Rose PW, Rubin G, Perera-Salazar R, *et al.* Explaining variation in cancer survival between 11 jurisdictions in the International Cancer Benchmarking Partnership: a primary care vignette survey. *BMJ Open* 2015;**5**:e007212. doi:10.1136/bmjopen-2014-007212
- 25 Epstein R, Street R. *Patient-centered communication in cancer care: promoting healing and reducing suffering*. Bethesda, MD, USA: : National Cancer Institute 2007. [http://outcomes.cancer.gov/areas/pcc/communication/pccm\\_flyer.pdf](http://outcomes.cancer.gov/areas/pcc/communication/pccm_flyer.pdf) (accessed 30 Apr2015).
- 26 Mishel MH, Germino BB, Lin L, *et al.* Managing uncertainty about treatment decision making in early stage prostate cancer: a randomized clinical trial. *Patient Educ Couns* 2009;**77**:349–59. doi:10.1016/j.pec.2009.09.009
- 27 Ayanian JZ, Zaslavsky AM, Arora NK, *et al.* Patients' Experiences With Care for Lung Cancer and Colorectal Cancer: Findings From the Cancer Care Outcomes Research and Surveillance Consortium. *J Clin Oncol* 2010;**28**:4154–61.
- 28

- 1 doi:10.1200/JCO.2009.27.3268
- 2
- 3 28 Harrison SE, Watson EK, Ward AM, *et al.* Cancer survivors' experiences of discharge
- 4 from hospital follow-up. *Eur J Cancer Care (Engl)* 2012;**21**:390–7. doi:10.1111/j.1365-
- 5 2354.2011.01312.x
- 6
- 7 29 Roberts KJ, Lepore SJ, Urken ML. Quality of life after thyroid cancer: an assessment of
- 8 patient needs and preferences for information and support. *J Cancer Educ*
- 9 2008;**23**:186–91.
- 10
- 11 30 Girgis A, Boyes A, Sanson-Fisher RW, *et al.* Perceived needs of women diagnosed with
- 12 breast cancer: rural versus urban location. *Aust N Z J Public Health* 2000;**24**:166–73.
- 13
- 14 31 Llewellyn CD, McGurk M, Weinman J. How satisfied are head and neck cancer (HNC)
- 15 patients with the information they receive pre-treatment? Results from the
- 16 satisfaction with cancer information profile (SCIP). *Oral Oncol* 2006;**42**:726–34.
- 17
- 18 32 Pollock K, Cox K, Howard P, *et al.* Service user experiences of information delivery
- 19 after a diagnosis of cancer: a qualitative study. *Support Care Cancer* 2008;**16**:963–73.
- 20
- 21 33 Griffiths P, Simon M, Richardson A, *et al.* Is a larger specialist nurse workforce in
- 22 cancer care associated with better patient experience? Cross-sectional study. *J Health*
- 23 *Serv Res Policy* 2013;**18**:39–46.
- 24
- 25 34 Armes J, Crowe M, Colbourne L, *et al.* Patients' supportive care needs beyond the end
- 26 of cancer treatment: a prospective, longitudinal survey. *J Clin Oncol* 2009;**27**:6172–9.
- 27
- 28 35 McIlmurray MB, Thomas C, Francis B, *et al.* The psychosocial needs of cancer patients:
- 29 findings from an observational study. *Eur J Cancer Care (Engl)* 2001;**10**:261–9.
- 30
- 31 36 Boberg EW, Gustafson DH, Hawkins RP, *et al.* Assessing the unmet information,
- 32 support and care delivery needs of men with prostate cancer. *Patient Educ Couns*
- 33 2003;**49**:233–42.
- 34
- 35 37 Hodgkinson K, Butow P, Hunt GE, *et al.* Breast cancer survivors' supportive care needs
- 36 2–10 years after diagnosis. *Support Care Cancer* 2007;**15**:515–23.
- 37
- 38 38 Wagland R, Recio-saucedo A, Simon M, *et al.* Development and testing of a text-
- 39 mining approach to analyse patients' comments on their experiences of colorectal
- 40 cancer care. *BMJ Qual Saf* 2015;:1–11. doi:10.1136/bmjqs-2015-004063
- 41
- 42 39 HS&DR - 14/156/15: PRESENT: Patient Reported Experience Survey Engineering of
- 43 Natural Text: developing practical automated analysis and dashboard representations
- 44 of cancer survey freetext answers.
- 45 2016.<http://www.nets.nihr.ac.uk/projects/hsdr/1415615> (accessed 26 Jan2016).
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58 29
- 59
- 60

Figure 1- Process of multi-stage coding.





## Appendix

Table 1 - Respondent demographics.<sup>7</sup>

<i>Demographic category</i>	<i>Demographic sub-category</i>	<i>Number of free-text respondents / as % of total free-text respondents (n = 4672)</i>	<i>Number of survey respondents / % of total survey respondents (n = 7352)</i>
<b>Tumour Group</b>	Breast	1168 / 25%	1717 / 23%
	Colorectal / Lower Gastrointestinal	722 / 15%	1112 / 15%
	Prostate	587 / 13%	954 / 13%
	Urological	455 / 10%	787 / 11%
	Haematological	405 / 9%	633 / 9%
	Gynaecological	340 / 7%	504 / 7%
	Lung	238 / 5%	427 / 6%
	Head and Neck	212 / 5%	332 / 5%
	Upper Gastrointestinal	210 / 4%	354 / 5%
	Other	161 / 3%	252 / 3%
	Skin	99 / 2%	163 / 2%
	Sarcoma	46 / 1%	64 / 1%
	Brain/Central Nervous System	29 / 1%	53 / 1%
<b>Health Board</b>	Velindre NHS Trust	1283 / 27%	2053 / 28%
	Betsi Cadwaladr University Health Board	1097 / 23%	1720 / 23%
	Abertawe Bro Morgannwg University Health Board	962 / 21%	1539 / 21%
	Hywel Dda University Health Board	699 / 15%	1069 / 15%
	Cardiff And Vale University Health Board	256 / 5%	384 / 5%
	Cwm Taf University Health Board	189 / 4%	307 / 4%
	Aneurin Bevan University Health Board	186 / 4%	280 / 4%
<b>Sex</b>	Male	2065 / 43%	3397 / 46%
	Female	2522 / 53%	3785 / 51%
	Data not available	85 / 2%	170 / 2%
<b>Approximate age range (years)</b>	95-99	9 / 0.2%	12 / 0.2%
	85-94	188 / 4%	332 / 5%
	75-84	1022 / 22%	1656 / 23%
	65-74	1509 / 32%	2315 / 31%
	55-64	920 / 13%	1386 / 19%
	45-54	428 / 11%	616 / 8%
	35-44	119 / 8%	178 / 2%
	25-34	41 / 6%	66 / 1%
	<=24	7 / 2%	15 / 0.2%
Data unavailable	429 / 9%	776 / 11%	
<b>Sexual Orientation</b>	Heterosexual / straight (opposite sex)	4275 / 92%	6595 / 90%
	Data not available	231 / 5%	459 / 6%
	Prefer not to answer	117 / 3%	222 / 3%
	Gay or Lesbian (same sex)	21 / 0.4%	27 / 0.4%

<sup>7</sup> Percentages for each demographic category may not add up to 100% due to rounding area.

	Bisexual (both sexes)	14 / 0.3%	19 / 0.3%
	Other	14 / 0.3%	30 / 0.4%
	Retired	2892 / 62%	4608 / 63%
	Full time employment	715 / 15%	1056 / 14%
	Part time employment	334 / 7%	497 / 7%
<b>Main employment status</b>	Unemployed – unable to work for health reasons	300 / 6%	494 / 7%
	Data not available	161 / 3%	266 / 4%
	Homemaker	127 / 3%	206 / 3%
	Other	108 / 2%	174 / 2%
	Unemployed – and seeking work	23 / 0.5%	36 / 0.5%
	Student (in education)	12 / 0.3%	15 / 0.2%
		Do not have long term condition	2631 / 56%
	Long-standing physical conditions	865 / 19%	1291 / 18%
<b>Longstanding conditions</b>	Long-standing illnesses, such as HIV diabetes, chronic heart disease, or epilepsy.	660 / 14%	1023 / 14%
	Deafness or severe hearing impairment	504 / 11%	852 / 12%
	Mental health conditions	117 / 3%	187 / 3%
	Blindness or partially sighted	106 / 2%	67 / 1%
	Learning disabilities	17 / 0.4%	31 / 0.4%
	White (British)	4467 / 96%	6991 / 95%
	White (Irish)	27 / 0.58%	44 / 1%
	White (other)	64 / 1%	102 / 1%
	Mixed (White and Black Carribean)	5 / 0.11%	7 / 0.1%
	Mixed (White and Black African)	1 / 0.02%	4 / 0.1%
	Mixed (White and Asian)	3 / 0.06%	4 / 0.1%
	Mixed (other)	2 / 0.04%	2 / 0.03%
<b>Ethnicity</b>	Indian	6 / 0.13%	7 / 0.1%
	Pakistani	3 / 0.06%	5 / 0.1%
	Bangladeshi	0 / 0.00%	1 / 0.01%
	Asian (other)	4 / 0.09%	6 / 0.1%
	Carribean	2 / 0.04%	3 / 0.04%
	African	4 / 0.09%	5 / 0.1%
	Black (other)	0 / 0.00%	0 / 0.00%
	Chinese	1 / 0.02%	5 / 0.1%
	Any other ethnic group	3 / 0.06%	4 / 0.1%
		Data unavailable	80 / 2%

**Table 2 – Results of Stage 1 analysis - no. of respondents providing positive and negative comments relating to categories of cancer patient experience.**

<i>Category</i>	<i>Negative respondents</i>	<i>Negative respondents as % of total category</i>	<i>Positive respondents</i>	<i>Positive respondents as % of total category</i>	<i>Total respondents</i> <sup>8</sup>	<i>Ratio (negative-to-positive comments, n : 1)</i>
Waiting for appointments	397	54%	342	46%	738	1.16
Communication between patients and staff (NOS <sup>9</sup> )	287	52%	287	52%	554	1
Surgery	181	33%	393	73%	541	0.46
General nursing	127	25%	402	78%	517	0.32
Hospital doctors	73	15%	411	86%	476	0.18
Investigations and diagnostic services	288	61%	198	42%	475	1.45
Consultants and specialists	72	15%	408	88%	465	0.18
Specialist nursing	108	25%	329	76%	433	0.33
GP	246	61%	161	40%	401	1.53
Chemotherapy	85	28%	233	77%	303	0.36
Aftercare (NOS)	199	69%	97	33%	290	2.05
Radiotherapy	67	27%	191	76%	251	0.35
Hospital environments	182	76%	53	22%	240	3.43
Communication between staff and/or institutions (NOS)	165	80%	44	21%	206	3.75
Waiting to be seen on the day	163	84%	31	16%	194	5.25
Travel-related issues during the cancer journey	122	76%	45	28%	161	2.71
Food and catering	128	84%	26	17%	153	4.94
Emotional, social, psychological support	94	71%	43	33%	132	2.19
Concerns about staffing levels	131	100%	-	#VALUE!	131	-
Oncology	31	26%	90	77%	117	0.34
Pain management	73	89%	10	12%	82	7.3
Out of hours and weekend care	60	88%	8	12%	68	7.50
Accident & Emergency	33	80%	8	20%	41	4.13

<sup>8</sup> As individual patients may have provided both a negative and a positive comment in a given area – the total number of respondents for a given category may be less than the sum of positive and negative respondents (i.e. each respondent would only be counted once for the area as a whole). Combined positive and negative percentage therefore may not equal 100%.

<sup>9</sup> Not otherwise specified – comments in these categories refer to generic aspects of experience (e.g. communication), which were not associated with a specific area of treatment or care (e.g. surgery).

33

Urology	7	18%	34	87%	39	0.21
Financial concerns	35	97%	3	8%	36	11.67
Physiotherapy	12	50%	12	50%	24	1
<b>Total</b>	<b>2313</b>	50%	<b>3818</b>	82%	<b>4672</b>	<b>0.60:1</b>

For peer review only

Table 3 - Results of Stage 3 analysis - number and percentages of respondents by theme/sub-theme

Theme	Sub-themes	Negative respondents (n)	Negative respondents (% of free-text respondents)	Positive respondents (n)	Positive respondents (% of free-text respondents)	Total respondents (n)	Total respondents (% of free-text respondents)	Total respondents (% of survey respondents)	Ratio (negative-to-positive, n : 1)
<b>Communication</b> <sup>10</sup>	Communication between patients and staff <sup>11</sup>	687	14.70%	809	17.32%	1472	31.51%	20.02%	0.85
	Communication between staff and/or institutions	208	4.45%	44	0.94%	252	5.39%	3.43%	4.73
	<i>Communication totals</i>	<i>854</i>	<i>18.28%</i>	<i>846</i>	<i>18.11%</i>	<i>1673</i>	<i>35.81%</i>	<i>22.76%</i>	<i>1.01</i>
<b>Waiting during the treatment and post-treatment phases</b>	Waiting for appointments	397	8.50%	342	7.32%	738	15.80%	10.04%	1.16
	Waiting on the day	163	3.49%	31	0.66%	194	4.15%	2.64%	5.26
	<i>Waiting totals</i>	<i>522</i>	<i>11.17%</i>	<i>372</i>	<i>7.96%</i>	<i>923</i>	<i>19.76%</i>	<i>12.55%</i>	<i>1.40</i>
<b>Staffing and resource levels</b>	Availability of aftercare <sup>12</sup>	217	4.64%	98	2.10%	312	6.68%	4.24%	2.21
	General comments about staffing levels (nursing and medical staff NOS)	266	5.69%	1	0.02%	267	5.71%	3.63%	266.00
	Availability and quality of staff on hospital wards at evening and weekends	62	1.33%	8	0.17%	70	1.50%	0.95%	7.75
	Availability of specialist nursing staff	63	1.35%	.	.	63	1.35%	0.86%	.
	<i>Staffing and resource levels totals</i>	<i>568</i>	<i>12.16%</i>	<i>107</i>	<i>2.29%</i>	<i>671</i>	<i>14.36%</i>	<i>0.86%</i>	<i>5.31</i>
<b>Speed and quality of diagnostic care</b>	GP role in diagnosis	159	3.40%	52	1.11%	211	4.52%	2.87%	3.06
	Investigations and diagnostic procedures <sup>13</sup>	173	3.70%	57	1.22%	193	4.13%	2.63%	3.04
	<i>Speed and quality of diagnostic care totals</i>	<i>314</i>	<i>6.72%</i>	<i>97</i>	<i>2.08%</i>	<i>411</i>	<i>8.80%</i>	<i>5.59%</i>	<i>3.24</i>
<b>Totals</b>		<b>1948</b>	<b>41.70%</b>	<b>1276</b>	<b>27.31%</b>	<b>3172</b>	<b>67.89%</b>	<b>43.14%</b>	<b>1.53</b>

<sup>10</sup> Sub-themes in this theme together all sub-categories of communication between patients and staff, whether NOS or belonging to a specific area of care or treatment (e.g. surgery).

<sup>12</sup> This sub theme collects all NOS and specific sub-categories relating to availability of aftercare.

<sup>13</sup> This sub-theme contains only comments relating to quality and speed of investigations and diagnostic services – other comments are excluded (see table 2).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

Category/sub-category	Respondents
Admin	117
AdminImprove	74
AdminPositive	43
Carers-Dependents-Family	27
CarDepFamImprove	24
CarDepFamPositive	2
Comorbidities	35
ComorbiditiesImprove	31
ComorbiditiesPositive	5
Facility closures	22
Finances	37
FinancesImprove	35
FinancesPositive	3
Travel	170
After care	312
AftercareImprove	217
AftercareE-S-MH-Improve	19
AftercareImproveNOS	170
InvestFollowImprove	28
AftercarePositive	98
AftercarePositiveNOS	83
InvestFollowPositive	17
BreastCare	157
Dressing-wound Care	32
Dress-woundCareImprove	29
Dress-woundCarePositive	3
EarlyDischarge	1
Emotional-Social-MH support	143
Emotional-Social-MHImprove	100
EmSocMHAAfterCareImprove	19
Emotional-Social-MHPositive	43
Environment	265
CleaningStaff	22
EnvironmentImprove	195
EnvBedLevelsImprove	60
EnvHospCleanImprove	22
EnvHospToiletImprove	16
EnvHospPrivacyImprove	22
EnvironmentPositive	53
BedLevelsPositive	1
EnvHospCleanPositive	18
Food-Catering	159
FoodCatImprove	132
FoodCatPositive	27
Hospital Infections	26
Nursing	1193
NursingImprove	421
NursAreas	102
NursAvailImprove(NOS)	32
NursComImprove	71
NursComImproveNOS	49
NursInfolImproveNOS	4
NursMannerImproveNOS	47
NursInfolImprove	7
NursMannerImprove	65
NursDisciplinesImprove	108
NursBreastImprove	19
NursBreastAvailImprove	8

1	NursCNSImprove	11
2	NursCNSAvailImprove	8
3	NursDistrictImprove	29
4	NursDistAvailImprove	11
5	NursKeyImprove	20
6	NursKeyAvailImprove	18
7	NursMacMilImprove	18
8	NursMacMilAvailImprove	9
9	NursSpecialImprove(NOS)	16
10	NursSpecialAvailImprove	13
11	NursImproveNOS	132
12	NursCareImproveNOS	72
13	NursOutOfHoursImprove	24
14	NursLevelsImprove	131
15	NursingPositive	811
16	NursComPositive	249
17	NursComPositiveNOS	2
18	NursInfoPositive	66
19	NursInfoPositiveNOS	38
20	NursMannerPositive	202
21	NursMannerPositiveNOS	153
22	NursDisciplinesPositive	329
23	NursBreastPositive	51
24	NursBreastManner-InfoPositive	18
25	NursChemoPositive	69
26	NursChemoInfoPositive	8
27	NursChemoMannerPositive	25
28	NursCNSPositive	45
29	NursCNSManner-InfoPositive	18
30	NursDistPositive	48
31	NursDistMannerPositive	10
32	NursKeyPositive	18
33	NursMacMilPositive	45
34	NursMacMilManner-InfoPositive	14
35	NursSpecialPositive(NOS)	70
36	NursSpecialInfoPositive(NOS)	6
37	NursSpecialMannerPositive(NOS)	19
38	NursPositiveNOS	410
39	Out of hours-Weekend(NOS)	71
40	OutofHours-WeekendImprove(NOS)	62
41	OutofHours-WeekendPositive(NOS)	8
42	Palliative Care	17
43	PalliativeCareImprove	4
44	PalliativeCarePositive	13
45	Staffing-Resource Levels	144
46	StaffingLevels(NOS)	142
47	StaffingLevelsImprove(NOS)	141
48	StaffingLevelsPositive(NOS)	1
49	Stoma	8
50	A&E	41
51	A&EImprove	33
52	A&EPositive	8
53	Anaesthesia	22
54	AnaesImprove	6
55	AnaesPositive	16
56	Chemotherapy	324
57	ChemolImprove	89
58	ChemoComImprove	28
59	ChemolInfoImprove	27
60		



1		
2	ChemolmproveNOS	61
3	ChemoPositive	236
4	ChemoComPositive	61
5	ChemoInfoPositive	15
6	ChemoMannerPositive	49
7	ChemoPositiveNOS	177
8	Choice-Treatment Options	64
9	Choice-TreatOpImprove	49
10	ChoiceTreatInfoDisImprove	27
11	Choice-TreatOpPositive	14
12	Clinical Trials	34
13	Communication	1377
14	CommunicationInterIntra-agency	252
15	CommunicationInterIntra-agencyImprove	208
16	ComInterIntraAgencyImproveNOS	171
17	NoteLossImproveNOS	15
18	CommunicationInterIntra-agencyPositiveNOS	44
19	CommunicationPatient-Provider	1171
20	CommunicationPatient-ProviderImprove	619
21	ComPatProvImproveNOS	315
22	InfoNOS	152
23	InfAfterNOS	8
24	InfDiagNOS	8
25	InfTreatNOS	62
26	MannerNOS	96
27	LangNOS	14
28	ManDiagNOS	57
29	CommunicationPatient-ProviderPositive	569
30	ComPatProvPosNOS	292
31	InfoPosNOS	77
32	InfoPosNOSMaterials	7
33	InfoPosNOSTelephone	12
34	ManDiagPosNOS	1
35	MannerPosNOS	217
36	MannerPersPosNOS	194
37	MannerProfPosNOS	74
38	Consultants-SpecialistsNOS	487
39	ConsultSpecialImprove	75
40	ConSpecAppSpeedImprove	11
41	ConSpecComImprove	47
42	ConSpecInfolImprove	25
43	ConSpecMannerImprove	22
44	ConsultSpecialPositive	415
45	ConSpecAppSpeedPositive	11
46	ConSpecComPositive	133
47	ConSpecAccessPositive	10
48	ConSpecInfoPositive	45
49	ConSpecMannerPositive	101
50	Dermatology	12
51	ENT	14
52	ENTImprove	6
53	ENTPositive	8
54	GP	422
55	GPIImprove	258
56	GPCarePDiagImprove	70
57	GPCondKnowImprove	18
58	GPServProvImprove	8
59	GPDiagImprove	159
60	GPDiagComImprove	13

1		
2	GPDiagSpeedImprove	39
3	GMisdiagImprove	35
4	GPreferralImprove	80
5	GPImproveNOS	33
6	GPPostive	165
7	GPDiagPositive	52
8	GPreferralPositive	41
9	GPPDiagCarePositive	43
10	GPPositiveNOS	70
11	Gynaecology	12
12	GynaelImprove	5
13	Gynaepositive	7
14	Haematology	27
15	HaemImprove	2
16	HaemPositive	25
17	HospitalDoctorsNOS	499
18	HospitalDoctorsImprove	79
19	HospDocComImprove	51
20	HospDocInfolImprove	26
21	HospDocLangImprove	4
22	HospDocMannerImprove	35
23	HospDocLevelsImprove	17
24	HospitalDoctorsPositive	422
25	HospDocComPositive	145
26	HospDocInfoPositive	11
27	HospDocMannerPositive	136
28	IntensiveCare	10
29	Investigations	502
30	InvestigationsImprove	306
31	InvestImproveNOS	104
32	InvestSpeedImprove	136
33	InvestInitialImprove	58
34	InvestMisImprove	37
35	WaitingResultsImprove	77
36	DiagWaitNOS	41
37	FollowResultsImproveNOS	4
38	TreatResultImproveNOS	9
39	WaitingResultsPositive	5
40	InvestigationsPositive	202
41	InvestPositiveNOS	81
42	InvestSpeedPositive	57
43	ScreeningPositive	65
44	ScreenBowelPos	25
45	ScreenBreastPos	28
46	Lymphodema	32
47	LympholImprove	19
48	LymphoPositive	15
49	Oncology	123
50	OncologyImprove	32
51	OncComImprove	20
52	OncInfolImprove	13
53	OncMannerImprove	7
54	OncologyPositive	94
55	OncComPositive	16
56	OncAccessPositive	2
57	OncInfoPositive	7
58	OncMannerPositive	10
59	OtherTreatments	50
60	OtherTreatImprove	35

1	OtherTreatPositive	15
2		
3	PainManagement	91
4	PainManagelImprove	82
5	PainChronicImprove	11
6	PainDischlImprove	8
7	PainWaitImprove	29
8	PainManagePositive	10
9	Pharmacy-Medication	80
10	PharmacyMedicationImprove	59
11	PharMedInfolImprove	5
12	PharmacyMedicationPositive	21
13	Physiotherapy	35
14	PhysiolImprove	12
15	PhysioPositive	13
16	PrivateTreatment-Care	89
17	PrivAfterCare	2
18	PrivDiag	59
19	PrivDiagAvoidDelay	45
20	PrivTreatCare	45
21	PrivNHSUnavail-Wait	18
22	PrivTreatCareImprove	4
23	PrivTreatCarePositive	23
24		
25	Radiotherapy	265
26	RadiotherapyImprove	71
27	RadComImprove	29
28	RadInfolImprove	24
29	RadMannerImprove	6
30	RadImproveNOS	42
31	RadiotherapyPositive	195
32	RadComPositive	74
33	RadInfoPositive	20
34	RadMannerPositive	65
35	RadPositiveNOS	121
36		
37	Respiratory	19
38	ResplImprove	6
39	RespPositive	14
40	Surgery	589
41	SurgeryImprove	190
42	SurgCancelDelayImprove	20
43	SurgComImprove	68
44	SurgInfolImprove	36
45	SurgLanglImprove	3
46	SurgMannerImprove	16
47	SurgeryImproveNOS	18
48	SurgFollowImprove	17
49	SurgPostOpImprove	54
50	PainPostOpImprove	16
51	SurgPostOpReclImprove	49
52	SurgPreOpImprove	18
53	SurgProcedImprove	15
54		
55	SurgeryPositive	407
56	SurgAppointSpeedPositive	78
57	SurgComPositive	63
58	SurgInfoPositive	19
59	SurgMannerPositive	47
60	SurgeryPositiveNOS	190
	SurgPostOpPositive	46
	SurgPostOpRecPositive	38
	SurgPreOpPositive	20

SurgProcedPositive	61
Urology	44
UrologyImprove	7
UrologyPositive	37
Waiting Times	930

For peer review only