



Fenton, G.E. and Halliday, D.M. and Mason, Rob and Bredy, Timothy W. and Stevenson, Carl W. (2016) Sex differences in learned fear expression and extinction involve altered gamma oscillations in medial prefrontal cortex. *Neurobiology of Learning and Memory*, 135 . pp. 66-72. ISSN 1095-9564

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/37029/8/Fenton%20et%20al%202016%20in%20press.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the Creative Commons Attribution Non-commercial No Derivatives licence and may be reused according to the conditions of the licence. For more details see: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

Sex differences in learned fear expression and extinction involve altered gamma oscillations in medial prefrontal cortex

†Georgina E. Fenton¹, David M. Halliday², Rob Mason³, Timothy W. Bredy^{4,5}, *Carl W. Stevenson¹

¹School of Biosciences, University of Nottingham, Sutton Bonington Campus, Loughborough, LE12 5RD, UK

²Department of Electronics, University of York, Heslington, York, YO10 5DD, UK

³School of Life Sciences, University of Nottingham, Queen's Medical Centre, Nottingham, NG7 2UH, UK

⁴Queensland Brain Institute, University of Queensland, Brisbane, Queensland 4072, Australia

⁵Department of Neurobiology and Behavior, University of California, Irvine, Irvine CA, 92697

Article type: Short Communication

Running title: Sex differences in learned fear & mPFC γ activity

Keywords: sex differences, fear conditioning, extinction, prelimbic, infralimbic, gamma oscillations

†Current Address:

Department of Neuroscience, Psychology and Behaviour
University of Leicester
University Road
Leicester
LE1 9HN
UK
Email: gef8@leicester.ac.uk

***Corresponding Author:**

Carl W. Stevenson
School of Biosciences
University of Nottingham
Sutton Bonington Campus
Loughborough
LE12 5RD
UK
Email: carl.stevenson@nottingham.ac.uk

Abstract

Sex differences in learned fear expression and extinction involve the medial prefrontal cortex (mPFC). We recently demonstrated that enhanced learned fear expression during auditory fear extinction and its recall is linked to persistent theta activation in the prelimbic (PL) but not infralimbic (IL) cortex of female rats. Emerging evidence indicates that gamma oscillations in mPFC are also implicated in the expression and extinction of learned fear. Therefore we re-examined our *in vivo* electrophysiology data and found that females showed persistent PL gamma activation during extinction and a failure of IL gamma activation during extinction recall. Altered prefrontal gamma oscillations thus accompany sex differences in learned fear expression and its extinction. These findings are relevant for understanding the neural basis of post-traumatic stress disorder, which is more prevalent in women and involves impaired extinction and mPFC dysfunction.

1. Introduction

Women show a greater prevalence of post-traumatic stress disorder (PTSD) than men [Glover et al. 2015; Maeng and Milad 2015], yet the neural mechanisms underlying this sex difference remain unclear. PTSD is characterized by impaired fear extinction [Milad et al. 2009; Jovanovic and Norrholm 2011]. This is the reduction in learned fear that results from repeated non-reinforced presentations of the conditioned stimulus (CS) [Herry et al. 2010]. PTSD is also associated with dysfunction of the medial prefrontal cortex (mPFC), a heterogeneous brain area important for mediating the expression and extinction of learned fear. Whereas the anterior cingulate cortex (ACC) is involved in learned fear expression, its extinction requires the ventromedial prefrontal cortex (vmPFC) [Linmann et al. 2012; Mueller et al. 2014]. Importantly, ACC and vmPFC dysfunction are observed in PTSD [Shin et al. 2009; Milad et al. 2009]. Accumulating evidence indicates sex differences in fear extinction that involve mPFC [Baran et al. 2010; Zeidan et al. 2011; Merz et al. 2012; Baker-Andresen et al. 2013; Rey et al. 2014]. However, the functional roles of different mPFC subregions in mediating sex differences in learned fear expression and extinction remain unclear.

Using *in vivo* electrophysiology, we have recently shown that sex differences in learned fear expression are linked to altered mPFC theta oscillations (4-12 Hz) in rats [Fenton et al. 2014]. Compared to males, females exhibited more fear during auditory fear extinction and its recall which was accompanied by persistent theta activation in prelimbic (PL) cortex, the rodent homolog of ACC. In contrast, we found no sex differences in theta activity in infralimbic (IL) cortex, the homologous area to vmPFC [Vidal-Gonzalez et al. 2006; Sierra-Mercado et al. 2011]. Compared to males, females also showed more contextual fear before extinction and extinction recall associated with persistent PL theta activation, whereas there was no accompanying sex difference in IL theta activation. Given the lack of sex differences in IL theta activity observed before and during extinction and its recall, we speculated that the enhanced fear shown by females was due to impaired contextual regulation of extinction rather than an extinction deficit *per se*. However, another possibility is that other types of oscillatory activity in mPFC are linked to sex differences in fear extinction.

Gamma oscillations (30-120 Hz) play a key role in prefrontal-dependent cognitive functions such as attentional processing [Benchenane et al. 2011]. Recent studies also indicate that both theta

and gamma synchrony between mPFC and other inter-connected areas are involved in various memory processes [Harris & Gordon 2015], including learned fear inhibition [Lesting et al. 2011; Courtin et al. 2014b; Stujenske et al. 2014; Wang et al. 2015]. Interestingly, emerging evidence indicates that oscillatory activity at lower gamma frequencies (~30-50 Hz) in mPFC plays a role in the extinction of learned fear. In mice, impaired fear extinction is associated with enhanced gamma activation in PL [Fitzgerald et al. 2014]. In humans, gamma activation in vmPFC accompanies the recall of fear extinction, while unsuccessful fear extinction recall is associated with a failure of vmPFC gamma activation [Mueller et al. 2014]. Therefore gamma oscillations in PL and IL might also be implicated in sex differences in learned fear expression and its extinction. Here we re-examined the local field potential (LFP) activity data from our recent study and analyzed PL and IL gamma oscillations before and during auditory fear extinction and its recall in males and females.

2. Material and methods

2.1. Animals and surgery

Young adult male and age-matched naturally cycling female Lister hooded rats (Harlan, UK) were used in these experiments, which were conducted with internal ethical approval and in accordance with the Animals (Scientific Procedures) Act 1986, UK. Electrodes were implanted into PL and IL under anesthesia and with analgesia as previously described [Fenton et al. 2014]. Rats were singly housed during recovery and behavioural testing, which started 10-14 days after surgery.

2.2. Behavioural testing

Auditory fear conditioning, extinction and extinction recall testing were conducted using two chambers described in detail elsewhere [Stevenson et al. 2009]. On Day 0 rats were habituated to contexts A and B (15 min each). On Day 1 rats underwent tone habituation (five tones alone; 30 s, 80 dB, 4 kHz, 2 min inter-trial interval (ITI)) followed by auditory fear conditioning (five tones co-terminating with footshock; 1 s, 0.5 mA, 2 min ITI) in context A. On Days 2 and 15 rats underwent extinction training and recall testing, respectively (30 tones alone; 30 sec ITI), in context B (Fig 1A). Freezing during tone alone presentations and tone-shock pairings on Day 1, and before (i.e. 2 min

prior to the first tone onset) and during tone presentations on Days 2 and 15, was scored manually and served as an index of learned fear.

2.3. In vivo electrophysiology

During behavioural testing LFP activity was recorded by connecting the electrodes to a preamplifier linked to a Plexon Recorder system (Plexon Inc, TX), via a headstage and a commutator. LFPs were band-pass filtered at 0.7-170 Hz and digitized at 1 kHz. All electrode placements in PL and IL were verified histologically as described previously [Fenton et al. 2014]. Only data from rats with confirmed electrode placements in PL and IL (Fig 2A) was included in the analysis.

2.4. Behavioural data analysis

The mean of freezing during two consecutive tones on Days 2 and 15 was used in the statistical analysis. Spontaneous fear recovery was calculated by dividing mean freezing during the first two tones on Day 15 by mean freezing during the last two tone-shock pairings on Day 1 and expressed as a percentage. Contextual fear before extinction and extinction recall was determined by assessing freezing before the first tone onset on Days 2 and 15. Freezing and spontaneous fear recovery were expressed as the mean + SEM. Sex differences in tone-induced freezing during conditioning, extinction and extinction recall were analyzed separately using two-way analysis of variance (ANOVA). Sex differences in spontaneous fear recovery were analyzed using an unpaired t-test. Differences in tone-induced freezing between extinction and extinction recall were analyzed separately in males and females using two-way ANOVA. Sex differences in contextual fear before extinction and extinction recall were analyzed using two-way ANOVA. Post-hoc comparisons were conducted using Bonferroni's testing. The significance level for all comparisons was set at $P < 0.05$.

2.5. In vivo electrophysiology data analysis

Using multi-taper spectral analysis [Fenton et al. 2013], spectral estimates of LFP activity in PL and IL (Fig 2B) were generated during early and late extinction and extinction recall by taking the mean of the first two and last two tones on Days 2 and 15 and pooling across males or females. Similarly, spectral estimates of LFP activity in the 2 min periods before tone onset on Days 2 and 15 were generated and pooled across males or females. Differences in LFP power during early vs late extinction, early vs late extinction recall, and before extinction vs before extinction recall were each

determined separately in males or females using the log ratio difference of spectra test [Diggle, 1990] and quantified statistically using 95% confidence intervals [Stevenson et al. 2007, 2008; Fenton et al. 2013, 2014]. The LFP analysis was restricted to the 30-45 Hz band, which coincides with the lower gamma frequencies recently implicated in fear extinction [Fitzgerald et al. 2014; Mueller et al. 2014].

3. Results

3.1. Females exhibit enhanced learned fear expression during extinction and extinction recall

Before presenting our mPFC gamma activity findings we first summarize the behavioural results from our recent study [Fenton et al. 2014]. We found no differences in freezing between males (n=9) and females (n=10) during the presentation of tones alone (data not shown) or tone-shock pairings during fear conditioning (Fig 1B). Females did show significantly greater tone-induced freezing during extinction (main effect of sex: $F_{(1,17)}=13.63$, $P<0.01$; Fig 1C) and extinction recall (main effect of sex: $F_{(1,17)}=27.41$, $P<0.001$; Fig 1D), indicating that females showed enhanced learned fear expression during extinction and its later recall. Females also showed enhanced spontaneous recovery of fear over time after extinction, as indicated by a significantly increased percentage of fear recovered during early extinction recall relative to late fear conditioning ($t_{17}=2.39$, $P<0.05$; Fig 1E). Despite this sex difference in spontaneous fear recovery both males (Fig 1F) and females (Fig 1G) showed savings of extinction, as indicated by a significant decrease in tone-induced freezing during extinction recall compared to extinction (males: day x block interaction: $F_{(14,112)}=5.89$, $P<0.0001$; Bonferroni's post-hoc test, Blocks 1-9: $P<0.05$) (females: day x block interaction: $F_{(14,126)}=2.48$, $P<0.001$; Bonferroni's post-hoc test: Blocks 2-11 and 15, $P<0.05$). Females also showed enhanced contextual fear before extinction and its recall, as indicated by a significant increase in freezing during the 2 min periods before tone presentations (main effect of sex: $F_{(1,17)}=8.70$, $P<0.01$; Fig 1H).

3.2. Females show persistent PL gamma activation during extinction and a failure of IL gamma activation during extinction retrieval

Changes in mPFC gamma activity during extinction are shown in Fig 3A. In males, gamma activity in PL was significantly decreased during late compared to early extinction ($P<0.05$), whereas IL gamma activity did not differ between late and early extinction. The differing patterns of gamma

activity in PL and IL, along with the low levels of PL-IL gamma synchrony that were observed throughout behavioural testing (coherence <0.1 ; data not shown), suggest that volume conduction of gamma oscillations between PL and IL was negligible. Females also showed no difference in gamma activity in IL between late and early extinction. However, in contrast to males, PL gamma activity did not change between late and early extinction in females. Sex differences were also observed in mPFC gamma activity during extinction recall (Fig 3B). In males, gamma activity in PL was significantly decreased during late compared to early extinction recall ($P<0.05$); conversely, IL gamma activity was significantly increased during late compared to early extinction recall ($P<0.05$). Females also showed a significant decrease in gamma activity in PL during late compared to early extinction recall ($P<0.05$). However, there was no difference in IL gamma activity between late and early extinction recall in females. Changes in mPFC gamma activity before extinction and extinction recall are shown in Fig 3C. In males, there was no difference in gamma activity in PL before extinction recall compared to before extinction, while IL gamma activity was significantly increased before extinction recall compared to before extinction ($P<0.05$). Females also showed no difference in PL gamma activity before extinction recall compared to before extinction, and a significant increase in gamma activity in IL before extinction recall compared to before extinction ($P<0.05$).

4. Discussion

We found similarities but also some notable differences between mPFC gamma and theta activation before and during extinction and its recall in males. As with theta activity, we found that PL gamma activity decreased during extinction and extinction recall and that gamma activity in IL increased during extinction recall, in keeping with the known roles of PL and IL in learned fear expression and extinction, respectively [Vidal-Gonzalez et al. 2006; Sierra-Mercado et al. 2011; Fenton et al. 2014]. However, we found no change in gamma activity in IL during extinction, which is different to the IL theta activation that we observed previously. This suggests that IL gamma oscillations might be preferentially involved in extinction memory rather than learning. We also found greater IL gamma activation before extinction recall compared to before extinction, which is similar to our observations for IL theta activity. In contrast, there were no differences in gamma activity in PL

before extinction recall compared to before extinction, whereas previously we found less PL theta activation before extinction recall compared to before extinction. This suggests that, in contrast to theta activity, PL gamma activity may not be involved in the contextual regulation of learned fear.

We also found sex differences in mPFC gamma activation during extinction and extinction recall. There was no change in PL gamma activation during extinction in females, which resembles our theta activation findings. In contrast to theta activity, females showed a small but significant decrease in PL gamma activity during extinction recall. As we observed in males, females showed no change in IL gamma activation during extinction, which again contrasts with the increased theta activation that we found previously. The most striking sex difference observed was related to IL gamma activity during extinction recall. In contrast to the gamma activation in males, there was no change in gamma activity in IL during extinction recall in females. This differs from the increase in IL theta activity that we observed previously during extinction recall in females. We also found no sex differences in PL or IL gamma activity before extinction recall compared to before extinction. Taken together, these results indicate that, compared to males, females showed sustained PL gamma activation during extinction and a failure of IL gamma activation during extinction recall.

Based on these new results and the emerging evidence implicating mPFC gamma oscillations in learned fear expression and extinction [Fitzgerald et al. 2014; Mueller et al. 2014], it is worth revisiting how we interpreted our behavioural results in light of our previous theta activity findings [Fenton et al. 2014]. We hypothesized that females showed enhanced learned fear expression during extinction and its recall as a result of impaired contextual regulation of extinction instead of resistance to extinction. This idea was based largely on our observation that sex differences in learned fear before and during extinction and extinction recall were associated with changes in PL, but not IL, theta activation. It also fits with evidence indicating a role for PL in the contextual renewal of fear after extinction [Orsini et al. 2011; Sharpe and Killcross 2015], together with known sex differences in contextual fear processing [Maren et al. 1994; Gupta et al. 2001]. Our finding of enhanced conditioned fear accompanied by sustained PL gamma activation during extinction in females adds to evidence indicating an important role for PL activity in learned fear expression [Vidal-Gonzalez et al. 2006; Sierra-Mercado et al. 2011; Fenton et al. 2014]. A recent study examining mouse strain

differences in fear extinction showed that enhanced gamma activation in PL is also associated with impaired extinction [Fitzgerald et al. 2014]. Although initial inspection of our behavioural results seems to agree with previous studies showing impaired fear extinction in females [Baran et al. 2009, 2010; Baker-Andresen et al. 2013], closer scrutiny of our data suggests that females exhibited intact extinction encoding. Both males and females showed extinction savings and there were no sex differences in IL activation during extinction. Females did exhibit enhanced spontaneous recovery of fear after extinction and a failure of IL gamma activation during extinction recall. Interestingly, the recall of fear extinction is linked to vmPFC gamma activation and unsuccessful extinction recall is associated with failed gamma activation in this region in humans [Mueller et al. 2014]. This suggests that female rats may have a specific impairment in the retrieval of extinction memory.

5. Conclusions

The present results confirm and extend our previous findings demonstrating sex differences in learned fear expression and extinction involving altered theta and gamma oscillations in mPFC (summarized in Table 1). Females showed enhanced conditioned fear and impaired extinction recall associated with persistent PL activation and a failure of IL activation, respectively, although the possibility remains that these sex differences may also involve changes in the contextual regulation of fear extinction. These findings have potential translational relevance given that PTSD is more prevalent in women, characterized by impaired inhibition of learned fear, and associated with mPFC dysfunction [Milad et al. 2009; Shin et al. 2009; Glover et al. 2015; Jovanovic and Norrholm 2011; Maeng and Milad 2015]. Further research is needed to determine the mechanisms underlying sex differences in learned fear inhibition and mPFC activity. Local GABA interneurons in mPFC, which regulate learned fear and its extinction, also modulate theta and gamma oscillations locally [Sangha et al. 2012; Courtin et al. 2014a; Glykos et al. 2015]. Moreover, functional connectivity within the neural circuitry mediating learned fear inhibition, which includes reciprocal projections between mPFC, amygdala and hippocampus [Herry et al. 2010], involves synchronized activity at theta and gamma frequencies [Lesting et al. 2011; Courtin et al. 2014b; Stujenske et al. 2014; Wang et al. 2015]. Importantly, sex differences in GABA signalling and interneuron expression in mPFC, as well

as hippocampus and amygdala function, have been reported [Maren et al. 1994; Gupta 2001; Blume et al. 2013; Cholanian et al. 2014; Marron Fernandez de Velasco et al. 2015]. Determining the role of gonadal hormones in mediating sex differences in the neural circuitry underpinning learned fear inhibition is another key issue to investigate in the future, given the evidence indicating their involvement in GABA signalling and neural circuit function during fear extinction [Zeidan et al. 2011; Merz et al. 2012; Cholanian et al. 2014; Rey et al. 2014; Mackenzie and Maguire, 2014]. Although we found sex differences in learned fear inhibition and mPFC gamma oscillations without accounting for variations in the phase of the females' estrous cycle, growing evidence indicates an important role for estrogen in modulating fear extinction [Glover et al. 2015; Maeng and Milad 2015].

Acknowledgements

This research was supported by the Australian Research Council (DP1096148; TWB) and a doctoral training grant from the University of Nottingham (GEF).

References

- Baker-Andresen D, Flavell CR, Li X, Bredy TW. 2013. Activation of BDNF signaling prevents the return of fear in female mice. *Learn Mem* 20:237-240.
- Baran SE, Armstrong CE, Niren DC, Conrad CD. 2010. Prefrontal cortex lesions and sex differences in fear extinction and perseveration. *Learn Mem* 17:267-278.
- Baran SE, Armstrong CE, Niren DC, Hanna JJ, Conrad CD. 2009. Chronic stress and sex differences on the recall of fear conditioning and extinction. *Neurobiol Learn Mem* 91:323-332.
- Benchenane K, Tiesinga PH, Battaglia FP. 2011. Oscillations in the prefrontal cortex: a gateway to memory and attention. *Curr Opin Neurobiol* 21:475-485.
- Blume SR, Hetzel A, Rosenkranz JA. 2013. Sex differences in the basolateral amygdala. *SFN Abstract* 673.01.
- Cholanian M, Lobzova A, Das B, Yelleswarapu C, Donaldson ST. 2014. Digital holographic microscopy discriminates sex differences in medial prefrontal cortex GABA neurons following amphetamine sensitization. *Pharmacol Biochem Behav* 124:326-332.
- Courtin J, Chaudun F, Rozeske RR, Karalis N, Gonzalez-Campo C, Wurtz H, Abdi A, Baufreton J, Bienvenu TC, Herry C. 2014a. Prefrontal parvalbumin interneurons shape neuronal activity to drive fear expression. *Nature* 505:92-96.
- Courtin J, Karalis N, Gonzalez-Campo C, Wurtz H, Herry C. 2014b. Persistence of amygdala gamma oscillations during extinction learning predicts spontaneous fear recovery. *Neurobiol Learn Mem* 113:82-89.
- Diggle PJ. 1990. Spectral analysis. In: Diggle PJ, editor. *Time series. A biostatistical introduction*. New York: Oxford University Press. p. 94–133.
- Fenton GE, Pollard AK, Halliday DM, Mason R, Bredy TW, Stevenson CW. 2014. Persistent prelimbic cortex activity contributes to enhanced learned fear expression in females. *Learn Mem* 21:55-60.
- Fenton GE, Spicer CH, Halliday DM, Mason R, Stevenson CW. 2013. Basolateral amygdala activity during the retrieval of associative learning under anesthesia. *Neuroscience* 13:146-156.

- Fitzgerald PJ, Whittle N, Flynn SM, Graybeal C, Pinard CR, Gunduz-Cinar O, Kravitz AV, Singewald N, Holmes A. 2014. Prefrontal single-unit firing associated with deficient extinction in mice. *Neurobiol Learn Mem* 113:69-81.
- Glover EM, Jovanovic T, Norrholm SD. 2015. Estrogen and extinction of fear memories: implications for posttraumatic stress disorder treatment. *Biol Psychiatry* 78:178-185.
- Glykos V, Whittington MA, LeBeau FE (2015) Subregional differences in the generation of fast network oscillations in the rat medial prefrontal cortex (mPFC) in vitro. *J Physiol* 593:3597-3615.
- Gupta RR, Sen S, Diepenhorst LL, Rudick CN, Maren S. 2001. Estrogen modulates sexually dimorphic contextual fear conditioning and hippocampal long-term potentiation (LTP) in rats. *Brain Res* 888:356-365.
- Harris AZ, Gordon JA. 2015. Long-range neural synchrony in behavior. *Annu Rev Neurosci* 38:171-94.
- Herry C, Ferraguti F, Singewald N, Letzkus JJ, Ehrlich I, Lüthi A. 2010. Neuronal circuits of fear extinction. *Eur J Neurosci* 31:599-612.
- Jovanovic T, Norrholm SD. 2011. Neural mechanisms of impaired fear inhibition in posttraumatic stress disorder. *Front Behav Neurosci* 5:44.
- Lesting J, Narayanan RT, Kluge C, Sangha S, Seidenbecher T, Pape HC. 2011. Patterns of coupled theta activity in amygdala-hippocampal-prefrontal cortical circuits during fear extinction. *PLoS One* 6:e21714.
- Linnman C, Zeidan MA, Furtak SC, Pitman RK, Quirk GJ, Milad MR. 2012. Resting amygdala and medial prefrontal metabolism predicts functional activation of the fear extinction circuit. *Am J Psychiatry* 169:415-423.
- MacKenzie G, Maguire J. 2014. The role of ovarian hormone-derived neurosteroids on the regulation of GABAA receptors in affective disorders. *Psychopharmacology* 231:3333-3342.
- Maeng LY, Milad MR. 2015. Sex differences in anxiety disorders: Interactions between fear, stress, and gonadal hormones. *Horm Behav* 76:106-117.

- Maren S, De Oca B, Fanselow MS. 1994. Sex differences in hippocampal long-term potentiation (LTP) and Pavlovian fear conditioning in rats: positive correlation between LTP and contextual learning. *Brain Res* 661:25-34.
- Marron Fernandez de Velasco E, Hearing M, Xia Z, Victoria NC, Luján R, Wickman K. 2015. Sex differences in GABABR-GIRK signaling in layer 5/6 pyramidal neurons of the mouse prelimbic cortex. *Neuropharmacology* 95:353-360.
- Merz CJ, Tabbert K, Schweckendiek J, Klucken T, Vaitl D, Stark R, Wolf OT. 2012. Neuronal correlates of extinction learning are modulated by sex hormones. *Soc Cogn Affect Neurosci* 7:819-830.
- Milad MR, Pitman RK, Ellis CB, Gold AL, Shin LM, Lasko NB, Zeidan MA, Handwerker K, Orr SP, Rauch SL. 2009. Neurobiological basis of failure to recall extinction memory in posttraumatic stress disorder. *Biol Psychiatry* 66:1075-1082.
- Mueller EM, Panitz C, Hermann C, Pizzagalli DA. 2014. Prefrontal oscillations during recall of conditioned and extinguished fear in humans. *J Neurosci* 34:7059-7066.
- Orsini CA, Kim JH, Knapska E, Maren S. 2011. Hippocampal and prefrontal projections to the basal amygdala mediate contextual regulation of fear after extinction. *J Neurosci* 31:17269-17277.
- Rey CD, Lipps J, Shansky RM. 2014. Dopamine D1 receptor activation rescues extinction impairments in low-estrogen female rats and induces cortical layer-specific activation changes in prefrontal-amygdala circuits. *Neuropsychopharmacology* 39:1282-1289.
- Sangha S, Ilenseer J, Sosulina L, Lesting J, Pape HC. 2012. Differential regulation of glutamic acid decarboxylase gene expression after extinction of a recent memory vs. intermediate memory. *Learn Mem* 19:194-200.
- Sharpe M, Killcross S. 2015. The prelimbic cortex uses contextual cues to modulate responding towards predictive stimuli during fear renewal. *Neurobiol Learn Mem* 118:20-29.
- Shin LM, Lasko NB, Macklin ML, Karpf RD, Milad MR, Orr SP, Goetz JM, Fischman AJ, Rauch SL, Pitman RK. 2009. Resting metabolic activity in the cingulate cortex and vulnerability to posttraumatic stress disorder. *Arch Gen Psychiatry* 66:1099-1107.

- Sierra-Mercado D, Padilla-Coreano N, Quirk GJ. 2011. Dissociable roles of prelimbic and infralimbic cortices, ventral hippocampus, and basolateral amygdala in the expression and extinction of conditioned fear. *Neuropsychopharmacology* 36:529-538.
- Stevenson CW, Spicer CH, Mason R, Marsden CA. 2009. Early life programming of fear conditioning and extinction in adult male rats. *Behav Brain Res* 205:505-510.
- Stevenson CW, Halliday DM, Marsden CA, Mason R. 2008. Early life programming of hemispheric lateralization and synchronization in the adult medial prefrontal cortex. *Neuroscience* 155:852-863.
- Stevenson CW, Halliday DM, Marsden CA, Mason R. 2007. Systemic administration of the benzodiazepine receptor partial inverse agonist FG-7142 disrupts corticolimbic network interactions. *Synapse* 61:646-663.
- Stujenske JM, Likhtik E, Topiwala MA, Gordon JA. 2014. Fear and safety engage competing patterns of theta-gamma coupling in the basolateral amygdala. *Neuron* 83:919-933.
- Vidal-Gonzalez I, Vidal-Gonzalez B, Rauch SL, Quirk GJ. 2006. Microstimulation reveals opposing influences of prelimbic and infralimbic cortex on the expression of conditioned fear. *Learn Mem* 13:728-733.
- Wang ME, Yuan RK, Keinath AT, Ramos Álvarez MM, Muzzio IA. 2015. Extinction of learned fear induces hippocampal place cell remapping. *J Neurosci* 35:9122-9136.
- Zeidan MA, Igoe SA, Linnman C, Vitalo A, Levine JB, Klibanski A, Goldstein JM, Milad MR. 2011. Estradiol modulates medial prefrontal cortex and amygdala activity during fear extinction in women and female rats. *Biol Psychiatry* 70:920-927.

Figure and Table Legends

Fig 1. Sex differences in learned fear before and during extinction and its later recall. (A) Schematic representation of the behavioural paradigm used. (B) There were no sex differences in freezing in response to tone-shock pairings during auditory fear conditioning. Females showed greater freezing in response to tone presentations during (C) extinction (** $P < 0.01$) and (D) extinction recall (** $P < 0.001$). (E) Females showed greater spontaneous fear recovery (i.e. freezing during early extinction recall relative to late fear conditioning; * $P < 0.05$). Compared to extinction, freezing during tone presentations was decreased during extinction recall in (F) males (* $P < 0.05$) and (G) females (* $P < 0.05$). (H) Females showed greater freezing before tone presentations during extinction and extinction recall (** $P < 0.01$) (figure adapted from [Fenton et al. 2014]).

Fig 2. (A) Representative example of electrode placements in PL and IL (indicated by the arrows). (B) Sample LFP traces recorded from PL and IL (figure adapted from [Fenton et al. 2014]).

Fig 3. Sex differences in mPFC gamma oscillations during extinction and its later recall. (A) Pooled gamma power spectra (left) and log ratio plots for pairwise comparisons of spectra (right) in PL (top) and IL (bottom) during the first and last tone blocks during extinction in males and females. Solid horizontal lines in the log ratio plots represent the upper and lower 95% confidence limits; positive log ratio values indicate increased power during the last compared to the first tone block, whereas negative values indicate decreased power during the last compared to the first tone block. In males, PL gamma power decreased during extinction, whereas gamma power in PL showed no change during extinction in females. There was no change in IL gamma power during extinction in males or females. (B) Pooled gamma power spectra (left) and log ratio plots (right) in PL (top) and IL (bottom) during the first and last tone blocks during extinction recall. Gamma power in PL decreased during extinction recall in males and, to a lesser extent, in females. In contrast, whereas IL gamma power increased during extinction recall in males, there was no change in gamma power in IL during extinction recall in females. (C) Pooled gamma power spectra (left) and log ratio plots (right) in PL (top) and IL (bottom) before tone presentations during extinction (Ext) and extinction recall (Ext Rec). Positive log ratio values indicate increased power before extinction recall compared to before extinction, whereas negative values indicate decreased power before extinction recall compared to

before extinction. There was no change in PL gamma power before extinction recall compared to before extinction in males or females. Gamma power in IL increased before extinction recall compared to before extinction in males and, to a lesser extent, in females.

Table 1. A summary of the changes in theta (θ) and gamma (γ) activation in PL and IL during fear extinction and its recall in male and female rats (\uparrow : increase; \downarrow : decrease; \leftrightarrow : no change).

Figure 1.




Figure 2.




Figure 3.




Table 1.

		Males		Females	
		θ	γ	θ	γ
During Extinction	PL	↓	↓	↑	↔
	IL	↑	↔	↑	↔
During Extinction Recall	PL	↓	↓	↔	↓
	IL	↑	↑	↑	↔