



UNITED KINGDOM · CHINA · MALAYSIA

England, Gary C.W. and Bright, L. and Pritchard, B. and Bowen, I. Mark and De Souza, M. and Silva, L.D.M. and Moxon, Rachel (2016) Canine reproductive ultrasound examination for predicting future sperm quality. *Reproduction in Domestic Animals*, 51 (Supp.3). pp. 1-6. ISSN 1439-0531

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/38759/1/Revised%20England%20Reproductive%20ultrasound%20assessment%2016%2008%2016.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see:

http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

1 **Canine reproductive ultrasound examination for predicting future sperm quality**

2

3

4 Running title: testis ultrasound to predict semen quality

5

6

7 Gary C.W. England^a, Lucy Bright^a, Beth Pritchard^a, I. Mark Bowen^a, Mírley Barbosa de

8 Souza^b, Lúcia Daniel Machado da Silva^b, Rachel Moxon^c

9 ¹School of Veterinary Medicine and Science, University of Nottingham, Sutton Bonington,

10 UK; ²Laboratory of Carnivore Reproduction, Veterinary School, Ceara State University,

11 Fortaleza, Brazil and ³National Breeding Centre, Guide Dogs, Leamington Spa, UK.

12

13

14 Corresponding author: gary.england@nottingham.ac.uk

15

16

17

18 **Contents**

19 The reproductive potential of male animals is commonly evaluated using a breeding
20 soundness examination incorporating B-mode ultrasound examination of the testes and
21 recently Doppler ultrasound examination of the testicular arteries. These techniques may
22 detect testicular normality or pathology, and while some measured parameters are associated
23 with semen quality at the time of ultrasound examination, few studies have investigated the
24 relationship with future semen quality.

25 We hypothesised that B-mode and Doppler ultrasound measurements would correlate with
26 future semen quality. Within two studies we investigated the relationship between ultrasound
27 measured testicular volume, testicular echogenicity, testicular homogeneity, subjective
28 assessment of the testicular parenchyma, testicular artery resistance index and pulsatility
29 index with subsequent semen quality. Fifty-five normal fertile dogs of which 29 had stable
30 semen quality and 26 had a subsequent decline in semen quality were examined during a six-
31 month period commencing 62 days after the ultrasound examination. Statistical analysis
32 showed that no ultrasound parameters were predictive of future total sperm output or
33 percentage live normal sperm. However, mean testicular echogenicity was positively related
34 to motility ($t = 2.202$, $P = 0.039$).

35 We conclude that quantitative ultrasound assessment of the appearance of the testicular
36 parenchyma has potential for evaluation of future semen quality in dogs.

37

38

39 **Keywords: B-mode, Doppler, ultrasound, dog, sperm motility, fertility**

40 **Introduction**

41 A breeding soundness examination is commonly performed to examine the breeding potential
42 of dogs. The procedure includes clinical examination of the reproductive tract, observation of
43 libido, examination of semen quality, and in some cases ultrasound examination of the
44 reproductive tract, and endocrine testing. More recently, Doppler ultrasonography has
45 allowed an additional evaluation of the reproductive organs, with useful information about
46 blood flow and velocity (Freitas et al., 2013; 2015). Our work to date has utilised B-mode
47 ultrasound to characterise the normal appearance of the testes and prostate gland and how
48 these are disturbed in cases of pathology (England, 1991; Souza et al., 2016). Recently, we
49 described digital image analysis of testicular and prostatic ultrasonographic echogenicity and
50 heterogeneity in dogs and their relation to semen quality (Moxon et al., 2015), and we
51 characterised differences of testicular artery blood flow measured using Doppler
52 ultrasonography in pre- and post-pubertal dogs (Souza et al., 2014; Souza et al., 2015a), and
53 in dogs with established infertility (Souza et al., 2015b). In these and other studies (Zelli et
54 al., 2013), measurements have generally been related to semen quality close to the time of
55 ultrasound examination rather than future semen quality as would be expected based upon the
56 time taken for spermatogenesis and sperm maturation. Interestingly, in a limited number of
57 studies in other species, some associations between the ultrasonographic appearance and
58 future semen quality have been established (Ahmadi et al., 2012; Brito et al., 2012; Arteaga
59 et al., 2015). Indeed, Arteaga et al. (2005) found that in bulls, testicular parenchymal pixel
60 intensity measured by ultrasound had a better association with future semen quality than with
61 present semen quality.

62

63 The aims of this retrospective clinical study were to examine whether testicular ultrasound
64 appearance and testicular artery blood flow measurements could be used to predict future

65 characteristics of semen quality. Two studies were performed with a total of 55 individual
66 dogs. In Study 1, testicular volume, testicular echogenicity, testicular homogeneity were
67 measured, and parenchymal appearance was subjectively scored in 24 normal fertile dogs of
68 which 11 had stable semen quality and 13 had a subsequent decline in semen quality of
69 unknown origin. In Study 2, testicular artery resistance index and pulsatility index were
70 measured in 31 dogs of which 18 had stable semen quality and 13 had a subsequent decline in
71 semen quality of unknown origin.

72

73

74 **Materials and methods**

75 *Study animals*

76 *Study 1: Testicular ultrasound appearance*

77 Twenty-four healthy stud dogs of two breeds (18 Labradors and six Golden retrievers) were
78 included. Dogs were aged 1.5 to 7.7 years (mean 4.3 ± 0.4 years) and weighed between 28.4
79 and 40.6 kg (mean 35.0 ± 0.5 kg). All dogs had proven normal semen quality and were fertile
80 based on achieving at least one pregnancy within six months prior to the study.

81

82 *Study 2: Testicular artery blood flow*

83 Thirty-one healthy stud dogs of two breeds (21 Labradors and 10 Golden retrievers) were
84 included. Dogs were aged 1.5 to 6.9 years (mean 4.3 ± 0.3 years) and weighed between 28.1
85 and 39.8 kg (mean 33.7 ± 0.6 kg). All dogs had proven normal semen quality and were fertile
86 in the six months prior to the study.

87

88 *Ultrasound examination*

89 *Study 1: Testicular volume and testicular ultrasound appearance*

90 Real time B-mode ultrasonography (Pandion 300s, Pie Data, UK) with a 10 MHz
91 mechanical-sector transducer was used to evaluate testicular volume and perform objective
92 and subjective assessment of testicular echogenicity and homogeneity. To measure testicular
93 volume, the testes were imaged in the sagittal, transverse and dorsal planes. The mediastinum
94 was used as a reference point for measuring the testicular length, width and height using
95 electronic callipers (Souza et al., 2014). Testicular volume was calculated using the formula
96 for an ellipse; volume = length x width x height x 0.5236 (Paltiel et al., 2002). Total testicular
97 volume (TTV) was calculated by adding together the volumes of each testis.

98

99 Mean testicular echogenicity (MTE) (calculated as the mean of nine values for testicular
100 echogenicity) and mean testicular homogeneity (MTH) (calculated as the standard deviation
101 of mean echogenicity) at the tissue level were determined using a semi-automated method as
102 previously described (Moxon et al., 2015). Briefly, two reference points (one in the near field
103 and one in the far field) were selected on the hyperechoic capsule of the testes (being selected
104 as the most echogenic structures identifiable) and a computer macro then randomly placed
105 nine sampling regions of interest (each 2.0 mm²) over the testicular parenchyma (avoiding the
106 central mediastinum). Within each region of interest, the mean pixel intensity (PI) was
107 measured. Of the two reference points the highest measurement of mean PI (most echogenic)
108 was used to calculate the echogenicity of the comparative region of interest as a percentage of
109 the highest mean pixel intensity using the following formula: Percentage echogenicity =
110 (Mean PI of capsule/Mean PI of testicular parenchyma) × 100. This methodology therefore
111 related all echogenicity measurements to a standard echogenic structure that would be
112 consistent between dogs. The mean of the nine echogenicity values for testicular echogenicity
113 were reported as mean echogenicity. Heterogeneity of testicular echogenicity (MTH) was
114 calculated as the standard deviation of the mean echogenicity of the regions of interest for

115 each organ. Using this method low values (low variation between regions of interest)
116 represented more homogenous tissues, whilst high values for heterogeneity (high variation
117 between the regions of interest) represented tissues that were less homogenous. These
118 measurements were at the tissue level rather than reporting visible gross changes.

119

120 Subjective assessment of testicular parenchymal echotexture was performed on recorded
121 digital images of the left and right testes. Each testis was classified into one of six categories
122 by the first author according to the echotexture of the testicular parenchyma. The image
123 classifications were; (1) hypoechoic, (2) normal echogenicity parenchyma with hypoechoic/
124 anechoic cysts, (3) normal echogenicity, (4) normal echogenicity parenchyma with echogenic
125 stippling, (5) hyperechoic parenchyma or (6) multiple changes. For data evaluation the
126 highest score from each testis was recorded as the testicular parenchymal echotexture for
127 each dog.

128

129 *Study 2: Testicular artery blood flow*

130 Doppler ultrasonography (M-Turbo, SonoSite[®], UK) with a 5 to 8 MHz micro-convex array
131 transducer was used to measure peak systolic velocity (PSV) and end diastolic velocity
132 (EDV). Dogs were restrained in the standing position and the transducer was placed at the
133 neck of the scrotum to identify the tortuous distal (looping) region of the supra-testicular
134 artery (Carrillo et al., 2012). The diameters of the testicular arteries were measured in the
135 longitudinal plane. The colour gain was adjusted to reduce any excess colour noise and the
136 pulsed Doppler gate was positioned within the lumen of the vessel. Three waves of a cardiac
137 cycle were used to measure mean testicular artery values for PSV and EDV, and these were
138 used by the machine software to calculate testicular artery resistance index (RI) and
139 pulsatility index (PI). This was repeated for each testis. The sample gate was 1 mm across all

140 regions. The angle of insonation used was less than 60°. The same operator performed each
141 examination. The mean of the two RI and two PI values from each testis were calculated and
142 used for analysis.

143

144 *Semen evaluation*

145 Immediately following ultrasound examination, semen was collected by digital manipulation
146 in the absence of a teaser bitch and evaluated using World Health Organisation standard
147 methods as described by England (1991). Only dogs with normal semen quality (Total sperm
148 output [TSO] $\geq 200 \times 10^6$ sperm; Percentage total live normal sperm [TLNS] $\geq 60\%$;
149 Percentage normal forward progressive motility [NFPM] $\geq 60\%$) were retained in the study.
150 Semen collection and evaluation was then undertaken between 1 and 7 further times for each
151 dog for the purposes of routine monitoring, for cryopreservation or insemination or when
152 there was concern that quality was declining. Samples collected within a six-month period
153 commencing 62 days after ultrasound examination were used for statistical analysis.

154

155 *Statistical analysis*

156 All data were examined for normality and were described as mean \pm one S.E. Semen quality
157 (TSO, TLNS, NFPM) after the initial assessment was calculated as a mean from all
158 subsequent collections (when there was more than one collection) and these values were used
159 for analysis. The difference in TLNS and NFPM between initial and subsequent semen
160 collections was determined using students t-tests or Wilcoxon signed-rank tests for paired
161 data.

162

163 For testicular ultrasound appearance, multiple regressions were used to determine whether
164 TTV, MTE, MTH and subjective appearance of the parenchyma were related to future semen

165 quality. Three multiple regressions were run; one for each semen quality measure (TSO,
166 TLNS, NFPM). Age and breed were included in the regressions and normality and co-
167 linearity were examined. For testicular artery blood flow, relationships between RI, PI, and
168 TSO, TLNS and NFPM were investigated using linear regression with TSO, TLNS and
169 NFPM as dependant variables. Data were analysed using XLStat (Addinsoft, USA) and SPSS
170 (version 20). Results were considered significant when $P < 0.05$.

171

172

173 **Results**

174 *Study 1: Testicular volume and testicular ultrasound appearance and future semen quality*

175 At the time of ultrasound examination TSO ranged from 200×10^6 to $2,470 \times 10^6$ (mean
176 $741.4 \pm 106.4 \times 10^6$); TLNS ranged from 64% to 87% (mean $75.7\% \pm 1.2\%$); NFPM ranged
177 from 70.0% to 95.0% (mean $87.3\% \pm 1.5\%$). At subsequent evaluations, semen quality was
178 stable for 11/24 dogs but declined for 13/24 dogs; of the latter, four dogs had poor quality
179 semen at subsequent evaluations (TLNS $< 60\%$; NFPM $< 60\%$). For the 13 dogs with semen
180 quality that declined, TNLS ($P = 0.04$) and NFPM ($P = 0.002$) were significantly lower at
181 subsequent evaluations than at the initial semen collected. In the monitoring period after
182 ultrasound examination, the range of the means for TSO was 336.0×10^6 to $3,234.0 \times 10^6$
183 (mean $1198.8 \pm 138.8 \times 10^6$), for TLNS was 46.0% to 95.0% (mean $76.1\% \pm 2.7\%$) and for
184 NFPM was 50.0% to 90.0% (mean $70.4\% \pm 1.7\%$).

185

186 TTV for the 24 dogs ranged from 36.1 to 60.5 cm^3 (mean $49.9 \pm 1.5 \text{ cm}^3$), MTE ranged from
187 45.3 to 96.9% (mean $61.3 \pm 1.9\%$) and MTH ranged from 4.3 to 9.8% (mean $6.6 \pm 0.3\%$).

188 Sixteen dogs had two testes that appeared normal and were classified as normal echogenicity

189 (3) and eight dogs were classified as not normal (scores 2, 4, 5 and 6) (Tables 1 and 2).

190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214

The multiple regression models for prediction of future semen quality (mean TSO, mean TLNS, mean NFPM) included all ultrasound parameters (TTV, MTE, MTH and subjective appearance). There were no ultrasound parameters that were predictive of future mean TSO or future mean TLNS. However, following model simplification, the model for NFPM predicted 27.9% of the variation ($F = 4.062$, $DF = 2$, $P = 0.032$), MTE was positively related to motility ($t = 2.202$, $P = 0.039$). Although the effect of age was not significant ($t = -1.949$, $p = 0.065$), age was retained in the final model as the effect of MTE on NFPM was lost on removal.

Study 2: Testicular artery blood flow and future semen quality

At the time of ultrasound examination TSO ranged from 200×10^6 to $2,448 \times 10^6$ (mean $986.5 \pm 95.3 \times 10^6$); TLNS ranged from 62.0% to 95.0% (mean $83.6\% \pm 1.4\%$); NFPM ranged from 60 to 90% (mean $79.2\% \pm 1.4\%$). At subsequent evaluations, semen quality was stable for 18/31 dogs but declined for 13/31 dogs; of the latter, four dogs had poor quality semen at subsequent evaluations (TLNS < 60%; NFPM < 60%). For the 13 dogs with semen quality that declined, TLNS ($P = 0.002$) and NFPM ($P = 0.002$) were significantly lower at subsequent evaluations than at the initial semen collected. In the monitoring period after ultrasound examination, the range of the means for TSO was 122.5×10^6 to $2,750.0 \times 10^6$ (mean 849.2 ± 86.5), for TLNS was 15.5% to 93.0% (mean $71.8\% \pm 3.4\%$) and for NFPM was 20.0% to 85.0% (mean $71.0\% \pm 2.5\%$).

Testicular artery RI ranged from 0.17 to 0.58 (mean 0.40 ± 0.02) and testicular artery PI ranged from 0.19 to 1.02 (mean 0.58 ± 0.04) (Table 3). Linear regression showed that there was no relationship between RI or PI at ultrasound examination and subsequent mean semen TSO (RI = $R^2 = 0.002$, $P = 0.796$, PI = $R^2 = 0.006$, $P = 0.687$) TLNS (RI = $R^2 = 0.008$, $P =$

215 0.635, $PI = R^2 = 0.010$, $P = 0.601$) or NFPM ($RI = R^2 = 0.001$, $P = 0.840$, $PI = R^2 = 0.002$, $P =$
216 0.819).

217

218 **Discussion**

219 B-mode and Doppler ultrasound examination of the reproductive tract of the dog provides
220 important and useful information about the state of the testes at the time of the examination
221 (Zelli et al., 2013; Moxon et al., 2015; Souza et al., 2015b). Studies to date have commonly
222 evaluated testicular echogenicity and heterogeneity, and testicular artery blood flow (Zelli et
223 al., 2013; Moxon et al., 2015; Souza et al., 2015b). The findings are perhaps not surprising
224 since previous work has shown that testicular parenchymal pixel intensity is associated
225 histologically with seminiferous tubule height, the proportion of tubules with a lumen, and
226 the size of the lumen (Evans et al., 1996; Giffin et al., 2009), and because blood flow through
227 the testicular artery is related to the rate of spermatogenesis (Kay et al., 1992). Interestingly,
228 in the present study we found a relationship between mean testicular echogenicity and future
229 semen quality in that higher testicular echogenicity at the tissue level was associated with
230 increased mean normal forward progressive motility in the subsequent monitoring period;
231 findings somewhat similar to those seen in the bull (Arteaga et al., 2005). The biological
232 reason for this is uncertain, however we postulate that this may be associated with a uniform
233 diameter of the seminiferous tubules rather than distortion of tubules sometimes seen cases of
234 pathology. There were however no associations between several ultrasonographic
235 characteristics and future semen quality; notably there was no relationship between total
236 testicular volume, mean testicular heterogeneity, subjective appearance of the testicular
237 parenchyma, testicular artery resistance index and testicular artery pulsatility index. The lack
238 of association between testicular volume and future semen quality is not surprising since
239 although this parameter is often purported to be predictive of current semen quality (Zelli et

240 al., 2013) this is not always the case (Souza et al., 2015b). Furthermore, relationships
241 between testicular artery blood flow are likely to relate more to current, rather than future,
242 semen quality since endothelial thickening and changes in blood flow occur secondarily to
243 testicular disease (Pinggera et al., 2008), while primary restriction of testicular artery
244 diameter results in rapid testicular changes (Kay et al., 1992). It was somewhat surprising that
245 in the present study no association was found between testicular parenchymal heterogeneity
246 or subjective appearance of the testicular parenchyma and subsequent semen quality, since
247 this has been observed in rams (Ahmadi et al., 2012). Interestingly, Ahmadi et al. (2012)
248 found stronger associations between epididymal echotexture and future semen quality than
249 they did between testicular echotexture and future semen quality. This is perhaps
250 unsurprising given the function of the epididymis as a sperm storage reservoir and site for
251 final sperm maturation, but importantly the study of Ahmadi et al. (2012) investigated semen
252 quality 60 days after ultrasound examination; an interval spanning one spermatogenic cycle
253 length. It is plausible that in the present study the method of calculating mean semen quality
254 throughout the six-month monitoring period (spanning 1-3 spermatogenic cycles [Soares et
255 al., 2009]) masked subtle changes in semen quality, although semen quality did decline in 26
256 of the 55 dogs across the two studies.

257 The present study is limited in scope in that the decline in semen quality was substantial in
258 only eight dogs, but offers a tantalising insight into the possibility that ultrasound can be used
259 to quantitatively assess pixel intensity representing physical properties of the testicular
260 parenchyma which are related to future semen quality. Further studies in this area and
261 extending to examination of epididymal appearance are warranted.

262

263

264 **Acknowledgements**

265 The authors thank Natasha White and Juliet Payne for animal technical assistance.

266

267

268 **References**

269

270 Arteaga AA, Barth AD, Brito LF, 2005; Relationship between semen quality and pixel-
271 intensity of testicular ultrasonograms after scrotal insulation in beef bulls. *Theriogenology*
272 **64**, 408-415.

273

274 Ahmadi B, Lau CP, Giffin J, Santos N, Hahnel A, Raeside J, Christie H, Bartlewski P, 2012;
275 Suitability of epididymal and testicular ultrasonography and computerized image analysis for
276 assessment of current and future semen quality in the ram. *Exp Biol Med* **237**, 186-193.

277

278 Brito LF, Barth AD, Wilde RE, Kastelic JP, 2012; Testicular ultrasonogram pixel intensity
279 during sexual development and its relationship with semen quality, sperm production, and
280 quantitative testicular histology in beef bulls. *Theriogenology* **78**, 69-76.

281

282 Carrillo JD, Soler M, Lucas X, Agut A, 2012; Colour and pulsed doppler ultrasonographic
283 study of the canine testis. *Reprod Domest Anim* **47**, 655–659.

284

285 England GCW, 1991; The relationship between ultrasonographic appearance, testicular size,
286 spermatozoal output and testicular lesions in the dog. *J Small Anim Pract* **32**, 306-311.

287

288 Evans AC, Pierson RA, Garcia A, McDougall LM, Hrudka F, Rawlings NC, 1996; Changes
289 in circulating hormone concentrations, testes histology and testes ultrasonography during
290 sexual maturation in beef bulls. *Theriogenology* **46**, 345-357.
291

292 Freitas LA, Pinto JN, Silva HVR, Uchoa DC, Mota Filho AC, da Silva LDM, 2013. Prostatic
293 and testicular doppler and echobiometry in Boxer dogs. *Acta Sci Vet* **41**, 1121-1129.
294

295 Freitas LA, Pinto JN, Silva HVR, da Silva LDM, 2015; Two-dimensional and Doppler
296 sonographic prostatic appearance of sexually intact French Bulldogs. *Theriogenology* **83**,
297 1140-1146.
298

299 Giffin JL, Franks SE, Rodriguez-Sosa JR, Hahnel A, Bartlewski PM, 2009; A study of
300 morphological and haemodynamic determinants of testicular echotexture characteristics in
301 the ram. *Exp Biol Med* **234**, 794-801.
302

303 Kay GW, Grobbelaar JA, Hattingh J, 1992; Effect of surgical restriction of growth of the
304 testicular artery on testis size and histology in bulls. *J Reprod Fertil* **96**, 549-553.
305

306 Moxon R, Bright L, Pritchard B, Bowen IM, de Souza MB, da Silva LD, England GC, 2015;
307 Digital image analysis of testicular and prostatic ultrasonographic echogenicity and
308 heterogeneity in dogs and the relation to semen quality. *Anim Reprod Sci* **160**, 112-119.
309

310 Paltiel HJ, Diamond DA, Di Canzio J, Zurakowski D, Borer JG, Atala A, 2002; Testicular
311 volume: comparison of orchidometer and US measurements in dogs. *Radiol* **222**, 114-119.
312

313 Pinggera GM, Mitterberger M, Bartsch G, Strasser H, Gradl J, Aigner F, Pallwein L,
314 Frauscher F, 2008; Assessment of the intratesticular resistive index by colour Doppler
315 ultrasonography measurements as a predictor of spermatogenesis. *Brit J Urol Intl* **101**, 722-
316 726.

317

318 Soares JM1, Avelar GF, França LR, 2009; The seminiferous epithelium cycle and its duration
319 in different breeds of dog (*Canis familiaris*). *J Anat* **215**, 462-471.

320

321 Souza MB, Barbosa, CC, Pereira, BS, Monteiro, CLB, Pinto, JN, Linhares, JCS, Silva, LDM,
322 2014; Doppler velocimetric parameters of the testicular artery in healthy dogs. *Res Vet Sci*
323 **96**, 533-36.

324

325 Souza MB, Barbosa CC, England GC, Mota Filho AC, Sousa CV, de Carvalho GG, Silva
326 HV, Pinto JN, Linhares JC, Silva LD, 2015a; Regional differences of testicular artery blood
327 flow in post pubertal and pre-pubertal dogs. *BMC Vet Res* **11**, 47-52.

328

329 Souza MB, England GC, Mota Filho AC, Ackermann CL, Sousa CV, de Carvalho GG, Silva
330 HV, Pinto JN, Linhares JC, Oba E, da Silva LD, 2015b; Semen quality, testicular B-mode
331 and Doppler ultrasound, and serum testosterone concentrations in dogs with established
332 infertility. *Theriogenology* **84**, 805-810.

333

334 Souza MB, Silva HV, Moxon R, Russo, M, England GC, 2016; B-mode, doppler and
335 contrast-enhanced ultrasonography of the prostate gland and testes of the dog. *In Practice* (in
336 press).

337

338 Zelli, R., Troisi, A., Ngonput, A.E., Cardinali, L. and Polisca, A., 2013; Evaluation of
339 testicular artery blood flow by Doppler ultrasonography as a predictor of spermatogenesis in
340 the dog. Res Vet Sci **95**, 632-637.

341 Table 1. The data for the 24 individual dogs showing each dog's age, breed, initial semen collection
 342 results (total sperm output [TSO], total live normal sperm [TLNS], normal forward progressive
 343 motility [NFPM]), total testicular volume (TTV) mean testicular echogenicity (MTE), mean testicular
 344 homogeneity (MTH), score for subjective appearance of the testes, number of subsequent semen
 345 collections and subsequent semen collection results (shown as a mean value where there was more
 346 than one subsequent collection in the six-month period). Four dogs were semen quality declined to a
 347 poor value are highlighted in bold.
 348

Dog		Initial semen collection								Subsequent semen collections				
Dog no.	Breed	Age	TSO (x 10 ⁶)	TLNS (%)	NFPM (%)	TTV (cm ³)	MTE (%)	MTH (%)	Subj. appearance score	N	Days from initial collection	TSO (x 10 ⁶)	TLNS (%)	NFPM (%)
1	Lab	5.42	360.0	87.0	80.0	36.26	56.69	4.65	2	1	219	375.0	92.0	70.0
2	Lab	1.65	749.0	74.0	90.0	40.67	52.24	7.68	3	1	128	720.0	82.0	75.0
3	Lab	2.28	390.0	75.0	90.0	48.68	61.99	5.85	6	1	224	480.0	89.0	90.0
4	GRet	5.56	660.0	75.0	95.0	49.21	59.98	4.32	3	1	171	912.0	52.0	50.0
5	Lab	5.17	2470.0	79.0	95.0	43.08	59.46	8.23	3	1	105	1417.0	56.0	70.0
6	Lab	5.30	592.5	77.0	95.0	54.26	45.27	7.82	4	1	171	336.0	60.0	50.0
7	Lab	4.30	272.5	84.0	85.0	54.51	54.69	6.27	4	1	133	625.0	95.0	70.0
8	Lab	1.52	200.0	64.0	80.0	56.12	63.67	6.69	3	1	181	510.0	85.0	65.0
9	Lab	2.06	301.0	71.0	85.0	60.45	53.35	5.71	4	1	186	1012.0	72.0	70.0
10	GRet	6.58	320.0	69.0	70.0	59.40	64.02	6.39	2	1	198	1060.0	46.0	70.0
11	Lab	1.81	580.0	68.0	90.0	37.05	64.95	7.15	3	1	171	1170.0	82.0	75.0
12	GRet	4.98	1425.0	78.0	95.0	58.52	54.51	8.43	3	1	174	2016.0	82.0	75.0
13	Lab	5.93	1029.0	70.0	70.0	59.59	96.90	7.63	3	1	95	897.0	71.0	80.0
14	Lab	5.62	1040.0	79.0	95.0	54.89	57.54	9.80	3	1	153	3234.0	82.0	75.0
15	Lab	5.09	465.0	73.0	80.0	44.80	62.52	4.52	3	2	161,238	1366.0	62.5	72.5
16	Lab	3.60	592.0	82.0	90.0	58.50	59.40	8.05	3	1	181	1008.0	85.0	75.0
17	Lab	4.56	551.0	86.0	95.0	36.06	63.65	8.42	3	2	139,165	832.5	89.0	67.5
18	Lab	5.60	629.0	70.0	85.0	50.57	66.48	6.15	3	2	152,173	1234.5	64.5	70.0
19	Lab	6.01	1662.5	75.0	90.0	53.25	57.76	5.52	4	2	145,243	1949.0	87.5	72.5
20	GRet	2.36	1020.0	72.0	90.0	50.84	66.85	5.62	3	3	167,177,184	2060.0	80.0	75.0
21	Lab	2.14	762.5	82.0	90.0	50.70	68.34	6.40	3	3	143,150,241	1182.3	80.7	71.7
22	GRet	7.68	390.0	80.0	90.0	49.58	62.44	5.29	3	4	169,175,176,183	1917.5	74.5	63.8
23	Lab	5.16	990.0	76.0	90.0	49.17	54.65	6.43	3	5	99,137,144,226,227	875.0	70.8	64.0
24	GRet	2.53	342.0	70.0	80.0	42.32	64.46	4.55	5	7	155,172,173,182,183,190,191	1583.1	86.4	73.6

349

350
351
352
353

Table 2. The results of subjective classification of the ultrasonographic appearance of the right and left testes of 24 fertile dogs with normal semen quality and the number of dogs with each overall score based on their highest (worst) score.

Classification	Description	Number of right testes scored	Number of left testes scored	Number of dogs with overall score
1	Hypoechoic	0	0	0
2	Normal echogenicity with hypoechoic/ anechoic cysts	2	2	2
3	Normal echogenicity	18	16	16
4	Normal echogenicity with echogenic stippling	4	4	4
5	Hyperechoic parenchyma	0	1	1
6	Multiple changes	0	1	1
Total		24	24	24

354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371

372 Table 3. The data for the 31 individual dogs showing each dog's age, breed, initial semen collection
 373 results (total sperm output [TSO], total live normal sperm [TLNS], normal forward progressive
 374 motility [NFPM]), testicular artery resistance index (RI) and pulsatility index (PI), number of
 375 subsequent semen collections and subsequent semen collection results (shown as a mean value where
 376 there was more than one subsequent collection in the six-month period). Four dogs were semen
 377 quality declined to a poor value are highlighted in bold.
 378

Dog		Initial semen collection						Subsequent semen collections				
Dog no.	Breed	Age	TSO (x 10 ⁶)	TLNS (%)	NFPM (%)	RI	PI	N	Days from initial collection	TSO (x 10 ⁶)	TLNS (%)	NFPM (%)
1	Lab	6.94	660.0	88.0	75.0	0.32	0.39	1	123	385.0	61.0	65.0
2	GRet	4.04	1600.0	91.0	80.0	0.46	0.69	1	245	768.0	74.0	70.0
3	GRet	4.04	424.0	87.0	70.0	0.30	0.41	1	161	570.0	72.0	75.0
4	Lab	3.04	330.0	89.0	80.0	0.32	0.40	1	216	802.0	93.0	70.0
5	Lab	5.69	865.0	87.0	85.0	0.36	0.52	1	156	1060.0	80.0	80.0
6	GRet	4.18	525.0	88.0	80.0	0.58	1.02	1	225	642.0	91.0	80.0
7	Lab	3.45	504.0	88.0	75.0	0.31	0.45	1	196	488.0	84.0	75.0
8	Lab	1.65	1230.0	87.0	85.0	0.38	0.51	1	207	180.0	77.0	80.0
9	Lab	6.07	2166.0	86.0	80.0	0.45	0.66	1	173	760.0	86.0	80.0
10	GRet	1.55	570.0	70.0	75.0	0.57	0.95	1	188	122.5	79.0	75.0
11	Lab	3.92	487.5	89.0	85.0	0.44	0.62	1	232	475.0	86.0	65.0
12	GRet	4.14	1000.0	62.0	70.0	0.36	0.52	1	233	357.5	22.0	20.0
13	Lab	4.25	1370.0	92.0	90.0	0.38	0.48	1	190	940.0	79.0	70.0
14	Lab	6.73	215.0	84.0	75.0	0.37	0.48	1	219	220.0	73.0	80.0
15	GRet	3.43	950.0	83.0	65.0	0.41	0.59	1	62	862.5	73.0	70.0
16	Lab	4.64	1515.0	85.0	80.0	0.55	0.81	1	217	941.0	89.0	85.0
17	Lab	4.49	1400.0	80.0	80.0	0.58	0.93	1	196	872.0	66.0	65.0
18	GRet	3.26	900.0	91.0	90.0	0.43	0.59	1	192	2750.0	81.0	85.0
19	Lab	2.24	1360.0	83.0	90.0	0.37	0.52	1	177	500.0	72.0	80.0
20	Lab	4.32	820.0	95.0	80.0	0.26	0.31	1	181	1100.0	91.0	80.0
21	Lab	4.71	2448.0	90.0	90.0	0.57	0.96	1	239	980.0	86.0	80.0
22	Lab	4.31	775.0	86.0	85.0	0.39	0.54	1	198	1025.0	68.0	70.0
23	GRet	5.53	650.0	62.0	65.0	0.42	0.61	2	152,203	740.0	37.0	35.0
24	GRet	4.47	200.0	84.0	75.0	0.39	0.53	2	133,235	1307.5	77.0	72.5
25	Lab	2.84	1295.0	86.0	85.0	0.17	0.19	2	78,221	1488.5	76.0	77.5
26	Lab	1.68	1455.0	85.0	85.0	0.26	0.31	2	133,195	710.0	39.0	67.5
27	Lab	4.13	985.0	80.0	90.0	0.47	0.72	2	65,102	1073.8	75.5	77.5
28	Lab	5.71	1297.5	74.0	70.0	0.36	0.50	2	83,211	804.0	81.5	70.0
29	Lab	4.08	1115.0	81.0	80.0	0.35	0.49	2	85,244	1270.5	81.5	82.5
30	Lab	5.92	875.5	82.0	80.0	0.48	0.73	3	131,159,182	1261.7	61.0	75.0
31	GRet	6.30	600.0	78.0	60.0	0.46	0.67	4	112,140,189,228	869.1	15.5	42.5

379