



The University of  
**Nottingham**

UNITED KINGDOM · CHINA · MALAYSIA

Moxon, Rachel and Bright, Lucy and Pritchard, Beth and Bowen, I. Mark and Souza, Mírley Barbosa de and Silva, Lúcia Daniel Machado da and England, Gary C.W. (2015) Digital image analysis of testicular and prostatic ultrasonographic echogenicity and heterogeneity in dogs and the relation to semen quality. *Animal Reproduction Science*, 160 . pp. 112-119. ISSN 1873-2232

**Access from the University of Nottingham repository:**

<http://eprints.nottingham.ac.uk/38918/1/Image%20analysis%20testes%20and%20prostate%20Animal%20Reproduction%20Science.pdf>

**Copyright and reuse:**

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the Creative Commons Attribution Non-commercial No Derivatives licence and may be reused according to the conditions of the licence. For more details see: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

**A note on versions:**

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact [eprints@nottingham.ac.uk](mailto:eprints@nottingham.ac.uk)

1 **Original article**

2

3 **Digital image analysis of testicular and prostatic ultrasonographic**  
4 **echogenicity and heterogeneity in dogs and the relation to semen quality.**

5

6 Rachel Moxon<sup>a</sup>, Lucy Bright<sup>b</sup>, Beth Pritchard<sup>b</sup>, Mark I Bowen<sup>b</sup>, Mirley Barbosa de  
7 Souza<sup>c</sup>, Lúcia Daniel Machado da Silva<sup>d</sup>, Gary C.W England<sup>b</sup>

8

9 <sup>a</sup> *Guide Dogs National Breeding Centre, Leamington Spa, Warwickshire, CV33*  
10 *9WF, UK.*

11 <sup>b</sup> *School of Veterinary Medicine and Science, The University of Nottingham,*  
12 *Sutton Bonington, Loughborough, LE12 5RD, UK.*

13 <sup>c</sup> *CAPES Foundation, Ministry of Education of Brazil, Brasilia – DF, CEP: 70.040-*  
14 *020, Brazil.*

15 <sup>d</sup> *Laboratory of Carnivore Reproduction, School of Veterinary Medicine, State*  
16 *University of Ceara 1700, Doutor Silas Munguba Avenue, CEP 60714-903,*  
17 *Fortaleza, CE, Brazil*

18

19

20 \*Corresponding author:

21 Tel: Tel.: +55.85.3101.9851, Fax: +55.85.3101.9840

22 Email address: [mirley.souza@gmail.com](mailto:mirley.souza@gmail.com)

23

24

25 **Abstract**

26 A semi-automated ultrasonographic method was developed to measure  
27 echogenicity and heterogeneity of the testes and prostate gland and relationships  
28 of these measures with semen quality were assessed in 43 fertile dogs. The  
29 relationship between animal age and body weight upon the volume of the testes,  
30 epididymal tail volume and prostate volume were also established.

31 Mean testicular echogenicity was negatively correlated with the percentage of  
32 morphologically normal live spermatozoa (more echogenic testes were  
33 associated with fewer normal sperm) but not with any other semen quality  
34 measure. Mean testicular heterogeneity was positively correlated with the total  
35 spermatozoal output (more heterogenous testes, being those with anechoic  
36 parenchyma and prominent echogenic stippling, were associated with greater  
37 sperm output) but not with any other semen quality measure. There was no  
38 relationship between either mean prostatic echogenicity or mean prostatic  
39 heterogeneity and any semen quality measure.

40 There was no relationship between age and any testicular or prostatic parameter;  
41 however bodyweight was significantly correlated with total testicular volume, total  
42 epididymal tail volume and total prostatic volume.

43 Testicular and prostatic ultrasonographic echogenicity and heterogeneity can be  
44 objectively assessed using digital image analysis and testicular echogenicity and  
45 heterogeneity may be useful adjunct measurements in a breeding soundness  
46 examination.

47

48 *Keywords:* canine; testes; prostate; semen quality; ultrasound

49

50 **Introduction**

51 Understanding reproductive function and fertility in the male is an essential  
52 element of breeding management in dogs. Commonly, reproductive potential is  
53 assessed by conducting a breeding soundness examination involving, amongst  
54 other things, clinical examination, ultrasound examination and semen collection  
55 and evaluation (Memon, 2007).

56  
57 B-mode real-time ultrasonography allows the accurate assessment of the size,  
58 shape, position, margination and internal architecture of the testes (England,  
59 1991; Eilts et al., 1993; Paltiel et al., 2002; Gouletsou et al., 2008; Souza et al.,  
60 2014) and prostate gland (Blum et al., 1985; Juniewicz et al., 1989; England,  
61 1991; Eilts et al., 1993; Ruel et al., 1998; Paltiel et al., 2002; Gouletsou et al.,  
62 2008; Freitas et al., 2013; Freitas et al., 2015). Ultrasonography also provides a  
63 valuable tool in assessing reproductive pathology (Cartee and Rowles, 1983;  
64 Feeney et al., 1987; Pugh and Konde, 1991; Cooney et al., 1992; England, 1995;  
65 Keenan, 1998; Nautrup and Tobias, 2001; Hecht, 2008).

66  
67 In clinical practice, ultrasound images are subjectively assessed and described in  
68 terms of their image texture; principally echogenicity and heterogeneity. A small  
69 number of studies have proposed a relationship between grossly detectable  
70 lesions within the testes and semen quality (England, 1991; Vencato et al., 2014).  
71 Objective analysis of echogenicity from measurements of pixel intensity is  
72 however possible using digital image analysis (Ivancic and Mai, 2008). This  
73 allows measurement of the characteristics of the tissue (Pierson and Adams,  
74 1995; Cardilli et al., 2010) and enables detection of changes in echogenicity

75 which may not be detected by the human eye (Rivers et al., 1996; Arteaga et al.,  
76 2005). Quantitative ultrasound measurement of ultrasonographic  
77 homogeneity/heterogeneity has also been previously assessed by calculation of  
78 the standard deviation of pixel intensity (Hershkovitz et al., 2010), and for testes  
79 ultrasound pixel heterogeneity has been directly correlated with tissue  
80 biochemical composition (Omer et al., 2012; Ahmadi et al., 2013). There are  
81 however only a few reports demonstrating a relationship between quantitative  
82 measurement of either testes echogenicity or heterogeneity and semen quality  
83 (Arteaga et al., 2005; Ahmadi et al., 2012). Kastelic and Brito (2012) proposed  
84 that the primary clinical use of ultrasonography was for grossly detectable lesions  
85 since quantitative pixel analysis was not predictive of semen quality in bulls. To  
86 date no quantitative ultrasonographic studies of the testes or prostate gland of  
87 the dog appear to have been published.

88  
89 The aim of this study was to measure testicular and prostatic ultrasonographic  
90 echogenicity and heterogeneity using digital image analysis, and investigate the  
91 relationships between these measures and semen quality in a group of known  
92 fertile dogs.

93

## 94 **Materials and Methods**

### 95 *Study animals*

96 Forty-three stud dogs (21 Labrador Retrievers, 12 Golden Retrievers, 6 German  
97 Shepherds, 1 Border Collie, 1 Flat Coated Retriever, 1 Irish Water Spaniel and 1  
98 Standard Poodle) with a mean weight of  $35.5 \pm 5.8$  kg (range 20.6 to 54.1 kg)  
99 aged between 1.1 and 9.3 years (mean  $4.2 \pm 2.0$  years) were examined. Dogs

100 were selected on the basis that they met the following inclusion criteria; (1)  
101 clinically healthy, (2) over one year of age, (3) **reproductively intact males** with no  
102 previous scrotal or prostatic surgery or exogenous hormone treatment, (4) proven  
103 fertility within the previous 6 months, and (5) having not ejaculated within the  
104 previous 48 hours.

105

#### 106 *Ultrasonographic measurements*

107 Ultrasound examinations of the testes, prostate **and epididymal tail** were  
108 undertaken once on each dog by the second and third authors respectively using  
109 a real time B-mode ultrasound machine (Pandion 300s, Pie Data UK Ltd.,  
110 Crawley, UK) with a 10MHz (testes) and 7.5MHz (prostate) mechanical-sector  
111 transducer. All machine settings including focal depth and gain settings were  
112 established at the first examination according to best image quality and remained  
113 unaltered for all remaining examinations which were performed over a **4-week**  
114 **period**.

115

116 The testes were imaged in the sagittal, transverse and dorsal planes and the  
117 prostate in the sagittal and transverse planes. Length, width and height of the  
118 testis and tail of the epididymis and of each lobe of the prostate were measured  
119 using the electronic callipers of the machine.

120

121 Testicular volume was calculated using the formula:  $\text{volume} = l \times w \times h \times 0.71$   
122 where  $l$  = sagittal diameter;  $w$  = transverse diameter and  $h$  = dorsal diameter  
123 (Hsieh et al., 2009; Gouletsou et al., 2008). Epididymal tail volume and the  
124 volume of each lobe of the prostate were calculated using the formula for an

125 ellipse: volume =  $l \times w \times h \times 0.523$  where  $l$  = length in a cranio-caudal direction  
126 (dorsal plane)  $w$  = sagittal diameter in a latero-medial direction (dorsal plane) and  
127  $h$  = sagittal diameter (sagittal plane).

128

#### 129 *Measurement of echogenicity and homogeneity*

130 Frozen digital images of the right and left testes in sagittal cross-section and of  
131 the prostate in the transverse plane were acquired onto a Laptop computer (Ergo  
132 Computing UK Ltd., Nottingham, UK) using video creation hardware (Dazzle  
133 Video Creator, Pinnacle Systems GmbH, Mountain View, California) and capture  
134 software ([www.virtualdub.org](http://www.virtualdub.org)). Images underwent semi-automated analysis using  
135 a macro developed with ImageJ software (National Institutes of Health, Bethesda,  
136 Maryland; <http://rsb.info.nih.gov/ij/>) to recover values for mean pixel intensity in the  
137 sampling window. Using this method the echogenicity of anechoic urine in the  
138 bladder and hepatic parenchyma of four normal 2 to 4 year old dogs were 6.7%  
139 and 78.0% respectively; higher percentage values representing structures that  
140 were more echogenic.

141

142 To measure echogenicity within the testes and prostate gland, two selected  
143 reference points (one in the near field and one in the far field) were selected on  
144 the hyperechoic capsule of the testes or the prostatic capsule (these being  
145 selected as the most echogenic structures identifiable). The computer macro then  
146 randomly placed nine sampling regions of interest (each 2.0 mm<sup>2</sup>) over the  
147 testicular parenchyma (avoiding the central mediastinum) (Figure 1), or three  
148 sampling regions within each lobe of the prostate gland. Within each region of  
149 interest the mean pixel intensity (PI) was measured. Of the two reference points

150 the highest measurement of mean PI (most echogenic) was used to calculate the  
151 echogenicity of the comparative region of interest as a percentage of the highest  
152 mean pixel intensity using the following formula:

153 Percentage echogenicity = (Mean PI of capsule / Mean PI of testicular or  
154 prostatic parenchyma) x 100. This methodology therefore related all echogenicity  
155 measurements to a standard echogenic structure that would be consistent  
156 between dogs.

157

158 The mean of the nine echogenicity values for testicular echogenicity and the  
159 mean of six echogenicity values for prostate echogenicity were reported as mean  
160 echogenicity for that organ.

161

162 Heterogeneity of testicular and prostatic echogenicity was calculated as the  
163 standard deviation of the mean echogenicity of the regions of interest for each  
164 organ. Using this method low values (low variation between regions of interest)  
165 represented more homogenous tissues, whilst high values for heterogeneity (high  
166 variation between the regions of interest) represented tissues that were less  
167 homogenous. These measurements were at the tissue level rather than reporting  
168 gross changes that would be observable by eye at the organ level.

169

#### 170 *Semen evaluation*

171 Semen was collected by digital manipulation in the absence of a teaser bitch  
172 (England, 1999a) immediately after the ultrasound examination. First and second  
173 fractions were collected and the combined volume recorded. Since the total  
174 duration of sexual excitement could vary between dogs, thus affecting the total



175 volume of the third (prostatic) fraction produced, prostatic fluid was collected over  
176 a defined time of 1-minute immediately following collection of the second  
177 fraction. Sperm motility (straight line velocity [VSL]) was objectively measured  
178 using computer image analysis (Hobson Tracking systems Ltd, Sheffield, UK) as  
179 previously described (Smith and England, 2001). The percentage of fast forward  
180 progressively motile sperm [PPFM] was determined as described by England  
181 (1999b). The percentage of morphologically normal live sperm (NLS) was  
182 assessed after examining 100 nigrosin-eosin stained spermatozoa at x 1000  
183 magnification, sperm concentration was measured using a haemocytometer, and  
184 total spermatozoal output (TSO) was calculated as previously described  
185 (England, 1999a).

186

#### 187 *Data analysis*

188 Measurements taken from the right and left testes, and the right and left prostatic  
189 lobes were compared using tests of difference. These measurements were then  
190 combined to provide values for total testicular volume (TTV), total epididymal tail  
191 volume (TEV) and total prostatic volume (TPV) volume.

192 Mean testicular echogenicity (MTE), mean testicular heterogeneity (MTH), mean  
193 prostatic echogenicity (MPE) and mean prostatic heterogeneity (MPH) were  
194 calculated for each dog.

195 Linear regression was used to determine whether age was related to the four  
196 ultrasound parameters (MTE, MTH, MPE and MPH). Relationships between  
197 MTE, MTH, MPE and MPH and semen quality were initially investigated using  
198 Spearman's rank correlation tests. Potential relationships were further  
199 investigated using linear regression. Relationships between age, bodyweight and

200 TTV, TEV and TPV and were investigated using linear regression with age and  
201 bodyweight as explanatory variables.

202 Statistical analysis was performed using XLStat software (Addinsoft, USA).

203 Values were considered significant when  $P < 0.05$ .

204

## 205 **Results**

206 There were no differences between measurements of the left compared with right  
207 testes, or left compared with right prostatic lobes.

208

209 There was substantial variation between the ultrasonographic measurements for  
210 the 43 dogs (Table 1). Mean straight line velocity (VSL) was  $16.7 \pm 1.2$  mm/s  
211 (range 8 to 39 mm/s), the mean percentage of fast forward progressively motile  
212 sperm was  $88 \pm 1.71\%$  (range 35 to 95%), - the mean percentage of  
213 morphologically normal live sperm (NLS) was  $76.0 \pm 1.1\%$  (range 57 to 88%),  
214 mean sperm concentration was  $463.9 \times 10^6 \pm 50.7 \times 10^6$  (range 95 to 1650 x  
215  $10^6$ ), and mean total spermatozoal output (TSO) was  $863.5 \times 10^6 \pm 74.1 \times 10^6$   
216 (range 180 to 1785 x  $10^6$ ).

217

218 Age was not related to MTE ( $r^2 = 0.007$ ,  $P = 0.583$ ), MTH ( $r^2 = 0.005$ ,  $P = 0.656$ ),  
219 MPE ( $r^2 = 0.033$ ,  $P = 0.240$ ) or MPH ( $r^2 = 0.012$ ,  $P = 0.483$ ). Mean testicular  
220 echogenicity was negatively correlated with the percentage morphologically  
221 normal live sperm ( $r = -0.455$ ,  $P = 0.003$ ,  $r^2 = 0.165$ ,  $P = 0.009$ ); more echogenic  
222 testes were associated with fewer normal sperm (Figure 2). There were no  
223 relationships between MTE and any other semen quality measure. Mean  
224 testicular heterogeneity was positively correlated with total spermatozoal output (r

225 = 0.416, P = 0.007,  $r^2 = 0.220$ , P = 0.002); more heterogenous testes at the  
226 tissue level were associated with greater sperm output (Figure 3). There were no  
227 relationships between MTH and any other semen quality measure. There were  
228 no relationships between mean prostatic echogenicity or mean prostatic  
229 heterogeneity and any measure of semen quality.

230

231 There was no relationship between age and any of the ultrasonographic  
232 measurements of testes, epididymis or prostate volume. Bodyweight was  
233 significantly related to TTV ( $r^2 = 0.27$ , P < 0.001), TEV ( $r^2 = 0.25$ , P = 0.001) and  
234 TPV ( $r^2 = 0.21$ , P = 0.002) (Figures 4a, 4b and 4c).

235

## 236 Discussion

237 This study assessed testicular and prostatic parenchymal echogenicity by  
238 measurement of pixel intensity in various regions of interest compared with an  
239 anatomically consistent echogenic reference point. This method has advantages  
240 over simple measurement of pixel intensity performed in the work of other  
241 authors (Pierson and Adams, 1995; Arteaga et al., 2005; Ivancic and Mai, 2008;  
242 Cardilli et al., 2010), where tissue echogenicity could be influenced by sound  
243 attenuation caused by subcutaneous tissue and tissue depth. This study also  
244 measured heterogeneity by examining the standard deviation of the mean pixel  
245 intensity of the regions of interest. A refinement of the methodology would be to  
246 sample a greater number of regions of a smaller area and perhaps consider  
247 statistical evaluation of the range of values for pixel intensity rather than to  
248 calculate the mean value, which by its nature tends to smooth the data.

249

250 Interestingly, within this population of recently fertile dogs there was substantial  
251 variation in semen quality and in many of the testicular and prostatic ultrasound  
252 measurements. Comparison between ultrasound parameters and semen quality  
253 showed that there was a significant negative correlation between mean testicular  
254 echogenicity and the percentage of morphologically normal live spermatozoa.  
255 Testes that were more echogenic were associated with fewer normal sperm in  
256 the ejaculate. Ahmandi et al. (2012) also studied fertile males but were unable  
257 to demonstrate a significant relationship between testicular echogenicity and  
258 semen quality, although their study included only six animals. We are uncertain of  
259 the histological variations that were present in our population of dogs that  
260 resulted in increased testicular echogenicity and were associated with poorer  
261 morphology, but not total sperm output. Presumably, in these dogs Sertoli cell  
262 numbers were normal, but subtle microstructural changes were present that  
263 affected sperm morphology. Induction of testicular pathology by scrotal insulation  
264 has been shown to cause changes in testicular echogenicity and associated  
265 changes in sperm morphology (Artega et al., 2005). That study found that  
266 hypoechoic testes were associated with poor morphology. We propose that either  
267 increased or decreased testicular echogenicity may reflect ultrastructural  
268 changes within the testes that are associated with altered sperm morphology.

269  
270 The present study also demonstrated that mean testicular heterogeneity was  
271 positively correlated with the total spermatozoal output; less heterogeneous  
272 testes were associated with reduced sperm output. It is important to recognise  
273 that heterogeneity was measured as the variation in pixel intensity within small  
274 sampling windows, such that testes which had an almost anechoic parenchyma

**Commented [RM1]:** Gary – we have said at the start of the discussion that we think their method could have led to echogenicity being affected by tissue depth etc, could that explain anything?

275 with prominent echogenic stippling were recorded as high heterogeneity.  
276 Presumably, focal regions of increased homogeneity (reduced variation between  
277 the anechoic parenchyma and the echogenic stippling) reflects a reduced density  
278 of fluid-containing seminiferous tubules accounting for reduced total sperm output  
279 but not changes in sperm morphology. Finding a relationship between testicular  
280 parenchymal heterogeneity and semen quality has only previously been reported  
281 in a small study in rams, where an inverse correlation was found between  
282 heterogeneity measurements and the percentage of sperm with normal  
283 morphology and progressive motility in samples collected 60 days after the  
284 ultrasound examination (Ahmadi et al., 2012). The present study found no  
285 relationship between sperm morphology or several objective measures of motility,  
286 but we collected and evaluated semen immediately after the ultrasound  
287 examination unlike the study of Ahmadi et al. (2012).

288  
289 An important clarification for clinical use of quantitative measurement of pixel  
290 intensity and calculation of the variation of pixel intensity is that these are  
291 measures at the level of the parenchyma, and are not a gross or overall  
292 assessment at the level of the organ which is a very different concept. Indeed,  
293 testes that have an overall heterogenous appearance characterised by irregular  
294 and diffuse echogenic structures within the parenchyma are often associated with  
295 low sperm output (Vencato et al., 2014).

296  
297 No relationships were found between echogenicity and heterogeneity of the  
298 prostate gland or any other measures of the prostate gland and semen quality.  
299 These findings are not surprising since the prostate gland solely contributes fluid

300 to the first and third fractions of the ejaculate (England et al., 1990), although the  
301 volumes of these fluids did not relate to any measurement of the prostate in the  
302 present study, unlike previous observations by Wheaton et al. (1979) who found  
303 that seminal volume was correlated to prostatic size. This may relate to the  
304 collection of a mixed first and second fraction and the time-restricted collection of  
305 third fraction in the present study, **since ejaculation may have been completed**  
306 **within 1 minute in some dogs but not others.**

307

308 Examination of the data collected from the left and right testes, and the left and  
309 right lobes of the prostate gland found no differences in any of the  
310 ultrasonographic measurements, similar to the findings of other authors (Pugh et  
311 al., 1990; England, 1991; Villaverde et al., 2014) but in contrast to the work of  
312 Souza et al. (2012). **In this study, ultrasonographic measurements of the left**  
313 **testis were higher than the right testis in two different breeds. However, a small**  
314 **number of dogs were used, in contrast with the present study, and a larger**  
315 **number of animals could demonstrate different results. Total testicular volume,**  
316 total epididymal tail volume and total prostatic volume were positively correlated  
317 ~~to~~with bodyweight, similar to the findings previously reported in other studies  
318 (Amann, 1986; Woodall and Johnstone, 1988; Ruel et al., 1998; Atalan et al.,  
319 1999a,b).

320

321 This study intentionally included dogs from a large age range and found no  
322 relationship between age and echogenicity or heterogeneity results from the  
323 testes or prostate, suggesting that age was not a contributing factor to the  
324 differences in ultrasonic appearance observed. No relationship was found

325 between age of the dog and total testicular volume, total epididymal tail volume  
326 and total prostatic volume, unlike the work of Mantziaras et al. (2014) who found  
327 a tendency for testes volume to increase until approximately 6 years of age and  
328 then to decrease. In contrast with the present study, prostatic volume has also  
329 been found to be related to age **by other authors** (Ruel et al., 1998; Atalan et al.,  
330 1999a,b; Lowseth et al., 1990; Mantziaras et al., 2014). It is possible that **the**  
331 inclusion criteria of the present study, requiring the dog to have been recently  
332 fertile, may have eliminated some older dogs with testicular disease or prostatic  
333 enlargement which may have been included in other studies. **In particular,**  
334 **previous work demonstration a link between age and prostatic size has frequently**  
335 **included dogs with prostate pathology (Brendler et al., 1983; Kay et al., 1989;**  
336 **Nielsen et al., 1990).**

337

### 338 **Conclusion**

339 Testicular and prostatic echogenicity and heterogeneity can be objectively  
340 measured by means of a semi-automated method using conventional ultrasound  
341 equipment. Relationships were apparent between mean testicular echogenicity  
342 and mean testicular heterogeneity with semen quality, such that superior semen  
343 samples was observed in testes that were less echogenic and had greater  
344 heterogeneity at the tissue level; corresponding to testes with an anechoic  
345 parenchyma with prominent echogenic stippling.

346 It is feasible that objective measurement of testicular echogenicity and  
347 heterogeneity may be useful adjunct measurements in a breeding soundness  
348 examination.

349

350 **Conflict of interest statement**

351 None of the authors has any financial or personal relationships that could  
352 inappropriately influence or bias the content of the paper.

353

354 **Acknowledgements**

355 The authors would like to acknowledge the work of Helen Freeman in support of  
356 this research. The authors ~~also~~ would also like to thank the funding agencies  
357 CNPq and CAPES, for the scholarship granted to M.B de Souza (process  
358 number BEX 3197/14-0).

359

360

361 **References**

362

363 Ahmadi, B., Lau, C.P., Giffin, J., Santos, N., Hahnel, A., Raesside, J., Christie,  
364 H., Bartlewski, P., 2012. Suitability of epididymal and testicular ultrasonography  
365 and computerized image analysis for assessment of current and future semen  
366 quality in the ram. *Exp. Biol. Med.* 237, 186-193.

367

368 Ahmadi, B., Mirshahi, A., Giffin, J., Oliveira, M.E., Gao, L., Hahnel, B., Bartlewski,  
369 P., 2013. Preliminary assessment of the quantitative relationships between  
370 testicular tissue composition and ultrasonographic image attributes in the ram.  
371 *Vet. J.* 198, 282-285.

372



373 Amann, R.P., 1986. Reproductive physiology and endocrinology of the dog. in:  
374 Morrow, D.A. (ed). Current Therapy in Theriogenology II. W. B. Saunders,  
375 Philadelphia, pp. 532-538.  
376

377 Arteaga, A.A., Barth, A.D., Brito, L.F.C., 2005. Relationship between semen  
378 quality and pixel-intensity of testicular ultrasonograms after scrotal insulation in  
379 beef bulls. Theriogenology. 64, 408-415.  
380

381 Atalan, G., Holt, P., Barr, F., Brown, P.J., 1999a. Ultrasonographic estimation of  
382 prostatic size in canine cadavers. Res. Vet. Sci. 67, 7-15.  
383

384 Atalan, G., Holt, P., Barr, F., 1999b. Ultrasonographic estimation of prostate size  
385 in normal dogs and relationship to bodyweight and age. J. Small Anim. Pract. 40,  
386 119-122.  
387

388 Brendler, C.B., Berry, S.J., Ewing, L.L., Mccollough, A.R., Cochran, R.C.,  
389 Strandberg, J.D., Zirkin, B.R., Coffey, D.S., Wheaton, L.G., Hiler, M.L., Bordy,  
390 M.J., Niswender, G.D., Scott, W.W., Walsh, P.C., 1983. Spontaneous benign  
391 prostatic hyperplasia in the Beagle. Age-associated changes in serum hormone  
392 levels, and the morphology and secretory function of the canine prostate. J Clin  
393 Invest. 71, 1114-1123.  
394

395 Blum, M.O., Bahnson, R.R., Lee, C., Deschler, J.W., Grayhack, J.T., 1985.  
396 Estimate of canine prostatic size by in vivo ultrasound and volumetric  
397 measurements. J. Urol. 133, 1082-1086.

398  
399 Cardilli, D.J., Toniollo, G.H., Pastore, A.A., Canola, J.C., Mercadante, M.E.Z.,  
400 Oliveira, J.A., 2010. Padrão ultrassonográfico do parênquima, mediastino, e  
401 túnicas testiculares em bovinos jovens da raça Nelore. Ci. Anim. Bras. 11, 899-  
402 905.  
403  
404 Cartee, R.E., Rowles, T., 1983. Transabdominal sonographic evaluation of the  
405 canine prostate. Vet. Radiol. Ultrasound. 24, 156-164.  
406  
407 Cooney, J.C., Cartee, R.E., Gray, B.W., Rumph, P.F., 1992. Ultrasonography of  
408 the canine prostate with histologic correlations. Theriogenology. 38, 877-895.  
409  
410 England, G.C.W., Allen, W.E. and Middleton, D.J., 1990. An investigation into the  
411 origin of the first fraction of the canine ejaculate. Res. Vet. Sci. 49, 66-70.  
412  
413 England, G.C.W., 1991. Relationship between ultrasonographic appearance,  
414 testicular size, spermatozoal output and testicular lesions in the dog. Journal of  
415 Small Animal Practice 32, 306-311.  
416  
417 England, G.C.W., 1995. Ultrasonographic diagnosis of non-palpable Sertoli cell  
418 tumours in infertile dogs. J. Small Anim. Pract. 36, 476-480.  
419  
420 England, G.C.W., 1999a. Semen quality in dogs and the influence of a short  
421 interval second ejaculation. Theriogenology. 52, 981-986.  
422

423 England, G.C.W., 1999b. Effect of clavulanic acid-potentiated amoxicillin on  
424 semen quality in dogs. *Theriogenology*. 52, 1117-1122.  
425

426 Eilts, B.E., Williams, D.B., Moser, E.B., 1993. Ultrasonic measurement of canine  
427 testes. *Theriogenology*. 40, 819-828.  
428

429 Feeney, D.A., Johnston, G.R., Klausner, J.S., Perman, V., Leninger, J.R.,  
430 Tomlinson M.J., 1987. Canine prostatic disease- comparison of ultrasonographic  
431 appearance with morphologic and microbiologic findings: 30 cases (1981-1985).  
432 *J. Am. Vet. Med. Assoc.* 190, 1027-1034.  
433

434 Freitas, L.A., Pinto, J.N., Silva, H.V.R., Uchoa, D.C., Mota Filho, A.C., Silva,  
435 L.D.M., 2013. Doppler e ecobiometria prostática e testicular em cães da raça  
436 Boxer. *Acta. Sci. Vet.* 41, 1121 – 1129.  
437

438 Freitas, L.A., Pinto, J.N., Silva, H.V.R., Silva, L.D.M., 2015. Two dimensional and  
439 Doppler sonographic prostatic appearance of sexually intact French Bulldogs.  
440 *Theriogenology*. 83, 1140-1146.  
441

442 Gouletsou, P., Galatos, A., Leontides, L., 2008. Comparison between  
443 ultrasonographic and caliper measurements of testicular volume in the dog.  
444 *Anim. Reprod. Sci.* 108, 1-12.  
445

446 Hecht, S., 2008. Male Reproductive Tract. in: Penninck, D., Anjou, M.A. (Eds).  
447 Atlas of small animal ultrasonography. Blackwell Publishing Inc., Iowa, pp. 417-  
448 444.  
449  
450 Hershkovitz, R., Amichay, K., Stein, G.Y., Tepper, R., 2010. The echogenicity of  
451 the normal fetal kidneys during different stages of pregnancy determined  
452 objectively. Arch. Gynecol. Obstet. 284, 807-811.  
453  
454 Hsieh, M.L., Huang, S.T., Huang, H.C., Chen, Y., Hsu, Y.C., 2009. The reliability  
455 of ultrasonographic measurements for testicular volume assessment: comparison  
456 of three common formulas with true testicular volume. Asian. J. Androl. 11, 261-  
457 265.  
458  
459 Ivancic, M., Mai, W., 2008. Qualitative and quantitative comparison of renal vs.  
460 hepatic ultrasonographic intensity in healthy dogs. Vet. Radiol. Ultrasound. 49,  
461 368-373.  
462  
463 Juniewicz, P.E., Lemp, B.M., Batzold, F.H., Reel, J.R., 1989. Transrectal  
464 ultrasonography as a method to monitor canine prostatic size in situ:  
465 measurements following endocrine manipulation and ejaculation. The Prostate.  
466 14, 265-277.  
467  
468 Kay, N.D., Ling, G.V., Nyland, T.G., Kennedy, P.C., Zinkl, J.G., 1989. Cytological  
469 diagnosis of canine prostatic disease using a urethral brush technique. J Am  
470 Anim Hosp Assoc 25, 517-526.

471

472 Kastelic, J.P., Brito, L.F., 2012. Ultrasonography for monitoring reproductive  
473 function in the bull. *Reprod. Domest. Anim. Suppl.* 3, 45-51.

474

475 Keenan, L.R.J., 1998. The infertile male. in: Simpson, G.M., England, G.C.W.,  
476 Harvey, M. (Eds). *Manual of Small Animal Reproduction and Neonatology*, 23 rd  
477 edn. Gloucester, British Small Animal Veterinary Association, London, p. 83-93.

478

479 Lowseth, L.A., Gerlach, R.F., Gillett, N.A., Muggenburg, B.A., 1990. Age-related  
480 Changes in the Prostate and Testes of the Beagle Dog. *Vet. Pathol.* 27, 347-353.

481

482 Mantziaras, G., Alonge, S., Luvoni, G.C., 2014. Ultrasonographic study of age-  
483 related changes on the size of prostate and testicles in healthy German  
484 Shepherd Dogs. *Proceedings of the 17th EVSSAR Congress, Wroclaw, Poland*  
485 pp.150.

486

487 Memon, M.A., 2007. Common causes of male dog infertility. *Theriogenology.* 68,  
488 322-328.

489

490 Nautrup, C.P., Tobias, R., 2001. Prostate gland. in: Nautrup, C.P., Tobias, R.  
491 (Eds). *An atlas and textbook of diagnostic ultrasonography of the dog and cat.*  
492 Manson Publisher Ltd, London, p.282-289.

493

494 Nielsen, S.W., Kennedy, P.C., 1990. Tumors of the genital systems. in: Moulton,  
495 J.E. (Ed). Tumors in Domestic Animals. University of California Press, Berkeley.  
496 pp 479-517.

497

498 Omer, R., Giffin, J., Hahnel, A., Bartlewski, P., 2012. Relationships of  
499 ultrasonographic and magnetic resonance image attributes to the  
500 histomorphology of ram testes. *Reprod. Biology.* 12, 355-361.

501

502 Paltiel, H.J., Diamond, D.A., Di Canzio, J., Zurakowski, D., Borer, J.G., Atala, A.,  
503 2002. Testicular volume: Comparison of orchidometer and US measurements in  
504 dogs. *Radiology.* 222, 114-119.

505

506 Pierson, R.A., Adams, G.P., 1995. Computer- assisted image analysis, diagnostic  
507 ultrasound and ovulation induction: strange bed fellows. *Theriogenology.* 43, 105-  
508 112.

509

510 Pugh, C.R., Konde, L.J., Park, R.D., 1990. Testicular ultrasound in the normal  
511 dog. *Vet. Radiology.* 31, 195-199.

512

513 Pugh, C.R., Konde, L.J., 1991. Sonographic evaluation of canine testicular and  
514 scrotal abnormalities - a review of 26 case-histories. *Vet. Radiology.* 32, 243-250.

515

516 Rivers, B.J., Walter, P.A., Holm, J.C., Letourneau, J.G., Finlay, D.E., Ritenour,  
517 E.R., King, V.L., O'Brien, T., Polzin, D.J., 1996. Gray-scale sonographic

518 characterization of aminoglycoside-induced nephrotoxicosis in a canine model.  
519 Invest. Radiol. 31, 639-651.  
520  
521 Ruel, Y.R, Barthez, P.Y., Mailles, A., Begon, D., 1998. Ultrasonographic  
522 evaluation of the prostate in healthy intact dogs. Vet. Radiol. Ultrasound. 39, 212-  
523 216.  
524  
525 Smith, S.C. and England, G.C.W., 2001. Effect of technical settings and semen  
526 handling upon motility characteristics of dog sperm measured using computer-  
527 aided sperm analysis. J. Reprod. Fertil. Suppl. 57, 151-159.  
528  
529 Souza, M.B., Barbosa, C.C., Pinto, J.N., Uchoa, D.C., Campello, C.C., Silva,  
530 L.D.M., 2012. Comparison of testicular volume between French Bulldog and  
531 Brazilian Terrier dogs. Proceedings of the 7th International Symposium on  
532 Canine and Feline Reproduction, Whistler, Canada.  
533  
534 Souza, M.B., Barbosa, C.C., Pereira, B.S., Monteiro, C.L.B., Pinto, J.N.,  
535 Linhares, J.C.S., Silva, L.D.M., 2014. Doppler velocimetric parameters of the  
536 testicular artery in healthy dogs. Res. Vet. Sci. 96, 533-536.  
537  
538 Vencato, J., Romagnoli, S., Stelletta, C., 2014. Trans-scrotal ultrasonography  
539 and testicular fine-needle aspiration cytology in the evaluation of ram sperm  
540 production. Small. Ruminant. Res. 120, 112-115.  
541

542 Villaverde, A.I.S.B., Fioratti, E.G., Ramos, R.S., Neves, R.C.F., Ferreira, J.C.P.,  
543 Cardoso, G.S., Padilha, P.M., Lopes, M.D., 2014. Blood and seminal plasma  
544 concentration of selenium, zinc and testosterone and their relationship to sperm  
545 quality and testicular biometry in domestic cats. Anim. Reprod. Sci. 150, 50-55.  
546 <http://dx.doi.org/10.1016/j.anireprosci.2014.08.004>  
547  
548 Wheaton, L.G., de Klerk, D.P., Strandberg, J.D., Coffey, D.S., 1979. Relationship  
549 of seminal volume to size and disease of the prostate in the beagle. Am. J. Vet.  
550 Res. 40, 1325-1328.  
551  
552 Woodall, P.F., Johnstone, I.P. 1988. Dimensions and allometry of testes,  
553 epididymes and spermatozoa in the domestic dog (*Canis familiaris*). J. Reprod.  
554 Fertil. 82, 603-609.  
555  
556  
557



558 **Legends to Figures**

559

560 Figure 1.

561 Software image representing the objective measurement of pixel intensity. Black  
562 rectangle and white rectangle are the near and far field echogenic reference  
563 points whilst the coloured squares represent the nine sampling regions over the  
564 parenchyma.

565

566 Figure 2.

567 Relationship between mean testicular echogenicity and percentage  
568 morphologically normal living sperm (NLS) for 43 dogs. Solid line shows  
569 regression analysis with 95% confidence limits.

570

571 Figure 3.

572 Relationship between mean testicular heterogeneity and total sperm output  
573 (TSO) for 43 dogs. Solid line shows regression analysis with 95% confidence  
574 limits.

575

576 Figure 4.

577 Relationship between body weight and (a) total testicular volume, (b) total  
578 epididymal tail volume and (c) total prostatic volume for 43 dogs. Solid line shows  
579 regression analysis with 95% confidence limits.