



The University of
Nottingham

UNITED KINGDOM · CHINA · MALAYSIA

Banerjee, Anirban and Hendrick, Paul and Bhattacharjee, P. and Blake, Holly (2016) A systematic review of outcome measures utilised to measure the change in self-management in clinical trials in non-cancer chronic pain. In: UK Society for Behavioural Medicine 12th Annual Scientific Meeting, 1-2 Dec. 2016, Cardiff, UK.

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/38996/1/2016%20Banerjee%20et%20al%20Systematic%20Review%20Pain%20UKSBM.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

Citation: Banerjee A, Hendrick P, Bhattacharjee P, Blake H. A systematic review of outcome measures utilised to measure the change in self-management in clinical trials in non-cancer chronic pain. In: Proceedings of the UK Society for Behavioural Medicine 12th Annual Scientific Meeting, Cardiff, Thursday 1st - Friday 2nd December 2016.

A systematic review of outcome measures utilised to measure the change in self-management in clinical trials in non-cancer chronic pain

Anirban Banerjee^{1,2}, Paul Hendrick¹, P Bhattacharjee², Holly Blake¹

¹University of Nottingham, Nottingham, UK, ²MSK Physiotherapy, Nottingham CityCare, Nottingham, UK

Background: There is lack of clarity and consistency in assessing the outcomes of self-management support programmes. Measures used often focus on particular constructs or behaviours such as pain, functional disability or self-efficacy.

Aim: Appraise and synthesise the evidence on outcome measures utilised to measure the effectiveness of self-management intervention in patients with non-cancer non-episodic chronic pain.

Methods: Medline, Embase, CINAHL, PsycINFO, the Cochrane Library and Google Scholar were searched in March 2015 (updated in February 2016). Included full text primary research reports of randomised or nonrandomised controlled trials, using SM outcome measure to examine effectiveness of nonsurgical interventions in adult participants. Study selection and risk of bias assessment was performed by two independent reviewers. Cochrane Handbook, the Cochrane Back Review Group and the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were followed.

Results: 2381 records were identified. 27 studies were included (24 in original 2015 search, further 4 in updated 2016 search). All were RCTs including 30-812 patients with chronic pain conditions including hip/knee osteoarthritis, rheumatoid arthritis, chronic low back pain, fibromyalgia and chronic fatigue syndrome. Most included studies had low risk of bias with the exception of blinding. 14 different validated measures were used in measuring SM and its change.

Conclusions: This systematic review presents the range of scales available for assessing change in SM in non-cancer chronic pain. Multi-construct scales are recommended for assessing SM alongside traditional pain and disability scales in chronic pain research and clinical practice.