



Fleming, Sarah A. and Blake, Holly and Gladman, John R.F. and Hart, Elizabeth and Lymbery, Mark and Dewey, Michael E. and McCloughry, Helen and Walker, Marion F. and Miller, Paul (2004) A randomised controlled trial of a care home rehabilitation service to reduce long-term institutionalisation for elderly people. *Age and Ageing*, 33 (4). pp. 384-390. ISSN 1468-2834

**Access from the University of Nottingham repository:**

<http://eprints.nottingham.ac.uk/39345/1/2004%20Fleming%20et%20al%20RCT%20Care%20Homes.pdf>

**Copyright and reuse:**

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: [http://eprints.nottingham.ac.uk/end\\_user\\_agreement.pdf](http://eprints.nottingham.ac.uk/end_user_agreement.pdf)

**A note on versions:**

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact [eprints@nottingham.ac.uk](mailto:eprints@nottingham.ac.uk)

**A randomised controlled trial of a care home rehabilitation service to reduce long-term institutionalisation for elderly people**

SARAH A. FLEMING<sup>1</sup>, HOLLY BLAKE<sup>1</sup>, JOHN R. F. GLADMAN<sup>1</sup>, ELIZABETH HART<sup>2</sup>, MARK LYMBERY<sup>3</sup>, MICHAEL E. DEWEY<sup>4</sup>, HELEN MCCLOUGHRY<sup>5</sup>, MARION WALKER<sup>1</sup>, PAUL MILLER<sup>6</sup>

<sup>1</sup>Division of Rehabilitation and Ageing, <sup>2</sup>School of Nursing, <sup>3</sup>School of Sociology and Social Policy, University of Nottingham, Nottingham NG7 2UH, UK

<sup>4</sup>Trent Institute for Health Services Research, Medical School, University Hospital, Nottingham NG7 2UH, UK

<sup>5</sup>Nottingham City Primary Care Trust, Gregory Boulevard, Nottingham NG7 5HY, UK

<sup>6</sup>Trent Institute for Health Services Research, University of Nottingham, Nottingham NG7 2UH, UK

Address correspondence to: J. R. F. Gladman, Division of Rehabilitation and Ageing, B Floor Medical School, University of Nottingham, Nottingham NG7 2UH, UK. Fax: (+44) 115 942 3618. Email: [john.gladman@nottingham.ac.uk](mailto:john.gladman@nottingham.ac.uk)

Received 27 January 2003; accepted in revised form 5 February 2004

Keywords: *health services for the aged, rehabilitation, care homes, randomised controlled trial, elderl*

## **Abstract**

**Objectives:** to evaluate the effect of a care home rehabilitation service on institutionalisation, health outcomes and service use.

**Design:** randomised controlled trial, stratified by Barthel ADL index, social service sector and whether living alone. The intervention was a rehabilitation service based in Social Services old people's homes in Nottingham, UK. The control group received usual health and social care.

**Participants:** 165 elderly and disabled hospitalised patients who wished to go home but were at high risk of institutionalisation (81 intervention, 84 control).

**Main outcome measures:** institutionalisation rates, Barthel ADL index, Nottingham Extended ADL score, General Health Questionnaire (12 item version) at 3 and 12 months, Health and Social Service resource use.

**Results:** the number of participants institutionalised was similar at 3 months (relative risk 1.04, 95% confidence intervals 0.65–1.65) and 12 months (relative risk 1.23, 95% confidence intervals 0.75–2.02). Barthel ADL Index, Nottingham Extended ADL score and General Health Questionnaire scores were similar at 3 and 12 months. The intervention group spent significantly fewer days in hospital over 3 and 12 months (mean reduction 12.1 and 27.6 days respectively,  $P < 0.01$ ), but spent a mean of 36 days in a care home rehabilitation service facility.

**Conclusions:** this service did not reduce institutionalisation, but diverted patients from the hospital to social services sector without major effects on activity levels or well-being.

## **Introduction**

Older people may move unnecessarily into long-term care because of inadequate rehabilitation after an acute illness [1]. Suitable rehabilitation is often limited because of a shortage of hospital beds [2]. In the UK, specific rehabilitation services, located in Social Services residential care homes, have been established to remedy this deficiency [3], especially since the promotion of Intermediate Care [4] as a means to deliver the National Service Framework for Older People [5]. Social Services residential care homes provide board, lodgings and personal care, without professional nursing or medical input.

Social Services departments are financially motivated to reduce the use of long-term institutional care, because they are responsible for funding it. Social Services care home rehabilitation services (CHRS) have the means to provide effective rehabilitation. Social Services care home staff are trained in the care of older people, Social Services occupational therapists can supervise and deliver rehabilitation, residents can have access to community-based rehabilitation services, and Social Services administer the provision of home care services. However, rehabilitation effectiveness is sensitive to organisation [6] and residential Intermediate Care services can inadvertently institutionalise [7]. Care home rehabilitation services could therefore, paradoxically, increase dependency and institutionalisation. Little is known about the effectiveness of such services.

A CHRS in Nottingham and Nottinghamshire was established and targeted at older people apparently destined for long-term care after an acute illness, who wished to go home, but where confidence and capability seemed to be a major factor in them doing so. Residential rehabilitation for up to 6 weeks in dedicated units within Social

Services old people's homes was provided.

We examined whether this CHRS had a major impact upon long-term care rates and improved rehabilitation out-comes (activity limitation and well-being), and the effect of the service on the use of health and social services.

## **Methods**

A pragmatic randomised controlled trial was performed. The local research ethics committee approved the study.

## **Intervention**

Over the recruitment period of the study (12 months from November 2000), the CHRS comprised 25 (rising to 40) beds in five (rising to six) units within Social Services old people's homes.

The CHRS units received input from 2.0 WTE Occupational Therapists, who assessed patients in the units and devised their treatment plans. There were 1.5 WTE Community Care Officers (Social Services employed staff with experience in the delivery of community care services for people with disability). Day to day staffing was by rehabilitation assistants: these were care assistants (workers without formal rehabilitation training) in the local authority homes in which the CHRS units were set, who had been trained by the Occupational Therapists. There were no dedicated physiotherapists: physiotherapy was provided by the existing community physiotherapy service. There was no dedicated medical cover: this was provided by the GP. There were no dedicated nurses: referrals were made to the District Nursing service.

Patients had single rooms, and had access to a dedicated rehabilitation kitchen. They were encouraged to practise the activities of daily living under the supervision of, or with the assistance of, the rehabilitation assistants. Home visits were encouraged, with the intention of increasing patients' confidence to return home. Treatment programmes were tailored to individual needs.

### **Study recruitment**

The referral criteria used and developed by the CHRS were for hospitalised patients who:

- were aged over 65
- lived in the Social Services districts served by the scheme
- wished to return to their own home
- no longer needed in-patient medical care
- were unable to return home due to activity limitation that might be improved for a period of short-term rehabilitation in a care home setting
- agreed to a period of rehabilitation in a care home setting
- met Social Services criteria for eligibility for residential home care.

The exclusion criteria were:

- Those with dementia, depression or distress that interfered with rehabilitation Those requiring two or more people to mobilise or perform personal activities of daily living, or with severe incontinence

All referrals were initially discussed with the referrer to confirm eligibility. The trial co-ordinator then obtained consent, completed baseline data collection and allocated the patient. A CHRS Occupational Therapist then assessed participants allocated to the CHRS, and arranged their transfer to the nearest unit to their home. When the study researcher was not available, all referrals were passed to the CHRS Occupational Therapist directly and were not included in this study.

## **Outcomes**

The outcomes of interest were:

- Place of residence
- Personal activities of daily living: Barthel ADL Index [8, 9]
- Instrumental activities of daily living: Nottingham Extended ADL (NEADL) scale [10]
- Psychological well-being : General Health Questionnaire (12 point version) (GHQ-12) [11]
- Hospital and CHRS bed days and re-admissions, use of day hospital and hospital out-patient departments, contacts with GPs and use of social services.

Health outcomes were recorded by post at 3 and 12 months from randomisation.

Ambiguous replies were clarified by telephone by a trained trial secretary who was independent of clinical services and masked to allocation. Participants who did not respond by post despite a telephone prompt and repeat mailing were visited at home by a researcher who was independent of clinical services and masked to allocation.

We have previously used this means of outcome assessment [12, 13] and have shown that observer bias is unlikely [14]. The use of health and social services resources were identified from routinely held service data, by a researcher who was independent of clinical services and masked to group allocation.

### **Sample size**

We set a target of 250 participants, to be recruited over 1 year. We calculated that this would be sufficient (power 80%, significance 5%, loss to follow-up 20%) to detect a reduction in the rate of placement in long-term residential and nursing home care from 60% to 30% (the latter being the level seen in pilot data).

### **Randomisation sequence generation**

A telephone randomisation service was used for allocation using computer generated balanced randomisation within strata. Stratification was by Social Services area (Nottingham City/ Nottinghamshire County), by Barthel Index at randomisation ( $\leq 14/20$ ,  $>14/20$ ) and by residential status (alone/not alone).

### **Statistical methods**

Categorical outcomes were analysed using contingency table analysis on an intention-to-treat basis. Health outcomes were analysed using multiple linear regression, adjusting for baseline characteristics and stratification variables (gender, age, baseline Barthel, location (city/county), living situation, cognitive impairment or language problem) in those with completed questionnaires only. Continuous service data were not normally distributed and were compared using non-parametric tests.

### **Results**



Figure 1 shows recruitment and patient flow through the trial. One hundred and sixty-five patients were recruited. The groups were well-matched at baseline for risk factors for institutionalisation, and the prevalence of these factors was high (Table 1).

There was no significant effect of allocation to the CHRS upon survival, rates of residential or nursing care, or the proportion living at home, at 3 or at 12 months from randomisation (Table 2). There was no significant effect of allocation to the CHRS upon the Barthel ADL Index, NEADL or GHQ-12 scores (Table 3).

Allocation to the CHRS reduced the time spent in hospital on the index admission (mean reduction 8.5 days), and non-significantly reduced re-admissions to hospital (Table 4). The mean number of hospital bed days saved rose from 12.1 by 3 months to 27.6 by 12 months. The CHRS group took significantly longer to return to their own homes after the index admission, and by 12 months had spent a mean of 19.1 more days in either a hospital or CHRS bed. There was no significant effect upon the use of other health resources. We were able to obtain limited data on Social Services resources only for those living within Nottinghamshire County Council's boundaries (51% of sample), where there was no significant impact of the intervention on the use of home care services.

## **Discussion**

This CHRS did not reduce placement rates in long-term residential and nursing homes, nor did it have a major impact upon activity levels or psychological well-being. It diverted patients from in-patient settings, but at the expense of a longer stay in a CHRS unit.

We designed our sample size to detect a moderate or large reduction in the rates of

use of institutional care but did not reach our target number, and one quarter of those allocated to the CHRS did not actually go to a CHRS unit. In fact, we observed a non-significant increase (RR 1.23, 95% CI 0.75–2.02) in the use of institutional care at 12 months in the group allocated to the CHRS. The 95% CI imply that an absolute reduction in the rates of institutional care of more than 6–8% was unlikely and on this basis we conclude that the study was adequately sized to exclude a clinically worthwhile benefit in terms of reducing institutional care rates. However, the study was inadequately sized to exclude the possibility of a clinically important increase in the rate of institutional care. More studies are needed to refute this possibility.

The means of outcome measurement used in this study are sensitive to rehabilitation intervention [13] and so the lack of effect of the CHRS upon activity levels and psycho- logical well-being is unlikely to be due to insensitivity. The CIs shown in Table 3 show that moderate to large benefits or hazards were unlikely.

The number of appropriate referrals was less than we had anticipated. Large numbers of people were referred, indicating that social workers and clinicians were aware of the service. There were many inappropriate referrals, indicating that referrers may have been uncertain of the referral criteria. Some appropriate patients may not have been referred.

The generalisability of our findings depends upon what the new service was compared with, as well as what that new service comprised. Table 4 shows that patients in the usual care group received a slightly longer initial period of hospital care than the CHRS group, but they did not receive prolonged rehabilitation or extensive rehabilitation in another facility such as a day hospital. The CHRS was not staffed like a hospital-based rehabilitation unit: the level of dedicated rehabilitation

staffing was low, and true multi- disciplinary teams did not exist. Levels of active rehabilitation were likely to be low in both groups. Similar conditions are likely to be found in many other parts of the UK and in other health care systems.

However, our study's findings relate to the CHRS operating in Nottingham during 2000, and do not necessarily apply to differently staffed or organised CHRSs elsewhere. In both institutional and community settings, there is ample evidence that organised active rehabilitation improves outcomes in people with a wide range of disabling illnesses [15]. One explanation for our findings is that the levels of rehabilitation in this CHRS were insufficient to affect health outcomes. If so, we would expect other CHRSs with similar staffing levels to have similar effects.

The CHRS units in this study were set in long-term care institutions, and the independence-promoting rehabilitation efforts of the CHRS staff may have been offset by other institutionalising influences, such as the expectation that they were in the home for care rather than rehabilitation. Our results do not necessarily apply to CHRSs that are dedicated units, rather than units within long stay institutions.

The randomised study did not assess patient satisfaction, nor could it examine further the clinical practice and organisational constraints that typified the CHRS, or identify possible means to improve outcomes. For the latter reasons, a qualitative study was also undertaken (reported elsewhere).

Although not its aim, the CHRS was successful at diverting old people from hospital, and doing so without doing harm other than delaying their return home. We saw a non-significant 10% short-term reduction in hospital readmission in the CHRS group, the consequence of which was an increasing number of hospital bed days saved over

the year of follow-up (mean number of bed days saved at 12 months = 27.6). Where there are shortages of publicly- funded hospital beds, this reduction in their use will be welcomed. However, the CHRS shifted resource use considerably from the health to the social services sector. An economic analysis is required to examine the cost-effectiveness of this arrangement to the health service, the social services and to society.

### **Key points**

- The care home service in this evaluation provided low levels of rehabilitation, which was delivered in dedicated units within existing long-term care institutions. Contrary to expectation and intention, it did not reduce unwanted institutionalisation or produce better health outcomes than ordinary hospital and Social Services aftercare.
- The care home rehabilitation service diverted patients from hospital to social services settings. Similar services may not improve the health of elderly people, but they may reduce the length of hospital stays and increase demands upon the social services.

### **Source of funding**

Trent NHS Executive.

### **Conflicts of interest**

Helen McCloughry was project manager for the CHRS during the study.

## References

1. Audit Commission. The Way to go Home. Rehabilitation and Remedial Services for Older People. London: Audit Commission, 2000. Available from URL: <http://www.audit-commission.gov.uk/reports/AC-REPORT.asp?CatID=&ProdID=40CDDD97-4563-47A9-B8F7-D4BC9AA1E44E> (accessed 25.6.03)
2. Department of Health. Shaping the Future NHS. Long Term Planning for Hospitals and Related Services. Consultation Document on the Findings of the National Beds Inquiry. London: Department of Health, 2000. Available from URL; <http://www.doh.gov.uk/pub/docs/doh/nationalbeds.pdf> (accessed 25.06.03)
3. Vaughan B, Lathlean J. Intermediate Care. Models in Practice. London: King's Fund Publishing, 1999.
4. Steiner A. Intermediate care—a good thing? *Age Ageing* 2001; 30 (Suppl 3): 33–39.
5. Department of Health. National Service Framework for Older People. London: Department of Health, 2001.
6. Stroke Unit Trialists' Collaboration. Organised inpatient (stroke unit) care for stroke (Cochrane Review). In *The Cochrane Library*, Issue 1, 2003. Oxford: Update Software.
7. Ishizaki T, Kobayashi Y, Tamiya N. The role of geriatric intermediate care facilities in long-term care for the elderly in Japan. *Health Policy* 1998; 43: 141–51.
8. Mahoney F, Barthel D. Functional evaluation: The Barthel Index. *MD State Med J* 1965; 14: 61–5.

- 9.** Collin C, Wade DT, Davies S, Horne V. The Barthel ADL Index: a reliability study. *Int Disabil Stud* 1988; 10: 61–3.
- 10.** Nouri FM, Lincoln NB. An extended activities of daily living scale for stroke patients. *Clin Rehabil* 1987; 1: 301–5.
- 11.** Goldberg D. *General Health Windsor: NFER-NELSON*, 1992.  
Questionnaire (GHQ-12).
- 12.** Parker CJ, Gladman JRF, Drummond AER *et al.* A multi-centre randomised controlled trial of leisure therapy and conventional occupational therapy after stroke. *Clin Rehabil* 2001; 15: 42–52.
- 13.** Cunliffe AL, Gladman JRF, Husbands SL, Miller P, Dewey ME, Harwood RH. Sooner and healthier: a randomised controlled trial and interview study of an early discharge rehabilitation service for older people. *Age Ageing* 2004; 33: (in press).
- 14.** Parker CJ, Dewey ME on behalf of the TOTAL study group. Assessing research outcomes by postal questionnaire with telephone follow-up. *Int J Epidemiol* 2000; 29: 1065–9.
- 15.** Turner-Stokes L ed. *The effectiveness of rehabilitation: a critical review of the evidence.* *Clin Rehabil* 1999; 13 (Supplement 1).




Figure 1. Recruitment and patient flow through the trial.

**Table 1. Baseline characteristics**

Characteristic	Intervention <i>n</i> = 81	Control <i>n</i> = 84	Both groups <i>n</i> = 165
Median age (IQR)	83 (78–89)	80 (76–87)	81 (77–88)
Female	55 (68%)	58 (69%)	113 (69%)
Living alone	72 (89%)	74 (88%)	146 (89%)
Pre-admission Oxford Handicap Scale			
No symptoms	19 (24%)	12 (14%)	31 (19%)
No significant disability	9 (11%)	3 (4%)	12 (7%)
Slight disability	9 (11%)	16 (19%)	25 (15%)
Moderate disability	41 (51%)	51 (61%)	92 (56%)
Moderately severe disability	3 (4%)	2 (2%)	5 (3%)
Median AMTS (IQR, mean) at randomisation*	7 (5–8, 6.7)	7 (6–9, 7)	7 (6–8, 6.9)
Median (IQR, mean) Barthel score at randomisation	14 (11–16, 13.8)	14 (12–17, 14.4)	14 (12–16, 14.1)
Median (IQR, mean) days in hospital at randomisation	20 (12–47, 37.2)	20 (12–41, 32.4)	20 (13–44, 34.8)
Principal diagnostic condition**			
Cardio-respiratory disorder	11 (14%)	15 (18%)	26 (16%)
Gastroenterology disorder	7 (9%)	4 (5%)	11 (7%)
Infection	1 (1%)	2 (2%)	3 (2%)
Neurological disorder	12 (15%)	11 (13%)	23 (14%)
Orthopaedic disorder	15 (19%)	14 (17%)	29 (18%)
Peripheral vascular disease	4 (5%)	1 (1%)	5 (3%)
Non-specific condition	28 (36%)	36 (43%)	64 (40%)

\*Adjusted Abbreviated Mental Test Score [total score = 9] (*n* = 151, 14 scores not obtained due to language problems).















\*\*Summary of medical notes classified into these empirical categories (by JRG, geriatrician) who was blind to group allocation.



**Table 2.** Overall outcomes at 3 and 12 months

	Intervention <i>n</i> = 81	Control <i>n</i> = 84	Relative risk (95% CI)
3 months			
Dead	6 (7%)	9 (11%)	0.69 (0.26–1.85)
In institution, including hospital	25 (31%)	25 (30%)	1.04 (0.65–1.65)
Dead or in institution	31 (38%)	34 (41%)	1.11 (0.84–1.46)
12 months			
Dead	22 (27%)	23 (27%)	0.99 (0.47–2.07)
In institution, including hospital	25 (31%)	21 (25%)	1.23 (0.75–2.02)
Dead or in institution	47 (58%)	44 (52%)	1.11 (0.84–1.46)

**Table 3.** Health outcomes at 3 and 12 months

Scale (range: worst to best score)	Mean difference [B] at 3 months (95% CI)	Difference: % of scale range at 3 months		Mean difference [B] at 12 months (95% CI)	Difference: % of scale range at 12 months	
		Intervention Worse ←	Intervention Better →		Intervention Worse ←	Intervention Better →
Barthel (0–20)	1.1 (–0.2 to 2.4)			0.2 (–1.5 to 1.9)		
Extended ADL total (0–66)	1.3 (–3.3 to 6.0)			0.8 (–5.4 to 7.1)		
Extended ADL sub-sections:						
Mobility (0–18)	0.2 (–1.1 to 1.6)			0.3 (–1.5 to 2.1)		
Kitchen (0–15)	0.8 (–0.8 to 2.4)			0.2 (–1.6 to 2.1)		
Domestic (0–15)	–0.2 (–1.5 to 1.2)			–0.6 (–2.2 to 1.0)		
Leisure (0–18)	0.4 (–0.8 to 1.7)			–0.1 (–1.5 to 1.7)		
GHQ-12 (36–0)	–0.2 (–2.6 to 2.2)			–1.1 (–3.8 to 1.7)		

Factors entered into all regression analyses included: Age at entry to trial, gender, stratifying variables [location at baseline (City or County), living arrangements at baseline (Alone/Not Alone), Barthel Index at baseline (cut-off ≤ 14)], cognitive deficit or language impairment at baseline (yes/no), randomisation group (Intervention/Control).

**Table 4.** Use of resources

	Intervention <i>n</i> = 81	Control <i>n</i> = 84	Comparison
<b>Hospital and CHRS use</b>			
Median (IQR, mean) days in hospital from randomisation to discharge	8 (7–15, 16.3)	18 (8–34, 24.8)	Median difference –7 (95% CI –11 to –2)
Median (IQR, mean) days in hospital from randomisation to 3 months	13 (7–25, 20.4)	26.5 (13–49, 32.5)	Median difference –10 (95% CI –17 to –5)
Median (mean) hospital bed days used from randomisation to 12 months	16 (8–35, 20.4)	34.5 (18–60, 48.0)	Median difference –12 (95% CI –20 to –5)
Number of patients re-admitted to hospital from randomisation to 3 months	22 (28%)	32 (38%)	RR 0.71 (95% CI 0.46–1.12)
Number of patients re-admitted to hospital from randomisation to 12 months	41 (51%)	46 (55%)	RR = 0.92 (95% CI 0.69–1.24)
Median (IQR, mean) days in CHRS facility from randomisation to 3 months	36 (0–54, 34.7)	0 (0–0, 0.0)	–
Median (IQR, mean) days in CHRS facility from randomisation to 12 months	38 (0–54, 35.0)	0 (0–0, 0.4)	–
Median (IQR, mean) days either in hospital or in CHRS facility from randomisation to 3 months	50 (31–78, 55.2)	26.5 (13–50, 32.9)	Median difference 22 (95% CI 13–31)
Median (IQR, mean) days either in hospital or in CHRS facility from randomisation to 12 months	60 (34–87, 67.6)	34.5 (18–63, 48.5)	Median difference 19 (95% CI 8–30)
<b>Community resource use</b>			
Median (IQR, mean) hospital out-patient visits from randomisation to 12 months	2 (1–6, 3.8)	4 (1–6, 4.6)	Median difference 0 (95% CI –1 to 0)
Number attending geriatric day hospital over 12 months	11 (14%)	13 (16%)	RR = 0.88 (95% CI 0.42–1.84)
Median (IQR) (mean (SD)) GP visits from randomisation to 3 months	1 (0–3, 1.9)	2 (0–3, 2.1)	Median difference 0 (95% CI –1 to 1)
Median (IQR) (mean (SD)) GP visits from randomisation to 12 months	3 (1–6, 4.0)	4 (0–6, 4.3)	Median difference –1 (95% CI –2 to 0)
Number in receipt of social services home care services from randomisation to 12 months	39 (93%) ( <i>n</i> = 42)	35 (83%) ( <i>n</i> = 42)	RR = 1.11 (0.95–1.31)

RR = relative risk, CI = confidence intervals.