



The University of
Nottingham

UNITED KINGDOM • CHINA • MALAYSIA

Rampioni, Giordano and Falcone, Marilena and Heeb, Stephan and Frangipani, Emanuela and Fletcher, Matthew P. and Dubern, Jean-Frédéric and Visca, Paolo and Leoni, Levi and Cámara, Miguel and Williams, Paul (2016) Unravelling the genome-wide contributions of specific 2-alkyl-4-quinolones and PqsE to quorum sensing in *Pseudomonas aeruginosa*. PLoS ONE, 12 (11). e1006029/1-e1006029/25. ISSN 1932-6203

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/39814/1/journal.ppat.1006029.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the Creative Commons Attribution licence and may be reused according to the conditions of the licence. For more details see:
<http://creativecommons.org/licenses/by/2.5/>

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

RESEARCH ARTICLE

Unravelling the Genome-Wide Contributions of Specific 2-Alkyl-4-Quinolones and PqsE to Quorum Sensing in *Pseudomonas aeruginosa*

Giordano Rampioni^{1,2*}, Marilena Falcone^{1a}, Stephan Heeb², Emanuela Frangipani^{1,2}, Matthew P. Fletcher², Jean-Frédéric Dubern², Paolo Visca¹, Livia Leoni¹, Miguel Cámara², Paul Williams²

1 Department of Science, University Roma Tre, Rome, Italy, **2** School of Life Sciences, Centre for Biomolecular Sciences, University of Nottingham, Nottingham, United Kingdom

^a Current address: Department of Biosciences, University of Milan, Milan, Italy

* giordano.rampioni@uniroma3.it



CrossMark

click for updates

OPEN ACCESS

Citation: Rampioni G, Falcone M, Heeb S, Frangipani E, Fletcher MP, Dubern J-F, et al. (2016) Unravelling the Genome-Wide Contributions of Specific 2-Alkyl-4-Quinolones and PqsE to Quorum Sensing in *Pseudomonas aeruginosa*. PLoS Pathog 12(11): e1006029. doi:10.1371/journal.ppat.1006029

Editor: Everett Pesci, East Carolina University School of Medicine, UNITED STATES

Received: June 9, 2016

Accepted: October 28, 2016

Published: November 16, 2016

Copyright: © 2016 Rampioni et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by: Biotechnology and Biological Sciences Research Council, UK (BB/F014392/1 to PW and MC, <https://www.bbsrc.ac.uk>); EU-FP7 collaborative Action (grant NABATIVI, Ref 223670 to MC and PW, https://ec.europa.eu/research/fp7/index_en.cfm); Italian Ministry for University and Research (RBF10LHD1 to GR, <http://www.istruzione.it>);

Abstract

The *pqs* quorum sensing (QS) system is crucial for *Pseudomonas aeruginosa* virulence both *in vitro* and in animal models of infection and is considered an ideal target for the development of anti-virulence agents. However, the precise role played by each individual component of this complex QS circuit in the control of virulence remains to be elucidated. Key components of the *pqs* QS system are 2-heptyl-4-hydroxyquinoline (HHQ), 2-heptyl-3-hydroxy-4-quinolone (PQS), 2-heptyl-4-hydroxyquinoline *N*-oxide (HQNO), the transcriptional regulator PqsR and the PQS-effector element PqsE. To define the individual contribution of each of these components to QS-mediated regulation, transcriptomic analyses were performed and validated on engineered *P. aeruginosa* strains in which the biosynthesis of 2-alkyl-4-quinolones (AQs) and expression of *pqsE* and *pqsR* have been uncoupled, facilitating the identification of the genes controlled by individual *pqs* system components. The results obtained demonstrate that *i*) the PQS biosynthetic precursor HHQ triggers a PqsR-dependent positive feedback loop that leads to the increased expression of only the *pqsABCDE* operon, *ii*) PqsE is involved in the regulation of diverse genes coding for key virulence determinants and biofilm development, *iii*) PQS promotes AQ biosynthesis, the expression of genes involved in the iron-starvation response and virulence factor production *via* PqsR-dependent and PqsR-independent pathways, and *iv*) HQNO does not influence transcription and hence does not function as a QS signal molecule. Overall this work has facilitated identification of the specific regulons controlled by individual *pqs* system components and uncovered the ability of PQS to contribute to gene regulation independent of both its ability to activate PqsR and to induce the iron-starvation response.

Italian Cystic Fibrosis Research Foundation (FFC 10/2013 to LL, <http://www.fibrosicisticaicerca.it>); Regione Lazio (LR 13/2008 - FILAS-RU-2014-1009 to PV, http://www.regione.lazio.it/ri_main/); Swiss National Science Foundation (PBLAP3-129398/1 to EF, <http://www.snf.ch/en/Pages/default.aspx>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Author Summary

Many bacterial pathogens control virulence gene expression and the development of antibiotic-resistant biofilms *via* intercellular communication through ‘quorum sensing’ (QS). QS systems depend on the synthesis, secretion and perception of diffusible signalling molecules that enable bacteria to synchronize their behaviour at the population level and are considered ideal targets for the development of anti-virulence drugs. *Pseudomonas aeruginosa* employs several overlapping QS circuits including the *pqs* system to control the expression of virulence determinants. The *pqs* QS system relies on multiple 2-alkyl-4-quinolones (AQs), including the *Pseudomonas* Quinolone Signal (PQS), as signal molecules. However, the individual contributions of key AQs and the effector proteins PqsR and PqsE within the auto-regulated *pqs* system have not been elucidated because of their inter-dependence. By constructing *P. aeruginosa* strains with multiple mutations in the *pqs* system and determining their transcriptomes in the presence or absence of PqsR, PqsE or exogenously supplied AQs, we define the distinct regulons involved and characterize a novel PQS signalling pathway independent of PqsR and the iron-starvation response.

Introduction

Pseudomonas aeruginosa is a multi-antibiotic resistant pathogen commonly responsible for hospital-acquired infections and is the main cause of morbidity and mortality in cystic fibrosis [1]. The pathogenicity of *P. aeruginosa* is multifactorial and host specific relying on the coordinated production of multiple virulence factors and the formation of antibiotic tolerant biofilms [2,3]. These are controlled by a quorum sensing (QS) intercellular communication network that integrates information on population structure/dynamics and the metabolic status of the cell with environmental cues [4–7]. Since *P. aeruginosa* QS mutants display attenuated pathogenicity, QS is a promising target for anti-virulence agents [8].

In *P. aeruginosa* QS involves three major inter-linked QS signalling pathways, namely the *las* and *rhl* systems that employ *N*-acylhomoserine lactones and the *pqs* QS system that uses 2-alkyl-4-quinolones (AQs) as QS signal molecules [5]. Data from expression studies and virulence factor profiling obtained by comparing wild type with different *pqs* mutants have revealed the extent of the *pqs* regulon and its relationship with the *las* and *rhl* regulons. For example, AQs are required for full transcription of genes coding for exoenzymes, exotoxins, lectins, secondary metabolites (*e.g.*, pyocyanin, hydrogen cyanide, rhamnolipids, pyochelin and pyoverdine) and biofilm development (reviewed in [9]). *P. aeruginosa* mutants defective in AQ biosynthesis or sensing are severely attenuated in plant and animal infection models [3,10,11]. Furthermore, AQs are detectable in sputum, blood and urine of individuals with cystic fibrosis and their presence correlates with clinical status [12].

The *pqs* system incorporates at least four transcriptional units, with *pqsABCDE* (PA0996-PA1000) and *pqsR* (PA1003) clustering at the same genetic locus, while *pqsH* (PA2587) and *pqsL* (PA4190) are distally located [13]. Our understanding of the molecular mechanisms governing *pqs*-dependent QS is however limited, largely because of the inter-dependent, auto-regulatory, multi-component nature of the system (Fig 1) [9].

P. aeruginosa produces >50 different AQs [14] of which 2-heptyl-3-hydroxy-4-quinolone (also known as the *Pseudomonas* Quinolone Signal, PQS) and its immediate precursor 2-heptyl-4-hydroxyquinoline (HHQ) are most closely associated with QS signalling. Most of the genes required for AQ biosynthesis are located in the *pqsABCDE* operon (Fig 1). PqsA converts anthranilic acid to anthraniloyl-CoA that is condensed with malonyl-CoA to form




Fig 1. The AQ biosynthetic pathway and *pqs* genes. Schematic representation of the AQ biosynthetic pathway and the *pqs* and *phn* genes in *P. aeruginosa* PAO1 and the isogenic Δ4AQ and Δ5AQ mutants. Main elements of the *pqs* QS system (HHQ, PQS, HQNO, PqsE, and PqsR) are in bold face. The PA number is indicated below the genes according to the *Pseudomonas* Genome Database [13]. Solid grey arrows represent biosynthesis; dashed grey arrows represent information flow; solid black arrow indicates activation (+); black T-line indicates negative regulation (-).

doi:10.1371/journal.ppat.1006029.g001

2-aminobenzoylacetyl-CoA (2-ABA-CoA) in a reaction catalysed by PqsD [15,16]. The thioesterase activity of PqsE converts 2-ABA-CoA into 2-aminobenzoylacetate (2-ABA) [17]; HHQ is formed through the condensation of octanoyl-coenzyme A and 2-ABA via the PqsBC heterodimer [18,19].

Additional enzymes are required for the biosynthesis of PQS and 2-heptyl-4-hydroxyquinoline *N*-oxide (HQNO) (Fig 1). Under aerobic conditions, HHQ is oxidized to PQS via the action of the monooxygenase PqsH [20]. A second monooxygenase, PqsL, is required together with the *pqsABCD* gene products for the synthesis of HQNO and related *N*-oxides [21].

The *pqs* system is subject to positive autoregulation, since the LysR-type transcriptional regulator PqsR (Mvfr), binds to the promoter region of *pqsABCDE* (*PpqsA*) and triggers

transcription once activated by HHQ or PQS (Fig 1) [22–24]. Therefore, by analogy with other QS systems, HHQ and PQS act as autoinducers by generating a positive feedback loop that accelerates their biosynthesis. Although both HHQ and PQS can function as QS signal molecules, it is not clear whether this regulatory effect is only exerted *via* PqsR, or also *via* PqsR-independent pathways. Although both HHQ and PQS activate PqsR, PQS has additional properties. For example, PQS promotes the formation of membrane vesicles (MVs) in which PQS is both bioactive and bioavailable [25], although it is not essential for MV formation [26]. The 3-hydroxy substituent also confers on PQS the ability to chelate ferric iron (Fe³⁺) [27]. Consequently, exogenous PQS triggers an iron-starvation response in *P. aeruginosa*, promoting the production of the siderophores, pyoverdine and pyochelin [27,28]. However, PQS cannot be considered as a siderophore *sensu stricto*, since it does not stimulate growth of a siderophore-defective *P. aeruginosa* mutant in iron-deficient growth conditions [27]. PQS appears instead to act as an iron trap associated with the outer membrane. In this context, the iron-chelating property of PQS may confer a survival advantage to *P. aeruginosa* in mixed bacterial populations by limiting the availability of iron to co-inhabitant species [29].

HQNO also contributes to the environmental competitiveness of *P. aeruginosa*, since it is a potent inhibitor of the cytochrome *bc₁* complex [30]. At present, the role played by HQNO in *P. aeruginosa* physiology and the mechanism by which HQNO self-poisoning is avoided, have not been determined.

Mutations in the *pqsA*, *pqsB*, *pqsC* or *pqsD* biosynthetic genes or in the regulatory gene *pqsR*, all abolish AQ production, while *P. aeruginosa* *pqsH* and *pqsL* mutants accumulate either HHQ and HQNO or HHQ and PQS, respectively [9,14,31]. Notably, while PqsE converts 2-ABA-CoA to 2-ABA, a mutation in *pqsE* does not affect AQ biosynthesis [11]. This is probably because the PqsE thioesterase functionality can be provided by alternative thioesterases [17]. PqsE over-expression however completely abrogates PqsA activity (Fig 1), and consequently AQ biosynthesis [11].

Although PqsE is dispensable for AQ biosynthesis, it is required for production of key virulence factors, such as pyocyanin, elastase, rhamnolipids, hydrogen cyanide, LecA lectin, and for biofilm maturation [11]. The activity of PqsE is also dependent on the *N*-butanoyl-homoserine lactone (C₄-HSL) receptor RhlR, which acts downstream but in synergy with PqsE [32]. Therefore, it is likely that PqsE has, as yet unidentified, functions in addition to its thioesterase activity [17]. Transcriptomic analyses have revealed that the expression of multiple genes requires *pqsE*, and that full virulence in plant and animal infection models is strongly dependent on this enzyme [6,11]. Although the crystal structure of PqsE has been solved and key active site residues identified, the mechanism by which it controls *P. aeruginosa* virulence gene expression is not understood [17,33].

Since the HHQ- and PQS-dependent activation of *pqsABCDE* transcription results in increased levels of both Aqs and PqsE, it is possible that functional effects previously considered to be HHQ- and/or PQS-dependent are mediated *via* PqsE. Alternatively, since PqsE over-expression abrogates AQ biosynthesis, some phenotypes altered as a consequence of increased *pqsE* expression may, at least in part, be under the control of HHQ and/or PQS. PqsE controls the expression of some virulence genes independent of contribution to AQ biosynthesis. The major reductions in LecA and pyocyanin production in an AQ-negative *pqsA* mutant for example could be restored fully by expressing *pqsE* from an inducible *tac* promoter [11]. Thus, the autoregulatory effect exerted by PqsE on its own transcription plays a homeostatic role in limiting AQ accumulation, thus impeding a clear understanding of the physiological role(s) played by PqsE.

Fig 1 shows how the *pqs* system components are interlinked. HHQ and PQS both induce transcription of the *pqsABCDE* operon *via* PqsR, increasing AQ biosynthesis and *pqsE*

expression. The latter in turn, exerts a repressive role on both AQ production and its own expression. This complexity has obscured comprehension of the physiological roles played by specific Aqs and PqsE. Characterization of the regulons controlled by individual components of the *pqs* system has not yet been reported. For example, the genes controlled via the *pqs* system have been investigated by comparing the transcriptional profiles of *P. aeruginosa* PA14 wild type and its *pqsH* isogenic mutant [23] by evaluating the effect of exogenous PQS on the *P. aeruginosa* PAO1 transcriptome [28] or by comparing the wild type PAO1 with *pqsA* or *pqsE* mutants [11]. In each case, numerous genes including those involved in virulence factor production, iron homeostasis and denitrification, appeared to be PQS-controlled. However, since, in the strains used, altered PQS levels led to dysregulation of HHQ and HQNO synthesis and *pqsE* expression, it is not possible to discriminate between the role(s) played by PQS from that of the other components of the *pqs* system. Similarly, it is not possible to determine whether PqsE-controlled genes in strains overexpressing this protein are controlled by PqsE itself or by the lack of Aqs resulting from *pqsE* overexpression [11].

To circumvent these limitations, a *P. aeruginosa* PAO1 mutant unable to synthesize Aqs or convert exogenously supplied Aqs was constructed. In this strain, termed $\Delta 4AQ$, *pqsE* expression is chemically inducible and uncoupled from the activity of the *PpqsA* promoter, thus exogenous AQ provision does not alter PqsE levels. Transcriptomic analyses were performed on the $\Delta 4AQ$ strain grown in the absence or in the presence of either HHQ, PQS or HQNO, or the exogenous inducer of PqsE expression (IPTG), thus enabling identification of the specific genes controlled by each *pqs* system component. Transcriptomic analyses were also performed on strains with *pqsR*-proficient or *pqsR*-deficient ($\Delta 5AQ$) genetic backgrounds to elucidate the physiological role(s) played by the transcriptional regulator PqsR.

Results and Discussion

Characterization of HHQ, PQS, HQNO and PqsE regulons

To identify the regulons controlled individually by HHQ, PQS, HQNO and PqsE, a quadruple mutant of *P. aeruginosa* PAO1, named $\Delta 4AQ$, was constructed. As depicted in Fig 1, this carries in frame deletions of *pqsA*, *pqsH* and *pqsL* genes, and incorporates an isopropyl β -D-L-thiogalactopyranoside (IPTG)-inducible *pqsE* gene. Preliminary experiments were performed to validate the $\Delta 4AQ$ strain. *P. aeruginosa* PAO1 wild type was grown in LB, while the isogenic $\Delta 4AQ$ mutant was grown in LB or in LB supplemented with either HHQ, PQS, or HQNO (40 μ M), or with IPTG (1 mM). All strains were grown to late exponential phase where the *pqs* system is maximally expressed [11]. Cell-free spent media and bacterial cells were respectively collected for determination of AQ levels by LC-MS/MS, and for quantification of *pqsE* mRNA levels by Real Time PCR. HHQ, PQS and HQNO were only recovered from the $\Delta 4AQ$ cultures if exogenously added, and were not converted into other Aqs (S1A Fig). Moreover, the *P. aeruginosa* $\Delta 4AQ$ strain grown in the absence of IPTG showed only basal levels of *pqsE* RNA ($\Delta 4AQ$ to wild type ratio ~ 0.2), irrespective of the presence or absence of Aqs while IPTG addition increased *pqsE* RNA levels by ~ 15 -fold relative to the parental strain (S1B Fig) [11]. Growth of the $\Delta 4AQ$ strain was not affected by exogenous provision of any AQ or IPTG (S1C Fig).

The transcriptional profiles of the $\Delta 4AQ$ strain grown with 40 μ M of HHQ, PQS or HQNO or with IPTG were compared by means of high-density oligonucleotide microarrays, using Affymetrix GeneChip for *P. aeruginosa* PAO1. This method was chosen to provide a reliable comparison with previously published data [6,10,11,23,28].

Following statistical validation of the dataset, only genes with a fold change > 2.5 and a *q*-value < 0.05 were considered for further analysis [34]. Table 1 lists the selected genes (see S1

Table 1. Selected genes whose transcription is controlled by HHQ, PQS and/or PqsE.

PA number ^a	Gene name ^a	HHQ ^b	PQS ^c	PqsE ^d	Product name ^a
PA0051	<i>phzH</i>			3.2 (3.6)	Potential phenazine-modifying enzyme
PA0083	<i>tssB1</i>			-5.1(-1.4)	TssB1
PA0084	<i>tssC1</i>			-3.5 (-1.3)	TssC1
PA0263	<i>hcpC</i>		-4.3		Secreted protein Hcp
PA0997* ^{f‡} ◇	<i>pqsB</i>	6.7	17.5		PqsB
PA0998* ^{f‡} ◇	<i>pqsC</i>	5.5	16.1		PqsC
PA0999* ^{f‡} ◇	<i>pqsD</i>	5.8	15.7		3-oxoacyl-[acyl-carrier-protein] synthase III
PA1000* ^f ◇	<i>pqsE</i>			22.8 (140.3)	Quinolone signal response protein
PA1001* ^f ◇	<i>phnA</i>			26.2 (ncd)	Anthranilate synthase component I
PA1002* ^f ◇	<i>phnB</i>			22.4 (ncd)	Anthranilate synthase component II
PA1245 ^{§f}	<i>aprX</i>		3.9		AprX
PA1706	<i>pcrV</i>		2.7		Type III secretion protein PcrV
PA1707	<i>pcrH</i>		3.1		Regulatory protein PcrH
PA1708	<i>popB</i>		5.6		Translocator protein PopB
PA1709	<i>popD</i>		3.0		Translocator outer membrane protein PopD precursor
PA1710	<i>exsC</i>		3.5		ExsC exoenzyme S synthesis protein C precursor
PA1711	<i>exsE</i>		3.1		ExsE
PA1712	<i>exsB</i>		2.6		Exoenzyme S synthesis protein B
PA1718	<i>pscE</i>		4.3		Type III export protein PscE
PA1901 ^f	<i>phzC2</i>			5.5 (6.3)	Phenazine biosynthesis protein PhzC
PA1902 [§]	<i>phzD2</i>			7.5 (9.8)	Phenazine biosynthesis protein PhzD
PA1903 ^f	<i>phzE2</i>			8.8 (9.6)	Phenazine biosynthesis protein PhzE
PA1904	<i>phzF2</i>			10.3 (9.9)	Probable phenazine biosynthesis protein
PA1905	<i>phzG2</i>			9.7 (9.6)	Probable pyridoxamine 5'-phosphate oxidase
PA2193*	<i>hcnA</i>			3.6 (2.1)	Hydrogen cyanide synthase HcnA
PA2194*	<i>hcnB</i>			3.1 (1.7)	Hydrogen cyanide synthase HcnB
PA2195*	<i>hcnC</i>			3.0 (1.6)	Hydrogen cyanide synthase HcnC
PA2300* ^f ◇	<i>chiC</i>			18.7 (8.2)	Chitinase
PA2426	<i>pvdS</i>		43.9		Sigma factor PvdS
PA2570* ^f ◇	<i>lecA</i>			26.3 (15.1)	LecA lectin
PA3361* [◇]	<i>lecB</i>			8.5 (10.4)	Fucose-binding lectin LecB
PA3391	<i>nosR</i>		-4.6	-4.1	Regulatory protein NosR
PA3478*	<i>rhlB</i>			3.6 (2.3)	Rhamnosyltransferase chain B
PA3479	<i>rhlA</i>			3.6 (2.3)	Rhamnosyltransferase chain A
PA3841	<i>exoS</i>		2.6		Exoenzyme S
PA3842	<i>spcS</i>		3.9		Specific <i>Pseudomonas</i> chaperone for ExoS, SpcS
PA4175	<i>prpL</i>		3.7		PrpL, protease IV
PA4205* ^f ◇	<i>mexG</i>			25.0 (47.2)	Hypothetical protein
PA4206* ^f ◇	<i>mexH</i>			16.4 (28.8)	Probable RND efflux membrane fusion protein precursor
PA4207* ^f ◇	<i>mexI</i>			18.5 (18.2)	Probable RND efflux transporter
PA4208* ^f ◇	<i>opmD</i>			11.6 (9.3)	Probable outer membrane protein precursor
PA4209* ^f ◇	<i>phzM</i>			4.1 (8.7)	Probable phenazine-specific methyltransferase
PA4210 [◇]	<i>phzA1</i>			10.2 (16.2)	Probable phenazine biosynthesis protein
PA4211* [◇]	<i>phzB1</i>			5.6 (10.1)	Probable phenazine biosynthesis protein
PA4217* [§] ◇	<i>phzS</i>			9.0 (9.3)	Flavin-containing monooxygenase
PA4227 ^{f‡}	<i>pchR</i>		11.2		Transcriptional regulator PchR
PA4468 [§]	<i>sodA</i>		89.4		Superoxide dismutase

(Continued)

Table 1. (Continued)

PA number ^a	Gene name ^a	HHQ ^b	PQS ^c	PqsE ^d	Product name ^a
PA4470 ^{§,j}	fumC1		114.7		Fumarate hydratase
PA4648	<i>cupE1</i>			3.0 (2.2)	Pilin subunit CupE1

^a PA number, gene name and product name are from the *Pseudomonas* Genome Database [13]. Genes previously reported as controlled by iron-starvation are in bold characters [40,41].

^{*}, genes whose transcription was altered in the $\Delta pqsR$ mutant with respect to the wild type strain [10]

[§], genes whose transcription was altered upon exogenous PQS provision [28]

^j, genes whose transcription was altered in the $\Delta pqsA$ mutant with respect to the wild type strain [11]

[‡], genes whose transcription was altered upon PqsE overexpression [11]

[◇], genes whose transcription was altered in the $\Delta pqsH$ mutant with respect to the wild type strain [23]. RND, Resistance-Nodulation-Cell division; MFS, major facilitator superfamily.

^b Fold change in gene expression in *P. aeruginosa* PAO1 $\Delta 4AQ$ grown in the LB supplemented with 40 μ M HHQ with respect to the same strain grown in LB.

^c Fold change in gene expression in *P. aeruginosa* PAO1 $\Delta 4AQ$ grown in the LB supplemented with 40 μ M PQS with respect to the same strain grown in LB.

^d Fold change in gene expression in *P. aeruginosa* PAO1 $\Delta 4AQ$ grown in the LB supplemented with 1 mM IPTG (to induce PqsE expression) with respect to the same strain grown in LB; in brackets is indicates the fold change in gene expression in *P. aeruginosa* PAO1 $\Delta pqsAHLE$ pUCPP. aeruginosa PAO1 $\Delta pqsAHLE$ pUCP18, both grown in LB. For the microarray analysis performed in the $\Delta pqsAHLE$ background, fold changes with a *q* value < 0.05 are indicated for selected virulence related genes, irrespective of the fold change. ncd, no change detected (*q* value > 0.05).

doi:10.1371/journal.ppat.1006029.t001

Table for complete list) satisfying this cut-off and hence significantly controlled by the Aqs and/or by PqsE. In brief, the RNA levels for 0, 3, 145, and 182 genes were significantly altered by HQNO, HHQ, PqsE and PQS respectively (Fig 2).

HQNO had no effect on the *P. aeruginosa* $\Delta 4AQ$ transcriptome indicating that this AQ does not function as a QS signal, implying an alternative role for HQNO in *P. aeruginosa* physiology. Since HQNO is a potent cytochrome *bc₁* complex inhibitor [9], it is likely that HQNO acts primarily as a secondary metabolite that increases the environmental competitiveness of *P. aeruginosa*.

Notably, only 3 genes were significantly controlled by HHQ, namely *pqsB*, *pqsC* and *pqsD* (Table 1; Fig 2). This suggests that HHQ controls only the *pqsABCDE* transcriptional unit so driving the positive feedback loop. The positive effect of HHQ on *PpqsA* activity is mediated by PqsR [22,24], such that the primary role of HHQ as a signal is to induce the PqsR-dependent expression of the *pqsABCDE* transcriptional unit, ultimately resulting in increased AQ biosynthesis and *pqsE* expression. As expected, the *pqsB*, *pqsC* and *pqsD* genes were also identified among the genes up-regulated by PQS (Table 1).

In the $\Delta 4AQ$ mutant background, PqsE emerges as a major effector of the *pqs* QS system, since the microarray analysis revealed it controls the expression of 145 genes in the $\Delta 4AQ$ strain, an AQ-negative background in which PqsE is unable to down-regulate AQ production. In particular, 72 genes were up-regulated and 73 down-regulated upon IPTG-induction of *pqsE* expression (Fig 2; S1 Table).

The 72 genes up-regulated by *pqsE* expression, included the pyocyanin biosynthetic genes (*phzA*, *phzB*, *phzC*, *phzD*, *phzE*, *phzF*, *phzG*, *phzM*, *phzS*), the *hcnABC* operon required for hydrogen cyanide biosynthesis, *rhlA* and *rhlB*, required for rhamnolipid biosynthesis, and *chiC*, coding for the extracellular chitinase ChiC. Moreover, PqsE exerted a positive effect on the transcription of genes involved in biofilm development e.g. *cupE1*, *lecA* and *lecB*, explaining the positive control of PqsE on biofilm formation [11], and on the *mexGHI-opmD* operon, coding for a Resistance-Nodulation-Cell division (RND) efflux pump involved in antibiotic




Fig 2. The AQ and PqsE regulons. Venn diagrams showing the number of genes controlled by HHQ, PQS, and PqsE in *P. aeruginosa* Δ 4AQ, and the overlap between the regulons.

doi:10.1371/journal.ppat.1006029.g002

resistance that is also essential for *pqs*-dependent QS. This is because *mexG* and *opmD* mutants are both avirulent in plant and rat infection models and fail to produce PQS, probably as a consequence of the intracellular accumulation of a toxic AQ metabolite [35]. Furthermore, pyocyanin functions as a signal in the *P. aeruginosa* QS network because it induced changes in the expression of over 50 genes (23 up-regulated and 29 down-regulated) [36]. Of these only *mexGHI-opmD*, PA2274 and PA3250 were also up-regulated via PqsE rather than PQS (Tables 1

and S1). Hence, although PqsE controls pyocyanin biosynthesis, it only regulates a sub-set of pyocyanin-dependent genes.

As shown in Fig 3A, many of PqsE up-regulated genes belong to the “Secreted Factors (toxins, enzymes, alginate)” and “Adaptation, Protection” functional classes (12.2% and 7.3%, respectively), highlighting the importance of PqsE in *P. aeruginosa* adaptive behaviour and virulence. However, most of the PqsE up-regulated genes (29.3%) are classified as “Hypothetical, unclassified, unknown”, limiting our comprehension of its physiological role.

With respect to the 73 genes down-regulated upon *pqsE* induction, they are mainly involved in energy metabolism and anaerobic respiration, including *gapA*, coding for glyceraldehyde 3-phosphate dehydrogenase, and almost all the *nir*, *nor*, *nar*, and *nos* genes (S1 Table). Indeed, the majority of the PqsE-repressed genes cluster in the “Energy metabolism” (24.3%), “Transport of small molecules” (17.1%), “Carbon compound catabolism” (11.7%) and “Biosynthesis of cofactors, prosthetic groups, and carriers” (9.0%) functional classes (Fig 3A). However, the physiological relevance of this repression is not clear, since the IPTG-mediated induction of PqsE does not affect bacterial growth, at least under aerobic conditions. It is also noticeable that two genes involved in type 6 secretion (T6SS; *tssB1* and *tssC1*) are down-regulated in response to *pqsE* induction (Table 1).

The global effect exerted by PqsE on the *P. aeruginosa* transcriptome is unlikely to be direct, since this protein does not possess a DNA-binding domain [37]. Moreover, PqsE activity is not exclusively a consequence of its thioesterase activity since *pqsE* expression is sufficient to restore pyocyanin in an AQ-deficient (*pqsA* mutant) background [6,11]. Hence, the multifunctional activity of PqsE may conceivably be a consequence of a *pqsE* regulatory RNA acting on the expression of *pqsE*-controlled genes. To investigate this possibility we quantified pyocyanin production in *P. aeruginosa* PAO1 $\Delta pqsA \Delta pqsE$ double mutant strains carrying plasmids for IPTG-inducible expression of wild type *pqsE*, or *pqsE* mutated variants lacking the first two codons (*pqsE* $\Delta 1-6$) or with a nucleotide insertion after the ATG to alter the protein frame (*pqsE*NoFrame). As shown in S2A Fig, pyocyanin production in the *P. aeruginosa* PAO1 $\Delta pqsA \Delta pqsE$ strains was restored only upon complementation with wild type *pqsE*, despite the presence of a *pqsE* transcript in the mutated variants (S2B Fig). These data suggest that the activity is not due to the *pqsE* RNA transcript but requires the PqsE protein, a finding that suggests PqsE has independent regulatory and thioesterase enzymatic functions.

When the $\Delta 4AQ$ strain was grown with IPTG, the *phnAB* operon was also up-regulated (Table 1). Knoten and co-workers reported that when *P. aeruginosa* PAO1 was grown in nutrient limiting conditions but not in LB, *pqsE* and *phnAB* were co-transcribed [38]. However, RT-PCR analysis of the PAO1 strain used in this study indicates that *pqsE* and *phnAB* are co-transcribed after growth in LB (S3 Fig), a finding consistent with the fold change increases quantified for *pqsE* (22.8), *phnA* (26.2) and *phnB* (22.4) upon addition of IPTG to the $\Delta 4AQ$ strain (Table 1). Although anthranilate is an AQ precursor [15], *pqsE*-overexpression results in the abrogation of AQ production via the strong repression of *PpqsA* promoter [11]. This repression is not apparent in our microarray analysis as a consequence of the lack of HHQ- and PQS-dependent *PpqsA* activation in the $\Delta 4AQ$ strain.

Anthranilate for AQ biosynthesis can be supplied by the anthranilate synthases TrpEG and PhnAB or via the kynurenine pathway that converts tryptophan into anthranilate [38]. The latter is the main source of anthranilate for PQS biosynthesis when tryptophan is present, and PhnAB appears to supply anthranilate only under nutrient-limiting conditions [38]. Consequently the increased transcription of *phnAB* following the IPTG-dependent induction of *pqsE* in the $\Delta 4AQ$ strain may increase intracellular anthranilate levels and so impact on gene expression independent of PqsE. To explore this possibility, we first quantified intracellular anthranilate levels in the $\Delta 4AQ$ strain grown in LB or in LB supplemented with 1 mM IPTG. S4A Fig




Fig 3. Functional classes of PqsE and PQS controlled genes. Histograms representing the distribution of (A) PqsE-controlled and (B) PQS-controlled genes according to their functional classification. Functional classes are from the *Pseudomonas* Genome Database [13].

doi:10.1371/journal.ppat.1006029.g003

shows that the IPTG-induced increase in *phnAB* expression did not result in higher levels of intracellular anthranilate. The slightly higher concentration of anthranilate in the $\Delta 4AQ$ strain in the absence of IPTG may be a consequence of the PqsE-mediated increases in the expression of genes such as *antR* and *catB* that are involved in the degradation of anthranilate (S1 Table). To confirm these data, we also compared the transcriptional profiles of a *P. aeruginosa* PAO1 quadruple mutant strain with deletions in *pqsA*, *pqsH*, *pqsL* and *pqsE* ($\Delta pqsAHLE$) and carrying a plasmid-borne copy of *pqsE* or the empty vector (pUCP18). The data obtained for selected virulence related genes are summarized in Table 1. The plasmid-mediated expression of *pqsE* in the $\Delta pqsAHLE$ genetic background did not affect *phnAB* expression. In addition, the data obtained was broadly consistent with that obtained for the inducible *pqsE* construct with respect to the genes involved in virulence factor production, biofilm formation and antibiotic resistance (Table 1). In addition, we validated the data with respect to the pyocyanin biosynthetic genes by introducing *PphzA1::lux* and *PphzA2::lux* transcriptional fusions onto the chromosome of the $\Delta pqsAHLE$ strain, since the microarray experiments cannot discriminate between the two *phz* operons as they are almost identical at the DNA level [39]. The results obtained with the reporter fusions confirm the microarray data and reveal that PqsE is responsible for driving the expression of *phzA1* but not *phzA2* (S4B Fig).

Despite the structural similarity between HHQ and PQS and their ability to activate PqsR via the same ligand binding site [24], the microarray data revealed that, in contrast to HHQ, PQS regulates the expression of 182 genes. In particular, 103 genes were up-regulated and 79 genes were down-regulated in response to exogenous PQS (Fig 2; S1 Table). The major proportion of PQS up-regulated genes (75%) are also induced by iron-starvation [40,41]. These consist of almost all the genes involved in the biosynthesis, uptake and response to the siderophores pyoverdine and pyochelin, including the regulatory genes *pvdS* and *pchR*. Moreover, metabolic and virulence genes previously shown to be induced by iron-starvation were strongly up-regulated by PQS, including fumarate hydratase (*fumC1*), superoxide dismutase (*sodA*) and two proteases (*prpL* and *aprX*) (Table 1). These findings are consistent with the iron-chelating activity of PQS inducing an iron-starvation response [27,28]. In addition to the iron-regulated genes, PQS increased the transcription of genes involved in Type 3 secretion (T3S; *pcrV*, *pcrH*, *popB*, *popD*, *exsC*, *exsE*, *exsB*, and *pscE*), and coding for both exotoxin ExoS (*exoS*) and its chaperone SpcS (*spcS*), indicating that PQS, independent of PqsE, contributes to *P. aeruginosa* virulence gene regulation (Table 1). PQS production has been indirectly linked to the regulation of T3S effector secretion in *P. aeruginosa* at the post-transcriptional level [42]. Interestingly, the *rhl* QS system that represses both *pqsA* and *pqsR* also negatively regulates *exoS* [43]. As anticipated, *pqsB*, *pqsC* and *pqsD* genes were all up-regulated by PQS (Table 1).

The 79 PQS-repressed genes mainly cluster in the “Energy metabolism” (42.7%), “Biosynthesis of cofactors, prosthetic groups, and carriers” (10.7%), and “Carbon compound catabolism” (7.8%) functional classes (Fig 3B). The repression exerted by PQS on certain metabolic genes could be due to its interaction with membranes and consequent perturbation of associated energy generation. Almost all the genes involved in denitrification (*nir*, *nor*, *nar*, and *nos* genes) are also down-regulated by PQS (S1 Table). These data are in line with previous work demonstrating that PQS represses anaerobic growth of *P. aeruginosa* by inhibiting denitrifying enzymes [44].

A comparison of the genes regulated by PQS and PqsE revealed that they control quite distinct regulons and up-regulate different sets of virulence genes (Table 1). The PQS and PqsE regulons only share 30 genes (Fig 2). Notably, 28 genes independently down-regulated by PQS and PqsE are all involved in denitrification (*nir*, *nor*, *nar*, and *nos* genes; S1 Table), indicating that there is some redundancy in the *pqs* system.




Fig 4. Validation of the microarray data by Real Time PCR. Relative mRNA levels of the genes indicated quantified by Real Time PCR in the *P. aeruginosa* $\Delta 4AQ$ strain grown in LB supplemented with 1 mM IPTG to induce PqsE expression (light-grey bars), or with 40 μ M PQS (white bars), HHQ (dark-grey bars), or HQNO (black bars), with respect to the same strain grown in LB. The average of two independent analyses each performed on three technical replicates is shown with standard deviations.

doi:10.1371/journal.ppat.1006029.g004

The reliability of the microarray data is supported by the observation that HHQ controls only one transcriptional unit, HQNO does not affect transcription under the growth conditions employed whereas PQS regulates 182 genes. Differential expression of selected genes by PQS or PqsE in the microarray experiment was validated by Real Time PCR analysis. A comparison between Table 1 and Fig 4 shows that the results obtained match the microarray data, since the mRNA levels of the *lecA*, *mexG* and *rhlA* genes increased upon IPTG-dependent induction of PqsE, while *nosR* decreased. Similarly, PQS increased the transcription of *sodA*, *pvdS*, *pchR*, and *aprX* but repressed *nosR* and *hcpC*. PqsE did not affect the transcript levels of PQS-controlled genes (i.e., *sodA*, *pvdS*, *pchR*, *aprX* and *hcpC*), and conversely PQS did not alter PqsE-regulated transcript levels (i.e., *lecA*, *mexG* and *rhlA*). Moreover, none of these transcripts were affected by HHQ or HQNO (Fig 4).

The potential effects of the Aqs and/or PqsE on *pqsH* or *pqsL* transcription cannot be inferred from the microarray analysis, since both genes were deleted in the $\Delta 4AQ$ strain. Therefore, the expression of chromosomal *pqsH* and *pqsL* *lux* promoter fusions was investigated in the *P. aeruginosa* $\Delta 4AQ$ strain. Neither promoter was influenced by exogenous HHQ, PQS or HQNO or by *pqsE* induction (S5 Fig). Thus, the autoregulatory activity of PQS is not directly exerted at the level of *pqsH* transcription, and HQNO has no effect in promoting its own biosynthesis. These data imply that a positive feedback loop exists in the *pqs* QS system only at the level of the *pqsABCDE-phnAB* transcriptional unit, with HHQ and PQS promoting their own biosynthesis by inducing PpqsA activity via PqsR.

Overall, our microarray experiments are consistent with previously published transcriptomic analyses highlighting the contributions of the *pqs* system to virulence factor production, ferric iron acquisition and energy metabolism [6,10,11,23,28]. However, our approach enabled us to discriminate between the physiological roles played by the distinct elements of the *pqs* QS system. For example, both PQS and PqsE were reported to affect iron-controlled genes, probably because in previous experimental settings, PqsE could control AQ biosynthesis [6,11]. However, our data demonstrate clearly that PQS but not PqsE, regulates the iron-regulated

genes. Similarly, certain PqsE-controlled virulence factors (e.g., pyocyanin, lectins, ChiC chitinase and the MexGHI-OpmD efflux pump) were reported to be PQS-controlled, probably because in previous experiments the addition of synthetic PQS or the abrogation of PQS synthesis (in *pqsR*, *pqsA* or *pqsH* mutants) led to dysregulation of *pqsE* expression [10,23,28]. Moreover, the use of the $\Delta 4AQ$ strain provides clear evidence that HHQ does not influence the *P. aeruginosa* transcriptome, and that HHQ exclusively regulates the *pqsABCDE-phnAB* transcriptional unit. Therefore, HHQ activity ultimately leads to the indirect control of specific physiological processes by increasing the expression of the effector protein PqsE and by acting as a substrate for PQS biosynthesis.

Comparison of the PQS and PqsR regulons reveals a PqsR-independent PQS regulon

Despite the structural similarity of the two AQs, HHQ controls the transcription of a single transcriptional unit (i.e., *pqsABCDE-phnAB*), while PQS regulates 182 genes. Since both AQs act as PqsR co-inducers [22,24], it is possible that the PqsR-HHQ complex only affects *PpqsA* activity, while the PqsR-PQS complex acts more globally as do other QS regulators such as LasR. However, PQS appears to influence gene expression via PqsR-dependent and PqsR-independent mechanisms, for example, by inducing an iron-starvation response [27,28].

To discriminate between PqsR-dependent and PqsR-independent PQS regulons, and to characterize the PqsR regulon itself, *pqsR* was deleted in the $\Delta 4AQ$ strain, generating the quintuple *P. aeruginosa* $\Delta 5AQ$ mutant (Fig 1). The transcriptomes of the $\Delta 4AQ$ and $\Delta 5AQ$ mutants, supplemented with PQS (40 μM), were compared. Only 4 genes were significantly down-regulated in the $\Delta 5AQ$ strain with respect to the $\Delta 4AQ$ mutant, namely *pqsR*, *pqsB*, *pqsC*, and *pqsD* ($-101.2 < \text{fold change} < -186.2$). An apparent strong down-regulation of *pqsR* was expected since this gene has been deleted from *P. aeruginosa* $\Delta 5AQ$. The down-regulation of *pqsB*, *pqsC*, and *pqsD* in *P. aeruginosa* $\Delta 5AQ$ strongly suggests that in PAO1 PqsR only triggers the transcription of the *pqsABCDE-phnAB* operon, and thus the *pqsA* promoter region is the only target for the PqsR-PQS complex.

Overall, these data imply that, apart from the *pqs* genes, the other 179 genes identified as PQS-regulated (S1 Table) are controlled by PQS via a PqsR-independent pathway(s). This regulatory activity is likely due, at least in part, to the iron-chelating activity of PQS, consistent with the finding that 77/100 genes up-regulated by PQS in a PqsR-independent manner are known to be induced by iron-starvation [40,41]. In contrast the 79 genes down-regulated by PQS have not previously been reported to be repressed in low-iron media. PQS could conceivably also control other phenotypes in both an iron- and a PqsR-independent manner through direct interactions with the outer membrane [25] or by acting as a pro- or anti-oxidant [45].

PQS activates *PpqsA* via both PqsR-dependent and PqsR-independent pathways

The transcriptome analysis performed on the $\Delta 4AQ$ strain indicates that PQS is more potent than HHQ in activating transcription of the *pqsABCDE-phnAB* operon in LB medium (Table 1). This is also consistent with a chromosomal reporter *PpqsA::lux* fusion in a *P. aeruginosa* *pqsAH* mutant, which cannot convert exogenously supplied HHQ to PQS, where EC_{50} values of $16.4 \pm 2.6 \mu M$ and $3.8 \pm 1.6 \mu M$ for HHQ and PQS respectively have been determined in LB medium [24]. However, HHQ activates the *PpqsA* promoter at a similar level to PQS when the bacteria are grown in an iron-deficient casamino acids (CAA) medium [27]. This suggests that PQS is more effective than HHQ in stimulating *PpqsA* activity because it induces an iron-starvation response. This hypothesis would be in agreement with iron-chelating

activity of PQS, with the evidence that high-iron concentrations negatively impact on *PpqsA* activity [6].

To investigate this possibility, the effect of HHQ, PQS and 2-heptyl-3-amino-4-quinolone (3-NH₂-PQS) on both *PpqsA* and *PpqsR* activation were compared in the Δ 4AQ (*pqsR*-proficient) and Δ 5AQ (*pqsR*-mutant) strains, grown in LB with or without 100 μ M FeCl₃. 3-NH₂-PQS is a potent PqsR agonist (EC₅₀ 0.4 \pm 0.1 μ M) isosteric with PQS but lacking iron-chelating activity [24]. In addition, although *pqsR* was not identified among the AQ-controlled genes in the transcriptome analysis (S1 Table) we included the *pqsR* promoter fusion experiments since it is not possible to exclude a regulatory effect below the fold change cut-off used (> 2.5) that has a significant effect on *PpqsA* activity.

Consistent with previous work, PQS was more effective than HHQ in up-regulating *PpqsA* in LB, while HHQ and 3-NH₂-PQS induced *PpqsA* activity at similar levels (Fig 5A). HHQ and 3-NH₂-PQS did not induce *PpqsA* in the *pqsR* mutant strain Δ 5AQ, while PQS exerted a positive, ~2 fold induction of *PpqsA* in this genetic background (Fig 5B). HHQ and 3-NH₂-PQS had no effect on *PpqsR* activity, while *PpqsR* was induced by PQS in both the Δ 4AQ and Δ 5AQ strains (~1.5 fold; Fig 5C and 5D). Interestingly, the PQS-dependent induction of *PpqsA* in the Δ 4AQ strain was strongly reduced when iron was added to the medium, showing the same activation level as that induced by HHQ and 3-NH₂-PQS. Iron supplementation had no effect on *PpqsA* activity when the promoter was induced with HHQ or 3-NH₂-PQS (Fig 5A). The reduced ability of PQS to induce *PpqsA* activity in the presence of 100 μ M FeCl₃ is likely due to its inability to induce *PpqsA* and *PpqsR* via the PqsR-independent pathway in the presence of high iron concentrations (Fig 5B–5D).

A plausible explanation for the above results is that the iron-chelating activity of PQS decreases the levels of available iron in LB medium, triggering an iron-starvation response with consequent activation of *PpqsA* and *PpqsR* via a PqsR-independent pathway. Indeed, the siderophore pyoverdine, which is only produced under iron limiting conditions [46], is detectable when PQS is added to the Δ 4AQ and Δ 5AQ cultures, but not when PQS is replaced with HHQ or 3-NH₂-PQS, or by excess iron (Fig 5), confirming that PQS triggers an iron-starvation response in LB. This is also in line with increased expression of the Fur-controlled iron-starvation sigma factor PvdS in the presence of PQS (Table 1).

The PqsR-independent activation exerted by PQS on *PpqsA* and *PpqsR* is not mediated by the iron-starvation response pathway

To determine whether the ability of PQS to induce the *PpqsA* and *PpqsR* promoters via a PqsR-independent pathway simply relies on its iron chelating properties, the effects of PQS and the iron chelators 2,2'-dipyridyl and deferiprone on *PpqsA* and *PpqsR* activity were compared in the *P. aeruginosa* strains Δ 4AQ and Δ 5AQ by means of transcriptional fusions. 2,2'-Dipyridyl chelates ferrous iron (Fe²⁺) [47], which is the prevalent intracellular iron species, while deferiprone chelates ferric iron (Fe³⁺), which prevails in extracellular environment, to form a 3:1 (deferiprone:Fe³⁺) complex [48], similar to the 3:1 ferric complexes formed by 2-hydroxy-3-alkyl-4-quinolones such as PQS [27]. Both 2,2'-dipyridyl and deferiprone induce iron-starvation in *P. aeruginosa* [49,50]. The results obtained show that 40 μ M PQS, 500 μ M 2,2'-dipyridyl or 160 μ M deferiprone all triggered similar levels of pyoverdine production in LB-grown cultures. However, neither 2,2'-dipyridyl nor deferiprone induce *PpqsA* or *PpqsR* activity, irrespective of the presence of PqsR (S6 Fig). These data strongly suggest that the PqsR-independent effect exerted by PQS on *PpqsA* and *PpqsR* does not depend on the ability of PQS to induce an iron-starvation response. Consistent with these findings, the activity of the *PpqsA* and *PpqsR* promoters was unchanged in the 4AQ and Δ 5AQ strains upon mutation of




Fig 5. Interplay of HHQ, PQS, PqsR and iron in controlling P_{pqSA} and P_{pqSR} activity. Maximal promoter activity quantified in the indicated strains carrying the transcriptional fusions $P_{pqSA}::lux$ (A and B) or $P_{pqSR}::lux$ (C and D). Strains were grown in LB or in LB supplemented with 40 μ M HHQ, PQS or 3-NH₂-PQS, as indicated below the graphs, in the absence (white bars) or presence (grey bars) of 100 μ M FeCl₃. Diamonds indicate the pyoverdine levels in the absence (white diamonds) or in the presence (grey

diamonds) of 100 μM FeCl_3 . Promoter activity and pyoverdine level are reported as Relative Light Units (RLU) and OD_{405} , respectively, normalized to cell density (OD_{600}). The average of three independent experiments is reported with standard deviations.

doi:10.1371/journal.ppat.1006029.g005

the *pvdS* gene that codes for the iron-starvation response sigma factor PvdS (S7A–S7D Fig); no differences in *PpqsA* and *PpqsR* activity were observed when these promoters were tested in *P. aeruginosa* PAO1 wild type and its isogenic $\Delta pvdS$ mutant (S7E Fig). Moreover, a Fur Titration Assay (FurTA) [51] revealed that the iron-response regulator Fur does not bind to the *PpqsA* or *PpqsR* promoter regions (S7F Fig).

Collectively, these data demonstrate that the ability of PQS to induce *PpqsA* and *PpqsR* via a PqsR-independent pathway does not rely on the capacity of PQS to induce an iron-starvation response via the master regulators PvdS and Fur. However, the effect of PQS on both promoters is inhibited when 100 μM FeCl_3 is added to the LB medium (Fig 5). To clarify this finding further, the ability of PQS to induce *PpqsA* activity was determined in *P. aeruginosa* $\Delta 4\text{AQ}$ grown in LB supplemented with increasing concentrations of FeCl_3 without Aqs or with either PQS or HHQ. In parallel, pyoverdine levels were determined in the culture supernatants of the $\Delta 4\text{AQ}$ strain grown in the presence of PQS to monitor the activation of the iron-starvation response. As shown in Fig 6, iron had no effect on the ability of HHQ to induce *PpqsA* activity or on the *PpqsA* basal level in the absence of Aqs. Conversely, increasing concentrations of FeCl_3 reduced the ability of PQS to promote *PpqsA* activity, and inhibited pyoverdine production, consistent with our previous data. Low FeCl_3 concentrations (from 0.4 μM to 1.6 μM) were sufficient to decrease the iron-starvation response, as indicated by reduced pyoverdine production, without affecting the PQS-dependent induction of *PpqsA*. The ability of HHQ and PQS to induce *PpqsA* was comparable when the medium iron concentration approximated to the theoretical PQS-saturating value, ranging from 12.5 to 25 μM (considering 40 μM PQS and 3:1 ratio of the PQS- Fe^{3+} complex).

Given that iron reduces the PqsR-independent expression of the *pqsA* or *pqsR* promoters in the absence of PvdS and that Fur does not bind to either promoter, our data suggest a regulatory




Fig 6. Effect of iron on the ability of PQS to stimulate *PpqsA* activity. Maximal *PpqsA* promoter activity measured in the *P. aeruginosa* $\Delta 4\text{AQ}$ strain carrying the transcriptional fusion *PpqsA*::lux, grown in LB (white bars) or in LB supplemented with 40 μM HHQ (light-grey bars) or 40 μM PQS (dark-grey bars), and FeCl_3 at the concentration indicated below the graph. White diamonds indicate the pyoverdine level in the supernatants of cultures grown in the presence of PQS with or without FeCl_3 . Promoter activity and pyoverdine are reported as Relative Light Units (RLU) and OD_{405} , respectively, normalized to cell density (OD_{600}). The average of three independent experiments is reported with standard deviation.

doi:10.1371/journal.ppat.1006029.g006

role for PQS in the absence of PqsR. This regulatory activity can however be abolished by increasing the medium iron content. However, of the 179 genes regulated by PQS *via* PqsR-independent pathways only 77 are controlled by the iron-starvation response. The remaining 102 genes (underlined in [S1 Table](#)) are regulated *via* an iron-starvation-independent and PqsR-independent PQS signalling pathway(s). Most of the repressed genes are involved in energy metabolism (49%) and include the *nir*, *nar*, *nos* genes involved in denitrification. Of the up-regulated genes, 39% are from the protein secretion/export functional class and include T3S genes such as *pcrV*, *exsC*, *exsE*, *exoS* and *spcS*. The mechanism by which these genes are regulated is not yet apparent but may be due to the anti-oxidant properties of PQS since these are likely to be inhibited by excess iron [45].

Since PQS promotes *PpqsA* activity in *P. aeruginosa* at different levels depending on the availability of iron, the expression of PqsE-controlled virulence factors in iron-poor environments is likely to be higher. In this context, it is tempting to speculate that the iron chelating ability of PQS may contribute to *P. aeruginosa* environmental fitness both by limiting the availability of iron to competing microorganisms and by increasing the expression of specific sets of genes important in challenging iron-poor environments. This process might be relevant during the colonization of the human host, when *P. aeruginosa* experiences iron starvation, and implies a new role for PQS as an extracellular iron sensor.

Conclusions

Although the central role of the *pqs* QS system in the control *P. aeruginosa* infection processes has been extensively studied, the precise role played by each individual element of this complex regulatory circuit remained to be defined. Here we have filled this knowledge gap by defining the specific genome-wide regulons for HHQ, PQS and HQNO and for the effector PqsE in the presence and absence of PqsR ([Fig 7](#)). Of 145 genes regulated *via* PqsE only 30 were co-regulated by PQS ([Fig 2](#)). Among the key genes controlled by PqsE in the absence of Aqs are those coding for the MexGHI-OpmD efflux pump and pyocyanin biosynthesis. Although biochemically PqsE functions as a thioesterase in AQ biosynthesis [17], the thioesterase-independent regulatory mechanism controlling gene expression requiring the PqsE protein remains to be elucidated. A striking feature of our transcriptome data is that signalling function of HHQ is simply to drive the expression of the *pqsABCDE-phnAB* transcriptional unit in a PqsR-dependent manner. These data highlight that unlike LuxR/AHL-based QS systems where the response regulator interacts with the promoters of multiple target genes, PqsR appears to target only one, the *pqsABCDE-phnAB* operon. Furthermore, HQNO, the *N*-oxide of HHQ, does not act as a signal molecule. In contrast to HHQ and HQNO, PQS is clearly a multi-functional molecule that operates *via* multiple PqsR-dependent and PqsR-independent pathways. In this it resembles *N*-(3-oxododecanoyl)homoserine lactone (3OC₁₂-HSL), which not only modulates the *P. aeruginosa* transcriptome *via* LasR and QscR, but also in the absence of any regulators incorporating an AHL-binding domain [52].

Materials and Methods

Bacterial strains and media

The bacterial strains used in this study are listed in [S2 Table](#). *E. coli* and *P. aeruginosa* strains were routinely grown at 37°C in Luria-Bertani (LB) broth with aeration. When required, LB was supplemented with synthetic 40 µM HHQ, PQS, or HQNO, or with 1 mM IPTG. FeCl₃, 2,2'-dipyridyl and deferiprone were used at the concentrations indicated. Aqs including 3-NH₂-PQS were synthesized as described previously [24]. Unless otherwise stated, antibiotics were added at the following concentrations: *E. coli*, 100 µg ml⁻¹ ampicillin (Ap), 10 µg ml⁻¹




Fig 7. Schematic representation of the *pqs* QS system in *P. aeruginosa*. The core of the *pqs* QS system is composed of the *pqsABCDE-phnAB* operon and the *pqsR* gene. Proteins coded by the *pqsABCDE-phnAB* operon synthesize HHQ that binds to and activates PqsR. The PqsR-HHQ complex promotes *Ppq*sA activity, thus increasing HHQ and PqsE levels. Notably, the *Ppq*sA promoter is the only target of the PqsR-HHQ complex. Apart from its contribution to HHQ biosynthesis, PqsE influences the *P. aeruginosa* transcriptome via a still uncharacterized AQ-independent pathway(s). In this way, PqsE up-regulates the expression of genes involved in virulence factor production, biofilm development, and antibiotic resistance. Conversely, PqsE down-regulates *Ppq*sA activity, AQ production and the expression of genes involved in denitrification and T6SS. The *pqsH* and *pqsL* genes are required for PQS and HQNO biosynthesis, respectively. HQNO did not affect the *P. aeruginosa* transcriptome, and probably contributes to environmental competition due to its cytochrome inhibitory activity. PQS chelates iron triggering the iron-starvation response and increasing the transcription of virulence factor genes coding for virulence factors such as pyoverdine, ExoS toxin and AprX protease. Moreover, PQS down-regulates genes involved in denitrification. Most of the regulatory effects exerted by PQS are PqsR-independent, since the PqsR-PQS (or PqsR-HHQ) complex only promotes *Ppq*sA activity. However, PQS also increases *Ppq*sA and *Ppq*sR expression via a PqsR-independent pathway(s) that is unrelated to the iron-starvation response, but is inhibited in the presence of high-iron concentrations. Dotted grey arrows indicate gene expression; solid grey arrows represent biosynthesis; solid black arrow indicates PqsR-dependent activation (+); dashed black arrows indicate PqsR-independent activation (+); black T-line indicates negative regulation (-); dashed grey arrows represent information flow.

doi:10.1371/journal.ppat.1006029.g007

tetracycline (Tc), or 30 $\mu\text{g ml}^{-1}$ chloramphenicol (Cm); *P. aeruginosa*, 200 $\mu\text{g ml}^{-1}$ tetracycline (Tc), 375 $\mu\text{g ml}^{-1}$ chloramphenicol (Cm), or 400 $\mu\text{g ml}^{-1}$ carbenicillin (Cb).

Recombinant DNA techniques

The plasmids and oligonucleotides used are listed in **S2** and **S3 Tables** respectively. Preparation of plasmid DNA, purification of DNA fragments, restrictions, ligations, and transformations in *E. coli* DH5 α or S17.1 λ pir competent cells were performed with standard procedures. DNA amplification was by Polymerase Chain Reaction (PCR) using the GoTaq Polymerase (Promega).

Construction of recombinant strains

The *P. aeruginosa* Δ 4AQ quadruple mutant strain was constructed by allelic exchange using the suicide vectors pDM4 Δ *pqsE*ind [11] and pDM4 Δ *pqsL* in the double mutant *P. aeruginosa*

PAO1 $\Delta pqsA \Delta pqsH$ [27]. The *P. aeruginosa* $\Delta 5AQ$ quintuple mutant was constructed using the suicide vector pDM4 $\Delta pqsR$ [24] in *P. aeruginosa* $\Delta 4AQ$. The *P. aeruginosa* PAO1 $\Delta pqsAHLE$ quadruple mutant strain was generated by allelic exchange using the suicide vectors pDM4 $\Delta pqsE$ [11] and pDM4 $\Delta pqsL$ in the double mutant *P. aeruginosa* PAO1 $\Delta pqsA \Delta pqsH$ [27]. pDM4 $\Delta pqsL$ was constructed by PCR amplifying the upstream and downstream fragments (~500 bp) of *pqsL* from PAO1 using the primers FW*pqsL*UP and RV*pqsL*UP, and FW*pqsL*DOWN and RV*pqsL*DOWN, respectively (S3 Table). The same procedures were used to introduce *pvdS* mutations into the *P. aeruginosa* wild type, $\Delta 4AQ$ and $\Delta 5AQ$ strains. In this case, the *E. coli* pEX $\Delta pvdS$ strain [53] was used as donor strain in the conjugation step.

For promoter activity studies, transcriptional fusions between the promoter regions of *pqsH*, *pqsL*, *pqsR*, *phzA1*, *phzA2* and the *luxCDABE* operon were constructed using the miniCTX-*lux* plasmid as previously described [27].

High-density oligonucleotide microarrays

Total RNA for the high-density oligonucleotide microarray experiments was extracted from 1 ml cultures of *P. aeruginosa* 4AQ, 5AQ or $\Delta pqsAHLE$ carrying the plasmid pUCP18 or pUCP*pqsE*, grown at 37°C with shaking at 200 rpm to an OD₆₀₀ 1.5 in LB or in LB supplemented with 40 μ M HHQ, PQS, HQNO, or 1 mM IPTG. Cells were mixed with 2 ml of RNA Protect Bacteria Reagent (Qiagen) the cells lysed and RNA was purified using RNeasy mini-columns (Qiagen), including the on-column DNase I digestion step. In addition, we treated the eluted RNA for 1 h at 37°C with TURBO DNase (0.1 units per μ g of RNA; Ambion). DNase I was removed with the RNeasy Column Purification Kit (Qiagen). RNA integrity was monitored by agarose gel electrophoresis, and the absence of contaminating chromosomal DNA was verified by PCR with primers pairs FW*pqsB*-RV*pqsB* and FW16SRT-RV16SRT (S3 Table).

Processing of the *P. aeruginosa* PAO1 Affymetrix GeneChip and statistical analysis of the dataset were performed at the Lausanne Genomic Technologies Facility, Center for Integrative Genomics, University of Lausanne, Switzerland. For each condition, two different pools of RNA were compared (biological duplicate), each containing RNAs from three independent extractions (technical triplicate). Fold changes > 2.5 with a *q*-value < 0.05 were considered as statistically significant. The *q*-value is the smallest False Discovery Rate (FDR) for which the test can be considered significant [34].

Reverse transcriptase and Real Time PCR analyses

For reverse transcriptase PCR (RT-PCR) and Real Time PCR analyses, RNA was extracted from *P. aeruginosa* PAO1 wild type, $\Delta 4AQ$ or $\Delta 5AQ$ grown to an OD₆₀₀ 1.5 in the same conditions as described above for the microarray experiments. cDNA synthesis was performed from 1 μ g of total purified RNA by using random hexamer primers and the iScript Reverse Transcription Supermix for RT-qPCR kit (BioRad). For RT-PCR, 50 ng of cDNA were PCR amplified with the GoTaq Polymerase (Promega) and primers FW*pqsERT* and RV*pqsERT* (for *pqsE*), FW*pqsE-phnA* and RV*pqsE-phnA* (for transcript spanning from *pqsE* to *phnA*), or FW*phnART* and RV*phnART* (for *phnA*) (S3 Table). After 5 min of denaturation at 95°C, the following reaction cycle was used for 30 cycles: 95°C for 30 s, 60°C for 30 s, and 72°C for 1 min. The PCR products were analysed on a 1% (w/v) agarose gel and stained with Midori Green DNA Stain (Nippon Genetis Europe GmbH).

Real-time PCRs were performed using the iTaq Universal SYBR Green Supermix (BioRad) and primers listed in S3 Table. Gene-specific primers employed in this analysis were designed using the Primer-Blast software (www.ncbi.nlm.nih.gov/tools/primer-blast) to avoid non-specific amplification of *P. aeruginosa* DNA. The reaction procedure involved incubation at 95°C

for 1 min and 40 cycles of amplification at 95°C for 10 s and 60°C for 45 s. Fluorescence was registered in the last 15 s of the 60°C step. 16S ribosomal RNA was chosen as the internal control to normalize the Real Time PCR data in each single run, and to calculate the relative fold change in gene expression by using the $2^{-\Delta\Delta C_t}$ method. The analysis was performed in duplicate on three technical replicates.

Measurements of promoter activity, pyoverdine, AQ and anthranilate concentrations

Bioluminescence was determined as a function of cell density using an automated luminometer-spectrometer (GENios Pro), as previously described [27]. Pyoverdine was quantified as OD₄₀₅ of culture supernatants appropriately diluted in 100 mM Tris-HCl (pH 8.0), and normalized for bacterial cell density (OD₆₀₀) [53]. AQs were quantified by LC-MS/MS after extracting cultures with acidified ethyl acetate [54]. Anthranilate levels were determined using quantitative LC-MS/MS following extraction of bacterial cell pellets with 80% (v/v) methanol. MS analysis was conducted under positive electrospray conditions (+ES) with the MS in MRM (multiple reaction monitoring) mode. The precursor-product ion mass transition used for the MRM detection was m/z 138.1–120.1. The relevant chromatographic peaks were compared to those of an anthranilate standard at a range of known concentrations.

For all the assays, the average data and standard deviations were calculated from at least three independent experiments.

FUR titration assay

The binding of Fur to the *PpqsA* and *PpqsR* promoter regions was investigated by transforming the miniCTX-*PpqsA::lux* [27] and miniCTX-*PpqsR::lux* plasmids into *E. coli* H1717 competent cells [49]. As positive and negative controls miniCTX-*PpchR::lux* and miniCTX-*lux* plasmids respectively were used. *PpchR::lux* was obtained by cloning a PCR fragment amplified from PAO1 with FW*PpchR* and RV*PpchR* (S3 Table). The resulting *E. coli* strains were grown for 16 h in LB broth supplemented with 10 $\mu\text{g ml}^{-1}$ Tc at 37°C, washed twice with saline, and then isolated on MacConkey agar supplemented with 10 $\mu\text{g ml}^{-1}$ Tc and 20 μM FeSO₄, as previously described [51]. Colony colour was checked after 24 h of incubation at 37°C.

Supporting Information

S1 Table. Genes whose transcription is controlled by HHQ, PQS and/or PqsE.
(PDF)

S2 Table. Bacterial strains and plasmids.
(PDF)

S3 Table. Oligonucleotides.
(PDF)

S1 Fig. Validation of the *P. aeruginosa* Δ 4AQ strain. (A) Levels of HHQ (white bars), PQS (light-grey bars) and HQNO (dark-grey bars) quantified by LC-MS/MS analysis in culture supernatants of *P. aeruginosa* Δ 4AQ grown in LB or in LB supplemented with 40 μM HHQ, PQS, or HQNO, or with 1 mM IPTG, as indicated. AQ levels were quantified in supernatants from three independent cultures grown to OD₆₀₀ 1.5. (B) Fold change in *pqsE* transcript levels measured by Real Time PCR in RNA extracted from Δ 4AQ strains grown as in (A). Data were normalized to the *pqsE* RNA level in the wild type. (C) Growth curves of *P. aeruginosa* Δ 4AQ grown in LB or in LB supplemented with 40 μM HHQ, PQS, or HQNO, or with 1 mM IPTG,

as indicated.

(PDF)

S2 Fig. The *pqsE* RNA transcript does not promote pyocyanin production. Pyocyanin production (A) and *pqsE* RNA levels measured by Real Time PCR (B) in *P. aeruginosa* $\Delta pqsA$ $\Delta pqsE$ double mutant strains carrying the pME6032 empty vector or pME6032-derivative plasmids for IPTG-inducible expression of wild type *pqsE* (pME-*pqsE*), or *pqsE* mutated variants lacking the first two codons (pME-*pqsE* $\Delta 1-6$) or with a nucleotide insertion after the ATG to alter the protein frame (pME-*pqsE*NoFrame). Culture supernatants and total RNAs are from the indicated strains grown to an OD₆₀₀ of 1.5 in LB supplemented with 1 mM IPTG. For the Real time PCR analysis, data are normalized to the *pqsE* RNA level measured in parallel in the *P. aeruginosa* PAO1 wild type.

(PDF)

S3 Fig. RT-PCR analysis showing co-transcription of *pqsE* and *phnA*. Amplification of cDNAs retro-transcribed from RNA extracted from (A) *P. aeruginosa* PAO1 wild type grown in LB and (B) *P. aeruginosa* $\Delta 4AQ$ grown in LB supplemented with 1 mM IPTG, to an OD₆₀₀ of 1.5. A 200 bp DNA region within the *pqsE* gene (*pqsE*), a 280 bp DNA region spanning from 97 bp upstream of the *pqsE* stop codon to 68 bp downstream of the *phnA* start codon (*pqsE-phnA*), and a 200 bp DNA region inside the *phnA* gene (*phnA*) were amplified from: 1, PAO1 genomic DNA (positive control); 2, cDNA; 3, the corresponding RNA (negative control). L, GeneRuler 100 bp DNA Ladder Plus (MBI Fermentas).

(PDF)

S4 Fig. PqsE does not affect anthranilate production and *PphzA2* activity, while it positively controls *PphzA1* activity. (A) Anthranilate was quantified by LC-MS/MS analysis in *P. aeruginosa* wild type and $\Delta 4AQ$ grown to an OD₆₀₀ of 1.5 in LB or in LB supplemented with 1 mM IPTG. Standard deviations are based on the mean values of three parallel cultures. (B) Maximal *PphzA1* and *PphzA2* promoter activity measured in *P. aeruginosa* PAO1 wild type and in the $\Delta pqsAHLE$ strain carrying the pUCP18 empty vector or the pUCP*pqsE* plasmid for constitutive expression of *pqsE*. Strains were grown in LB or in LB supplemented with 40 μ M HHQ or PQS. Promoter activity is reported as Relative Light Units (RLU)/OD₆₀₀.

(PDF)

S5 Fig. Aqs and PqsE do not affect *PpqsH* and *PpqsL* activity. Maximal promoter activity measured in *P. aeruginosa* $\Delta 4AQ$ strains carrying the transcriptional fusions (A) *PpqsH::lux* or (B) *PpqsL::lux*. Strains were grown in LB or in LB supplemented with 40 μ M Aqs or 1 mM IPTG, as indicated below the graphs. Promoter activity is reported as Relative Light Units (RLU)/OD₆₀₀.

(PDF)

S6 Fig. The iron-chelators 2,2'-dipyridyl and deferiprone do not increase *PpqsA* activity. Maximal promoter activity in strains carrying the transcriptional fusions *PpqsA::lux* (A and B) or *PpqsR::lux* (C and D). Strains were grown in LB or in LB supplemented with 40 μ M PQS, 500 μ M 2,2'-dipyridyl (DIP), or 160 μ M deferiprone (DEF), as indicated. Diamonds indicate the pyoverdine levels measured in parallel in culture supernatants. Promoter activity is reported as Relative Light Units (RLU)/OD₆₀₀; pyoverdine levels are reported as OD₄₀₅ normalized to cell density (OD₆₀₀).

(PDF)

S7 Fig. Impact of Fur and PvdS on the *PpqsA* and *PpqsR* promoter regions. (A-D) Maximal promoter activity in the strains carrying the transcriptional fusions *PpqsA::lux* (A and B) or

PpqSR::lux (C and D). White bars indicate the *pvdS*-proficient genetic backgrounds ($\Delta 4AQ$ and $\Delta 5AQ$); grey bars indicate the *pvdS*-mutant genetic backgrounds ($\Delta 4AQ\Delta pvdS$ and $\Delta 5AQ\Delta pvdS$). Strains were grown in LB or in LB supplemented with 40 μM HHQ or PQS, as indicated. Diamonds indicate the pyoverdine levels measured in culture supernatants in the *pvdS*-proficient (white diamonds) or *pvdS*-mutant (grey diamonds) genetic backgrounds. Promoter activity is reported as Relative Light Units (RLU)/OD₆₀₀; pyoverdine levels are reported as OD₄₀₅ normalized to cell density (OD₆₀₀). (E) Maximal *PpqSA::lux* and *PpqSR::lux* promoter activity in the wild type (white bars) and $\Delta pvdS$ (grey bars) strains grown in LB. Promoter activity is reported as Relative Light Units (RLU)/OD₆₀₀. (F) *E. coli* H1717 cells containing the plasmids indicated and grown for 24 h at 37°C on McConkey agar supplemented with 10 $\mu\text{g ml}^{-1}$ Tc and 20 μM FeSO₄. Red-staining indicates the ability to ferment lactose and hence the binding of Fur to the target promoter. miniCTX-*PpchR::lux*, positive control (red colonies); miniCTX-*lux*, negative control (white colonies). (PDF)

Acknowledgments

We thank Catherine A. Ortori (Centre for Analytical Sciences, School of Pharmacy, University of Nottingham) and Nigel Halliday for the LC-MS/MS analyses, Siri Ram Chhabra and Alex Truman (Centre for Biomolecular Sciences, University of Nottingham) for AQ synthesis, and the Lausanne Genomic Technologies Facility staff (Center for Integrative Genomics, University of Lausanne, Switzerland), in particular Keith Harshman, Alexandra Paillusson, Leonore Wigger, Sylvain Pradervand and Mélanie Dupasquier, for bioinformatics assistance with the microarray analysis.

Author Contributions

Conceived and designed the experiments: GR SH PV LL MC PW.

Performed the experiments: GR MF EF JFD MPF.

Analyzed the data: GR SH EF MPF PV LL MC PW.

Contributed reagents/materials/analysis tools: GR PV LL MC PW.

Wrote the paper: GR PV LL MC PW.

References

1. El Zowalaty ME, Al Thani AA, Webster TJ, El Zowalaty AE, Schweizer HP, Nasrallah GK, et al. *Pseudomonas aeruginosa*: arsenal of resistance mechanisms, decades of changing resistance profiles, and future antimicrobial therapies. *Future Microbiol.* 2015; 10:1683–706. doi: [10.2217/fmb.15.48](https://doi.org/10.2217/fmb.15.48) PMID: [26439366](https://pubmed.ncbi.nlm.nih.gov/26439366/)
2. Gellatly SL, Hancock RE. *Pseudomonas aeruginosa*: new insights into pathogenesis and host defences. *Pathog Dis.* 2013; 67:159–73. doi: [10.1111/2049-632X.12033](https://doi.org/10.1111/2049-632X.12033) PMID: [23620179](https://pubmed.ncbi.nlm.nih.gov/23620179/)
3. Dubern JF, Cigana C, De Simone M, Lazenby J, Juhas M, Schwager S, et al. Integrated whole-genome screening for *Pseudomonas aeruginosa* virulence genes using multiple disease models reveals that pathogenicity is host specific. *Environ Microbiol.* 2015; 17:4379–93. doi: [10.1111/1462-2920.12863](https://doi.org/10.1111/1462-2920.12863) PMID: [25845292](https://pubmed.ncbi.nlm.nih.gov/25845292/)
4. Kirisits MJ, Parsek MR. Does *Pseudomonas aeruginosa* use intercellular signalling to build biofilm communities? *Cell Microbiol.* 2006; 8:1841–9. doi: [10.1111/j.1462-5822.2006.00817.x](https://doi.org/10.1111/j.1462-5822.2006.00817.x) PMID: [17026480](https://pubmed.ncbi.nlm.nih.gov/17026480/)
5. Williams P, Cámara M. Quorum sensing and environmental adaptation in *Pseudomonas aeruginosa*: a tale of regulatory networks and multifunctional signal molecules. *Curr Opin Microbiol.* 2009; 12:182–91. doi: [10.1016/j.mib.2009.01.005](https://doi.org/10.1016/j.mib.2009.01.005) PMID: [19249239](https://pubmed.ncbi.nlm.nih.gov/19249239/)

6. Hazan R, He J, Xiao G, Dekimpe V, Apidianakis Y, Lesic B, et al. Homeostatic interplay between bacterial cell-cell signaling and iron in virulence. *PLoS Pathog.* 2010; 6(3):e1000810. doi: [10.1371/journal.ppat.1000810](https://doi.org/10.1371/journal.ppat.1000810) PMID: [20300606](https://pubmed.ncbi.nlm.nih.gov/20300606/)
7. Lee J, Zhang L. The hierarchy quorum sensing network in *Pseudomonas aeruginosa*. *Protein Cell.* 2015; 6:26–41. doi: [10.1007/s13238-014-0100-x](https://doi.org/10.1007/s13238-014-0100-x) PMID: [25249263](https://pubmed.ncbi.nlm.nih.gov/25249263/)
8. Rampioni G, Leoni L, Williams P. The art of antibacterial warfare: deception through interference with quorum sensing-mediated communication. *Bioorg Chem.* 2014; 55:60–8. doi: [10.1016/j.bioorg.2014.04.005](https://doi.org/10.1016/j.bioorg.2014.04.005) PMID: [24823895](https://pubmed.ncbi.nlm.nih.gov/24823895/)
9. Heeb S, Fletcher MP, Chhabra SR, Diggle SP, Williams P, Cámara M. Quinolones: from antibiotics to autoinducers. *FEMS Microbiol Rev.* 2010; 35:247–74.
10. Déziel E, Gopalan S, Tampakaki AP, Lépine F, Padfield KE, Saucier M, et al. The contribution of MvfR to *Pseudomonas aeruginosa* pathogenesis and quorum sensing circuitry regulation: multiple quorum sensing-regulated genes are modulated without affecting *lasRI*, *rhlRI* or the production of *N*-acyl-L-homoserine lactones. *Mol Microbiol.* 2005; 55:998–1014. doi: [10.1111/j.1365-2958.2004.04448.x](https://doi.org/10.1111/j.1365-2958.2004.04448.x) PMID: [15686549](https://pubmed.ncbi.nlm.nih.gov/15686549/)
11. Rampioni G, Pustelny C, Fletcher MP, Wright VJ, Bruce M, Rumbaugh KP, et al. Transcriptomic analysis reveals a global alkyl-quinolone-independent regulatory role for PqsE in facilitating the environmental adaptation of *Pseudomonas aeruginosa* to plant and animal hosts. *Environ Microbiol.* 2010; 12:1659–73. doi: [10.1111/j.1462-2920.2010.02214.x](https://doi.org/10.1111/j.1462-2920.2010.02214.x) PMID: [20406282](https://pubmed.ncbi.nlm.nih.gov/20406282/)
12. Barr HL, Halliday N, Cámara M, Barrett DA, Williams P, Forrester DL, et al. *Pseudomonas aeruginosa* quorum sensing molecules correlate with clinical status in cystic fibrosis. *Eur Respir J.* 2015; 46:1046–54. doi: [10.1183/09031936.00225214](https://doi.org/10.1183/09031936.00225214) PMID: [26022946](https://pubmed.ncbi.nlm.nih.gov/26022946/)
13. Winsor GL, Lam DK, Fleming L, Lo R, Whiteside MD, Yu NY, et al. *Pseudomonas* Genome Database: improved comparative analysis and population genomics capability for *Pseudomonas* genomes. *Nucleic Acids Res.* 2011; 39:596–600.
14. Déziel E, Lépine F, Milot S, He J, Mindrinos MN, Tompkins RG, et al. Analysis of *Pseudomonas aeruginosa* 4-hydroxy-2-alkylquinolines (HAQs) reveals a role for 4-hydroxy-2-heptylquinoline in cell-to-cell communication. *Proc Natl Acad Sci USA.* 2004; 101:1339–44. doi: [10.1073/pnas.0307694100](https://doi.org/10.1073/pnas.0307694100) PMID: [14739337](https://pubmed.ncbi.nlm.nih.gov/14739337/)
15. Coleman JP, Hudson LL, McKnight SL, Farrow JM 3rd, Calfee MW, Lindsey CA, et al. *Pseudomonas aeruginosa* PqsA is an anthranilate-coenzyme A ligase. *J Bacteriol.* 2008; 190:1247–55. doi: [10.1128/JB.01140-07](https://doi.org/10.1128/JB.01140-07) PMID: [18083812](https://pubmed.ncbi.nlm.nih.gov/18083812/)
16. Zhang YM, Frank MW, Zhu K, Mayasundari A, Rock CO. PqsD is responsible for the synthesis of 2,4-dihydroxyquinoline, an extracellular metabolite produced by *Pseudomonas aeruginosa*. *J Biol Chem.* 2008; 283:28788–94. doi: [10.1074/jbc.M804555200](https://doi.org/10.1074/jbc.M804555200) PMID: [18728009](https://pubmed.ncbi.nlm.nih.gov/18728009/)
17. Drees SL, Fetzner S. PqsE of *Pseudomonas aeruginosa* acts as pathway-specific thioesterase in the biosynthesis of alkylquinolone signaling molecules. *Chem Biol.* 2015; 22:611–8. doi: [10.1016/j.chembiol.2015.04.012](https://doi.org/10.1016/j.chembiol.2015.04.012) PMID: [25960261](https://pubmed.ncbi.nlm.nih.gov/25960261/)
18. Dulcey CE, Dekimpe V, Fauvelle DA, Milot S, Groleau MC, Doucet N, et al. The end of an old hypothesis: the *Pseudomonas* signalling molecules 4-hydroxy-2-alkylquinolines derive from fatty acids, not 3-ketofatty acids. *Chem Biol.* 2013; 20:1481–91. doi: [10.1016/j.chembiol.2013.09.021](https://doi.org/10.1016/j.chembiol.2013.09.021) PMID: [24239007](https://pubmed.ncbi.nlm.nih.gov/24239007/)
19. Drees SL, Li C, Prasetya F, Saleem M, Drevény I, Williams P, et al. PqsBC, a condensing enzyme in the biosynthesis of the *Pseudomonas aeruginosa* Quinolone Signal: crystal structure, inhibition, and reaction mechanism. *J Biol Chem.* 2016; 291:6610–24. doi: [10.1074/jbc.M115.708453](https://doi.org/10.1074/jbc.M115.708453) PMID: [26811339](https://pubmed.ncbi.nlm.nih.gov/26811339/)
20. Schertzer JW, Brown SA, Whiteley M. Oxygen levels rapidly modulate *Pseudomonas aeruginosa* social behaviors via substrate limitation of PqsH. *Mol Microbiol.* 2010; 77:1527–38. doi: [10.1111/j.1365-2958.2010.07303.x](https://doi.org/10.1111/j.1365-2958.2010.07303.x) PMID: [20662781](https://pubmed.ncbi.nlm.nih.gov/20662781/)
21. Lépine F, Milot S, Déziel E, He J, Rahme LG. Electrospray/mass spectrometric identification and analysis of 4-hydroxy-2-alkylquinolines (HAQs) produced by *Pseudomonas aeruginosa*. *J Am Soc Mass Spectrom.* 2004; 15:862–9. doi: [10.1016/j.jasms.2004.02.012](https://doi.org/10.1016/j.jasms.2004.02.012) PMID: [15144975](https://pubmed.ncbi.nlm.nih.gov/15144975/)
22. Wade DS, Calfee MW, Rocha ER, Ling EA, Engstrom E, Coleman JP, et al. Regulation of *Pseudomonas* quinolone signal synthesis in *Pseudomonas aeruginosa*. *J Bacteriol.* 2005; 187:4372–80. doi: [10.1128/JB.187.13.4372-4380.2005](https://doi.org/10.1128/JB.187.13.4372-4380.2005) PMID: [15968046](https://pubmed.ncbi.nlm.nih.gov/15968046/)
23. Xiao G, Déziel E, He J, Lépine F, Lesic B, Castonguay MH, et al. MvfR, a key *Pseudomonas aeruginosa* pathogenicity LTTR-class regulatory protein, has dual ligands. *Mol Microbiol.* 2006; 262:1689–99.
24. Ilangovan A, Fletcher M, Rampioni G, Pustelny C, Rumbaugh K, Heeb S, et al. Structural basis for native agonist and synthetic inhibitor recognition by the *Pseudomonas aeruginosa* quorum sensing

- regulator PqsR (MvfR). *PLoS Pathog.* 2013; 9(7):e1003508. doi: [10.1371/journal.ppat.1003508](https://doi.org/10.1371/journal.ppat.1003508) PMID: [23935486](https://pubmed.ncbi.nlm.nih.gov/23935486/)
25. Mashburn-Warren L, Howe J, Garidel P, Richter W, Steiniger F, Roessle M, et al. Interaction of quorum signals with outer membrane lipids: insights into prokaryotic membrane vesicle formation. *Mol Microbiol.* 2008; 69:491–502. doi: [10.1111/j.1365-2958.2008.06302.x](https://doi.org/10.1111/j.1365-2958.2008.06302.x) PMID: [18630345](https://pubmed.ncbi.nlm.nih.gov/18630345/)
 26. Macdonald IA, Kuehn MJ. Stress-induced outer membrane vesicle production by *Pseudomonas aeruginosa*. *J Bacteriol.* 2013; 195:2971–81. doi: [10.1128/JB.02267-12](https://doi.org/10.1128/JB.02267-12) PMID: [23625841](https://pubmed.ncbi.nlm.nih.gov/23625841/)
 27. Diggle SP, Matthijs S, Wright VJ, Fletcher MP, Chhabra SR, Lamont IL, et al. The *Pseudomonas aeruginosa* 4-quinolone signal molecules HHQ and PQS play multifunctional roles in quorum sensing and iron entrapment. *Chem Biol.* 2007; 14:87–96. doi: [10.1016/j.chembiol.2006.11.014](https://doi.org/10.1016/j.chembiol.2006.11.014) PMID: [17254955](https://pubmed.ncbi.nlm.nih.gov/17254955/)
 28. Bredenbruch F, Geffers R, Nimtz M, Buer J, Häussler S. The *Pseudomonas aeruginosa* quinolone signal (PQS) has an iron-chelating activity. *Environ Microbiol.* 2006; 8:1318–29. doi: [10.1111/j.1462-2920.2006.01025.x](https://doi.org/10.1111/j.1462-2920.2006.01025.x) PMID: [16872396](https://pubmed.ncbi.nlm.nih.gov/16872396/)
 29. Nguyen AT, Jones JW, Ruge MA, Kane MA, Oglesby-Sherrouse AG. Iron depletion enhances production of antimicrobials by *Pseudomonas aeruginosa*. *J Bacteriol.* 2015; 197:2265–75. doi: [10.1128/JB.00072-15](https://doi.org/10.1128/JB.00072-15) PMID: [25917911](https://pubmed.ncbi.nlm.nih.gov/25917911/)
 30. Voggu L, Schlag S, Biswas R, Rosenstein R, Rausch C, Götz F. Microevolution of cytochrome *bd* oxidase in *Staphylococci* and its implication in resistance to respiratory toxins released by *Pseudomonas*. *J Bacteriol.* 2006; 188:8079–86. doi: [10.1128/JB.00858-06](https://doi.org/10.1128/JB.00858-06) PMID: [17108291](https://pubmed.ncbi.nlm.nih.gov/17108291/)
 31. Gallagher LA, McKnight SL, Kuznetsova MS, Pesci EC, Manoil C. Functions required for extracellular quinolone signaling by *Pseudomonas aeruginosa*. *J Bacteriol.* 2002; 184:6472–80. doi: [10.1128/JB.184.23.6472-6480.2002](https://doi.org/10.1128/JB.184.23.6472-6480.2002) PMID: [12426334](https://pubmed.ncbi.nlm.nih.gov/12426334/)
 32. Farrow JM 3rd, Sund ZM, Ellison ML, Wade DS, Coleman JP, Pesci EC. PqsE functions independently of PqsR-*Pseudomonas* quinolone signal and enhances the *rhl* quorum-sensing system. *J Bacteriol.* 2008; 190:7043–51. doi: [10.1128/JB.00753-08](https://doi.org/10.1128/JB.00753-08) PMID: [18776012](https://pubmed.ncbi.nlm.nih.gov/18776012/)
 33. Folch B, Déziel E, Doucet N. Systematic mutational analysis of the putative hydrolase PqsE: toward a deeper molecular understanding of virulence acquisition in *Pseudomonas aeruginosa*. *PLoS One.* 2013; 8(9):e73727. doi: [10.1371/journal.pone.0073727](https://doi.org/10.1371/journal.pone.0073727) PMID: [24040042](https://pubmed.ncbi.nlm.nih.gov/24040042/)
 34. Storey JD, Tibshirani R. Statistical significance for genome-wide studies. *Proc Natl Acad Sci USA.* 2003; 100:9440–5. doi: [10.1073/pnas.1530509100](https://doi.org/10.1073/pnas.1530509100) PMID: [12883005](https://pubmed.ncbi.nlm.nih.gov/12883005/)
 35. Aendekerck S, Diggle SP, Song Z, Høiby N, Cornelis P, Williams P, et al. The MexGHI-OpmD multidrug efflux pump controls growth, antibiotic susceptibility and virulence in *Pseudomonas aeruginosa* via 4-quinolone-dependent cell-to-cell communication. *Microbiology.* 2005; 151:1113–25. doi: [10.1099/mic.0.27631-0](https://doi.org/10.1099/mic.0.27631-0) PMID: [15817779](https://pubmed.ncbi.nlm.nih.gov/15817779/)
 36. Dietrich LE, Price-Whelan A, Petersen A, Whiteley M, Newman DK. The phenazine pyocyanin is a terminal signalling factor in the quorum sensing network of *Pseudomonas aeruginosa*. *Mol Microbiol.* 2006; 61:1308–21. doi: [10.1111/j.1365-2958.2006.05306.x](https://doi.org/10.1111/j.1365-2958.2006.05306.x) PMID: [16879411](https://pubmed.ncbi.nlm.nih.gov/16879411/)
 37. Yu S, Jensen V, Seeliger J, Feldmann I, Weber S, Schleicher E, et al. Structure elucidation and preliminary assessment of hydrolase activity of PqsE, the *Pseudomonas* quinolone signal (PQS) response protein. *Biochemistry.* 2009; 48:10298–307. doi: [10.1021/bi900123j](https://doi.org/10.1021/bi900123j) PMID: [19788310](https://pubmed.ncbi.nlm.nih.gov/19788310/)
 38. Knoten CA, Wells G, Coleman JP, Pesci EC. A conserved suppressor mutation in a tryptophan auxotroph results in dysregulation of *Pseudomonas* quinolone signal synthesis. *J Bacteriol.* 2014; 196:2413–22. doi: [10.1128/JB.01635-14](https://doi.org/10.1128/JB.01635-14) PMID: [24748618](https://pubmed.ncbi.nlm.nih.gov/24748618/)
 39. Recinos DA, Sekedat MD, Hernandez A, Cohen TS, Sakhtah H, Prince AS, Price-Whelan A, Dietrich LE. Redundant phenazine operons in *Pseudomonas aeruginosa* exhibit environment-dependent expression and differential roles in pathogenicity. *Proc Natl Acad Sci USA.* 2012; 109:19420–5. doi: [10.1073/pnas.1213901109](https://doi.org/10.1073/pnas.1213901109) PMID: [23129634](https://pubmed.ncbi.nlm.nih.gov/23129634/)
 40. Ochsner UA, Wilderman PJ, Vasil AI, Vasil ML. GeneChip expression analysis of the iron starvation response in *Pseudomonas aeruginosa*: identification of novel pyoverdine biosynthesis genes. *Mol Microbiol.* 2002; 45:1277–87. PMID: [12207696](https://pubmed.ncbi.nlm.nih.gov/12207696/)
 41. Palma M, Worgall S, Quadri LE. Transcriptome analysis of the *Pseudomonas aeruginosa* response to iron. *Arch Microbiol.* 2003; 180:374–9. doi: [10.1007/s00203-003-0602-z](https://doi.org/10.1007/s00203-003-0602-z) PMID: [14513207](https://pubmed.ncbi.nlm.nih.gov/14513207/)
 42. Singh G, Wu B, Baek MS, Camargo A, Nguyen A, Slusher NA, et al. Secretion of *Pseudomonas aeruginosa* type III cytotoxins is dependent on *Pseudomonas* quinolone signal concentration. *Microb Pathog.* 2010; 49:196–203. doi: [10.1016/j.micpath.2010.05.013](https://doi.org/10.1016/j.micpath.2010.05.013) PMID: [20570614](https://pubmed.ncbi.nlm.nih.gov/20570614/)
 43. Hogardt M, Roeder M, Schreff AM, Eberl L, Heesemann J. Expression of *Pseudomonas aeruginosa* *exoS* is controlled by quorum sensing and RpoS. *Microbiology.* 2004; 150:843–51. doi: [10.1099/mic.0.26703-0](https://doi.org/10.1099/mic.0.26703-0) PMID: [15073294](https://pubmed.ncbi.nlm.nih.gov/15073294/)

44. Toyofuku M, Nomura N, Kuno E, Tashiro Y, Nakajima T, Uchiyama H. Influence of the *Pseudomonas* quinolone signal on denitrification in *Pseudomonas aeruginosa*. *J Bacteriol.* 2008; 190:7947–56. doi: [10.1128/JB.00968-08](https://doi.org/10.1128/JB.00968-08) PMID: [18931133](https://pubmed.ncbi.nlm.nih.gov/18931133/)
45. Häussler S, Becker T. The *Pseudomonas* quinolone signal (PQS) balances life and death in *Pseudomonas aeruginosa* populations. *PLoS Pathog.* 2008; 4(9):e1000166. doi: [10.1371/journal.ppat.1000166](https://doi.org/10.1371/journal.ppat.1000166) PMID: [18818733](https://pubmed.ncbi.nlm.nih.gov/18818733/)
46. Visca P. Iron regulation and siderophore signaling in *Pseudomonas aeruginosa* virulence. In: Ramos JL, editor. *Pseudomonas*, vol. 2. New York: Kluwer Academic/Plenum Publishers; 2004. p. 69–124.
47. Bergh A, Offenhardt P, George P, Haight GP. Complexes of 2,2', 2'', 2-quaterpyridine with ferrous and ferric ions. *J Chem Soc.* 1964; 1:1533–8.
48. Boelaert JR, Van Cutsem J, de Loch M, Schneider YJ, Crichton RR. Deferoxamine augments growth and pathogenicity of *Rhizopus*, while hydroxypyridinone chelators have no effect. *Kidney Int.* 1994; 45:667–71. PMID: [8196268](https://pubmed.ncbi.nlm.nih.gov/8196268/)
49. Liu Y, Yang L, Molin S. Synergistic activities of an efflux pump inhibitor and iron chelators against *Pseudomonas aeruginosa* growth and biofilm formation. *Antimicrob Agents Chemother.* 2010; 54:3960–3. doi: [10.1128/AAC.00463-10](https://doi.org/10.1128/AAC.00463-10) PMID: [20566773](https://pubmed.ncbi.nlm.nih.gov/20566773/)
50. Thompson MG, Corey BW, Si Y, Craft DW, Zurawski DV. Antibacterial activities of iron chelators against common nosocomial pathogens. *Antimicrob Agents Chemother.* 2012; 56:5419–21. doi: [10.1128/AAC.01197-12](https://doi.org/10.1128/AAC.01197-12) PMID: [22850524](https://pubmed.ncbi.nlm.nih.gov/22850524/)
51. Stojiljkovic I, Bäumlér AJ, Hantke K. Fur regulon in Gram-negative bacteria. Identification and characterization of new iron-regulated *Escherichia coli* genes by a fur titration assay. *J Mol Biol.* 1994; 236:531–45. doi: [10.1006/jmbi.1994.1163](https://doi.org/10.1006/jmbi.1994.1163) PMID: [8107138](https://pubmed.ncbi.nlm.nih.gov/8107138/)
52. Chugani S, Greenberg EP. LuxR homolog-independent gene regulation by acyl-homoserine lactones in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci USA.* 2010; 107:10673–8. doi: [10.1073/pnas.1005909107](https://doi.org/10.1073/pnas.1005909107) PMID: [20498077](https://pubmed.ncbi.nlm.nih.gov/20498077/)
53. Imperi F, Massai F, Facchini M, Frangipani E, Visaggio D, Leoni L, et al. Repurposing the antimycotic drug flucytosine for suppression of *Pseudomonas aeruginosa* pathogenicity. *Proc Natl Acad Sci USA.* 2013; 110:7458–63. doi: [10.1073/pnas.1222706110](https://doi.org/10.1073/pnas.1222706110) PMID: [23569238](https://pubmed.ncbi.nlm.nih.gov/23569238/)
54. Ortori CA, Dubern JF, Chhabra SR, Cámara M, Hardie K, Williams P, et al. Simultaneous quantitative profiling of *N*-acyl-L-homoserine lactone and 2-alkyl-4(1*H*)-quinolone families of quorum-sensing signaling molecules using LC-MS/MS. *Anal Bioanal Chem.* 2011; 399:839–50. doi: [10.1007/s00216-010-4341-0](https://doi.org/10.1007/s00216-010-4341-0) PMID: [21046079](https://pubmed.ncbi.nlm.nih.gov/21046079/)