

1 Group W invasive meningococcal disease (MenW IMD) has steadily increased in England
2 since 2009.^{1,2} This increase has occurred in all regions and age groups, and is due to the rapid
3 expansion of a single endemic hyper-virulent strain belonging to the sequence type 11 clonal
4 complex (ST-11cc). MenW ST-11cc IMD is associated with more severe and atypical
5 presentations compared to group B disease.¹ In response, the Joint Committee on Vaccination
6 and Immunisation (JCVI) recommended the introduction of meningococcal ACWY
7 (MenACWY) conjugate vaccine to the national immunisation programme in England.^{1,3}
8 Public Health England (PHE) initiated this programme on the 26th June 2015. As a result,
9 MenACWY vaccine replaced the existing time-limited ‘freshers’ programme from August
10 2015, and was substituted directly for MenC vaccine in the routine adolescent school
11 programme for 14-15 year olds. Currently, a catch-up campaign is also being implemented to
12 offer MenACWY vaccine to all 14-18 year olds.^{3,4}

13 The rationale for targeting the vaccine solely at older adolescent and young adults is that this
14 group represents the main reservoir of meningococcal carriage.⁵ Introduction of the MenC
15 conjugate vaccine previously reduced carriage of MenC strains in this population^{6,7} and
16 evidence suggests that the quadrivalent vaccine may have a similar effect on carriage of
17 MenW strains.⁸ Reduced carriage in this population should lead to other age groups being
18 protected indirectly.⁹

19 In this study, we report MenACWY vaccine coverage in first year students arriving at the
20 University of Nottingham (UoN) in September 2015. We also report the uptake of
21 MenACWY vaccine offered to unvaccinated students via a campus-based mass vaccination
22 campaign as part of a local initiative by the University of Nottingham Health Service
23 (UNHS), in liaison with UoN, during the registration period.

24 Following approval by the Research Ethics Committee at the UoN, students were recruited to
25 assess the uptake of MenACWY conjugate vaccination prior to arrival at the university.
26 During registration (17-23rd September 2015), as part of standard clinical practice,
27 interviews were conducted with each student by UNHS healthcare professionals and their
28 vaccination history was confirmed and recorded on the UNHS registration database. Students
29 who reported that they had not received MenACWY vaccine during the interview were
30 offered immediate pro gratis vaccination with Menveo. Searches of the UNHS registration
31 database (EMIS Web software; EMIS Health, Leeds, UK software) were later performed to
32 determine vaccine coverage in the registered first year student population prior to arrival at
33 the UoN and in a subgroup of international students. Uptake of MenACWY vaccine during
34 registration was also assessed by UNHS database search and separately, to provide
35 corroborative data, by a self-reported questionnaire from 134 students recruited post-
36 registration on 28th and 29th September 2015 in two separate halls of residence on the main
37 campus of the UoN.

38 During the registration period, 7049 first year students (aged 17-25 years) registered with the
39 UNHS. On searching the UNHS database, 2160 (31%) of the 7049 registered students were
40 recorded as having received MenACWY conjugate vaccine prior to arrival at the UoN. Of the
41 remaining 4889 students who were eligible for vaccination, 2809 (57%) accepted the offer of
42 vaccination and 2080 (43%) declined vaccination. Overall, following registration,
43 MenACWY vaccine coverage in first year students at the UoN as assessed by UNHS
44 database search increased from 31% to 71%. From the follow-up questionnaire post-
45 registration on 28th and 29th September 2015, MenACWY vaccine coverage in 134 students
46 was found to be 64% confirming the significant rise in coverage following the UNHS
47 campaign. Students' self-reported vaccination status agreed with that in their health record.

48 In a subset of 804 international students aged 17-25 years, who registered with UNHS on 17th
49 and 18th September 2015, only 7 (1%) confirmed they had already received the MenACWY
50 vaccine prior to arrival. Of the remaining 797, 572 (72%) accepted and 225 (28%) declined
51 vaccination. International students, compared to UK-based students, are less likely to have
52 received MenACWY vaccine in their home country prior to arriving at university in UK.
53 Nevertheless, uptake of MenACWY vaccine when offered was equally as good in this group.
54 In the absence of a local mass vaccination campaign, it is likely that coverage would be
55 particularly low in international students, placing them at risk of MenW disease.
56 The overall vaccine coverage of 31% in this population of first year students on arrival at the
57 UoN is unsurprisingly low but did confirm that the national vaccination campaign was
58 successful in reaching a proportion of prospective English university students. PHE have
59 used a temporary general practice-based sentinel surveillance system to provide a provisional
60 estimate of vaccination coverage for the first cohort offered MenACWY vaccine from August
61 2015. An evaluation at the end of January 2016 suggested that coverage was 33.7%¹⁰, which
62 is consistent with our findings. The slightly lower MenACWY vaccine coverage found in our
63 student population is likely to be due to the inclusion of a proportion of international students
64 who were much less likely to have received vaccination before arrival.

65 High rates of transmission of meningococcal strains occur in students during the first year of
66 university attendance. As students also exhibit high mobility and wide geographical
67 distribution, they represent a major vehicle for spread of meningococcal strains into
68 communities throughout the UK. Targeting of this age group with the MenACWY vaccine
69 has the potential to significantly perturb spread of the virulent MenW ST-11cc strain. If
70 representative of coverage in this age group as a whole in England, ca. 30% MenACWY
71 vaccine coverage may be too low to significantly reduce meningococcal carriage and
72 transmission, accepting that the level of coverage needed to achieve such an effect is not

73 known. Consequently, a herd protection effect leading to a fall in MenW IMD may not be
74 achieved in the short term. In contrast, the significant boost in coverage (from 31% to 71%)
75 following the vaccine campaign at registration at the UoN demonstrates the importance of
76 offering vaccination at enrolment in tertiary educational establishments and suggests that a
77 more general implementation of this strategy may significantly improve vaccine coverage in
78 this age group. As the cost of the MenACWY vaccine is funded centrally for first year
79 university students as part of the national campaign, and an administration fee is also
80 provided (currently £9.80 per vaccine administered), the campus-based mass vaccination
81 campaign was essentially cost-neutral as advertising and the necessary facilities were
82 arranged and provided in liaison with the host institution as part of the pastoral care for new
83 students. The campus-based mass vaccination campaign delivered during the registration
84 period, also provided significant logistical advantages compared with offering a large number
85 of individual appointments for students to receive vaccination. However, a substantial
86 proportion of unvaccinated students still declined vaccination (due to a lack of perceived
87 need or benefit) suggesting that further advertising of the national campaign will be necessary
88 to raise awareness and acceptance of the MenACWY vaccine in this demographic.
89

90 References

- 91 1. Ladhani SN, Beebeejaun K, Lucidarme J et al. Increase in endemic *Neisseria meningitidis*
92 capsular group W sequence type 11 complex associated with severe invasive disease in
93 England and Wales. *Clin Infect Dis* 2015;**60**:578–85.
- 94 2. Public Health England (2016). Health Protection Report 9(7). 27 February 2015.
- 95 3. Public Health England, Meningococcal ACWY (MenACWY) vaccination programme:
96 [https://www.gov.uk/government/collections/meningococcal-acwy-menacwy-vaccination-](https://www.gov.uk/government/collections/meningococcal-acwy-menacwy-vaccination-programme)
97 programme; 2016 [accessed 04/05/16].
- 98 4. Public Health England (2016). Health Protection Report 10(8). 26 February 2016.
- 99 5. Christensen H, May M, Bowen L, Hickman M, Trotter CL. Meningococcal carriage by
100 age: a systematic review and meta-analysis. *Lancet Infect Dis* 2010;**10**:853–61.
- 101 6. Maiden MCJ, Stuart JM, for The UK Meningococcal Carriage Group. Carriage of
102 serogroup C meningococci 1 year after meningococcal C conjugate polysaccharide
103 vaccination. *The Lancet* 2002;**359**:1829–30.
- 104 7. Maiden MCJ, Ibarz-Pavón AB, Urwin R et al. Impact of meningococcal serogroup C
105 conjugate vaccines on carriage and herd immunity. *J Infect Dis* 2008;**197**:737–43.
- 106 8. Read RC, Baxter D, Chadwick DR et al. Effect of a quadrivalent meningococcal ACWY
107 glycoconjugate or a serogroup B meningococcal vaccine on meningococcal carriage: an
108 observer-blind, phase 3 randomised clinical trial. *The Lancet* 2014;**384**:2123–31.
- 109 9. Vetter V, Baxter R, Denizer G et al. Routinely vaccinating adolescents against
110 meningococcus: targeting transmission & disease. *Expert Rev Vaccines* 2016;
111 DOI:10.1586/14760584.2016.1130628.
- 112 10. Public Health England (2016). Health Protection Report 10(9). 4 March 2016.

113

114