

Hens, Bart and Corsetti, Maura and Spiller, Robin C. and Marciani, Luca and Vanuytsel, Tim and Tack, Jan and Talattof, Arjang and Amidon, Gordon L. and Koziolek, Mirko and Weitschies, Werner and Wilson, Clive G. and Bennink, Roelof J. and Brouwers, Joachim and Augustijns, Patrick (2017) Exploring gastrointestinal variables affecting drug and formulation behavior: methodologies, challenges and opportunities. International Journal of Pharmaceutics, 519 (1-2). pp. 79-97. ISSN 1873-3476

Access from the University of Nottingham repository: http://eprints.nottingham.ac.uk/39960/1/FinalVersion-Revised-FOR_EPRINTS_1_YEAR_EMBARGO_ELSEVIER.pdf

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the Creative Commons Attribution Non-commercial No Derivatives licence and may be reused according to the conditions of the licence. For more details see: http://creativecommons.org/licenses/by-nc-nd/2.5/

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

Exploring Gastrointestinal Variables Affecting Drug and Formulation Behavior: Methodologies, Challenges and Opportunities

Bart Hens^{1,5}, Maura Corsetti^{2,3}, Robin Spiller², Luca Marciani², Tim Vanuytsel³, Jan Tack³, Arjang Talattof^{4*}, Gordon L. Amidon⁵, Mirko Koziolek^{6,7}, Werner Weitschies⁷, Clive G. Wilson⁸, Roelof J. Bennink⁹, Joachim Brouwers¹, Patrick Augustijns¹

¹ Drug Delivery & Disposition, KU Leuven, Leuven, Belgium

² Nottingham Digestive Diseases Centre and NIHR Biomedical Research Unit in Gastrointestinal and Liver Diseases at Nottingham University Hospitals NHS Trust and the University of Nottingham

³ Translational Research Center for Gastrointestinal Disorders (TARGID), KU Leuven, Leuven, Belgium

⁴ Office of Generic Drugs, Center for Drug Evaluation and Research, U.S. Food and Drug Administration, Silver Spring, Maryland, USA.

⁵ College of Pharmacy, University of Michigan, Ann Arbor, Michigan, USA

⁶ Drug Delivery, Disposition and Dynamics, Monash Institute of Pharmaceutical Sciences, Monash University, Parkville, Australia

⁷ Center of Drug Absorption and Transport, Department of Pharmaceutical Technology and Biopharmacy, University of Greifswald, Greifswald, Germany

⁸ Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, United Kingdom

⁹ Academic Medical Center Amsterdam, Department of Nuclear Medicine, Amsterdam, The Netherlands

* Disclaimer: The views expressed in this article are those of the authors and not necessarily those of the Food and Drug Administration (FDA).

Corresponding author: Patrick Augustijns – patrick.augustijns@kuleuven.be

Address: Drug Delivery & Disposition, Campus Gasthuisberg O&N 2, Box 921, Herestraat 49, 3000 Leuven, Belgium. Tel.: +32 16 330301 ; fax: +32 16 330305.

Keywords: intestinal absorption, MRI, scintigraphy, manometry, telemetry, intraluminal profiling

1. Abstract

Various gastrointestinal (GI) factors affect drug and formulation behavior after oral administration, including GI transfer, motility, pH and GI fluid volume and composition. An in-depth understanding of these physiological and anatomical variables is critical for a continued progress in oral drug development. In this review, different methodologies (invasive versus non-invasive) to explore the impact of physiological variables on formulation behavior in the human GI tract are presented, revealing their strengths and limitations. The techniques mentioned allow for an improved understanding of the role of following GI variables: gastric emptying (magnetic resonance imaging (MRI), scintigraphy, acetaminophen absorption technique, ultrasonography, breath test, intraluminal sampling and telemetry), motility (MRI, small intestinal/colonic manometry and telemetry), GI volume changes (MRI and ultrasonography), temperature (telemetry) and intraluminal pH (intraluminal sampling and telemetry).

2. Introduction

In 2015, the FDA's Center for Drug Evaluation and Research (CDER) approved 45 new drugs for clinical use (U.S. Food and Drug Administration, 2016). Of these new drugs, 55% were commercialized as oral formulations either as tablets (16), capsules (8) or granules (1), indicating that the oral route of administration is still of major interest. For the last couple of decades, efforts are underway to develop and validate biorelevant *in vitro* and *in silico* tools to predict the oral absorption of new drug products prior to the start of animal studies or clinical trials (Lennernäs et al., 2014). Reliable models are essential to reduce the occurrence of failures in a late phase, resulting in time- and cost-saving drug development.

After oral intake, a drug formulation is challenged by several gastrointestinal (GI) barriers. Complex variables such as pH, GI secretions and GI transit/motility along the GI tract can affect drug release and absorption in a positive or negative way depending on the physicochemical properties of the drug compound (e.g. pH/pKa interplay for ionizable drugs) and/or the formulation characteristics (e.g. pH-sensitive coating). In addition, inter-subject variability in characteristics of the GI environment may cause variable drug absorption and systemic drug availability (Riethorst et al., 2015). Determining the median and range for specific GI variables, and understanding how they influence oral drug behavior along the GI tract, will be helpful in the field of drug development.

The present review provides an overview of different methodologies that can be applied to study physiological variables of the human GI tract and their impact on oral drug absorption in healthy and patient populations. The methodologies have been subdivided into two groups: (i) noninvasive and (ii) invasive methodologies. Although some of the methodologies are more historical than operational, they have been of paramount importance for innovations in drug and formulation development; in addition, they have created the basis for several novel methodologies, leading to an in-depth comprehension of different GI variables: gastric emptying (magnetic resonance imaging (MRI), scintigraphy, acetaminophen absorption technique, ultrasonography, breath test, intraluminal sampling and telemetry), motility (MRI, small intestinal/colonic manometry and telemetry), GI volume changes (MRI and ultrasonography), temperature (telemetry) and GI pH (intraluminal sampling and telemetry).

3. Noninvasive methodologies

a. Scintigraphy

Disturbances in the normal movement of food along the GI tract can be associated with abdominal pain, early satiety and nausea. Measurement of GI transit, especially gastric emptying (GE), therefore has become a routine and important test in clinical physics and gastroenterology clinics for patients with dyspeptic symptoms. Techniques previously used by physiologists include gastric intubation to sample dye dilution, sampling of gastric meal volumes (recoverable after different periods; Hunt, 1963), or the movement of radio-opaque markers using X-rays. The principle of incorporating gamma-emitting radiopharmaceuticals into a food matrix or a formulation allowed the non-invasive visualization of the ligand with a much reduced dose compared to X-rays and less interference with normal function (Digenis et al., 1977).

Studies of GE using scintigraphic techniques with radiolabeled standard meals were introduced by Griffith and colleagues in 1966 (Griffith et al., 1966) and further refined in the seventies with the introduction of simultaneous measurement of GE of both solids and liquids in 1976 (Heading et al., 1976). This was facilitated by labeling the phases with radionuclides of different energies that could be distinguished by gating in separate channels. GE scintigraphy has been used for more than 50 years for clinical and investigative purposes, and is usually ordered to confirm or exclude whether gastroparesis (delayed GE) is a cause of patient's symptoms (Maurer, 2015). Scintigraphy employing a gamma camera

has become the gold standard for the evaluation of GE of solids and liquids in all types of GI disorders and for assessing the efficacy of gastroprokinetic drugs and surgical procedures (Cuomo et al., 2001). Besides planar scintigraphy, single-photon emission computed tomography (SPECT) can be performed. The extra dimension of this technique is the ability to detect changes in gastric accommodation after intravenous injection of the radiopharmaceutical ^{99m}Tc-pertechnetate which shows uptake in the gastric mucosa (Bennink et al., 2004). It was demonstrated that dynamic gastric scintigraphy allows visualization and characterization of antral contractions and can also be used to evaluate the distribution of food inside the stomach and to quantify the emptying of a radiolabeled test meal from each compartment (Bennink et al., 1998). In the scintigraphic determination of GE, there are many parameters which can affect the final result. Factors such as meal size and composition, subject age and weight and measurement technique are known to influence GE. It is therefore of paramount importance that, as much as possible, standardized conditions are used in performing scintigraphic GE studies. Until recently, there were no accepted standards for performing GE scintigraphy. This problem raised concerns about the continued acceptance of GE scintigraphy without consistent methodology (Maurer, 2008). As a result, in 2007 a consensus recommendation was published jointly by the Gastrointestinal Council of the Society of Nuclear Medicine and Molecular Imaging and the American Neurogastroenterology and Motility Society (Abell et al., 2008). The consensus group recommended a solid-meal GE test "using readily available technology and normative data, which can provide clinicians with standardized results". This consensus recommendation was adopted by the Society of Nuclear Medicine and Molecular Imaging (Donohoe et al., 2009) and was included in a joint practice guideline from the American College of Radiology/Society for Pediatric Radiology and the Society of Nuclear Medicine and Molecular Imaging. In interpreting GE studies, one needs to understand the multiple factors that affect GE, particularly the separate roles of the fundus and antrum. Visual inspection of early distribution of a solid meal in the stomach has become increasingly recognized as important. Although liquids rapidly disperse throughout the stomach, solids will initially localize predominantly in the fundus until slow, sustained fundal contractions move them to the antrum (Bennink et al., 1998). The normal values for scintigraphic GE studies are usually based on male controls or a mixed control population. Historically it has been assumed that men and women have identical GE but gender differences were

occasionally noted in studies (Datz et al., 1987). The effect of gender on GE remains controversial though there is more and more evidence that women have slower GE rates (Wang et al., 2015). In female volunteers, "delayed" GE of solids seems to be related to a prolonged lag phase during which no emptying is observed, decelerating the emptying rate (Bennink et al., 1998). Moreover, GE seemed to be delayed in elderly compared to young subjects (Soenen et al., 2015). It has been demonstrated that motor dysfunction of the upper GI tract and/or colon is common in patients with functional (nonorganic) upper and/or lower GI symptoms. An association has been reported between delayed GE of solids and liquids and symptoms of postprandial fullness, early satiety, nausea and vomiting by several groups (Sarnelli et al., 2003; Stanghellini et al., 1996), although not confirmed by other studies (Talley et al., 2006).

Following the behavior of oral formulations in the upper GI tract

Early studies exploring the behavior of drug formulations utilized the expertise developed in partnerships between pharmaceutical sciences and radionuclide imaging since the potential for understanding the *in vivo* release of drugs and relating this to absorption was unsurpassed by any other imaging technology (Hardy and Wilson, 1981). The availability of technetium-99m eluted as [^{99m}Tc] sodium pertechnetate from a portable Mo-99 generator further strengthened the acceptance of scintigraphy as a clinical tool. The isotope has ideal properties: a half-life of 6.03 h and an energy of 140 keV, without beta-radiation resulting in a low radiation burden. For the study of the behavior of drug formulations in the GI lumen, it was important to determine the gut compartment in which the formulation disintegrated. The use of non-absorbed marker materials was essential, as otherwise the view of the gut would be swamped by hepatic blood flow. For this reason, chelates such as [^{99m}Tc]-labeled diethylene-triaminepentaacetic acid ([^{99m}Tc]-DTPA) were widely used as solution markers and Amberlite resins as material to mimic undissolved particulate (e.g. in a hard gelatin capsule). An example image of late dispersion and emptying in a volunteer who received a capsule formulation containing indium-111- labeled Amberlite IR120 (presented by the highlighted colour) is shown in Figure 1.



Figure 1: Emptying following dispersion of a capsule containing micronised Amberlite labeled with Indium-111.

Early studies conducted by the Nottingham group investigated factors such as differential GE of drugs relative to meal according to size of the matrix (Davis et al., 1984). It has been established in dogs that large objects were more likely to be retained in the fed state. This retention was also seen in man, accentuated by intake of a heavy meal with larger enteric coated dosage forms (Wilson et al., 1989), whereas systems that disintegrated or dispersed into component particulates emptied with the liquid phase. Results for swelling dosage forms were inconsistent due to the variable initiation of the housekeeper sequence (migrating motor complex or MMC) relative to the rate of swelling. Similarly, floating dosage forms needed sufficient contents to be present and were very sensitive to posture (Bennett et al., 1984). For Biopharmaceutics Classification System (BCS) class I compounds, the rate of emptying of the stomach determines the absorption profile, since dissolution may occur in the stomach and drug uptake occurs as soon as it enters the small intestine. Based on this principle, the acetaminophen absorption technique was introduced as an alternative technique for measuring GE (discussed in the next section: Acetaminophen Absorption Technique). In the development of a novel fast acting formulation containing paracetamol (FD-APAP), the granulate was labeled by the addition of [¹¹¹In]-labeled DTPA (Wilson et al., 2011). Because scintigraphy allows a more rapid sampling than taking blood, the effects of small changes in dissolution rate of a drug compound become more evident in gastric scintigraphy compared to the acetaminophen absorption technique (Figure 2).



Figure 2: Composite of the pharmacokinetic data and the illustrated images for standard acetaminophen 1000 mg dose (2 x 500 mg tablets) and fast acting formulation containing paracetamol 1000 mg dose (FD-APAP; 2 x 500 mg tablets. Adopted from Wilson et al. (2011). Copyright Taylor & Francis 2011.

Regarding small intestinal transit times (SITT), scintigraphy demonstrated reasonably constant transit times of approximately 3 h (3 ± 1 h; mean \pm SEM) in 201 normal subjects, irrespective of prandial state (ranging from fasted state to heavy breakfast) and formulation type (solutions, small pellets, and single units (matrix tablets and osmotic pumps)).

Gastro-retention is a principle which can be used to deliver, for instance, weakly basic drug compounds which have good acid solubility but which precipitate upon entry in the upper intestine. Such materials have a pH-based absorption window in the proximal intestine. By retaining a formulation in the stomach, the drug can be allowed to percolate into the duodenum at a slow rate maintaining the systemic exposure more effectively than an immediate release formulation for instance. There are several mechanisms including flotation, swelling, and folding and coating (Lopes et al., 2016). Scintigraphy can be used to measure the residence time and dispersion of the formulation in the stomach to evaluate the extent of coating. For example, sucralfate is used to treat and prevent ulcers by forming a precipitate on the exuded proteins lost from the ulcer site. The compound will form strong complexes with indium – labeled radiopharmaceuticals. A formulation of [^{113m}In] -labeled sucralfate was administered to volunteers and a typical image is shown in Figure 3.



Figure 3: Scintiscan of [^{113m}Tn]-labeled sucralfate after swallowing 20 ml labeled suspension.

In a related fashion, gastro-retentive dosage forms can be radiolabeled by incorporation of a nonabsorbed radiopharmaceutical. This is useful for evaluating swelling dosage forms and floating systems. Swelling, by changing the geometry, could sustain delivery by a physicochemical rather than solely a physiological mechanism once it had left the stomach. Scintigraphy assists by unambiguously demonstrating that the sustaining action is in part due to residence in the stomach.

b. Acetaminophen absorption technique

Since the seventies, new insights have been generated in the variability of therapeutic drug outcomes among patient populations. Different physiological factors such as intestinal motility, GE rate, splanchnic blood flow and volume, composition and pH of GI fluids were explored in detail in animal experiments. Their clinical relevance, however, remained uncertain as only a few studies were carried out in man. In 1973, Heading et al. investigated the relationship between gastric half time ($T_{1/2, G}$) and the plasma C_{max} and T_{max} of acetaminophen (paracetamol) in patients (Heading et al., 1973). Acetaminophen is classified as a highly soluble and highly permeable drug (BCS class 1) (Amidon et al., 1995). Based on these properties, it can be hypothesized that GE directly influences paracetamol absorption (as measured from the plasma concentration-time profile) by controlling the rate at which the drug is delivered to the small intestine. Within 6 days following acetaminophen absorption study, the GE rate (expressed as gastric half-time; $T_{1/2, G}$) was measured by a sequential scintigraphic technique using the chelate indium-113m DTPA, the gold standard technique to measure GE in those days. Maximum plasma concentrations (C_{max}) and time to maximum plasma concentrations (T_{max}) of acetaminophen were strongly correlated to $T_{1/2, G}$, as depicted in Figure 4a and Figure 4b, respectively.



Figure 4: (a) Relationship between $T_{1/2, G}$ and the peak plasma acetaminophen (i.e. paracetamol) concentration (r= -0.77, p < 0.005). (b) Relationship between GE half time ($T_{1/2, G}$) and the time of the peak plasma acetaminophen concentration (r= 0.76, p < 0.005). Adopted from Heading et al. (1973). Copyright Wiley 1973.

Although statistically significant correlations were observed for the entire population of 14 patients, the GE of 5 out of 14 patients did not correlate with the plasma C_{max} nor T_{max} of acetaminophen. In this study, the authors claimed that interindividual variation in the rate of drug absorption was due to differences in the rate of GE, which in turn depends on (i) disease state, (ii) endocrine and autonomic activity and (iii) GI variables such as volume, temperature, pH and composition of the stomach contents. Willems and colleagues reviewed studies carried out from 1973 until 1996 and concluded that the lack of a standardized protocol confounded the pharmacokinetic parameter (C_{max} , T_{max} , AUC, etc.) that was able to describe the process of GE appropriately (Sanaka et al., 1997; Willems et al., 2001). Moreover, caloric differences between test meals and the use of different dosage forms (tablet versus solution) make it difficult to interpret and compare all studies with each other (Sanaka et al., 1998). Therefore, because of these drawbacks, other techniques are more preferred to study gastric emptying.

c. Radio-opaque markers

Radio-opaque markers (i.e. barium, platinum, tantalum, gold) are high-density particles of metals or compounds that will absorb radiation and can be visualized by radiography (X-rays). Back in 1969, Hinton et al. explored the use of radio-opaque markers to measure GI transit times in healthy subjects and in ileostomy subjects (Hinton et al., 1969). Twenty solid cylindrical pellets of polythene with various thickness (2, 3 or 5 mm), containing 20% (w/w) of barium sulphate, were orally administered before breakfast.

Based on radiographs of the stool of 25 healthy volunteers, pellets could be visualized and data were presented as (a) the time when the first markers appeared and (b) when 80% of the markers appeared in the feces. The first marker(s) passed within 30 hours (range: 9-59 h) and 80% of the markers passed in 45 hours (range: 25-84 h, except for one volunteer: 150 h). In ileostomy subjects, the average GI transit time was 2.5 h, ranging from 0.75 to 4.75 h.

Rather than working with stool samples, abdominal radiographs can estimate transit times in the different colonic segments. In 1987, Metcalf and colleagues measured colonic transit times with the use of distinctive markers, ingested by 24 healthy subjects on 3 consecutive days (Metcalf et al., 1987). Daily abdominal X-rays were taken to visualize the position of these markers. Transit times were calculated by the following equation:

$$T = \Delta t \times \frac{n}{N}$$
 (Equation 1)

Where T is the transit time in a given colonic segment, Δt is the time between two marker ingestions (in this case 24 h), n is the number of markers counted in a given colonic segment and N is the number of markers ingested daily. A mean colonic transit time of 35.0 h was calculated in this study for healthy volunteers. In case of patients complaining of functional constipation, Pomerri and colleagues investigated colonic transit times in 50 patients who were asked to ingest 10 identical cylindrical radio-opaque markers every day during 10 days (Pomerri et al., 2008). In addition to the radio-opaque markers, 8 to 10 ml of a commercial thick 113% (w/v) barium paste was co-administered as a colonic tracer. On day 11, fluoroscopic images were obtained in order to visualize the exact place of the different

administered doses of radio-opaque markers and, subsequently, colonic transit times were calculated by Equation 1.

Figure 5 depicts the fluoroscopic image of a constipated patient at day 11 of the study. The paste of the barium formulation that was orally ingested clearly identifies the different regions of the colon. Colonic segments of interest were the right colon (Figure 5A; ileocecal junction together with the ascending colon and right part of the transverse colon), the left colon (Figure 5B; left part of the transverse colon and the descending colon) and the recto-sigmoid colon (Figure 5C).



Figure 5: Fluoroscopic image showing small radio-opaque particles present in the different anatomical segments of the colon; a: right colonic segment; b: left colonic segment; c: recto-sigmoid colon. Adapted from Pomerri et al. (2008). Copyright Elsevier 2008.

Results indicated a transit time of 30.5 ± 24.4 h for the right colonic segment, a transit time of 42.9 ± 34.9 h for the left colonic segment and 23.6 ± 21.4 h for the recto-sigmoid colon, resulting in a total colonic transit time of 96.9 ± 62.0 h. Focusing on colonic drug delivery, important lessons need to be kept in mind. Despite the reasonably constant small intestinal transit time (cfr. supra: Scintigraphy), the food or drug matrix spends a considerable time at the ileocecal junction, which functions mechanistically as a valve allowing fecal material to become more concentrated before transfer to the colon (Davis et al., 1986). For time-dependent colonic drug delivery systems, which will release drug after a predetermined lag time, movement through this junction can be a significant variable (Jose et al., 2009;

Kinget et al., 1998; Pišlar et al., 2015; Wilson, 2010). Moreover, in case of GI disorders (ulcerative colitis and Crohn's disease), episodes of diarrhea can reduce colonic transit times in such a way that drug delivery at the site of the colon will be inefficient (van der Sijp et al., 1993).

Comparing different studies regarding GI transit times is quite challenging when using radio-opaque markers: most studies do not report appropriately what happened after the intake of the markers (mealtimes, medication use, personal activities, etc.). Therefore, standardized protocols are necessary and indispensable in order to further rely on this technique and to allow comparison between results with other studies.

d. <u>Ultrasonography</u>

As already mentioned, scintigraphy is the gold standard for measurement of GE (Wilding, 2002). The emitted radiation and high costs, however, are still well-argued reasons to look for alternative techniques. Ultrasonography (also known as echography) is a non-invasive, low cost and safe methodology to study GE in patients and healthy persons (Gilja et al., 1999). A transducer is placed on the abdominal skin, with contact gel to enhance wave conduction, and generates an 'ultrasound' with such a high frequency that is not audible to humans. These ultrasound waves will travel through the body and will be echoed back to the interface between hard and soft structures of tissues to make a distinction across the different organs. These reflected waves will be displayed as images (ultrasonograms) to the operator (Bateman and Whittingham, 1982).

In 1984, King et al. applied transabdominal two-dimensional (2D) ultrasonography in order to study (i) terminal antral contractions, (ii) duodenal contractions, (iii) forward flow movement of gastric content and (iv) retrograde flow movements through the pylorus (King et al., 1984). Seventeen healthy volunteers were given a test meal of 500 ml of water flavored with orange juice and warmed to 37°C, to which 0.5 g chopped and sieved bran was added. Subsequently, volunteers were sitting in an upright position and the upper abdomen was scanned directly after ingestion of the meal up to 30 min (Figure 6).



Figure 6: (a) Ultrasonographic image of the upper abdomen after ingestion of the test meal. Stomach and proximal duodenum are distended and bran particles can be seen as small white specks suspended in gastric contents. Concentric antral contraction, which is throwing an acoustic shadow, is travelling along the mid portion of the antrum towards the pylorus. (b) Peristaltic wave has progressed along the antrum and is now approaching the pylorus. (c) An overview of all different organs that can be detected after ultrasonography. Adopted from King et al. (1984). Copyright British Medical Journal 1984.

In the same study, four volunteers were selected to investigate antroduodenal motility and transpyloric

fluid movement (Figure 7).



Figure 7: Comparison of terminal antral contractions, duodenal contractions, forward flow and retrograde flow through pylorus with their corresponding duration of time. Adopted from King et al. (1984). Copyright British Medical Journal 1984.

Duodenal contractions can be seen to occur after the midpoint of the antral contractions. Forward flow of gastric content was mostly observed immediately before or after terminal antral contractions, whereas retrograde flow of intestinal contents was observed immediately before duodenal contractions take place. Accurate assessment of intragastric volumes, however, was not possible with this twodimensional technique. A three-dimensional (3D) view of the stomach can assess the volumes of the different gastric compartments and their emptying characteristics. In addition to the ultrasound transducer, a commercially available magnetometer-based position and orientation measurement (POM) was added to the system in order to obtain 3D images (Gilja et al., 1997). Examples of 3D wireframes of the stomach after intake of a 500 ml soup meal are depicted in Figure 8. The average compartmental postprandial gastric volumes in 14 healthy male volunteers are shown in Figure 9.



Figure 8: 3D wireframes of the stomach of a healthy subject reconstructed from ultrasound images obtained by magnetic scanhead tracking. Panel A presents fasting stomach volumes, whereas panels B, C and D present postprandial stomach volumes after 5, 10 and 15 min of intake of a soup meal, respectively. Adopted from Gilja et al. (1997). Copyright Elsevier 1997.



Figure 9: Gastric volumes in the distal, proximal and total part of the stomach as a function of time after administration of 500 ml of soup to 14 male volunteers. Gastric volumes were measured in the distal and proximal compartments of the stomach. Adopted from Gilja et al. (1997). Copyright Elsevier 1997.

The observed $T_{1/2, G}$ was 22.1 ± 3.8 min. Intragastric volume distribution, expressed as the ratio between proximal and distal gastric volumes, varied from 3.6 ± 2.1 (5 min after meal) to 2.7 ± 1.9 (30 min after meal), indicating progressive redistribution of the meal towards the distal stomach. This technique is easy to handle, non-invasive and does not involve the use of radiation. The success of this technique, however, is totally dependent on the expertise of the researcher.

e. Breath test

The gastric emptying breath test is a non-invasive, non-operator dependent technique, based on stable (e.g. 13 C) or radioactive (e.g. 14 C) isotope testing in the breath after ingestion of a labeled solid and/or liquid meal. In a typical protocol, 13 C or 14 C octanoic acid (i.e. medium chain fatty acid) is solubilized in egg yolk and cooked in an omelet, which is eaten by the subject together with a piece of toast and a glass of water (Ghoos et al., 1993; Perri et al., 2005). Once the omelet has reached the duodenum, rapid disintegration of the labeled solid phase occurs with subsequent absorption and oxidation of 13 C/ 14 C octanoic acid to 13 CO₂/ 14 CO₂, respectively, which can be measured in breath samples of the volunteer/patient. Since gastric emptying to the duodenum is the rate-limiting step in this entire process, the rate of 13 CO₂/ 14 CO₂ appearing in the breath as a function of time is reflecting the GE rate. The stable isotope 13 C is preferred over radioactive 14 C to reduce radiation burden to patients, volunteers and the environment. When both solid and liquid GE are assessed simultaneously, dual 13 C and 14 C labeling of

the respective solid and liquid parts of the meal is needed. The metabolic pathway is depicted in Figure 10.



Figure 10: Subsequent metabolic steps of ¹³C-octanoic acid in egg yolk after oral administration. Adopted from Perri et al. (2005). Copyright European Review for Medical and Pharmacological Sciences 2005.

Ghoos and colleagues collected exhaled ¹³CO₂ from 36 subjects (16 healthy volunteers and 20 patients with dyspeptic symptoms) after eating a scrambled egg of which the yolk was doped with 100 mg ¹³C octanoic acid and the egg white was labeled with 3 mCi of ^{99m}Tc albumin colloid (Ghoos et al., 1993). Half-life of gastric emptying ($T_{1/2, G}$) was estimated based on the ¹³CO₂ breath excretion (% dose/h) and gamma radiation (% retention of ^{99m}Tc albumin colloid) by making use of mathematical derivations (Figure 11).



Figure 11: (a) Breath test: ${}^{13}CO_2$ breath excretion in (% dose/h) as a function of time after ingestion of ${}^{13}C$ -octanoic acid. Tlag_b stands for the time to maximum ${}^{13}CO_2$ excretion; $t_{1/2b}$ is the breath test determined GE time. (b) Scintigraphic test: % gastric radioactivity retention as a function of time after ingestion of ${}^{99m}Tc$. $T_{1/2s}$ is the time at which half of the γ -emitting labeled meal is retained in the stomach and Tlag_s is the time needed for the antral contractions to grind solids into particles small enough to pass through the pylorus. Adopted from Perri et al. (2005). Copyright European Review for Medical and Pharmacological Sciences 2005.

 $T_{1/2, G}$ derived from the breath test (t_{1/2b}) and from the scintigraphy study (t_{1/2s}) showed a correlation coefficient of 0.89. A mean value of 72 ± 22 min was observed for t_{1/2b}. This study demonstrated that the breath test can be a valid alternative to scintigraphy to assess GE.

As many factors (i.e. disease, drugs, gastric surgery, endocrine disorders, etc.) can accelerate or delay GE, these should be taken into account in drug and formulation development for these specific populations. To assess acceleration and delay in GE, Maes and colleagues performed a study in nine healthy volunteers using ¹⁴C octanoic acid breath test for solids (Maes et al., 1994). On three consecutive days, volunteers were tested in three different conditions: (i) without medication, (ii) 30 minutes after intravenous administration of 200 mg of erythromycin (motilin-receptor agonist with strong prokinetic properties) and (iii) 60 minutes after oral administration of 30 mg of propantheline (an anti-muscarinic, transit-inhibiting agent). In all nine subjects, $T_{1/2, G}$ decreased after intravenous administration of erythromycin (p = 0.0168), as depicted in Figure 12.



Figure 12: Comparison of T_{1/2, G} in each individual in normal conditions and after administration of erythromycin and propantheline. Adopted from Maes et al. (1994). Copyright British Medical Journal 1994.

For all three test conditions, $T_{1/2, G}$ were 37.44 ± 21.32 min (erythromycin), 71.22 ± 26.68 min (basal), and 141.44 ± 87.74 min (propantheline). These values can serve as reference data for optimization of *in vitro/in silico* models. These half-emptying times can easily be programmed in *in vitro* models that are frequently applied to predict the *in vivo* performance of a drug in postprandial conditions. For example, standard implemented $T_{1/2, G}$ in TIM-1 model is set on 80 min, mimicking the *in vivo* situation in fed state conditions (Brouwers et al., 2011). In case of an accelerated or delayed state of GE, the $T_{1/2, G}$ can easily be reprogrammed (Hens et al., 2014). Which technique to use for measuring GE (scintigraphy, acetaminophen absorption technique, breath test, ultrasonography, etc.), will depend on the impact of the technique for the volunteer/patient, ethical considerations, costs and facilities available for performing the study.

f. Magnetic resonance imaging (MRI)

Magnetic resonance imaging (MRI) images the hydrogen protons whose signal characteristics change depending on their physicochemical environment, mobility and concentration. It has multi-planar capability, good imaging speed and resolution, and superb soft tissue contrast, which is tunable to different tissues and fluids. After an initial slow development, MRI of gastrointestinal function has developed substantially. MRI has been demonstrated to complement other techniques by providing unique data on the undisturbed bowel environment and therefore has the ability to improve the in vitro/in vivo relevance of drug disposition modeling. MRI is inherently suited to image fluids and organ volumes. Gastric volumes and GE have been measured since the inception of GI MRI (Evans et al., 1993; Schwizer et al., 1992; Stehling et al., 1989). Gastric secretion has been investigated by monitoring the dilution of a gadolinium (Gd) contrast agent (Curcic et al., 2015; Hoad et al., 2015) and gastric volume data are increasingly being handled using semi-automated data processing (Bharucha et al., 2014). More recently, the attention is shifting to monitoring the *in vivo* environment of standard drug testing conditions. A recent study investigated gastric fluid contents under the standard fasting and fed oral dosage form testing conditions (Mudie et al., 2014, Koziolek et al., 2014). This work has been extended to the small bowel environment. MRI showed that liquid in the small bowel is present in discrete liquid "pockets" (Schiller et al., 2005); these have been quantified under conditions representing bioavailability/bioequivalence (BA/BE) studies at fasted baseline and after a 240 ml glass of water, using methods validated against naso-duodenal infusion (Hoad et al., 2007). The method is based on a heavily T2 (i.e. time when 63% of the transverse magnetization has decayed) weighted sequence that images only freely mobile water. The signal from less mobile water molecules, for instance associated with mucus, will have decayed before the imaging time and is therefore not included (Figure 13).



Figure 13: MRI images of the stomach and small bowel of a healthy volunteer after ingestion of 240 ml water. (A) stomach; (B) abdomen. (C) Maximum intensity projection small bowel water pockets. Separate water pockets are rendered in different colors. Adopted from Mudie et al. (2014). Copyright ACS 2014.

MRI has been used in a limited number of studies to directly visualize a drug product. In one study, the intragastric distribution of a colloidal carrier based on Gd-labeled liposomes was studied (Faas et al., 2001). In another study, MRI was used to monitor the gastric position and residence time of gastric-retentive tablets labeled with iron particles or Gd chelates (Steingoetter et al., 2003). In addition, MRI has been used to indirectly visualize the effect of an oral formulation on the GI system. These include

imaging the effect of a polyethylene glycol bowel preparation on bowel fluids distribution and colonic motility (Marciani et al., 2014) and the formation and time courses of sodium alginates formulations for gastroesophageal reflux (Marciani et al., 2002; Sweis et al., 2013). Cine-MRI can image the GI tract with sufficient speed to visualize motility. This can be done on the stomach looking at antral motility frequency and amplitude (Kunz et al., 1998; Marciani et al., 2001; Schwizer et al., 1994, 1996; Wright et al., 1999). More recently, the technique has been extended to the small bowel (Hahnemann et al., 2015; Menys et al., 2014; Odille et al., 2012) and to the colon (Buhmann et al., 2005; Menys et al., 2014). Small bowel motility is most commonly acquired under MRI enterography (Patak et al., 2001) whereby the subjects are asked to ingest a large amount of fluid to distend the bowel walls. Respiratory motion is a problem for quantitation. Either breath hold acquisition (Froehlich et al., 2005) or postprocessing data registration techniques (Hamy et al., 2014) can be employed to overcome this issue and enable quantitation. Another parameter that MRI can measure is bowel transit. This has been done using 'transit capsule markers' filled with MRI-visible fluid. The imaging was based either on standard ¹H MRI (Chaddock et al., 2014; Schiller et al., 2005) but also on ¹⁹F MRI (Hahn et al., 2011, 2012, 2014), an interesting approach though the availability of the necessary hardware is scarce. The position of the MRI markers can be visualized and transit determined in a fashion similar to the radiological methods (Chaddock et al., 2014). A limitation is the relatively large size of the capsules which are likely to stay inside the stomach until a migrating motor complex moves them into the small bowel. Hence their transit time may differ from that of small tablets but this limitation could be removed by the development of smaller markers or by making use of magnetic marker monitoring (MMM). In this approach, an oral dosage form will be marked with ferromagnetic material to create a magnetic dipole that can be traced in the gastrointestinal tract as a function of time after magnetization in a strong magnetic field. As such, the transit of these dosage forms can accurately be visualized as 3D pathways, enabling the evaluation of, for instance, gastric residence and emptying of an oral dosage form. This is demonstrated in Figure 14, where the passage of a non-disintegrating capsule through the stomach of a healthy male subject is shown in four repeated experiments (Weitschies et al., 2010).



Figure 14: Passages of a non-disintegrating capsule through the stomach monitored in four repeated experiments in a healthy volunteer. (a) Experiment 1 (gastric emptying time: 24 min), (b) Experiment 2 (gastric emptying time: 20 min), (c) Experiment 3 (gastric emptying time: 14 min) and (d) Experiment 4 (gastric emptying time: 25 min). Adopted from Weitschies et al., 2010. Copyright Elsevier 2010.

4. Invasive methodologies

a. Intraluminal sampling methodology

Intraluminal sampling and subsequent characterization of the aspirated human gastrointestinal fluids is a versatile approach to investigate different aspects of drug and formulation behavior. The use of intestinal catheters to explore drug and formulation behavior dates back to the eighties, starting with perfusion of intestinal segments to evaluate *in vivo* drug permeability, stability and dissolution. Afterwards, the sampling technology was applied to study the intraluminal behavior of orally administered drug products by measuring drug concentrations along the GI tract (Figure 15).



Figure 15: Schematic of tube placement with aspiration channels in the stomach (1), duodenum (2), proximal jejunum (3), and distal jejunum (4).

In 1985, Jobin et al. perfused different regions of the GI tract with a solution of metoprolol. Disappearance of the drug over time in the aspirated intestinal fluid and appearance in the plasma indicated intestinal absorption. Approximately 60% of drug that was emptied from the stomach was absorbed from the duodenum, whereas 50% of metoprolol that left from the duodenum was absorbed from the proximal jejunum (Jobin et al., 1985). A more refined technique to study human permeability for different drug compounds is the Loc-I-Gut[®] perfusion technique. A multilumen catheter is equipped with two balloons which are inflated to isolate a specific region of interest along the intestinal fluid, the net human effective permeability can be calculated (Dahan et al., 2012; Lennernäs, 1998). Moreover, the Loc-I-Gut[®] perfusion technique has also been widely applied to explore intestinal drug metabolism (Petri et al., 2003; Tannergren et al., 2003) and *in vivo* dissolution of drugs in the intestinal tract when the segment of interest will be perfused with a suspension of the drug (Bønløkke et al., 1997).

Aside from intestinal perfusion, intubation techniques allow exploring intraluminal drug concentrations after oral administration of drug formulations. In combination with the simultaneous assessment of systemic drug concentration, this allows the evaluation of (i) how a drug and its corresponding formulation behave after oral intake, (ii) how this behavior affects drug absorption and (iii) how certain GI variables (e.g. GE, pH, secretions, transit times, etc.) influence oral drug disposition in man (Brouwers and Augustijns, 2014). Many drug compounds nowadays exhibit very poor aqueous solubility. Their fate depends on the formulation or dosing strategy in order to achieve sufficient therapeutic systemic concentrations. Enabling formulations such as solid dispersions, nano- and microparticles and cyclodextrin/co-solvent based solutions are examples of methods used to increase solubilization or achieve supersaturated solution of the drug, with the ultimate goal of enhancing oral absorption. Aspiration of gastric and duodenal fluids in parallel with blood sampling resulted in new insights into how these enabling strategies behave after oral intake (Brouwers et al., 2006; Geboers et al., 2015; Hens et al., 2015a; Stappaerts et al., 2015). Moreover, drug solubility measurements in these aspirated fluids can reveal whether the drug was supersaturated at the time of aspiration or directly started to precipitate (Hens et al., 2015b; Psachoulias et al., 2011; Van Den Abeele et al., 2015).

In addition to being instrumental in exploring gastrointestinal formulation behavior, the sampling technique can also be used to collect blank gastrointestinal fluids. These fluids can then be used as solvent system in *in vitro* assays (e.g. solubility determination (Augustijns et al., 2014; Kostewicz et al., 2014b), permeability assays (Wuyts et al., 2015a, 2015b) and supersaturation evaluation (Bevernage et al., 2013)), or, upon characterization, as reference for the optimization of simulated gastrointestinal fluids. Characterization of intestinal fluids collected from healthy volunteers in terms of pH, bile salts, phospholipids, cholesterol and lipid degradation products indicate large differences in composition which potentially implies differences in solubilizing capacity that may eventually account for inter-individual variations in systemic exposure after oral administration of lipophilic drugs (Kalantzi et al., 2006; Lindahl et al., 1997; Riethorst et al., 2015). Unraveling the composition of these intestinal fluids has led to the development of simulated intestinal fluids (fasted and fed state simulated intestinal fluids) and have been further optimized throughout the years. Substantial inter-individual differences have also been demonstrated by electron microscopy imaging of the colloidal structures present in intestinal fluids, especially after intake of a meal (Fatouros et al., 2009; Müllertz et al., 2012; Riethorst et al., 2016). Characterization of the contents of the distal ileum and cecum revealed major differences in terms of

liquid content, buffer capacity and pH compared to the present fluids in the upper small intestine (Reppas et al., 2015). These differences may significantly influence the performance of drugs during their residence in the lower parts of the intestine. Unraveling the composition of colonic fluids led to better insights in colonic drug delivery strategies, resulting in pH-responsive and bacterially-triggered drug delivery technology (Basit et al., 2004, 2009; Ibekwe et al., 2008; McConnell et al., 2008; Vertzoni et al., 2010a), as well as optimized dissolution media to assess the performance of for instance colontargeted drug formulations in the lower intestine. These findings have led to the development of biorelevant fasted and fed state simulated colonic fluid (FaSSCoF and FeSSCoF, respectively), developed to mimic the fluids collected from the ascending colon in healthy adults (Vertzoni et al., 2010b). Thinking beyond the intestine, a lot of interest goes out to the dynamic yet complex environment of the stomach (Van Den Abeele et al., 2016b). The acidic pH as reported in literature may have a major impact on oral drug behavior along the GI tract in case of weakly basic compounds. Reported median values ranging from 1.55 to 1.8 may enhance drug's gastric dissolution, depending on the dissociation constant(s) of the drug compound. Besides gastric pH, gastric motility has shown to affect drug distribution along the stomach (Van Den Abeele et al., 2016a). Further studies should focus on how gastric motility will affect drug appearance in plasma in terms of C_{max} and T_{max} which may explain the failure of bioequivalence studies in some cases due to variability in gastrointestinal motility among subjects when an oral drug is administered (Talattof et al., 2016a). Characterization of gastric fluids has led to the development of simulated gastric media, referred to as fasted and fed state simulated gastric fluid (FaSSGF and FeSSGF, respectively). Biorelevant implementation of pH, bile salts and phospholipids is reflected in FaSSGF, whereas FeSSGF is composed of equal amounts of FaSSGF and a liquid meal, as frequently applied in intraluminal profiling studies in fed state conditions (Vertzoni et al., 2007).

Site-specific aspiration has led to the exploration of formulation behavior in the more distal parts of the GI tract, especially when dealing with delayed/controlled release formulations (Hens et al., 2016). Measuring drug concentrations at multiple site-specific locations along the GI tract will expand our knowledge about the life cycle of the drug after oral administration. A recent study implemented a

redesigned aspiration catheter (3 meters in length with 4 aspiration ports located in the stomach, duodenum/proximal jejunum, mid jejunum, and distal jejunum/ileum, depicted in Figure 15) to quantify mesalamine concentrations along the various regions of the GI tract following ingestion correlating pharmacokinetics with local availability of medications within the GI tract (Brown et al., 2014). Previously, Marathe et al. showed that the systemic availability of metformin, which is primarily absorbed in the small intestine, is significantly improved with delayed GE (Marathe et al., 2000). Oberle and Amidon linked the double peak phenomenon in plasma levels of cimetidine with the variable gastric emptying and transit along the intestinal tract combined with a short plasma elimination half-life and poor absorption (Oberle and Amidon, 1987). As stated by Brouwers and Augustijns, gastrointestinal profiling has moved forward last decades to evaluate intraluminal drug and formulation behavior in humans. Intensified use of this technique will guide us to the next level in optimization of *in vitro* and *in silico* tools for intestinal absorption (Brouwers and Augustijns, 2014).

b. Small intestinal manometry

GI motility and the associated migrating motor or myoelectric complex (MMC) play a crucial role in transporting ingested material from the stomach through the intestine and into the colon by means of segmental and peristaltic contractions (Cannon, 1912; Deloose et al., 2012). The propagating wave of peristalsis is regulated by hormones, paracrine signaling, and the autonomic nervous system, while segmentation is carried out by longitudinal muscle relaxation and circular muscle contraction thereby mixing GI contents with digestive enzymes and ensuring composition uniformity and sufficient epithelial contact for absorption (Culen et al., 2013). The GE rate is controlled by gastric distention promoting emptying and intestinal stimuli slowing emptying (Hunt, 1963). Contractile activity propels matter from the stomach into the small bowel where segmental contractions beginning in the duodenum reach the terminal ileum in approximately 2 hours (Grivel and Ruckebusch, 1972). The MMC is defined by three distinct phases: phase I is an inert period with little activity; phase II features sporadic contractions gradually ascending in magnitude but with little net forward movement of gastric contents; and phase III is characterized by powerful, high frequency contractile bursts that promote emptying of contents where peak flow rates are observed (Deloose et al., 2012; Kerlin et al., 1982). As the contractile activity propagates, it becomes less spatiotemporally organized resulting in slower propulsion rates in

the distal small bowel (Sarna and Otterson, 1989). A typical phase I through III tracing is shown in Figure 16, adapted from Hansen (Hansen, 2002).



Figure 16: Tracing showing phase I, phase II, and phase III of the fasting cycle in the antrum (A1 & A2), duodenum (D1 & D2), and jejunum (J1 & J2). Adopted from Hansen (2002). Copyright Physiological Research 2002.

In the context of drug products, GI motility has become the focus of an increasing number of studies, especially in how it affects formulations with respect to transit and dissolution (Chen et al., 2013; Greiff and Rowbotham, 1994). Recent modeling approaches have been undertaken to incorporate variable GI transit times (Hénin et al., 2012) as well as simulate such physiological effects on oral drug products and their GI residence times, which can impact the extent of absorption and ultimately bioavailability/bioequivalence (Talattof et al., 2016b). Motility can be measured by means of manometry, employing catheters perfused with deionized water to record intestinal pressure in the duodenum, jejunum, and ileum. The spacing of the ports on motility catheters serves not only for recording of local pressure but also for determining the propagation velocities of contractile activities. The pressure is measured with a capillary infusion pump system with pressure transducers transforming intraluminal pressure into electrical signals via the strain gauge transducers (Figure 17). With improving technologies, high resolution manometric studies can potentially reveal new insights into of GI motility. It is thus important to acknowledge the complexities of GI motility as a source of tremendous intrinsic variability that needs further investigation, perhaps linked with higher-level physiology of slow wave propagation that is potentially an underlying phenomenon driving motility but whose exact relationships

remain unclear (Myers et al., 2002; O'Grady et al., 2010).



Figure 17: A 16-channel pressure recording of an anterograde antroduodenal phase III of the MMC (upper panel) and two sequences shown with high temporal resolution (lower panel). The start point of the duodenal pressure waves is indicated by a black dot and the direction of propagation of the duodenal pressure waves indicated. A, antrum; D, duodenum; S, pyloric sleeve. Adopted from Castedal and Abrahamsson (2001). Copyright Blackwell Science Ltd 2011.

c. <u>Colonic manometry</u>

The colon represents the distal part of the GI tract and plays an important role in the maintenance of electrolyte and fluid balance, in the metabolism of carbohydrates and bile acids, in the absorption of fatty acids and as a reservoir for feces. It is generally believed that it achieves these functions through the prolongation of the residence of its contents inside the lumen, which explains the longer colonic residence time as compared to small bowel transit time (approximately 35 hours vs 4 hours, respectively) in healthy subjects (Phillips, 1984).

Colonic motility is an important determinant of colonic transit and can be studied by different techniques including colonic manometry. As reported by Dinning and colleagues, the study of colonic motility was performed with catheters applying different recording systems (water-perfused or solid state) but generally allowing to measure the pressure created by colonic contractions on a limited number of pressure sensors (from 4 to 16) spaced 7-12 cm one from each other (conventional or low-resolution manometry) (Dinning et al., 2013). Manometric recordings of the relatively inaccessible human colon

are complicated by a number of technical difficulties that typically necessitate partial or total removal of colonic contents in order to position a recording catheter. Therefore, most of these studies have been conducted in pre-cleaned colon applying a colonoscopy-assisted retrograde intubation via the anus. Only a limited number of studies have been conducted using retrograde intubation in non-prepared colon (Rao et al., 2001). Even fewer studies have been performed by means of anterograde intubation via naso-colonic intubation (Bampton et al., 2001).

In the case of 24-hours conventional manometry, it has been demonstrated that colonic motility mainly consists of non-propagated activity as opposed to propagated sequences of low amplitude (5-60 mmHg) and high amplitude (more than 100 mmHg) occurring only 2.4 ± 0.1 and 0.4 ± 0.1 per hour, respectively (Bampton et al., 2001). Both anterograde and retrograde propagating sequences have been described as well as simultaneous pan-colonic low-amplitude pressurization events (Bampton et al., 2001; Rao et al., 2001). In healthy subjects, colonic motility in general is reduced during sleeping time and significantly increases at awakening and during meals (Rao et al., 2001). It has been reported that anterograde propagating sequences occur more frequently during the awaking period, simultaneous sequences significantly increase at awakening and after the meal, retrograde sequences represent a minority of the colonic motor patterns, and high-amplitude propagating sequences occur at awakening and about 1-2 hours after the meal (Bampton et al., 2001; Rao et al., 2001).

Recently, the introduction of the high-resolution manometry (HRM) has revealed new aspects of the colonic motility in healthy volunteers (Dinning et al., 2013). Two types of catheters have been applied: a fiber-optic catheter with up to 90 recording sensors spaced 1 cm from each other (Dinning et al., 2013) and a solid state catheter with up to 40 pressure sensors spaced 1-2.5 cm from each other (Chen et al., 2014; Corsetti et al., 2016). In a study with a fiber-optic catheter, the majority of activity in the postprandial period were retrograde propagating sequences, mainly occurring in the sigmoid region (Dinning et al., 2013). The relevance of simultaneous pressurization sequences, previously observed in conventional manometry studies, was confirmed in several studies using a HRM solid state catheter (Chen et al., 2014; Corsetti et al., 2016). Based on preliminary data in healthy adults, this activity seems to represent the most frequent colonic motor pattern, which increases significantly during the meal and

decreases afterwards, in concert with increased occurrence of retrograde propagating colonic sequences in the left colon (Chen et al., 2014; Corsetti et al., 2016).

With respect to the role of different colonic motor patterns on transport of colonic contents, few data are available in the literature. In studies combining the conventional manometry with scintigraphic assessment of bolus movements, it has been demonstrated that colonic propulsion is relatively infrequent (only 30% of propagating sequences are propulsive) and is associated with propagating sequences originating in the more proximal part of the colon, which have a higher amplitude and lower velocity of propagation (Cook et al., 2000). In studies focusing on the pre- and postprandial period, it has been observed that before the meal, the presence of simultaneous sequences associates with almost no movement of colonic contents (Bazzocchi et al., 1990). During the first 30 minutes after the meal, the presence of simultaneous and retrograde sequences in the sigmoid colon is associated with movement of colonic contents in retrograde direction (Bazzocchi et al., 1990). After this period, the colonic contents move both in anterograde and retrograde direction until the occurrence of the high-amplitude propagating sequences, which normally move the majority of colonic contents from the proximal part (transverse) to the distal colon, inducing the sensation and, in a next step, the activation of defecation (Bazzocchi et al., 1990; Herbst et al., 1997). Anterograde and retrograde low-amplitude propagating sequences are believed to play a major role in the mixing function, which allows the colon to absorb water and nutrients and slow the progression of luminal contents (Phillips, 1984). The increase of both simultaneous and retrograde activity, observed by both conventional and high resolution manometry in the left colon during the postprandial period, has been hypothesized to play a role in facilitating the retrograde transport of colonic contents to the transverse colon, producing the necessary distension to generate high-amplitude propagating sequences occurring later after the meal (Bazzocchi et al., 1990).

Regarding colonic motility alterations in bowel disorders, all but one study used conventional manometry. Overall, these studies have demonstrated that the colonic response to a meal (i.e. the number of retrograde propagating sequences and the number of high-amplitude propagating sequences) is reduced in adults with functional constipation as compared to healthy subjects (Dinning et al., 2013). In adults with functional diarrhea, the number of high and low amplitude anterograde propagating

sequences was increased, while simultaneous and retrograde activity was less frequent (Herbst et al., 1997). A similar increase in anterograde propagating sequences has been observed in patients with moderately active ulcerative colitis, but not in patients in remission (Bassotti et al., 2014).

When compared with the small bowel, the colon in general plays a minor role in the absorption of oral drugs. However, its role is relevant for the absorption of certain drugs specifically delivered to the colon or metabolized to their active form in the colon (e.g. sulfasalazine) for local treatment (i.e. inflammatory bowel disease). Moreover, numerous studies have recently shown the importance of the colon as a target for oral absorption: large drug molecules such as proteins and peptides that are known to degrade in the acidic environment of the stomach may show moderate colonic absorption, if delivered intact in the colon (Amidon et al., 2015). Finally, the colon may contribute to the absorption of drugs with extended or delayed absorption kinetics, including highly dosed, low solubility drugs and modified release dosage forms (Markopoulos et al., 2015).

Several factors have been reported to influence the absorption and activity of colon delivered drugs, including low volume and high viscosity and low volume of colonic fluids (challenging drug dissolution), colonic pH (the colonic pH varies across the different segments of the colon and this can affect the pharmacokinetic and pharmacodynamics of drugs), the presence of bacteria (which produce different enzymes able to metabolize drugs and to induce the formation of active or inactive metabolites) and the colonic transit time (which can modify the bioavailability of drugs as such) (Amidon et al., 2015; Kostewicz et al., 2014a).

Considering that colonic manometry needs bowel preparation and positioning assisted by colonoscopy, this technique could be less attractive as compared to other non-invasive methodologies (e.g. cine-MRI) to study the influence of colonic motility on oral absorption and activity of drugs. However, it can be considered in studies aiming to assess the influence of colonic motility on colonic and systemic profiling of an orally administered drug targeting the colon. In these cases, catheters to collect colonic fluids can be advanced together with the manometry probe in the colon and maintained during the entire duration of colonic motility recording.

d. <u>Telemetry</u>

The interest in telemetric capsules in the field of oral biopharmaceutics has increased in recent years due to significant progress in developing supporting technologies and the improved accessibility of certified commercial products. Telemetric capsules are used by physicians to examine the GI tract for organic pathologies and to characterize transit conditions in health and disease. For the pharmaceutical scientist, the spectrum of applications is wider as both the sensor and the delivery technologies are useful. These allow identification of the luminal conditions, visualization of disintegration/drug release in the gut, and calibration of regional drug absorption. Apart from the marketed products that are presented in Table 1 and discussed below, recent publications describe prototypes developed to overcome limitations of the currently available devices, incorporating functions such as anchoring and repositioning, steerable devices and improved control on release (Goffredo et al., 2015; Munoz et al., 2014).

	Dimensions	Application in oral	Domostra
	(mm x mm)	biopharmaceutics	Kemarks
Heidelberg pH capsule	21 x 8	Investigation of luminal pH profiles	Extensive experience (> 100 studies)
SmartPill [®]	26 x 13	Investigation of luminal pH, temperature and pressure profiles	CE* certification
Enterion [®]	32 x 11	Regional drug absorption studies	Drug actively released as bolus Tracking by scintigraphy
InteliSite [®]	35 x 10	Regional drug absorption studies	Drug passively released as bolus Tracking by scintigraphy
IntelliCap [®]	27 x 11	Regional drug absorption studies Investigation of luminal pH and temperature profiles	Drug actively released either in controlled manner or as bolus CE certification
PillCam[®]	26 x 11	Visualization of <i>in vivo</i> behavior of oral dosage forms	CE certification

Table 1: Selection of telemetric capsules used in the field of oral biopharmaceutics.

*CE: Conformité Européene/ European Conformity

The GI pH value is one of the major parameters determining intraluminal solubility of the majority of orally administered drugs, but unfortunately, the number of techniques to measure GI pH values is limited. In contrast to commonly applied aspiration techniques and pH catheters, telemetric capsules represent a comfortable alternative. As they are ingested in the same way as capsules or tablets and move freely through the gut, they generate profiles that provide valuable information about the physiological transit conditions of solid oral dosage forms. A major drawback of pH-sensing telemetric capsules is the pH drift of the commonly applied ion-selective field effect transistor (ISFET) sensors (Khanna, 2013). This effect is typically minimized by drift correction based on post-calibration results

(Abbas et al., 2014). One of the first systems used in biopharmaceutics, was the Heidelberg pH capsule. In 1990, Dressman and co-workers used this system to measure GI pH profiles in the stomach and duodenum after a high-fat meal (Dressman et al., 1990). These data provided important insights into fed state human GI physiology and were the basis for the development of biorelevant dissolution tests.

In addition to high-temporal pH and temperature monitoring, the SmartPill[®] system offers the unique opportunity to measure pressure within the GI tract (Cassilly et al., 2008). The GI motility can be crucial for oral drug delivery as high pressures can cause unwanted drug release profiles such as dose dumping. In contrast to catheters used for manometry, SmartPill[®] is freely moving within the GI tract; thus, the generated pressure profiles are representative for large non-digestible dosage forms such as hydrogel matrix tablets. The transit conditions after ingestion of the high-caloric, high-fat FDA standard breakfast used for fed state BA/BE studies were determined by use of SmartPill[®] in a recent study by Koziolek and co-workers (Koziolek et al., 2015) (Figure 18). It was shown in this study that the maximum pressures in the human GI tract amount to 293 ± 109 mbar and typically arise shortly before or during GE (Koziolek et al., 2015).



Figure 18: Individual pH (black), pressure (red) and temperature (blue) profiles over time measured by administering the SmartPill system 30 min after beginning of the intake of the high-caloric, high-fat standard breakfast recommended for food-effect studies. (GE: gastric emptying, CA: colonic arrival). Adopted from Koziolek et al. (2015). Copyright Elsevier 2015.

Telemetric capsule systems can further be used to study regional drug absorption in the GI tract. As such, they may aid development and characterization of new formulations as the presence of absorption windows or the potential for colon targeting can be evaluated. Drug release is initiated either passively by physical or chemical triggers (e.g. HF capsule, InteliSite[®]) or actively by different mechanisms (e.g.

Enterion[®], IntelliCap[®]). The volume that can be released is typically below 1 ml (Goffredo et al., 2015). The IntelliCap[®] system is currently the most advanced system. It can be programmed to release a drug either continuously or as a bolus by a miniaturized pump (Becker et al., 2014; van der Schaar et al., 2013). The system consists of a body and a cap (Figure 19).



Figure 19: The orally swallowable IntelliCap[®] system to quantify regional drug absorption in the human GI tract. Adopted from Becker et al., 2014. Copyright Springer Link 2014.

The cap is clipped onto the body and consists of a medication container that can be filled with a drug liquid. Based on real-time pH and temperature profiles, the drug can be released at a certain region of the GI tract making the system a valuable tool for regional drug absorption studies (Becker et al., 2014). Söderlind and co-workers investigated the value of the IntelliCap[®] system for regional drug absorption studies by comparing the plasma concentration profiles of metoprolol (BCS class I) after three different release profiles with the administration of an oral solution. They showed that the system was suitable to deliver the drug in a controlled manner over a period of up to 14 h. However, the IntelliCap[®] version used in this study was not able to deliver uniform drug pulses, which would limit its suitability for studies aiming at the immediate release of a drug at a certain place (e.g. colon targeting) (Söderlind et al., 2015). Recently, an updated version of the IntelliCap (IntelliCap FR) was developed to ensure the fast release of a drug in either liquid or solid form; however, clinical data proving the usefulness of this modification have not been published yet. In GI investigations, wireless capsule endoscopy (WCE) has

revolutionized investigative procedures which were uncomfortable for the patient. Therefore, for example, the following studies attempted to visualize the *in vivo* behavior of solid oral dosage forms within the GI tract by making use of WCE: Pedersen and co-workers attached a PillCam[®] with a filament to the teeth in order to keep the system inside the stomach. By this, they could record the disruption and dispersion of an oil-filled capsule and a supersaturating nano-emulsifying drug delivery system (SNEDDS) (Pedersen et al., 2014). Another interesting approach was published recently by Keller and colleagues. They managed to control the position of a PillCam[®] with the aid of magnets (Keller et al., 2011).

Although telemetric capsules are already valuable tools in oral biopharmaceutics, their field of application is limited to certain biopharmaceutical questions. In particular, the high price of the systems as well as the lack of tracking tools and localization control is often limiting the usage of telemetric capsules. In addition, the large dimensions (similar to capsule size 000) of most systems can alter gastric transit times, especially in the fed state (Koziolek et al., 2015), and may pose a risk to subjects or patients with GI obstruction. Nonetheless, the current progress in technology will most probably overcome these limitations and contribute to expanded functionality (e.g. active locomotion, sampling capability) of telemetric capsules. Advances in micro-robotic technology including better power usage, signal transmission, delivery technologies and new sensors ensure that their importance for oral biopharmaceutics will further increase in the future (Amoako-Tuffour et al., 2014; Goffredo et al., 2015).

5. Future perspectives: multidisciplinary approach

As there is currently no 'ideal' technique able to provide a holistic view on the interplay between GI physiology and oral drug delivery, the combination of different techniques seems critical for further progress in understanding this interplay. For the next years to come, research may therefore focus on intraluminal profiling of a drug along the GI tract in parallel with examining specific GI variables (e.g. GE, motility, residual volumes, pH, etc.) to evaluate their influence on intraluminal concentrations of the drug and, ultimately, oral drug absorption. This implies that collaborations among physicians and pharmaceutical scientists are indispensable to reveal the dynamic nature of the GI tract by the above-

mentioned techniques and how this dynamic nature helps to explain the behavior of an orally administered drug. The acquired multi-dimensional information of drug concentrations and measured impact of the dynamic physiology on these concentrations is crucial to open discussion on whether or not we need to consider implementing certain GI variables in predictive yet simple *in vitro* and *in silico* models, depending on the type of drug and/or formulation.

6. Acknowledgments

This work has received support from the Innovative Medicines Initiative Joint Undertaking (http://www.imi.europa.eu) under Grant Agreement No. 115369, resources of which are composed of financial contribution from the European Union's Seventh Framework Program and EFPIA companies' in kind contribution. Bart Hens acknowledges the Institute for the Promotion of Innovation through Science and Technology in Flanders (IWT Vlaanderen) for a PhD grant. Tim Vanuytsel is a senior clinical investigator of the Research Foundation Flanders (FWO). This work was also supported by Award # HHSF223201510157C by the U.S. Food and Drug Administration (FDA); this review represents the position of the authors and not necessarily that of the FDA.

7. <u>References</u>

- Abbas, A., Wilding, G.E., Sitrin, M.D., 2014. Does Colonic Transit Time Affect Colonic pH ? J Gastroenterol Hepatol Res 3, 1103–1107. doi:10.6051/j.issn.2224-3992.2014.03.399
- Abell, T.L., Camilleri, M., Donohoe, K., Hasler, W.L., Lin, H.C., Maurer, A.H., McCallum, R.W., Nowak, T., Nusynowitz, M.L., Parkman, H.P., Shreve, P., Szarka, L.A., Snape, W.J., Ziessman, H.A., American Neurogastroenterology and Motility Society and the Society of Nuclear Medicine, 2008. Consensus recommendations for gastric emptying scintigraphy: a joint report of the American Neurogastroenterology and Motility Society and the Society of Nuclear Medicine. J Nucl Med Technol 36, 44–54. doi:10.2967/jnmt.107.048116
- Amidon, G.L., Lennernäs, H., Shah, V.P., Crison, J.R., 1995. A theoretical basis for a biopharmaceutic drug classification: the correlation of in vitro drug product dissolution and in vivo bioavailability. Pharm. Res. 12, 413–420.
- Amidon, S., Brown, J.E., Dave, V.S., 2015. Colon-targeted oral drug delivery systems: design trends and approaches. AAPS PharmSciTech 16, 731–741. doi:10.1208/s12249-015-0350-9
- Amoako-Tuffour, Y., Jones, M.L., Shalabi, N., Labbé, A., Vengallatore, S., Prakashm, S., 2014. Ingestible Gastrointestinal Sampling Devices : State-of-the-Art and Future Directions. Crit Rev Biomed Eng 42, 1–15.
- Augustijns, P., Wuyts, B., Hens, B., Annaert, P., Butler, J., Brouwers, J., 2014. A review of drug solubility in human intestinal fluids: implications for the prediction of oral absorption. Eur J Pharm Sci 57, 322–332. doi:10.1016/j.ejps.2013.08.027
- Bampton, P.A., Dinning, P.G., Kennedy, M.L., Lubowski, D.Z., Cook, I.J., 2001. Prolonged multi-point recording of colonic manometry in the unprepared human colon: providing insight into potentially relevant pressure wave parameters. Am. J. Gastroenterol. 96, 1838–1848. doi:10.1111/j.1572-0241.2001.03924.x

- Basit, A.W., Podczeck, F., Newton, J.M., Waddington, W.A., Ell, P.J., Lacey, L.F., 2004. The use of formulation technology to assess regional gastrointestinal drug absorption in humans. Eur J Pharm Sci 21, 179–189.
- Basit, A.W., Short, M.D., McConnell, E.L., 2009. Microbiota-triggered colonic delivery: robustness of the polysaccharide approach in the fed state in man. J Drug Target 17, 64–71. doi:10.1080/10611860802455805
- Bassotti, G., Antonelli, E., Villanacci, V., Baldoni, M., Dore, M.P., 2014. Colonic motility in ulcerative colitis. United European Gastroenterol J 2, 457–462. doi:10.1177/2050640614548096
- Bateman, D.N., Whittingham, T.A., 1982. Measurement of gastric emptying by real-time ultrasound. Gut 23, 524–527.
- Bazzocchi, G., Ellis, J., Villanueva-Meyer, J., Jing, J., Reddy, S.N., Mena, I., Snape, W.J., 1990.
 Postprandial colonic transit and motor activity in chronic constipation. Gastroenterology 98, 686–693.
- Becker, D., Zhang, J., Heimbach, T., Penland, R.C., Wanke, C., Shimizu, J., Kulmatycki, K., 2014. Novel Orally Swallowable IntelliCap[®] Device to Quantify Regional Drug Absorption in Human GI Tract Using Diltiazem as Model Drug. AAPS PharmSciTech DOI: 10.12. doi:10.1208/s12249-014-0172-1
- Bennett, C.E., Hardy, J.G., Wilson, C.G., 1984. The influence of posture on the gastric emptying of antacids. International Journal of Pharmaceutics 21, 341–347. doi:10.1016/0378-5173(84)90192-3
- Bennink, R., Peeters, M., Van den Maegdenbergh, V., Geypens, B., Rutgeerts, P., De Roo, M.,
 Mortelmans, L., 1998. Comparison of total and compartmental gastric emptying and antral motility between healthy men and women. Eur J Nucl Med 25, 1293–1299.
- Bennink, R.J., van den Elzen, B.D., Kuiken, S.D., Boeckxstaens, G.E., 2004. Noninvasive measurement of gastric accommodation by means of pertechnetate SPECT: limiting radiation dose without losing image quality. J. Nucl. Med. 45, 147–152.
- Bevernage, J., Brouwers, J., Brewster, M.E., Augustijns, P., 2013. Evaluation of gastrointestinal drug supersaturation and precipitation: strategies and issues. Int J Pharm 453, 25–35. doi:10.1016/j.ijpharm.2012.11.026
- Bharucha, A.E., Karwoski, R.A., Fidler, J., Holmes, D.R., Robb, R.A., Riederer, S.J., Zinsmeister, A.R., 2014. Comparison of manual and semiautomated techniques for analyzing gastric volumes with MRI in humans. Am. J. Physiol. Gastrointest. Liver Physiol. 307, G582-587. doi:10.1152/ajpgi.00048.2014
- Bønløkke, L., Christensen, F.N., Knutson, L., Kristensen, H.G., Lennernäs, H., 1997. A New Approach for Direct In Vivo Dissolution Studies of Poorly Soluble Drugs. Pharm Res 14, 1490–1492. doi:10.1023/A:1012149511768
- Brouwers, J., Anneveld, B., Goudappel, G.-J., Duchateau, G., Annaert, P., Augustijns, P., Zeijdner, E., 2011. Food-dependent disintegration of immediate release fosamprenavir tablets: In vitro evaluation using magnetic resonance imaging and a dynamic gastrointestinal system. European Journal of Pharmaceutics and Biopharmaceutics 77, 313–319. doi:10.1016/j.ejpb.2010.10.009
- Brouwers, J., Augustijns, P., 2014. Resolving intraluminal drug and formulation behavior:
 Gastrointestinal concentration profiling in humans. European Journal of Pharmaceutical
 Sciences, Theme Issue 5th World Conference on Drug Absorption, Transport and Delivery 61, 2–10. doi:10.1016/j.ejps.2014.01.010
- Brouwers, J., Tack, J., Lammert, F., Augustijns, P., 2006. Intraluminal drug and formulation behavior and integration in in vitro permeability estimation: a case study with amprenavir. J Pharm Sci 95, 372–383. doi:10.1002/jps.20553
- Brown, S.W., Bly, M.J., Baker, J., May, K., Fioritto, A.F., Hasler, W.L., Sun, D., 2014. Su1081 A Novel Method to Investigate Local Concentrations of Mesalamine in the Gastrointestinal Tract of Healthy Volunteers. Gastroenterology 146, S-367. doi:10.1016/S0016-5085(14)61330-5

- Buhmann, S., Kirchhoff, C., Wielage, C., Mussack, T., Reiser, M.F., Lienemann, A., 2005. Assessment of large bowel motility by cine magnetic resonance imaging using two different prokinetic agents: a feasibility study. Invest Radiol 40, 689–694.
- Cannon, W.B., 1912. Peristalsis, segmentation, and the myenteric reflex. American Journal of Physiology 30, 114–128.
- Cassilly, D., Kantor, S., Knight, L.C., Maurer, A.H., Fisher, R.S., Semler, J., Parkman, H.P., 2008. Gastric emptying of a non-digestible solid: assessment with simultaneous SmartPill pH and pressure capsule, antroduodenal manometry, gastric emptying scintigraphy. Neurogastroenterol Motil 20, 311–319. doi:NMO1061 [pii]10.1111/j.1365-2982.2007.01061.x
- Castedal, M., Abrahamsson, H., 2001. High-resolution analysis of the duodenal interdigestive phase III in humans. Neurogastroenterol. Motil. 13, 473–481.
- Chaddock, G., Lam, C., Hoad, C.L., Costigan, C., Cox, E.F., Placidi, E., Thexton, I., Wright, J., Blackshaw, P.E., Perkins, A.C., Marciani, L., Gowland, P.A., Spiller, R.C., 2014. Novel MRI tests of orocecal transit time and whole gut transit time: studies in normal subjects. Neurogastroenterol. Motil. 26, 205–214. doi:10.1111/nmo.12249
- Chen, J.-H., Yu, Y., Yang, Z., Yu, W.-Z., Huizinga, J.D., 2014. Sa2007 Colonic Motor Patterns in Patients With Chronic Constipation Assessed by High Resolution Manometry. Gastroenterology 146, S-353. doi:10.1016/S0016-5085(14)61275-0
- Chen, M.-L., Sadrieh, N., Yu, L., 2013. Impact of osmotically active excipients on bioavailability and bioequivalence of BCS class III drugs. AAPS J 15, 1043–1050. doi:10.1208/s12248-013-9509-z
- Cook, I.J., Furukawa, Y., Panagopoulos, V., Collins, P.J., Dent, J., 2000. Relationships between spatial patterns of colonic pressure and individual movements of content. Am. J. Physiol. Gastrointest. Liver Physiol. 278, G329-341.
- Corsetti, M., Pagliaro, G., Desmedts, I., Deloose, E., Gevers, A., Scheerens, C., Rommel, N., Tack, J., 2016. Pan-colonic pressurizations associated with relaxation of the anal sphincter in health and disease: a new colonic motor pattern identified using high-resolution manometry. Am J gastroenterol.
- Culen, M., Rezacova, A., Jampilek, J., Dohnal, J., 2013. Designing a dynamic dissolution method: a review of instrumental options and corresponding physiology of stomach and small intestine. J Pharm Sci 102, 2995–3017. doi:10.1002/jps.23494
- Cuomo, R., Sarnelli, G., Grasso, R., Bruzzese, D., Pumpo, R., Salomone, M., Nicolai, E., Tack, J., Budillon, G., 2001. Functional dyspepsia symptoms, gastric emptying and satiety provocative test: analysis of relationships. Scand. J. Gastroenterol. 36, 1030–1036.
- Curcic, J., Sauter, M., Schwizer, W., Fried, M., Boesiger, P., Steingoetter, A., 2015. Validation of a golden angle radial sequence (GOLD) for abdominal T1 mapping during free breathing: demonstrating clinical feasibility for quantifying gastric secretion and emptying. J Magn Reson Imaging 41, 157–164. doi:10.1002/jmri.24530
- Dahan, A., Lennernäs, H., Amidon, G.L., 2012. The fraction dose absorbed, in humans, and high jejunal human permeability relationship. Mol. Pharm. 9, 1847–1851. doi:10.1021/mp300140h
- Datz, F.L., Christian, P.E., Moore, J., 1987. Gender-related differences in gastric emptying. J. Nucl. Med. 28, 1204–1207.
- Davis, S.S., Hardy, J.G., Fara, J.W., 1986. Transit of pharmaceutical dosage forms through the small intestine. Gut 27, 886–892.
- Davis, S.S., Hardy, J.G., Taylor, M.J., Whalley, D.R., Wilson, C.G., 1984. A comparative study of the gastrointestinal transit of a pellet and tablet formulation. International Journal of Pharmaceutics 21, 167–177. doi:10.1016/0378-5173(84)90091-7
- Deloose, E., Janssen, P., Depoortere, I., Tack, J., 2012. The migrating motor complex: control mechanisms and its role in health and disease. Nat Rev Gastroenterol Hepatol 9, 271–285. doi:10.1038/nrgastro.2012.57

- Digenis, G.A., Beihn, R.M., Theodorakis, M.C., Shambhu, M.B., 1977. Use of 99mTc-labeled triethylenetetramine-polystyrene resin for measuring the gastric emptying rate in humans. J Pharm Sci 66, 442–443.
- Dinning, P.G., Wiklendt, L., Gibbins, I., Patton, V., Bampton, P., Lubowski, D.Z., Cook, I.J., Arkwright, J.W., 2013. Low-resolution colonic manometry leads to a gross misinterpretation of the frequency and polarity of propagating sequences: Initial results from fiber-optic highresolution manometry studies. Neurogastroenterol. Motil. 25, e640-649. doi:10.1111/nmo.12170
- Donohoe, K.J., Maurer, A.H., Ziessman, H.A., Urbain, J.-L.C., Royal, H.D., Martin-Comin, J., Society for Nuclear Medicine, American Neurogastroenterology and Motility Society, 2009. Procedure guideline for adult solid-meal gastric-emptying study 3.0. J Nucl Med Technol 37, 196–200. doi:10.2967/jnmt.109.067843
- Dressman, J.B., Berardi, R.R., Dermentzoglou, L.C., Russell, T.L., Schmaltz, S.P., Barnett, J.L., Jarvenpaa, K.M., 1990. Upper gastrointestinal (GI) pH in young, healthy men and women. Pharm Res 7, 756–761.
- Evans, D.F., Lamont, G., Stehling, M.K., Blamire, A.M., Gibbs, P., Coxon, R., Hardcastle, J.D., Mansfield,
 P., 1993. Prolonged monitoring of the upper gastrointestinal tract using echo planar
 magnetic resonance imaging. Gut 34, 848–852.
- Faas, H., Schwizer, W., Feinle, C., Lengsfeld, H., de Smidt, C., Boesiger, P., Fried, M., Rades, T., 2001. Monitoring the intragastric distribution of a colloidal drug carrier model by magnetic resonance imaging460. Pharm. Res. 18, 460–466.
- Fatouros, D.G., Walrand, I., Bergenstahl, B., Müllertz, A., 2009. Colloidal structures in media simulating intestinal fed state conditions with and without lipolysis products. Pharm. Res. 26, 361–374. doi:10.1007/s11095-008-9750-9
- Froehlich, J.M., Patak, M.A., von Weymarn, C., Juli, C.F., Zollikofer, C.L., Wentz, K.-U., 2005. Small bowel motility assessment with magnetic resonance imaging. J Magn Reson Imaging 21, 370– 375. doi:10.1002/jmri.20284
- Geboers, S., Haenen, S., Mols, R., Brouwers, J., Tack, J., Annaert, P., Augustijns, P., 2015. Intestinal behavior of the ester prodrug tenofovir DF in humans. Int J Pharm 485, 131–137. doi:10.1016/j.ijpharm.2015.03.002
- Ghoos, Y.F., Maes, B.D., Geypens, B.J., Mys, G., Hiele, M.I., Rutgeerts, P.J., Vantrappen, G., 1993.
 Measurement of gastric emptying rate of solids by means of a carbon-labeled octanoic acid breath test. Gastroenterology 104, 1640–1647.
- Gilja, O.H., Detmer, P.R., Jong, J.M., Leotta, D.F., Li, X.N., Beach, K.W., Martin, R., Strandness, D.E., 1997. Intragastric distribution and gastric emptying assessed by three-dimensional ultrasonography. Gastroenterology 113, 38–49.
- Gilja, O.H., Hausken, T., Ødegaard, S., Berstad, A., 1999. Gastric emptying measured by ultrasonography. World J Gastroenterol 5, 93–94. doi:10.3748/wjg.v5.i2.93
- Goffredo, R., Accoto, D., Guglielmelli, E., 2015. Swallowable smart pills for local drug delivery: present status and future perspectives. Expert Rev Med Devices 12, 585–599. doi:10.1586/17434440.2015.1061933
- Goffredo, R., Accoto, D., Guglielmelli, E., 2015. Swallowable smart pills for local drug delivery: present status and future perspectives. Expert Rev Med Devices 12, 585–599. doi:10.1586/17434440.2015.1061933
- Greiff, J.M., Rowbotham, D., 1994. Pharmacokinetic drug interactions with gastrointestinal motility modifying agents. Clin Pharmacokinet 27, 447–461. doi:10.2165/00003088-199427060-00004
- Griffith, G.H., Owen, G.M., Kirkman, S., Shields, R., 1966. Measurement of rate of gastric emptying using chromium-51. Lancet 1, 1244–1245.
- Grivel, M.L., Ruckebusch, Y., 1972. The propagation of segmental contractions along the small intestine. J. Physiol. (Lond.) 227, 611–625.

- Hahn, T., Kozerke, S., Schwizer, W., Fried, M., Boesiger, P., Steingoetter, A., 2014. Real-time multipoint gastrointestinal 19-fluorine catheter tracking. Magn Reson Med 71, 302–307. doi:10.1002/mrm.24654
- Hahn, T., Kozerke, S., Schwizer, W., Fried, M., Boesiger, P., Steingoetter, A., 2012. 19F MR imaging golden angle-based capsule tracking for intestinal transit and catheter tracking: initial in vivo experience. Radiology 265, 917–925. doi:10.1148/radiol.12120297
- Hahn, T., Kozerke, S., Schwizer, W., Fried, M., Boesiger, P., Steingoetter, A., 2011. Visualization and quantification of intestinal transit and motor function by real-time tracking of 19F labeled capsules in humans. Magn Reson Med 66, 812–820. doi:10.1002/mrm.22822
- Hahnemann, M.L., Nensa, F., Kinner, S., Gerken, G., Lauenstein, T.C., 2015. Motility mapping as evaluation tool for bowel motility: initial results on the development of an automated colorcoding algorithm in cine MRI. J Magn Reson Imaging 41, 354–360. doi:10.1002/jmri.24557
- Hamy, V., Dikaios, N., Punwani, S., Melbourne, A., Latifoltojar, A., Makanyanga, J., Chouhan, M., Helbren, E., Menys, A., Taylor, S., Atkinson, D., 2014. Respiratory motion correction in dynamic MRI using robust data decomposition registration - application to DCE-MRI. Med Image Anal 18, 301–313. doi:10.1016/j.media.2013.10.016
- Hansen, M.B., 2002. Small intestinal manometry. Physiol Res 51, 541–556.
- Hardy, J.G., Wilson, C.G., 1981. Radionuclide imaging in pharmaceutical, physiological and pharmacological research. Clin Phys Physiol Meas 2, 71–121.
- Heading, R.C., Nimmo, J., Prescott, L.F., Tothill, P., 1973. The dependence of paracetamol absorption on the rate of gastric emptying. British Journal of Pharmacology 47, 415–421. doi:10.1111/j.1476-5381.1973.tb08339.x
- Heading, R.C., Tothill, P., McLoughlin, G.P., Shearman, D.J., 1976. Gastric emptying rate measurement in man. A double isotope scanning technique for simultaneous study of liquid and solid components of a meal. Gastroenterology 71, 45–50.
- Hénin, E., Bergstrand, M., Standing, J.F., Karlsson, M.O., 2012. A mechanism-based approach for absorption modeling: the Gastro-Intestinal Transit Time (GITT) model. AAPS J 14, 155–163. doi:10.1208/s12248-012-9324-y
- Hens, B., Brouwers, J., Anneveld, B., Corsetti, M., Symillides, M., Vertzoni, M., Reppas, C., Turner, D.B., Augustijns, P., 2014. Gastrointestinal transfer: in vivo evaluation and implementation in in vitro and in silico predictive tools. Eur J Pharm Sci 63, 233–242. doi:10.1016/j.ejps.2014.07.008
- Hens, B., Brouwers, J., Corsetti, M., Augustijns, P., 2015a. Gastrointestinal behavior of nano- and microsized fenofibrate: In vivo evaluation in man and in vitro simulation by assessment of the permeation potential. Eur J Pharm Sci 77, 40–47. doi:10.1016/j.ejps.2015.05.023
- Hens, B., Brouwers, J., Corsetti, M., Augustijns, P., 2015b. Supersaturation and Precipitation of Posaconazole upon Entry in the Upper Small Intestine in Humans. J Pharm Sci. doi:10.1002/jps.24690
- Hens, B., Corsetti, M., Brouwers, J., Augustijns, P., 2016. Gastrointestinal and Systemic Monitoring of Posaconazole in Humans After Fasted and Fed State Administration of a Solid Dispersion. J Pharm Sci 105, 2904–2912. doi:10.1016/j.xphs.2016.03.027
- Herbst, F., Kamm, M.A., Morris, G.P., Britton, K., Woloszko, J., Nicholls, R.J., 1997. Gastrointestinal transit and prolonged ambulatory colonic motility in health and faecal incontinence. Gut 41, 381–389.
- Hinton, J.M., Lennard-Jones, J.E., Young, A.C., 1969. A new method for studying gut transit times using radioopaque markers. Gut 10, 842–847. doi:10.1136/gut.10.10.842
- Hoad, C.L., Marciani, L., Foley, S., Totman, J.J., Wright, J., Bush, D., Cox, E.F., Campbell, E., Spiller, R.C., Gowland, P.A., 2007. Non-invasive quantification of small bowel water content by MRI: a validation study. Phys Med Biol 52, 6909–6922. doi:10.1088/0031-9155/52/23/009
- Hoad, C.L., Parker, H., Hudders, N., Costigan, C., Cox, E.F., Perkins, A.C., Blackshaw, P.E., Marciani, L., Spiller, R.C., Fox, M.R., Gowland, P.A., 2015. Measurement of gastric meal and secretion

volumes using magnetic resonance imaging. Phys Med Biol 60, 1367–1383. doi:10.1088/0031-9155/60/3/1367

Hunt, J.N., 1963. GASTRIC EMPTYING IN RELATION TO DRUG ABSORPTION. Am J Dig Dis 8, 885-894.

- Ibekwe, V.C., Khela, M.K., Evans, D.F., Basit, A.W., 2008. A new concept in colonic drug targeting: a combined pH-responsive and bacterially-triggered drug delivery technology. Aliment. Pharmacol. Ther. 28, 911–916. doi:10.1111/j.1365-2036.2008.03810.x
- Jobin, G., Cortot, A., Godbillon, J., Duval, M., Schoeller, J.P., Hirtz, J., Bernier, J.J., 1985. Investigation of drug absorption from the gastrointestinal tract of man. I. Metoprolol in the stomach, duodenum and jejunum. Br J Clin Pharmacol 19 Suppl 2, 97S–105S.
- Jose, S., Dhanya, K., Cinu, T., Litty, J., Chacko, A., 2009. Colon targeted drug delivery: Different approaches. Journal of Young Pharmacists 1, 13–19. doi:http://dx.doi.org.kuleuven.ezproxy.kuleuven.be/10.4103/0975-1483.51869
- Kalantzi, L., Goumas, K., Kalioras, V., Abrahamsson, B., Dressman, J.B., Reppas, C., 2006.
 Characterization of the human upper gastrointestinal contents under conditions simulating bioavailability/bioequivalence studies. Pharm. Res. 23, 165–176. doi:10.1007/s11095-005-8476-1
- Keller, J., Fibbe, C., Volke, F., Gerber, J., Mosse, A.C., Reimann-Zawadzki, M., Rabinovitz, E., Layer, P., Schmitt, D., Andresen, V., Rosien, U., Swain, P., 2011. Inspection of the human stomach using remote-controlled capsule endoscopy: a feasibility study in healthy volunteers (with videos). Gastrointest Endosc 73, 22–28. doi:10.1016/j.gie.2010.08.053
- Kerlin, P., Zinsmeister, A., Phillips, S., 1982. Relationship of motility to flow of contents in the human small intestine. Gastroenterology 82, 701–706.
- Khanna, V.K., 2013. Remedial and adaptive solutions of ISFET non-ideal behaviour. Sensor Rev 33, 228–237. doi:10.1108/02602281311324681
- King, P.M., Adam, R.D., Pryde, A., McDicken, W.N., Heading, R.C., 1984. Relationships of human antroduodenal motility and transpyloric fluid movement: non-invasive observations with real-time ultrasound. Gut 25, 1384–1391.
- Kinget, R., Kalala, W., Vervoort, L., van den Mooter, G., 1998. Colonic drug targeting. J Drug Target 6, 129–149. doi:10.3109/10611869808997888
- Kostewicz, E.S., Aarons, L., Bergstrand, M., Bolger, M.B., Galetin, A., Hatley, O., Jamei, M., Lloyd, R., Pepin, X., Rostami-Hodjegan, A., Sjögren, E., Tannergren, C., Turner, D.B., Wagner, C., Weitschies, W., Dressman, J., 2014a. PBPK models for the prediction of in vivo performance of oral dosage forms. Eur J Pharm Sci 57, 300–321. doi:10.1016/j.ejps.2013.09.008
- Kostewicz, E.S., Abrahamsson, B., Brewster, M., Brouwers, J., Butler, J., Carlert, S., Dickinson, P.A., Dressman, J., Holm, R., Klein, S., Mann, J., McAllister, M., Minekus, M., Muenster, U., Müllertz, A., Verwei, M., Vertzoni, M., Weitschies, W., Augustijns, P., 2014b. In vitro models for the prediction of in vivo performance of oral dosage forms. Eur J Pharm Sci 57, 342–366. doi:10.1016/j.ejps.2013.08.024
- Koziolek, M., Grimm, M., Garbacz, G., Kühn, J.-P., Weitschies, W., 2014. Intragastric volume changes after intake of a high-caloric, high-fat standard breakfast in healthy human subjects investigated by MRI. Mol. Pharm. 11, 1632–1639. doi:10.1021/mp500022u
- Koziolek, M., Schneider, F., Grimm, M., Modeβ, C., Seekamp, A., Roustom, T., Siegmund, W., Weitschies, W., 2015. Intragastric pH and pressure profiles after intake of the high-caloric, high-fat meal as used for food effect studies. J Control Release 220, 71–78. doi:10.1016/j.jconrel.2015.10.022
- Koziolek, M., Schneider, F., Grimm, M., Modeβ, C., Seekamp, A., Roustom, T., Siegmund, W., Weitschies, W., 2015. Intragastric pH and pressure profiles after intake of the high-caloric, high-fat meal as used for food effect studies. J Control Release 220, 71–78. doi:10.1016/j.jconrel.2015.10.022
- Kunz, P., Crelier, G.R., Schwizer, W., Borovicka, J., Kreiss, C., Fried, M., Boesiger, P., 1998. Gastric emptying and motility: assessment with MR imaging--preliminary observations. Radiology 207, 33–40. doi:10.1148/radiology.207.1.9530296

Lennernäs, H., 1998. Human intestinal permeability. J. Pharm. Sci. 87, 403–410. doi:10.1021/js970332a

- Lennernäs, H., Aarons, L., Augustijns, P., Beato, S., Bolger, M., Box, K., Brewster, M., Butler, J.,
 Dressman, J., Holm, R., Julia Frank, K., Kendall, R., Langguth, P., Sydor, J., Lindahl, A.,
 McAllister, M., Muenster, U., Müllertz, A., Ojala, K., Pepin, X., Reppas, C., Rostami-Hodjegan,
 A., Verwei, M., Weitschies, W., Wilson, C., Karlsson, C., Abrahamsson, B., 2014. Oral
 biopharmaceutics tools time for a new initiative an introduction to the IMI project OrBiTo.
 Eur J Pharm Sci 57, 292–299. doi:10.1016/j.ejps.2013.10.012
- Lindahl, A., Ungell, A.L., Knutson, L., Lennernäs, H., 1997. Characterization of fluids from the stomach and proximal jejunum in men and women. Pharm. Res. 14, 497–502.
- Lopes, C.M., Bettencourt, C., Rossi, A., Buttini, F., Barata, P., 2016. Overview on gastroretentive drug delivery systems for improving drug bioavailability. International Journal of Pharmaceutics 510, 144–158. doi:10.1016/j.ijpharm.2016.05.016
- Maes, B.D., Hiele, M.I., Geypens, B.J., Rutgeerts, P.J., Ghoos, Y.F., Vantrappen, G., 1994. Pharmacological modulation of gastric emptying rate of solids as measured by the carbon labelled octanoic acid breath test: influence of erythromycin and propantheline. Gut 35, 333–337.
- Marathe, P.H., Wen, Y., Norton, J., Greene, D.S., Barbhaiya, R.H., Wilding, I.R., 2000. Effect of altered gastric emptying and gastrointestinal motility on metformin absorption. Br J Clin Pharmacol 50, 325–332.
- Marciani, L., Garsed, K.C., Hoad, C.L., Fields, A., Fordham, I., Pritchard, S.E., Placidi, E., Murray, K., Chaddock, G., Costigan, C., Lam, C., Jalanka-Tuovinen, J., De Vos, W.M., Gowland, P.A., Spiller, R.C., 2014. Stimulation of colonic motility by oral PEG electrolyte bowel preparation assessed by MRI: comparison of split vs single dose. Neurogastroenterol. Motil. 26, 1426– 1436. doi:10.1111/nmo.12403
- Marciani, L., Little, S.L., Snee, J., Coleman, N.S., Tyler, D.J., Sykes, J., Jolliffe, I.G., Dettmar, P.W., Spiller, R.C., Gowland, P.A., 2002. Echo-planar magnetic resonance imaging of Gaviscon alginate rafts in-vivo. J. Pharm. Pharmacol. 54, 1351–1356. doi:10.1211/002235702760345428
- Marciani, L., Young, P., Wright, J., Moore, R., Coleman, N., Gowland, P.A., Spiller, R.C., 2001. Antral motility measurements by magnetic resonance imaging. Neurogastroenterol. Motil. 13, 511–518.
- Markopoulos, C., Vertzoni, M., Symillides, M., Kesisoglou, F., Reppas, C., 2015. Two-Stage Single-Compartment Models to Evaluate Dissolution in the Lower Intestine. J Pharm Sci 104, 2986– 2997. doi:10.1002/jps.24485
- Maurer, A., 2008. Consensus report on gastric emptying: what's needed to prevent tarnishing a gold standard? J. Nucl. Med. 49, 339. doi:10.2967/jnumed.106.037507
- Maurer, A.H., 2015. Gastrointestinal Motility, Part 1: Esophageal Transit and Gastric Emptying. J. Nucl. Med. 56, 1229–1238. doi:10.2967/jnumed.112.114314
- McConnell, E.L., Short, M.D., Basit, A.W., 2008. An in vivo comparison of intestinal pH and bacteria as physiological trigger mechanisms for colonic targeting in man. J Control Release 130, 154– 160. doi:10.1016/j.jconrel.2008.05.022
- Menys, A., Hamy, V., Makanyanga, J., Hoad, C., Gowland, P., Odille, F., Taylor, S.A., Atkinson, D., 2014. Dual registration of abdominal motion for motility assessment in free-breathing data sets acquired using dynamic MRI. Phys Med Biol 59, 4603–4619. doi:10.1088/0031-9155/59/16/4603
- Metcalf, A.M., Phillips, S.F., Zinsmeister, A.R., MacCarty, R.L., Beart, R.W., Wolff, B.G., 1987. Simplified assessment of segmental colonic transit. Gastroenterology 92, 40–47.
- Mudie, D.M., Murray, K., Hoad, C.L., Pritchard, S.E., Garnett, M.C., Amidon, G.L., Gowland, P.A., Spiller, R.C., Amidon, G.E., Marciani, L., 2014. Quantification of gastrointestinal liquid volumes and distribution following a 240 mL dose of water in the fasted state. Mol. Pharm. 11, 3039–3047. doi:10.1021/mp500210c

- Müllertz, A., Fatouros, D.G., Smith, J.R., Vertzoni, M., Reppas, C., 2012. Insights into intermediate phases of human intestinal fluids visualized by atomic force microscopy and cryo-transmission electron microscopy ex vivo. Mol. Pharm. 9, 237–247. doi:10.1021/mp200286x
- Munoz, F., Alici, G., Li, W., 2014. A review of drug delivery systems for capsule endoscopy. Adv. Drug Deliv. Rev. 71, 77–85. doi:10.1016/j.addr.2013.12.007
- Munoz, F., Alici, G., Li, W., 2014. A review of drug delivery systems for capsule endoscopy. Adv Drug Deliv Rev 71, 77–85. doi:10.1016/j.addr.2013.12.007
- Myers, A.G., Bradshaw, L.A., McDowell, J.G., Richards, W.O., 2002. Magnetic vector analysis of gastrointestinal electrical control activity, in: Engineering in Medicine and Biology, 2002. 24th Annual Conference and the Annual Fall Meeting of the Biomedical Engineering Society EMBS/BMES Conference, 2002. Proceedings of the Second Joint. Presented at the Engineering in Medicine and Biology, 2002. 24th Annual Conference and the Annual Fall Meeting of the Biomedical Engineering Society EMBS/BMES Conference, 2002. Proceedings of the Second Joint, pp. 1728–1729 vol.2. doi:10.1109/IEMBS.2002.1106623
- Oberle, R.L., Amidon, G.L., 1987. The influence of variable gastric emptying and intestinal transit rates on the plasma level curve of cimetidine; an explanation for the double peak phenomenon. J Pharmacokinet Biopharm 15, 529–544.
- Odille, F., Menys, A., Ahmed, A., Punwani, S., Taylor, S.A., Atkinson, D., 2012. Quantitative assessment of small bowel motility by nonrigid registration of dynamic MR images. Magn Reson Med 68, 783–793. doi:10.1002/mrm.23298
- O'Grady, G., Du, P., Cheng, L.K., Egbuji, J.U., Lammers, W.J.E.P., Windsor, J.A., Pullan, A.J., 2010. Origin and propagation of human gastric slow-wave activity defined by high-resolution mapping. American Journal of Physiology - Gastrointestinal and Liver Physiology 299, G585– G592. doi:10.1152/ajpgi.00125.2010
- Patak, M.A., Froehlich, J.M., von Weymarn, C., Ritz, M.A., Zollikofer, C.L., Wentz, K., 2001. Noninvasive distension of the small bowel for magnetic-resonance imaging. Lancet 358, 987–988.
- Pedersen, P.B., Bar-Shalom, D., Baldursdottir, S., Vilmann, P., Müllertz, A., 2014. Feasibility of capsule endoscopy for direct imaging of drug delivery systems in the fasted upper-gastrointestinal tract. Pharm Res 31, 2044–53. doi:10.1007/s11095-014-1306-6
- Perri, F., Pastore, M.R., Annese, V., 2005. 13C-octanoic acid breath test for measuring gastric emptying of solids. Eur Rev Med Pharmacol Sci 9, 3–8.
- Petri, N., Tannergren, C., Holst, B., Mellon, F.A., Bao, Y., Plumb, G.W., Bacon, J., O'Leary, K.A., Kroon, P.A., Knutson, L., Forsell, P., Eriksson, T., Lennernas, H., Williamson, G., 2003.
 Absorption/metabolism of sulforaphane and quercetin, and regulation of phase II enzymes, in human jejunum in vivo. Drug Metab. Dispos. 31, 805–813.
- Phillips, S.F., 1984. Functions of the large bowel: an overview. Scand. J. Gastroenterol. Suppl. 93, 1– 12.
- Pišlar, M., Brelih, H., Mrhar, A., Bogataj, M., 2015. Analysis of small intestinal transit and colon arrival times of non-disintegrating tablets administered in the fasted state. Eur J Pharm Sci 75, 131– 141. doi:10.1016/j.ejps.2015.03.001
- Pomerri, F., Zandonà, M., Barusco, A., Razzoli, M., Muzzio, P.C., 2008. Comparison of fluoroscopy and radiography in the assessment of colonic transit time using radio-opaque markers. Clinical Radiology 63, 1019–1025. doi:10.1016/j.crad.2008.05.002
- Psachoulias, D., Vertzoni, M., Goumas, K., Kalioras, V., Beato, S., Butler, J., Reppas, C., 2011. Precipitation in and supersaturation of contents of the upper small intestine after administration of two weak bases to fasted adults. Pharm. Res. 28, 3145–3158. doi:10.1007/s11095-011-0506-6
- Rao, S.S., Sadeghi, P., Beaty, J., Kavlock, R., Ackerson, K., 2001. Ambulatory 24-h colonic manometry in healthy humans. Am. J. Physiol. Gastrointest. Liver Physiol. 280, G629-639.
- Reppas, C., Karatza, E., Goumas, C., Markopoulos, C., Vertzoni, M., 2015. Characterization of Contents of Distal Ileum and Cecum to Which Drugs/Drug Products are Exposed During

Bioavailability/Bioequivalence Studies in Healthy Adults. Pharm. Res. 32, 3338–3349. doi:10.1007/s11095-015-1710-6

- Riethorst, D., Baatsen, P., Remijn, C., Mitra, A., Tack, J., Brouwers, J., Augustijns, P., 2016. An In-Depth View into Human Intestinal Fluid Colloids: Intersubject Variability in Relation to Composition. Mol. Pharm. 13, 3484–3493. doi:10.1021/acs.molpharmaceut.6b00496
- Riethorst, D., Mols, R., Duchateau, G., Tack, J., Brouwers, J., Augustijns, P., 2015. Characterization of Human Duodenal Fluids in Fasted and Fed State Conditions. J Pharm Sci. doi:10.1002/jps.24603
- Sanaka, M., Koike, Y., Yamamoto, T., Mineshita, S., Yamaoka, S., Hirama, S., Tanaka, H., Kuyama, Y., Yamanaka, M., 1997. A reliable and convenient parameter of the rate of paracetamol absorption to measure gastric emptying rate of liquids. Int J Clin Pharmacol Ther 35, 509– 513.
- Sanaka, M., Kuyama, Y., Yamanaka, M., 1998. Guide for judicious use of the paracetamol absorption technique in a study of gastric emptying rate of liquids. J. Gastroenterol. 33, 785–791.
- Sarna, S.K., Otterson, M.F., 1989. Small intestinal physiology and pathophysiology. Gastroenterol. Clin. North Am. 18, 375–404.
- Sarnelli, G., Caenepeel, P., Geypens, B., Janssens, J., Tack, J., 2003. Symptoms associated with impaired gastric emptying of solids and liquids in functional dyspepsia. Am. J. Gastroenterol. 98, 783–788. doi:10.1111/j.1572-0241.2003.07389.x
- Schiller, C., Fröhlich, C.-P., Giessmann, T., Siegmund, W., Mönnikes, H., Hosten, N., Weitschies, W.,
 2005. Intestinal fluid volumes and transit of dosage forms as assessed by magnetic resonance imaging. Aliment. Pharmacol. Ther. 22, 971–979. doi:10.1111/j.1365-2036.2005.02683.x
- Schwizer, W., Fraser, R., Borovicka, J., Asal, K., Crelier, G., Kunz, P., Boesiger, P., Fried, M., 1996.
 Measurement of proximal and distal gastric motility with magnetic resonance imaging. Am. J.
 Physiol. 271, G217-222.
- Schwizer, W., Fraser, R., Borovicka, J., Crelier, G., Boesiger, P., Fried, M., 1994. Measurement of gastric emptying and gastric motility by magnetic resonance imaging (MRI). Dig. Dis. Sci. 39, 1015–103S.
- Schwizer, W., Maecke, H., Fried, M., 1992. Measurement of gastric emptying by magnetic resonance imaging in humans. Gastroenterology 103, 369–376.
- Sitti, M., Ceylan, H., Hu, W., Giltinan, J., Turan, M., Yim, S., Diller, E., 2015. Biomedical Applications of Untethered Mobile Milli/Microrobots. Proc IEEE 103, 217–230. doi:10.1109/JPROC.2014.2385105
- Söderlind, E., Abrahamsson, B., Erlandsson, F., Wanke, C., Iordanov, V., von Corswant, C., 2015. Validation of the IntelliCap® system as a tool to evaluate extended release profiles in human GI tract using metoprolol as model drug. J Control Release 217, 300–307. doi:10.1016/j.jconrel.2015.09.024
- Soenen, S., Rayner, C.K., Horowitz, M., Jones, K.L., 2015. Gastric Emptying in the Elderly. Clinics in Geriatric Medicine, Nutrition in Older Adults 31, 339–353. doi:10.1016/j.cger.2015.04.003
- Stanghellini, V., Tosetti, C., Paternico, A., Barbara, G., Morselli-Labate, A.M., Monetti, N., Marengo,
 M., Corinaldesi, R., 1996. Risk indicators of delayed gastric emptying of solids in patients with functional dyspepsia. Gastroenterology 110, 1036–1042.
- Stappaerts, J., Geboers, S., Snoeys, J., Brouwers, J., Tack, J., Annaert, P., Augustijns, P., 2015. Rapid conversion of the ester prodrug abiraterone acetate results in intestinal supersaturation and enhanced absorption of abiraterone: in vitro, rat in situ and human in vivo studies. Eur J Pharm Biopharm 90, 1–7. doi:10.1016/j.ejpb.2015.01.001
- Stehling, M.K., Evans, D.F., Lamont, G., Ordidge, R.J., Howseman, A.M., Chapman, B., Coxon, R., Mansfield, P., Hardcastle, J.D., Coupland, R.E., 1989. Gastrointestinal tract: dynamic MR studies with echo-planar imaging. Radiology 171, 41–46. doi:10.1148/radiology.171.1.2928545

- Steingoetter, A., Weishaupt, D., Kunz, P., M\u00e4der, K., Lengsfeld, H., Thumshirn, M., Boesiger, P., Fried, M., Schwizer, W., 2003. Magnetic resonance imaging for the in vivo evaluation of gastric-retentive tablets. Pharm. Res. 20, 2001–2007.
- Sweis, R., Kaufman, E., Anggiansah, A., Wong, T., Dettmar, P., Fried, M., Schwizer, W., Avvari, R.K., Pal, A., Fox, M., 2013. Post-prandial reflux suppression by a raft-forming alginate (Gaviscon Advance) compared to a simple antacid documented by magnetic resonance imaging and pHimpedance monitoring: mechanistic assessment in healthy volunteers and randomised, controlled, double-blind study in reflux patients. Aliment. Pharmacol. Ther. 37, 1093–1102. doi:10.1111/apt.12318
- Talattof, A., Price, J.C., Amidon, G.L., 2016a. Gastrointestinal Motility Variation and Implications for Plasma Level Variation: Oral Drug Products. Mol. Pharm. 13, 557–567. doi:10.1021/acs.molpharmaceut.5b00774
- Talattof, A., Price, J.C., Amidon, G.L., 2016b. Gastrointestinal Motility Variation and Implications for Plasma Level Variation: Oral Drug Products. Mol. Pharm. doi:10.1021/acs.molpharmaceut.5b00774
- Talley, N.J., Locke, G.R., Lahr, B.D., Zinsmeister, A.R., Tougas, G., Ligozio, G., Rojavin, M.A., Tack, J., 2006. Functional dyspepsia, delayed gastric emptying, and impaired quality of life. Gut 55, 933–939. doi:10.1136/gut.2005.078634
- Tannergren, C., Knutson, T., Knutson, L., Lennernäs, H., 2003. The effect of ketoconazole on the in vivo intestinal permeability of fexofenadine using a regional perfusion technique. Br J Clin Pharmacol 55, 182–190.
- U.S. Food and Drug Administration, 2016. Drug Innovation New Molecular Entity and New Therapeutic Biological Product Approvals for 2015 [WWW Document]. URL http://www.fda.gov/Drugs/DevelopmentApprovalProcess/DrugInnovation/ucm430302.htm (accessed 1.12.16).
- Van Den Abeele, J., Brouwers, J., Mattheus, R., Tack, J., Augustijns, P., 2015. Gastrointestinal Behavior of Weakly Acidic BCS Class II Drugs in Man - Case Study Diclofenac Potassium. J Pharm Sci. doi:10.1002/jps.24647
- Van Den Abeele, J., Brouwers, J., Tack, J., Augustijns, P., 2016a. Exploring the link between gastric motility and intragastric drug distribution in man. European Journal of Pharmaceutics and Biopharmaceutics.
- Van Den Abeele, J., Rubbens, J., Brouwers, J., Augustijns, P., 2016b. The dynamic gastric environment and its impact on drug and formulation behaviour. Eur J Pharm Sci 96, 207–231. doi:10.1016/j.ejps.2016.08.060
- van der Schaar, P.J., Dijksman, J.F., Broekhuizen-de Gast, H., Shimizu, J., van Lelyveld, N., Zou, H., Iordanov, V., Wanke, C., Siersema, P.D., 2013. A novel ingestible electronic drug delivery and monitoring device. Gastrointest Endosc 78, 520–528. doi:http://dx.doi.org/10.1016/j.gie.2013.03.170
- van der Sijp, J.R., Kamm, M.A., Nightingale, J.M., Britton, K.E., Mather, S.J., Morris, G.P., Akkermans, L.M., Lennard-Jones, J.E., 1993. Radioisotope determination of regional colonic transit in severe constipation: comparison with radio opaque markers. Gut 34, 402–408.
- Vertzoni, M., Diakidou, A., Chatzilias, M., Söderlind, E., Abrahamsson, B., Dressman, J.B., Reppas, C., 2010a. Biorelevant media to simulate fluids in the ascending colon of humans and their usefulness in predicting intracolonic drug solubility. Pharm. Res. 27, 2187–2196. doi:10.1007/s11095-010-0223-6
- Vertzoni, M., Diakidou, A., Chatzilias, M., Söderlind, E., Abrahamsson, B., Dressman, J.B., Reppas, C., 2010b. Biorelevant media to simulate fluids in the ascending colon of humans and their usefulness in predicting intracolonic drug solubility. Pharm. Res. 27, 2187–2196. doi:10.1007/s11095-010-0223-6
- Vertzoni, M., Pastelli, E., Psachoulias, D., Kalantzi, L., Reppas, C., 2007. Estimation of intragastric solubility of drugs: in what medium? Pharm. Res. 24, 909–917. doi:10.1007/s11095-006-9209-9

- Wang, Y.T., Mohammed, S.D., Farmer, A.D., Wang, D., Zarate, N., Hobson, A.R., Hellström, P.M., Semler, J.R., Kuo, B., Rao, S.S., Hasler, W.L., Camilleri, M., Scott, S.M., 2015. Regional gastrointestinal transit and pH studied in 215 healthy volunteers using the wireless motility capsule: influence of age, gender, study country and testing protocol. Aliment. Pharmacol. Ther. 42, 761–772. doi:10.1111/apt.13329
- Weitschies, W., Blume, H., Mönnikes, H., 2010. Magnetic marker monitoring: high resolution realtime tracking of oral solid dosage forms in the gastrointestinal tract. Eur J Pharm Biopharm 74, 93–101. doi:10.1016/j.ejpb.2009.07.007
- Wilding, I., 2002. Bioequivalence testing for locally acting gastrointestinal products: what role for gamma scintigraphy? J Clin Pharmacol 42, 1200–1210.
- Willems, M., Quartero, A.O., Numans, M.E., 2001. How useful is paracetamol absorption as a marker of gastric emptying? A systematic literature study. Dig. Dis. Sci. 46, 2256–2262.
- Wilson, C.G., 2010. The transit of dosage forms through the colon. International Journal of Pharmaceutics 395, 17–25. doi:10.1016/j.ijpharm.2010.04.044
- Wilson, C.G., Clarke, C.P., Starkey, Y.Y.L., Clarke, G.D., 2011. Comparison of a novel fast-dissolving acetaminophen tablet formulation (FD-APAP) and standard acetaminophen tablets using gamma scintigraphy and pharmacokinetic studies. Drug Dev Ind Pharm 37, 747–753. doi:10.3109/03639045.2010.538058
- Wilson, C.G., Washington, N., Greaves, J.L., Kamali, F., Rees, J.A., Sempik, A.K., Lampard, J.F., 1989.
 Bimodal release of drug in a sustained release Ibuprofen formulation: A scintigraphic and pharmacokinetic open study in healthy volunteers under different conditions of food intake. Int J Pharmaceut 50, 155–161.
- Wright, J., Evans, D., Gowland, P., Mansfield, P., 1999. Validation of antroduodenal motility measurements made by echo-planar magnetic resonance imaging. Neurogastroenterol. Motil. 11, 19–25.
- Wuyts, B., Riethorst, D., Brouwers, J., Tack, J., Annaert, P., Augustijns, P., 2015a. Evaluation of fasted state human intestinal fluid as apical solvent system in the Caco-2 absorption model and comparison with FaSSIF. Eur J Pharm Sci 67, 126–135. doi:10.1016/j.ejps.2014.11.010
- Wuyts, B., Riethorst, D., Brouwers, J., Tack, J., Annaert, P., Augustijns, P., 2015b. Evaluation of fasted and fed state simulated and human intestinal fluids as solvent system in the Ussing chambers model to explore food effects on intestinal permeability. Int J Pharm 478, 736–744. doi:10.1016/j.ijpharm.2014.12.021