

Gladman, John R.F. and Buckell, John and Young, John and Smith, Andrew and Hulme, Claire and Saggu, Satti and Godfrey, Mary and Enderby, Pam and Teale, Elizabeth and Longo, Roberto and Gannon, Brenda and Holditch, Claire and Eardley, Heather and Tucker, Helen (2016) Understanding the models of community hospital rehabilitation activity (MoCHA): a mixed method study. BMJ Open . ISSN 2044-6055 (In Press)

Access from the University of Nottingham repository:

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: http://eprints.nottingham.ac.uk/end user agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk



This is an author produced version of *Understanding the Models of Community Hospital rehabilitation Activity (MoCHA): a mixed method study.*

White Rose Research Online URL for this paper: http://eprints.whiterose.ac.uk/103835/

Article:

Gladman, J, Buckell, J, Young, J et al. (11 more authors) (2016) Understanding the Models of Community Hospital rehabilitation Activity (MoCHA): a mixed method study. BMJ Open. ISSN 2044-6055 (In Press)

© 2016, The Authors. Published by BMJ Publishing Group. This is an author produced version of a paper accepted for publication in BMJ Open. Uploaded in accordance with the publisher's self-archiving policy



Understanding the Models of Community Hospital rehabilitation Activity (MoCHA): a mixed method study

Corresponding author: John Gladman. Postal address: Room B111, Division of Rehabilitation and Ageing, B Floor Medical School, Queen's Medical Centre, Nottingham NG7 2UH UK. Affiliation: Division of Rehabilitation and Ageing, School of Medicine, University of Nottingham, Nottingham, UK. Email: john.gladman@nottingham.ac.uk. Telephone +44 115 823 0242. Fax +44 115 823 0239

Name(s) of co-author(s) with complete affiliations (in addition to institutions where appropriate), institutions, city and country:

John Buckell, School of Public Health, Yale University, New Haven USA

John Young, Academic Unit of Elderly Care and Rehabilitation, University of Leeds, Leeds UK

Andrew Smith, Academic Unit of Health Economics, University of Leeds, Leeds UK

Clare Hulme, Academic Unit of Health Economics, University of Leeds, Leeds UK

Satti Saggu, Academic Unit of Elderly Care and Rehabilitation, University of Leeds, Leeds UK

Mary Godfrey, Academic Unit of Elderly Care and Rehabilitation, University of Leeds, Leeds UK

Pam Enderby, ScHARR, University of Sheffield, Sheffield, UK

Elizabeth Teale, Academic Unit of Elderly Care and Rehabilitation, University of Leeds, Leeds UK

Roberto Longo, Academic Unit of Health Economics, University of Leeds, Leeds UK

Brenda Gannon, Academic Unit of Health Economics, University of Leeds, Leeds UK

Claire Holditch, NHS Benchmarking Network, Manchester, UK

Heather Eardley, The Patients Association, London, UK

Helen Tucker, The Community Hospitals Association, Ilminster, UK

Keywords (up to five):

- 1. Community hospitals
- 2. Mixed method research
- 3. Health economics
- 4. Geriatric medicine
- 5. Health services administration and management

Strengths and limitations: 88 words

Abstract: 296 words.

Main text word count, excluding Figure 1: 4705 words

Authors, funding and competing interests statements: 48 words

STRENGHTHS AND LIMITATIONS OF THIS STUDY

Strengths

This mixed methods study will use economic analyses of large, existing, up-to-date
databases together with new empirical observational data to provide evidence of
value to commissioners and providers about how to optimize the performance in
community hospitals which would not be possible using other methods

Weaknesses

- The nature of the economic findings on relative performance will be limited by the size and content of the existing databases which are to be analysed
- The findings may not generalise beyond the UK setting

ABSTRACT

Introduction

To understand the variation in performance between community hospitals, our objectives are: to measure the relative performance (cost efficiency) of rehabilitation services in community hospitals; to identify the characteristics of community hospital rehabilitation that optimise performance; to investigate the current impact of community hospital in-patient rehabilitation for older people on secondary care and the potential impact if community hospital rehabilitation was optimised to best practice nationally; to examine the relationship between the configuration of intermediate care and secondary care bed use; and to develop toolkits for commissioners and community hospital providers to optimise performance.

Methods and analysis

Four linked studies will be performed.

Study 1: Cost efficiency modelling will apply econometric techniques to datasets from the NHS Benchmarking Network surveys of community hospital and intermediate care. This will identify community hospitals' performance and estimate the gap between high and low performers. Analyses will determine the potential impact if the performance of all community hospitals nationally was optimised to best performance, and examine the association between community hospital configuration and secondary care bed use.

Study 2: A national community hospital survey gathering detailed cost data and efficiency variables will be performed.

Study 3: In-depth case studies of three community hospitals, two high and one low performing, will be undertaken. Case studies will gather routine hospital and local health economy data. Ward culture will be surveyed. Content and delivery of treatment will be observed. Patients and staff will be interviewed.

Study 4: Co-designed web-based quality improvement toolkits for commissioners and providers will be developed, including indicators of performance and the gap between local and best community hospitals performance.

Ethics and dissemination

Publications will be in peer reviewed journals, reports will be distributed through stakeholder organisations.

Ethical approval was obtained from the Bradford Research Ethics committee (reference: 15/YH/0062).

INTRODUCTION

There are 10 million people in the UK who are over 65 years old. Between 2010 and 2035, the proportion of the population aged 65 and over is expected to rise from around 16% to around 23%; and the proportion of the population aged 85 and over from around 2% to around 5%.[1]. A growing older population will be accompanied by an increasing impact on health and social care services. Decision makers across the NHS and local government are attempting to reconfigure services in response.

Rehabilitation lies at the heart of best practice for older people. Rising emergency admissions and reductions in acute hospital beds, leading to shorter lengths of stay, limits the scope for rehabilitation in general hospitals.[2] Intermediate care provides short-term, community based support and rehabilitation, delivered using a range of service models including community hospitals, care homes and home-based. The UK National Audit of Intermediate Care [3] has implied an under-provision of intermediate care services and insufficient whole system impact. There is uncertainty both about what type of intermediate care works and what configuration of intermediate care services offers a comprehensive repertoire of provision.

Community hospitals are part of this uncertainty about intermediate care. Community hospitals are local hospitals providing a range of health care facilities and resources that usually does not include emergency or acute medical in-patient care, intensive care or major surgery, but very commonly includes in-patient rehabilitation for older people. Community hospitals are long established in the UK and internationally.[4-7]. High quality evidence supports community hospitals as effective bed-based rehabilitation services for older people when compared to general hospitals. [8,9,10] The care experience reported by patients in community hospital wards is favoured over that provided in general hospital wards.[8] As there are nearly 300 community hospitals in the UK,[11] it is important that community hospitals nationally are organised and configured to deliver these superior outcomes.

Three key findings are apparent from two national surveys, the NHS Benchmarking Network Community Hospitals Project and the National Audit of Intermediate Care which between them provide information on 180 community hospitals, approximately two-thirds of UK community hospitals. First, these studies confirm that a core function of the contemporary community hospital is rehabilitation, largely for older people: 97% of community hospitals provide rehabilitation. Second, community hospital wards are extremely variable. Examples of variability include: bed provision per 100,000 weighted population (range <10 to 70);

clinical leadership (50% nurse led; 50% consultant led); average length of stay (11 to 58 days); cost per admission (£3,700 to £17,500); and cost per day (£140 to £450). Third, there is potential for improvement. If a community hospital with 20 beds and 90% occupancy with a length of stay at the 75% quartile (31 days) improved to the 25% quartile (21 days), it would be able to treat approximately 100 more patients per year - a 48% increase. Alternatively, if a 20 bed community hospital with a cost per occupied bed day at the 75% quartile (£200 per day) improved to the 25% quartile (£110 per day) the annual savings would be approximately £650,000.

The reasons behind the variations, and therefore the steps needed to improve, are speculative as no detailed study has been designed and conducted systematically to investigate this issue. There is also a paucity of information available to service planners about the staffing levels, comparative outcomes and efficiencies of community hospitals in relation to alternative forms of community rehabilitation services. The Models of Community Hospital Activity (MoCHA) study (01/04/2014 to: 31/03/2017) will address these deficiencies in the evidence base. Its objectives are:

- 1. To measure the relative performance (cost efficiency) of rehabilitation services in community hospitals
- 2. To identify the characteristics of community hospital in-patient rehabilitation for older people that optimise performance
- 3. To investigate the current impact of community hospital in-patient care for older people on secondary care and the potential impact if community hospital rehabilitation was optimised to best practice nationally
- 4. To examine the relationship between the configuration of intermediate care and secondary care bed use
- 5. To develop toolkits for commissioners and community hospital providers to optimise performance.

METHODS AND ANALYSIS

A mixed methods approach will be taken, combining established quantitative and qualitative

techniques, consisting of four interlinked studies. This approach employs the use of existing

data alongside complementary prospective data to produce insights that could not be

achieved by the study of the components alone, and represents an efficient and effective

research design.

Study 1. An analysis of cost efficiency amongst community hospitals

Study 2. A national survey of community hospitals

Study 3. A multi- method comparative case study of purposively selected community hospital

wards providing rehabilitation to older people

Study 4. The development of web-based quality improvement tools for community hospitals

and commissioners.

Figure 1, the Community Hospital Study Map, illustrates the relationship of the four studies

to the five research objectives.

<insert Figure 1 here>

Legend: Figure 1. The Community Hospital Study Map

Study 1: an analysis of cost efficiency amongst community hospitals

Objective1 - to measure the relative performance (cost efficiency) of rehabilitation

services in community hospitals

Figure 2 illustrates the various components of Study1.

<insert Figure 2 here>

Legend: Figure 2. The components of the cost efficiency study (Study 1)

Previous economic evidence on community hospitals derives from a cost-effectiveness

analysis embedded within a multi-centre randomised controlled trial.[9] The efficiency

analysis we will adopt in this study is different in as far as it provides a framework to assess

6

the extent to which resources that have already been allocated to health services are optimally deployed. A service or process is said to be productively efficient if it produces a given output at the least possible cost.[12] The aim of this analysis is to identify the performance range for community hospital wards and to explain variations in costs for rehabilitation of older people.

In broad terms, the inputs required for the efficiency analysis include costs (operating costs), output (e.g., occupied bed days), and quality (e.g. the frequency of multidisciplinary team (MDT) meetings) data, and the outputs are community hospital-specific efficiency scores; an assessment of the scope for efficiency gains across the sector; and information for how costs vary with important variables such as scale and quality. The data sources used in this study are from the National Audit of Intermediate Care and the Community Hospitals NHS Benchmarking Network datasets, and Hospital Episode Statistics (HES) as described in Table 1. Both the National Audit of Intermediate Care and NHS Benchmarking Network Community Hospitals data are collected directly from commissioner and provider organisations via an online data collection tool. Data are held by the NHS Benchmarking Network in a SQL database. The NHS Benchmarking Network undertakes validation checks to ensure data quality. Data collection is annual. The data will be provided to the research project in CSV files in anonymised format. A sub-contract is in place with East London NHS Foundation Trust, the NHS Benchmarking Network's host organisation, for re-imbursement to the Network of the costs of the Community Hospitals data collection.

Data are available at the whole hospital level which includes the full range of services and at the ward level which is on rehabilitation services only. These services constitute a large proportion (circa 70%) of the total activity of community hospitals. Analysis at this level allows for greater comparability between community hospitals because services are very similar. Therefore more reliable estimates of efficiency are derived. To support this claim, we have analysed hospital level data in preliminary modelling, but the results were implausible. This was not the case when the ward level data was analysed. Ward level analysis seems more appropriate empirically. Also, it may be of greater use to managers who have a clearer idea as to the source of inefficiency.

Table 1 Data sources

Data	Data Source(s)			Features		Collection	
National Audit		Geriatrics	,		Commissioners (62) and providers (112) in Round 1, Round 2 has	,	from

Intermediate Care	Services; AGILE (Chartered physiotherapists working with older people); the Royal College of Physicians; the Royal College of Nursing; the Royal College of Speech & Language Therapists; The Patients Association; and NHS Benchmarking Network	higher participation (92 commissioners, provider numbers need validation); 370 IC services, both home- and bed-based; information on demography (age, gender, pre-admission accommodation, place of referral), level of required care, clinical outcomes, service outcomes, patient reported experience measure (PREM)	
Community Hospitals NHS Benchmarking Network (NHSBN) datasets	NHS Benchmarking Network	Opt-in scheme for NHSBN members; around 180 community hospitals; two years of data; information on workforce, activity, investment levels, organisational features, services provided and quality measures	Survey of community hospitals by NHSBN
Hospital Episode Statistics (HES)	Health and Social Care Information Centre (HSCIC)	Patient level data of hospital admissions; collected in episodes of care; range of episode- and patient-specific information available	Recorded at all secondary care providers

We will investigate the variation in efficiency across community hospitals wards using a range of well-established but analytically complex methods including corrected ordinary least squares (COLS) regression analysis, Stochastic Frontier Analysis (SFA) and Data Envelopment Analysis (DEA). Corrected ordinary least squares (COLS) regression models enable relationships between (in this case) costs and cost drivers to be estimated, thus revealing the extent to which changing cost drivers will impact on cost. SFA is a similar approach that likewise investigates relationships between costs and the cost drivers. However, in this model, the estimated relationship is interpreted as an efficiency frontier, and community hospital ward inefficiency is measured against that frontier. SFA is widely used in the academic literature [13] and is also used by some economic regulators in the UK and internationally.[14] SFA has been used recently in the NHS setting.[15, 16] DEA is a method which uses mathematical-programming techniques to produce an efficiency score for each unit analysed.[17] These techniques have been extensively applied to the health sector in general and to the hospital sector in particular.[18]

COLS and SFA are described as below.

c=f(y,w,q,z)+e(1)e=u+v(2)

Where.

- c costs: ward level operating costs; exclusive of capital
- y outputs: a measure of the services provided, e,g. occupied bed days, admissions, discharges (these are commonly used in health cost analyses [18]). For modelling we prefer occupied bed days; other measures are used for sensitivity analysis.
- w input prices: as we have operating costs, we have input prices for labour and materials.
- q quality: measures to capture the quality of services, e.g. physio/OT ratio (as recommended by our clinicians; a unique measure for community hospitals), frequency of staff meetings, whether the community hospital is engaged in research. We also apply measures to capture unobserved heterogeneity for any quality that is not captured by these variables or is imperfectly measured.
- z observable heterogeneity: differences between services for which we have data, including case mix, risk factors and hospital characteristics, e.g. service variables (as a proxy for case mix), admission criteria for the community hospital, length of stay (a proxy for risk factors), the age of the community hospital, inter alia. We also apply measures to capture unobserved heterogeneity for any differences between services that are not captured by these variables or are imperfectly measured.
- f(...) represents the cost function. This is the relationship between costs and drivers of costs. That is, the factors in the brackets in equation (1) explain differences in costs between community hospitals; these factors are removed from our measure of inefficiency.
- e error term or model residual the remainder of costs that are not explained by the cost function. In the COLS model, this is used to compute the inefficiency
- u inefficiency in the stochastic frontier model (the SFA model is an extension of COLS that allows the separation of inefficiency from random noise [19]).
- v random statistical noise (e.g. measurement error, untoward events).

It is important to deal with variations in costs for which data are not available (or variables that are imperfectly measured; for example, service quality, which is difficult to define and to measure). This is termed unobserved heterogeneity in economic jargon. It is possible to use econometric methods to do this in four broad ways, which have been used in health care settings [15, 16]:

- 1. The stochastic frontier model can accommodate random noise, which, in part, comprises unobserved heterogeneity
- 2. Using panel data, allowing for community hospital-specific effects to control for unobserved heterogeneity
- 3. The use of the Mundlak approach to dealing with unobserved heterogeneity
- 4. The latent class stochastic frontier model

The specific approach is an empirical issue. Statistical testing is used for model specification.

In sum, econometric methods allow us to: measure the relationship between costs and cost drivers (e.g. output), including measuring economies of scale; control for observed and unobserved differences between community hospitals; separate out random noise from efficiency estimates; and estimate the relative cost efficiency of rehabilitation services for older people provided in the community hospital.

We anticipate some missing data and propose to address this as follows. We first categorise our variables into groups according to economic theory: costs, outputs, input prices, environmental variables and quality.[13] We then make an assessment of the extent of missing data and decide on variables to be used based on initial statistical testing and consultation with the research group members. Next, we conduct modelling based on the base data set (i.e. unmodified data) and on an imputed data set as modified by a range of standard imputation methods.[20] We conduct appropriate sensitivity analysis and select our preferred model. We therefore set out the following procedure for dealing with missing data items:

- (i) Categorise variables according to economic taxonomy
- (ii) Analyse and summarise missing data
- (iii) Decide on final set of variables for modelling
- (iv) Run a range of models on base data (no imputation)
- (v) Run a range of models on imputed data
- (vi) Conduct statistical testing/sensitivity analysis
- (vii) Selection of preferred model

Objective 2 - To identify the characteristics of community hospital in-patient rehabilitation for older people that optimise performance

The analyses described for Objective 1 identify, after taking account of differences between hospitals as captured by the range of cost drivers, the community hospitals that perform best and provides an estimate for the gap between the higher performing community hospital wards and the others. Thus, "best" care in this health economic context is defined by describing the features of those hospitals that have the highest relative efficiency.

Objective 3 - to investigate the current impact of community hospital in-patient care for older people on secondary care and the potential impact if community hospital rehabilitation was optimised to best practice nationally

A multiple regression analysis will be used to investigate the relationship between secondary care utilisation, identified in the HES data, and relevant variables characterising the quantum and nature of community hospital care within the relevant Clinical Commissioning Group (CCG) area. We use the standard regression approach rather than frontier-based approaches as the focus is on how the explanatory variables impact on the dependent variable, secondary care utilisation, rather than on the efficiency of certain providers. The community hospital variables of interest are the number of community hospital beds used for the rehabilitation of older people and parameters of efficiency and quality of care as identified in Study 1.

Objective 4 - to examine the relationship between the configuration of intermediate care and secondary care bed use

We will adopt a whole system perspective in order to determine if there is an association between the configuration of intermediate care (community-based rehabilitation) services and secondary care utilisation by older people. This analysis will use the National Audit of Intermediate Care (NAIC) and Hospital Episode Statistics (HES) datasets. A standard regression model will be used to estimate the relationship between secondary care utilisation (sourced from HES) and key variables sourced from NAIC capturing the configuration of community based rehabilitation services (the capacity of intermediate care: community hospitals, home-based rehabilitation, care home rehabilitation, and reablement services). The way in which community based rehabilitation services interact with each other, and with the wider secondary care system, is complex and a key part of the project will be to determine how to capture this complexity into a set of measures for inclusion in the model.

Study 2: National survey of community hospitals

Objective 3 - to investigate the current impact of community hospital in-patient care for older people on secondary care and the potential impact if the community hospital rehabilitation was optimised to best practice nationally

Two existing data sets for this research programme, the NAIC and the NHS Benchmarking Network Community Hospital datasets, comprise only 2/3 of UK community hospitals. The purpose of this national survey of community hospitals is to draw on the findings of Study 1 and to use a brief survey instrument to obtain a complete description UK community hospital rehabilitation practice. The survey will focus on those variables most strongly associated with community hospital efficiency (for example, total occupied bed days, input prices, staff mix and bed occupancy), allowing inferences to be drawn about the scale of the work to optimise community hospital ward care nationally. The results will help Clinical Commissioning Groups to plan strategic changes to community rehabilitation services. The survey instrument will be posted with telephone reminders and will use existing or modified questions from the previous NHS Benchmarking Network community hospital surveys that align to the key performance features identified in the efficiency models in Study 1.

Participants to the survey will be recruited by the NHS Benchmarking Network through advertising the project to its 337 Network member organisations and an inventory of UK community hospitals recently updated by the Birmingham Community Hospital Study Group and the Community Hospitals Association. Ward staff supported by the audit departments of participating organisations will complete the survey, on-line – as is usual practice in NHS Benchmarking Network surveys, completing one survey form for each rehabilitation ward in the hospital. A second round using a shorter survey instrument including only the variables required to calculate the cost efficiency model will be directed at the non-responders.

The preferred performance (cost efficiency) model will be applied to the results of study 2 which, being more up to date and complete than the historical data used to produce the models, will enable the most complete and accurate estimate of the performance of community hospitals and enable the best estimate of the likely consequence across the country of optimising the performance of community hospitals (objective 3).

Study 3: In-depth case studies

Objective 2 - to identify the characteristics of community hospital rehabilitation that optimise performance

Using a comparative case study design,[21-23] we will draw on multiple methods and perspectives better to understand current in-patient community hospital care and contribute

to an understanding of what best cost-efficient performance looks like in practice. This method was previously employed successfully in a study of intermediate care.[24]

Drawing on Study 1, we will purposively select 3 case studies of community hospital wards providing rehabilitation to older people. We will identify two high performers and one low performer based on relative efficiency from Study 1, to enable in-depth exploration of how cases that differ in respect of performance vary in terms of care delivery and user-centred outcomes.

Within case studies, we will employ maximum variation sampling [25] of staff to ensure inclusion of staff members from different disciplines and seniority levels. We will also purposively select a sample of older patients and their caregiver (5/6 patient/caregiver dyads in each case study), and follow them from admission to discharge via observation, conversations/interviews and medical records. Selection will be based on typical and critical case sampling strategies - patients that are typical of those on community hospital wards and critical in that they pose particularly difficult challenges for delivery, such as cognitive impairment.

We will employ multiple quantitative and qualitative data collection methods to build up a picture of each community hospital case, the design and execution of which will incorporate processes for ensuring rigor and quality.[26] This will include ward patient profiles from routinely collected aggregate data and an assessment of care culture (shared philosophy, leadership, mutual support and team working) using a structured questionnaire with staff.[27] We will observe practice, including the work of therapy, nursing and medical staff, interactions between professionals and between professionals and patients, decision-making and discharge planning (30-40 hours in each case study). Detailed descriptions of settings, events, interactions and activities will be maintained in field notes. Emerging categories about the data will be tested through more focused observation.

Qualitative interviews with staff (6-7 in each ward) will include how the work of rehabilitation and care for older people is understood, what makes it work and for whom, the resources available and the professional, organisational, cultural and other contextual factors affecting delivery from their different perspectives. Through discussion of anonymised cases we will explore the kinds of patients perceived as best suited to the service and those most likely to benefit.

Data will be collected about the experience of care from a patient and caregiver perspective via observation of care delivery on the ward and of multi-disciplinary team and family meetings and contemporaneous conversations with patients. This approach is particularly valuable for example for patients with cognitive problems or dementia, since data are collected in real time it does not require either verbal facility or ability to recall.

We will interview patients selected for observation and their caregivers, in each case study site shortly before discharge from the community hospital to reduce problems of recall. As data analysis will proceed simultaneously with data collection, we will leave open the possibility of undertaking a small number of additional interviews to pursue promising lines of enquiry not anticipated in advance. The interviews will assess if and how the care they received facilitated recovery from their perspective.

Data analysis will be in six stages.

Stage 1: With the ward as the unit of analysis, we will construct a narrative description of the structure (bed base, staffing, patient profile), activities (throughput) and care culture. Quantitative data from the hospital admission systems (age, sex, reason for admission, type of residence, length of inpatient stay, discharge destination, hospital mortality) will be analysed to provide descriptive statistics, while qualitative staff interview data and observational data will be drawn upon to examine how beliefs and values are translated into practice in each organisation.

Stage 2: Employing grounded theory analytical techniques,[28] such as simultaneous data collection and analysis, constant comparison and search for negative cases, we will examine the process of delivery of rehabilitation care to patients with different characteristics and needs in the real life context of the ward environment by drawing on the observational data and conversations with staff and patients. Such grounded theory techniques provide a robust approach to analysis and direct attention on conditions, processes and consequences pertinent to this study.

Stage 3: With the patient as the unit of analysis, we will similarly employ grounded theory analytical techniques to compare and contrast experiences and outcomes across patients similar to, and different from, each other in terms of the nature of the event that precipitated admission and their prior characteristics (such as cognitive impairment). We will refer to the recovery trajectories developed in research on intermediate care [29] which take account of

the diversity of patient characteristics and what recovery means from the perspective of the patient.

Stage 4: Using analytic induction, we will compare and contrast cases in their structure, culture and process of delivery drawing on the narrative descriptions of each case from Stages 1 & 2. We will specifically focus on features of structure, culture and delivery processes that differentiate between high and low performing cases.

Stage 5: Involves synthesising and simplifying the conditions and delivery processes from Stage 4 to begin to explore the relationship between those conditions, delivery processes and patient outcomes from Stage 3. This will involve the use of Qualitative Comparative Analysis (QCA) techniques [30,31] to identify which key factors are logically necessary and sufficient to result in these recovery trajectories or outcomes.

Stage 6: We will review the different causal paths resulting from the QCA analysis and then test them out through further perusal of the case narratives. The outcomes of the case studies will be examined alongside the cost efficiency analysis to explore and seek to account for, discrepancies in the qualitative and quantitative studies. The final part of the analysis will focus on locating the community hospital rehabilitation within the broader intermediate care system for older people.

Study 4 - to develop toolkits for commissioners and community hospital providers to optimise performance (objective 5)

We will develop two web-based interactive toolkits for use by local commissioners and community hospital teams respectively that support operational changes to optimise their community hospital wards. The web pages for the online toolkits will be built in asp.net and linked to the Network's SQL Server database which contains the NAIC and the NHSBN Community Hospital Programme data. The toolkits will be securely accessible via the NHSBN website. The toolkits will be based on previous similar work conducted by NHSBM for other clinical service areas and will include three core elements: local performance indicators benchmarked against best performance; a calculation of local potential performance gains; and case studies.

The key indicators of community hospitals that optimise performance will be presented in a series of dashboards. The dashboards will be customised to suit the different needs of

commissioners and providers. The dashboards will show the national performance range, the performance level of the best performing community hospitals and local performance against each key indicator. The dashboards will show summary information for all community hospitals in the particular health economy with the ability to drill down to the performance of individual community hospitals and additional performance metrics.

A Quality, Innovation, Productivity, and Prevention (QIPP) calculator will show the gap between local and best performing community hospitals, the investment required to meet best practice levels of community hospital capacity and potential savings from meeting best practice efficiency values. The potential impact on local secondary care of optimising community hospital capacity and performance will be modelled.

Descriptions of the in-depth case studies generated in Study 3 will be available for download as part of the toolkit. The case studies will explain how the best performing community hospitals achieved high levels of performance.

The content of the toolkits will be co-produced and iteratively modified with a group of 3 to 6 community hospital teams, the Community Hospitals Association and The Patients Association. The toolkits will be launched at workshops within the final conference event and delegate suggestions for further modification considered.

Initial testing will be conducted with the group of 3 to 6 community hospital teams by the NHSBN analytics team. The web pages will then be published to the testing area of the Network's servers. A further round of testing will take place with 3 to 6 new community hospital sites (commissioners and providers). Amendments can be made at this stage to ensure the toolkits function as specified and meet user requirements.

Once testing is complete, the toolkit will be published to Network's live web and database environments.

ETHICS AND DISSEMINATION

Study 3 required ethical approval as patient consent is necessary to obtain information from medical records about individual outcomes, to undertake specific ward observations relating to their care, and to carry out qualitative interviews/conversations with them and their caregivers about their experiences of care in the ward. Ethical approval was obtained via the Health Research Authority that governs research ethics in the UK (Reference: 15/YH/0062). The main ethical issue is one of consent to participate among patients who may lack capacity on account of dementia or delirium. To exclude such patients from the study risks the loss of voices of those for whom care delivery may be particularly problematic. The consent procedures will adhere to the Mental Capacity Act (2005) and accompanying Code of Practice. First, where an individual has been identified by staff as having a condition that could affect their mental capacity, we will consider how best to approach the person, with advice from ward staff (for example, times when the person may be more alert. or when their relative is present). Second, the researcher will explain the study in clear terms and ascertain whether the person understands the information. Third, if the person is deemed unable to make a decision about participation in the research, we will identify a suitable personal consultee either with the patient or with a staff member (close relative or friend). If there is no suitable personal consultee, a nominated consultee will be consulted to enable patients without next of kin to participate. If the consultee advises that a patient who lacks capacity would be willing to take part in the study then that person will be included in the research, providing that they show no signs of unwillingness to participate (for example becoming distressed, upset, or anxious in the presence of the researcher or when discussing the study). Consent will be an ongoing process. The researcher will repeatedly check with participants that they are happy to continue and will be sensitive to any signs of distress or unwillingness to proceed. If any such signs verbal or nonverbal are present, we will discontinue.

The academic outputs of the overall study will be published both as a report to the funder and in peer reviewed papers. Dissemination of these and associated reports and other outputs such as the toolkits will be disseminated through our stakeholder networks including the NHS Benchmarking Agency and the Community Hospitals Association.

AUTHORS CONTRIBUTIONS

All authors contributed to writing the protocol and this manuscript.

FUNDING STATEMENT

This work was supported by NIHR Health Services and Delivery Research Project: 12/177/04.

COMPETING INTERESTS STATEMENT

Dr Helen Tucker is employed by the Community Hospitals Association, which receives financial support from UK community hospitals.

REFERENCES

- [1] Office for National Statistics.2012. Population Ageing in the United Kingdom, its Constituent Countries and the European Union. London: ONS. Available online at http://www.ons.gov.uk/ons/rel/mortality-ageing/focus-on-older-people/population-ageing-in-the-united-kingdom-and-europe/rpt-age-uk-eu.html Accessed 21/02/2015
- [2] Young J, Robinson J, Dickinson E. Rehabilitation for older people BMJ 1998;316: 1108.
- [3] Young J, Gladman JRF, Forsyth DR, et al. The Second National Audit of Intermediate Care. Age Ageing 2015;44(2):182-184
- [4] Tucker, H. Discovering integrated care in community hospitals. Journal of Integrated Care 2013:21:336-346.
- [5] Van Oosterbos, H. The future of integrated primary care: community health centres at the heart of the neighbourhood. International Journal of Integrated Care 2006;6: 1-2.
- [6] Bernabei R, Landi F, Gambassi G, et al. Randomised trial of impact of model of integrated care and case management for older people living in the community. BMJ 1998; 316:1348-1351.
- [7] Lappegard, Ø and Hjortdahl, P. Perceived quality of an alternative to acute hospitalization: An analytical study at a community hospital in Hallingdal, Norway. Social Science & Medicine 2014;119:27-35.

- [8] Young J, Green J, Forster A, et al. Postacute care for older people in community hospitals: a multicentre randomized, controlled trial. J American Geriatr Assoc 2007;55: 1995-2002.
- [9] O'Reilly J, Lowson K, Green J, et al. Post-acute care for older people in community hospitals a cost-effectiveness analysis within a multi-centre randomised controlled trial. Age and Ageing 2008;37:513-520.
- [10] Garasen H, Winspoll, Johnsen R. Long term outcomes after intermediate care at a community hospital for elderly patients: 12 months follow up of a randomised controlled trial Scan J Public Health 2008;36:197.
- [11] Tucker H, Moore B, Jones S, et al. Profiling Community Hospitals in England 1998-2008. Community Hospitals Association: Department of Health and Community Health Partnership 2008.

http://www.communityhealthpartnerships.co.uk/index.php?ob=1&id=321

- [12] Farrell, MJ. The Measurement of Productive Efficiency. Journal of the Royal Statistical Society (A, general) 1957;120: 253–281.
- [13] Coelli T, Rao DSP, O'Donnell CJ, et al An Introduction to Efficiency and Productivity Measurement. Springer: New York, 2005.
- [14] Smith, ASJ (2008), International Benchmarking of Network Rail's Maintenance and Renewal Costs: An Econometric Study Based on the LICB Dataset (1996-2006), Report for the Office of Rail Regulation, October 2008.
- [15] Buckell J, Smith A, Longo R, et al. Efficiency, heterogeneity and cost function analysis: empirical evidence from pathology services in the National Health Service in England. Applied Economics 2015;47:3311-3331.
- [16] Smith ASJ, Buckell J, Wheat P, et al. Hierarchical performance and unobservable heterogeneity in health: A dual-level efficiency approach applied to NHS pathology in England. In: Greene W, Sickles RC, Khalaf L, Veall M. & Voia M. (eds.) Productivity and Efficiency Analysis. New York: Springer Proceedings in Business and Economics 2015.

- [17] Cooper WW, Seiford LM, Zhu J. Handbook on Data Envelopment Analysis. Kluwer Academic Publishers: Massachusetts, USA 2004.
- [18] Hollingsworth B. The measurement of efficiency and productivity of health care delivery. Health Economics 2008;17:1107–1128.
- [19] Parmeter CF & Kumbhakar SC. 2014. Efficiency Analysis: A Primer on Recent Advances. Foundations and Trends in Econometrics, 7, 191-385.
- [20] White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. Statistics in Medicine 2011;30:377-399.
- [21] Ragin C. 'Casing' and the process of social inquiry. In Charles C Ragin and Howard S Becker (Eds.). What is a Case: Exploring the Foundations of Social Inquiry. Cambridge: Cambridge University Press, 1992, pp 217-226.
- [22] Yin R. Case Study Research: Design and Methods (4th Ed.). Thousand Oaks, CA: Sage Publications, 2009.
- [23] Byrne D. Case-based methods: why we need them; what they are; how to do them. In David Byrne and Charles C Ragin, (Eds.). The Sage Handbook of Case-based Methods. London: Sage Publications, 2012; pp 1-10.
- [24] Godfrey M, Keen J, Townsend J et al. Evaluation of intermediate care as a system of care for older people, Report to the Department of Health Policy Research Programme. Leeds: University of Leeds, 2005.
- [25] Patton MQ. Qualitative research & evaluation methods 3rd ed. London: Sage, 2002.
- [26] Long AF and Godfrey M. An evaluation tool to assess the quality of qualitative research studies in systematic reviews, International Journal of Social Research Methodology 7 (2): 181-196, 2004.
- [27] Patterson M, Nolan M, Musson G. From Metrics to Meaning: Culture Change and Quality of Acute Hospital Care for Older People. London: HMSO, 2011.

- [28] Charmaz K. Constructing Grounded Theory. 2nd Edition. London: Sage Publications, 2014.
- [29] Godfrey M and Townsend J. Older people in transition from illness to health: trajectories of recovery. Qual. Health Research 2008; 18: 939-951.
- [30] Ragin CC. Fuzzy-set Social Science. Chicago: University of Chicago Press, 2000.
- [31] Ragin CC, Rihoux B. Qualitative comparative analysis (QCA): state of the art and prospects. Qualitative Methods 2004;2(2):3-13.