

RESEARCH

Open Access

A palatable hyperlipidic diet causes obesity and affects brain glucose metabolism in rats

Debora Estadella¹, Lila M Oyama¹, Allain A Bueno², Carlos A Habitante³, Gabriel I Souza¹, Eliane B Ribeiro¹, Caio SM Motoyama¹ and Claudia M Oller do Nascimento^{1*}

Abstract

Background: We have previously shown that either the continuous intake of a palatable hyperlipidic diet (H) or the alternation of chow (C) and an H diet (CH regimen) induced obesity in rats. Here, we investigated whether the time of the start and duration of these feeding regimens are relevant and whether they affect brain glucose metabolism.

Methods: Male Wistar rats received C, H, or CH diets during various periods of their life spans: days 30-60, days 30-90, or days 60-90. Experiments were performed the 60th or the 90th day of life. Rats were killed by decapitation. The glucose, insulin, leptin plasma concentration, and lipid content of the carcasses were determined. The brain was sliced and incubated with or without insulin for the analysis of glucose uptake, oxidation, and the conversion of [1-¹⁴C]-glucose to lipids.

Results: The relative carcass lipid content increased in all of the H and CH groups, and the H30-60 and H30-90 groups had the highest levels. Groups H30-60, H30-90, CH30-60, and CH30-90 exhibited a higher serum glucose level. Serum leptin increased in all H groups and in the CH60-90 and CH30-90 groups. Serum insulin was elevated in the H30-60, H60-90, CH60-90, CH30-90 groups. Basal brain glucose consumption and hypothalamic insulin receptor density were lower only in the CH30-60 group. The rate of brain lipogenesis was increased in the H30-90 and CH30-90 groups.

Conclusion: These findings indicate that both H and CH diet regimens increased body adiposity independent treatment and the age at which treatment was started, whereas these diets caused hyperglycemia and affected brain metabolism when started at an early age.

Introduction

Glucose is considered to be the major nutrient for cells of the adult nervous system. A small portion of glucose is used for biosynthesis pathways. Dhopeswarkar and Subramanian [1] showed that intracranially administered D- [U-¹⁴C] glucose was incorporated in saturated fatty acids and, to a lesser extent, in monounsaturated fatty acids. Poor glucose metabolism has also been related to memory problems [2].

In a previous review, it was reported that high-fat diets, especially diets enriched with saturated fatty acids, impaired the learning and memory of rodents and that these effects could be related to insulin resistance and

impaired glucose metabolism [3,4]. However, polyunsaturated fatty acids improve cognitive function and are beneficial for the prevention of cognitive decline [5]. Kaiyala et al. [6] reported a reduction of insulin uptake and clearance by the central nervous system in dogs that were fed a hyperlipidic diet.

Insulin receptors are expressed in several regions of the central nervous system [7,8]. Experimental evidence has demonstrated the important role of the insulin central receptors as normal energy balance controllers [9-11] in addition to their role in neuronal growth and differentiation during development [12,13]. According to Carnevali et al. [14], insulin modulates the leptin signal transduction pathway in the hypothalamus, playing a synergistic role in regulating food intake and controlling weight.

Previous experiments have shown that rats fed a high-fat diet had an increased fat mass percentage even when

* Correspondence: claudia.oller@unifesp.br

¹Disciplina de Fisiologia da Nutrição, Departamento de Fisiologia, Universidade Federal de São Paulo, São Paulo, SP, Brazil

Full list of author information is available at the end of the article

the energy intake was not elevated [15,16]. Likewise, we previously demonstrated that the continuous intake of a hyperlipidic palatable diet and the alternation of a high-fat intake with periods of chow intake for 8 weeks caused obesity and affected the lipid metabolism of rats in a similar way, although the energy intake remained unchanged (KJ) [17].

In these experimental models, obesity was accompanied by hyperleptinemia, normoglycemia, hyperinsulinemia and insulin resistance. The age, gender weight, and period on high-fat diets of the animals influenced their metabolism [16,18-20].

There are controversial results regarding the effect of hyperlipidic diets on the expression of leptin receptors (Ob-Rb and Ob-Ra) in the central nervous system. Madiehe et al. [21] suggested that a decrease in the levels of both the long and short form of the hypothalamus leptin receptors could induce leptin resistance in hyperlipidic-fed rats. In mice given a high-fat diet, Lin et al. [22] reported an increase in mRNA leptin receptor expression after 8 weeks and a reduction of this expression after 19 weeks. Boado et al. [23] detected OB-Ra up-regulation in the blood-brain-barrier of the rats after 14 weeks of high-fat feeding.

The suckling-weaning transition of rats is accompanied by marked dietary changes. Suckling rats eat a very high-fat, low-carbohydrate diet, and this is replaced by a low-fat, high-carbohydrate diet at the time of weaning [24]. The utilization of substrates by the brain depends on several factors: 1) the plasma concentrations of the substrate, 2) the permeability of the blood-brain barrier to the substrates, and 3) the intrinsic capability of the brain to utilize the substrates [25].

The present study was designed to investigate whether the continuous feeding of a palatable hyperlipidic diet or cycling a hyperlipidic diet with a chow diet induced obesity. We evaluated whether the time of the start and duration of these feeding regimens were relevant and whether they affected brain glucose metabolism.

Materials and Methods

Animals

The Experimental Research Committee of the Federal University of São Paulo approved the procedures used in the experiments presented here (CEP n°0446/03).

Male Wistar rats supplied by the animal care facility of the Physiology Department of the Federal University of São Paulo were assigned to 8 groups according to diet composition: (C) receiving a chow diet, (H) receiving a palatable hyperlipidic diet and (CH) receiving a cycled diet (cycles alternating weekly between a chow diet and a palatable hyperlipidic diet). The time of the start and duration of these feeding regimens were also studied using the following groups: diet received from the 30th to

60th day of life (30-60), diet received from the 30th to 90th day of life (30-90) and diet received from the 60th to 90th day of life (60-90).

All groups were maintained in a room at 23°C ± 1°C with light from 7:00 AM to 7:00 PM and received food and water *ad libitum*. Body weight and food intake were measured weekly. Spillage was minimized due to the use of specially designed covered food cups. All the animals were sacrificed in the fed state in the early morning to avoid chronobiological variations. The animals were acclimatized in a quiet room next to the laboratory under subdued light for 1 h and were taken to the laboratory individually to minimize stress.

Preparation of the palatable hyperlipidic diet

The palatable hyperlipidic diet consists of commercial rat chow plus peanuts, milk chocolate, and sweet biscuits in the proportion of 3:2:2:1. All components were powdered and mixed. This diet was composed of 20% protein, 20% fat, and 40% carbohydrate. The control diet contained 20% protein, 4.5% fat, and 55% carbohydrate. The caloric densities of the diets were determined using an IKA-C400 adiabatic calorimeter. The caloric density was 21.40 kJ/g (35% of calories from fat) for the palatable hyperlipidic diet and 17.03 kJ/g for the chow diet.

Fatty acid composition of the diets

Five hundred milligrams of the diets were treated with 2.0 mL of methanol:benzene (4:1, v/v) followed by 200 µL of acetyl chloride under light agitation [26]. Fatty acid methyl esters were separated (SP2560 column, Supelco, Bellefonte, PA, USA) and quantified using gas-liquid chromatography with an ionizable flame detector (Perkin Elmer, Wellesley, MA, USA) and hydrogen as the carrier gas. The injection and detection temperatures were 260°C and 280°C, respectively. The run temperature started at 135°C and increased to 195°C during a run time of 45 min. The fatty acid composition of the diets is presented in Table 1.

Experimental Procedure

Experiments were performed on the 60th or 90th day of the rats' lives, and the rats were killed by decapitation. Trunk blood was collected, serum was obtained via centrifugation, and aliquots were taken to measure the concentration of glucose using commercial kits from Labtest Diagnostic S.A (Minas Gerais, Brazil) and the concentrations of insulin and leptin using ELISA (Linco Research, Inc., USA).

The brain was removed, the cerebellum and brain stem were discarded, and the remaining brain was sliced as described by Liu et al. (2005) [27] and immediately incubated to determine the brain glucose consumption, glucose oxidation, and lipid synthesis. Another set of

Table 1 Fatty acid profile of the diets, as percent of total lipid content

Fatty Acid	Total Fatty Acid (%)	
	C	H
C14:0	0.11	0.53
C16:0	13.37	17.85
C16:1n7	0.21	0.15
C17:0	0.09	0.12
C18:0	2.95	11.05
C18:1n9	24.13	36.59
C18:2n6	49.45	27.27
C18:3n3	3.73	0.49
C20:0	0.42	1.19
C22:0	0.04	2.14
C22:2n6	0.04	0.04
C23:0	0.07	0.04
C22:5n3	0.22	0.14
C24:0	0.34	1.15
SFA	17.4	34.06
MUFA	25.59	38.00
PUFA	56.97	27.95

Control diet (C) and palatable hyperlipidic diet (H)

SFA. saturated fatty acids; MUFA. monounsaturated fatty acids; PUFA. polyunsaturated fatty acids. * Data are means \pm SEMs of three determinations.

rats was used for the Western blotting analysis. The carcasses were weighed and stored in a freezer (-20°C) for lipid content analysis.

Carcass lipid content determination

The carcasses were eviscerated, weighed, and stored at -20°C. The lipid content was measured as described by Stansbie et al. [28] and standardized using the method described by Oller do Nascimento and Williamson [29]. Briefly, the eviscerated carcass was autoclaved at 120°C for 90 min and then homogenized with double its mass of water. Triplicate aliquots of this homogenate were weighed and digested in 3 ml of 30% KOH and 3 ml of ethanol for at least 2 h at 70°C in capped tubes. After cooling, 2 ml of 12 N H₂SO₄ were added, and the sample was washed three times with petroleum ether for lipid extraction. The results are expressed as grams of lipid/100 g of carcass.

Brain incubation

The total brain was sliced and used for the incubation procedure (~100 mg). Incubations were performed at 37°C in 25 ml Erlenmeyer flasks equipped with a central well. The incubation medium consisted of 2.0 ml of Krebs-Henseleit buffer containing 5 mM glucose plus 0.04 μ Ci ¹⁴C-glucose with or without 0.5 U/ml insulin, as it has been previously demonstrated that chronic hyperinsulinemia could alter the glucose utilization of several brain regions [30]. The flasks were continuously shaken

and flushed with carbogenium (O₂/CO₂, 95/5%) during the incubation period.

After 1 hour, incubation was stopped by the addition of 0.5 ml of 4 N H₂SO₄ to the main well, and 0.3 ml of NaOH (1 N) was then added to the central well for ¹⁴CO₂ collection. After two hours, the brain slices were removed, and the total lipids were extracted with 10 ml of chloroform:methanol (2:1). The radioactivity of this extract represented the conversion of [1-¹⁴C]-glucose to lipids. The rates of conversion of [1-¹⁴C]-glucose to ¹⁴CO₂ and incorporation into the lipid fraction were expressed as μ mol/h \times g of tissue. The incubation medium was used for the determination of the brain glucose consumption using commercial kits from Labtest Diagnostic S.A. These methods have been previously described [31].

Western blot analysis

The rat cranium was opened, and the hypothalamus was quickly removed, minced coarsely and immediately homogenized in 1 mL of buffer [100 mM Tris (pH 7.6), 10 mM Na₃VO₄, 2 mM PMSE, 10 mM EDTA, 100 mM NaF, 10 mM Na₄P₂O₇, 0.1 mg/mL \times aprotinin] using an ULTRA TURAX IKA T-18 BASIC generator operated at maximum speed for 30 s. Then, 10% Triton X-100 was added, and the solution was clarified by centrifugation (12,000 rpm, 20 min, at 4°C).

The total protein content from the entire tissue extraction was determined using standard BSA, following previously described methods [32]. One hundred micrograms of hypothalamus total protein was loaded onto a sodium dodecyl sulfate-polyacrylamide gel (5% stacking gel; 10% running gel), separated by electrophoresis and then electroblotted onto nitrocellulose membranes (Hybond-C Extra, Amersham) using a wet electroblotter (Bio-Rad, CA, USA). After blotting, the membranes were blocked in Tris-buffered saline (TBS)-Tween buffer, pH 7.5 (20 mM Tris/500 mM NaCl/0.05% Tween-20) containing 10% skimmed milk powder for 2 h and then exposed to either anti-insulin receptor (IR) or anti-leptin receptor (OB-R) at a dilution of 1/200 in TBS-Tween buffer (pH 7.5) containing 1% BSA for 2 h. The membranes then were washed and incubated with anti-rabbit Ig or anti-mouse Ig conjugated to horseradish peroxidase and diluted to 1/1000 in the same buffer for 1 h. After a series of washes in TBS-Tween buffer, the protein bands were visualized by chemiluminescence with an ECL luminescence kit (Amersham) and exposure to Hyper-film ECL (Amersham). The size of the protein bands was determined using electrophoresis color markers. The antibodies against IR (sc-711) and OB-R (sc-8391) were obtained from Santa Cruz Biotechnology (Santa Cruz, Calif., USA), and the anti-rabbit Ig and anti-mouse Ig conjugated to horseradish peroxidase were

obtained from Sigma (USA). A quantitative analysis of the blots was performed using Image J software.

Statistical Analysis

The results are expressed as the mean \pm standard error of means. Statistical comparisons were carried out using analysis of variance (one-way ANOVA) following by a post-hoc analysis (Tukey test) to compare the effects of the different diets among animals of the same age. The level of significance was set at $p < 0.05$.

Results

Body Weight, Energy Intake and Carcass lipid content

The increment in body weight was not significantly different among the C30-60, H30-60 and CH30-60 groups. When the animals were fed a palatable hyperlipidic diet or the cycled diet from the 30th to 90th or from the 60th to 90th day of life, we observed a significant increase in body weight compared to the respective C groups (Table 2).

The total caloric intake per body weight gain (kJ/100 g) was significantly lower in the H60-90, CH60-90 and CH30-90 groups compared to the C60-90 and C30-90 groups (Table 2).

The palatable hyperlipidic diet and the food cycle treatments increased the relative carcass lipid content in all of the experimental groups compared to the C groups. In addition, the H30-60 group had a higher carcass lipid content than the CH30-60 group (Table 2).

Glucose, leptin and insulin serum concentrations

Rats exposed to the palatable hyperlipidic diet (groups H30-60, CH30-60, H30-90 and CH30-90) exhibited higher serum glucose concentrations compared to the respective chow diet-fed rats. When given the palatable hyperlipidic diet, either continuously or cycled with

chow beginning on the 60th day of life, the serum glucose levels were not significantly affected.

The serum leptin levels increased in all of the H groups and in the CH60-90 and CH30-90 groups compared to the respective C groups.

Both diet regimens and the different periods of treatment were effective in increasing the serum insulin concentrations in the H30-60, H60-90, CH60-90 and CH30-90 groups compared to the respective chow diet-fed rats (Table 3).

Brain glucose consumption, glucose oxidation and brain lipogenesis rate

The basal brain glucose consumption was lower in the CH30-60 group than in the C30-60 and H30-60 groups, and the insulin addition to the media normalized this parameter. Brain glucose consumption was similar between the H groups and the C groups with or without insulin addition (Table 4).

The brain lipogenesis rate from ¹⁴C-glucose, both basal levels and when stimulated by insulin, was increased in the H30-90 and CH30-90 groups compared with the C30-90 group (Table 5).

The diet regimens did not alter the brain glucose oxidation in any of the experimental groups compared to the control groups (Table 6).

Quantification of the Insulin and leptin receptors

(Figure 1A, B and 1C shows the quantification of the insulin receptor (IR) from the extracted tissue of the entire hypothalamus of the rats. The cycled diet from the 30th to 60th day of life significantly decreased the insulin receptor density compared to the C group.

No significant difference in leptin receptor protein levels was observed among the groups (Figure 1D, E and 1F).

Table 2 Increment in body weight (g), total caloric intake (kJ/100 g b.w gain) and carcass relative lipid content (g/100 g)

		C	H	CH
Increment in body weight (g)	30-60	118.1 \pm 4.8 ^a	124.7 \pm 8.2 ^a	126.8 \pm 6.8 ^a
	60-90	77.1 \pm 3.1 ^a	91.9 \pm 3.2 ^b	103.9 \pm 3.3 ^c
	30-90	198.1 \pm 9.9 ^a	207.8 \pm 7.4 ^{ab}	237.2 \pm 13.5 ^b
Total caloric intake (kJ/100 g b.w)	30-60	7017.9 \pm 656.0 ^a	6882.7 \pm 464.7 ^a	6875.1 \pm 357.0 ^a
	60-90	15282.7 \pm 656.0 ^a	12662.3 \pm 408.1 ^b	11057.6 \pm 359.7 ^b
	30-90	10289.9 \pm 642.9 ^a	9675.2 \pm 401.6 ^{ab}	8162.0 \pm 369.9 ^b
Carcass Relative lipid content(g/100 g)	30-60	2.32 \pm 0.18 ^a	5.48 \pm 0.35 ^b	3.94 \pm 0.29 ^c
	60-90	2.65 \pm 0.17 ^a	4.26 \pm 0.34 ^b	3.75 \pm 0.32 ^b
	30-90	2.65 \pm 0.17 ^a	3.94 \pm 0.26 ^b	3.49 \pm 0.21 ^b

Control rats (C), rats treated with palatable hyperlipidic diet (H) and rats treated with food cycles alternating chow and palatable hyperlipidic diets (CH). The animals received these diets from 30th to 60th days of life (30-60), from 30th to 90th days of life (30-90), or from 60th to 90th days of life (60-90). Values are means \pm standard error of the mean of 10 animals. Values in the same row with different superscript letters are significantly different from one another at $p < 0.05$ (Tukey's test)

Table 3 Serum glucose (mg/dL), insulin (μU/mL) and leptin (ng/mL) concentrations

		C	H	CH
Glucose(mg/dL)	30-60	98.7 ± 3.6 ^a	116.9 ± 3.8 ^b	116.5 ± 3.6 ^b
	60-90	95.5 ± 4.1 ^a	98.3 ± 2.8 ^a	104.0 ± 5.4 ^a
	30-90	95.5 ± 4.1 ^a	115.1 ± 4.4 ^b	115.6 ± 4.7 ^b
Leptin(μU/mL)	30-60	1.57 ± 0.19 ^a	7.41 ± 1.31 ^b	3.54 ± 0.32 ^a
	60-90	3.41 ± 0.24 ^a	6.67 ± 0.99 ^b	6.41 ± 1.08 ^b
	30-90	3.41 ± 0.24 ^a	8.27 ± 1.01 ^b	9.68 ± 1.76 ^b
Insulin(ng/mL)	30-60	15.54 ± 1.19 ^a	21.94 ± 1.81 ^b	16.95 ± 1.10 ^a
	60-90	19.10 ± 1.45 ^a	25.93 ± 1.57 ^b	29.81 ± 2.52 ^b
	30-90	19.10 ± 1.45 ^a	19.63 ± 2.18 ^a	25.48 ± 1.09 ^b

Control rats (C), rats treated with palatable hyperlipidic diet (H) and rats treated with food cycles alternating chow and palatable hyperlipidic diets (CH). The animals received these diets from 30th to 60th days of life (30-60), from 30th to 90th days of life (30-90), or from 60th to 90th days of life (60-90). Values are means ± standard error of the mean of 10 animals. Values in the same row with different superscript letters are significantly different from one another at p < 0.05 (Tukey's test)

Discussion

In the present investigation, we found that both hyperlipidic diet regimes increased the carcass lipid content in all of the groups studied. We also found that when the special diet began on the 30th day of life, the serum glucose level and brain glucose metabolism were more affected.

The palatable hyperlipidic diet induced a more pronounced body weight gain in the H60-90, CH60-90, and CH30-90 groups, regardless of the regimen of administration, i.e., either continuously administered or cycled with chow, although significant differences were observed based on the age and treatment period. Because caloric intake was lower in these animals, the palatable hyperlipidic diet may have increased their metabolic efficiency, as reported by others [33-35]. Similarly, successive caloric restriction and high-fat re-feeding cycles have been shown to increase metabolic efficiency and promote obesity [36].

Experimental data on the effects of palatable hyperlipidic diets are somewhat contradictory with respect to body weight gain. The collective data are not clear regarding the effects of hyperlipidic diets, also referred to

as cafeteria diets, on the body weight gain of animals. Some authors have not reported an increase in body weight [19,37], whereas others have found an elevated body weight [38,39].

Both hyperlipidic diet regimes tested here increased the relative carcass lipid content in all of the studied groups. Previously, we have shown that a hyperlipidic diet administered for 8 weeks induced a more pronounced body weight gain and adiposity compared to chow diet [17]. In the present study, only 4 weeks on a hyperlipidic diet, both continuously and alternated with a chow diet, caused an enhancement in adiposity that was accompanied by an increase in serum leptin concentration, which shows that short periods of hyperlipidic diets can promote obesity. In previous studies, several obese animal models have also developed elevated circulating leptin levels [15,40].

A number of reports have demonstrated that high-fat diets may promote hyperglycemia [26,41,42]. We found higher serum glucose levels in the H30-60, CH30-60, H30-90 and CH30-90 groups than in the respective control groups. This result may suggest the occurrence of glucose intolerance. In the H60-90 and CH60-90 groups, glucose intolerance appeared to be compensated for by hyperinsulinemia, which adjusted serum glucose levels.

In our study, we did not find changes in the levels of hypothalamic leptin receptor protein, despite the observed hyperleptinemia. This finding is consistent with the results of Sahu et al. [43] and Peiser et al. [44] who also did not verify an alteration in the hypothalamus leptin receptor protein levels of rats treated with a hyperlipidic diet. In contrast, Munzberg et al. [45] studied several regions of the hypothalamus using STAT3 to map leptin-responsive cells in the brain and observed resistance to leptin action in the arcuate nucleus of the rats fed a high-fat diet.

Glucose is the major energy substrate for neurons. The major objective of our study was to analyze the effect of a hyperlipidic diet, administered either continuously or cycled with a chow diet, on brain glucose metabolism in different periods of life in the rat. There are data to

Table 4 Brain glucose consumption (μmol/g tissue/h) of control rats (C)

		C	H	CH
Glucose(μmol/g tissue/h)	30-60	22.52 ± 1.53 ^a	20.84 ± 2.19 ^a	14.59 ± 1.34 ^b
	60-90	21.27 ± 1.47 ^a	20.47 ± 1.97 ^a	15.77 ± 2.10 ^a
	30-90	21.27 ± 1.47 ^a	22.46 ± 2.19 ^a	19.73 ± 1.81 ^a
Glucose + Insulin(μmol/g tissue/h)	30-60	21.56 ± 2.15 ^a	19.87 ± 1.44 ^a	18.76 ± 1.28 ^a
	60-90	21.56 ± 1.93 ^a	18.47 ± 1.43 ^a	15.54 ± 1.78 ^a
	30-90	21.56 ± 1.93 ^a	20.06 ± 1.44 ^a	24.35 ± 2.12 ^a

Control rats (C), rats treated with palatable hyperlipidic diet (H) and rats treated with food cycles alternating chow and palatable hyperlipidic diets (CH). The animals received these diets from 30th to 60th days of life (30-60), from 30th to 90th days of life (30-90), or from 60th to 90th days of life (60-90). Values are means ± standard error of the mean of 10 animals. Values in the same row with different superscript letters are significantly different from one another at p < 0.05 (Tukey's test)

Table 5 Brain lipogenesis rate (¹⁴C-glucose incorporated in lipid/g tissue/h)

		C	H	CH
Glucose (¹⁴ C-glucose incorporated in lipid/g tissue/h)	30-60	4.91 ± 0.22 ^a	4.74 ± 0.14 ^a	4.79 ± 0.14 ^a
	60-90	4.26 ± 0.17 ^a	4.96 ± 0.23 ^a	4.68 ± 0.29 ^a
	30-90	4.26 ± 0.17 ^a	6.05 ± 0.20 ^b	5.79 ± 0.14 ^b
Glucose + insulin (¹⁴ C-glucose incorporated in lipid/g tissue/h)	30-60	4.77 ± 0.16 ^a	4.76 ± 0.17 ^a	4.46 ± 0.17 ^a
	60-90	4.53 ± 0.18 ^a	4.61 ± 0.25 ^a	4.63 ± 0.24 ^a
	30-90	4.53 ± 0.18 ^a	5.66 ± 0.18 ^b	5.37 ± 0.10 ^b

Control rats (C), rats treated with palatable hyperlipidic diet (H) and rats treated with food cycles alternating chow and palatable hyperlipidic diets (CH). The animals received these diets from 30th to 60th days of life (30-60), from 30th to 90th days of life (30-90), or from 60th to 90th days of life (60-90). Values are means ± standard error of the mean of 10 animals. Values in the same row with different superscript letters are significantly different from one another at p < 0.05 (Tukey's test)

suggest a that high-fat diet, especially one enriched in saturated fatty acids, decreases the learning capacity and memory of rodents and is related to insulin resistance and impaired glucose metabolism [3,4].

The administration of successive cycles of chow and high-fat diets for 30 d (CH30-60) elevated the plasma glucose levels and diminished the brain glucose consumption in relation to the C30-60 group; this was accompanied by a decrease in the levels of insulin receptor protein in the hypothalamus. A non-significant reduction (25.8%) was also observed in brain glucose consumption in the CH60-90 group compared with the C60-90 group. However, this result was not observed in the H animals or the CH30-90 group compared to the respective controls. In addition, the presence of insulin in the incubation media elevated brain glucose consumption in the CH30-90 group compared with the CH60-90 group (24.35 ± 2.12 vs. 15.54 ± 1.78, p < 0.05), which suggests that long-term treatment caused an adaptation in brain glucose metabolism.

It was reported in a review [46] that arachidonic acid, a polyunsaturated fatty acid, stimulates glucose uptake in cerebral cortical astrocytes and thus plays a role in the regulation of energy metabolism in the cerebral cortex. In the present study, the high-fat diet that was used was rich in saturated and monounsaturated fatty acids and poor in polyunsaturated fatty acids compared to the chow diet, which could promote a decrease in the

incorporation of polyunsaturated fatty acids in the neuronal cell membrane and cause a decrease in glucose uptake or a defect in the expression or function of insulin receptors in the brain. Yu et al. [47] showed a positive correlation between glucose uptake by astrocytes and arachidonic acid concentration.

As described by Girard et al. [25], the concentration of GLUT1 and the rate of glucose transport in the rat brain during the suckling period are lower than the levels in adult rats. It is important to note that these changes in brain metabolism correspond to the animal's diet: milk during the suckling period, which is high in fat, or adult solid food, which is high in carbohydrates.

Duelli et al. [48] have reported a decrease in GLUT1 protein density during chronic hyperglycemia followed by an increase in brain glucose utilization. However, existing data are inconsistent with respect to the effects of hyperglycemia on brain glucose consumption. Some authors have demonstrated a decrease in brain glucose consumption [49], whereas others have shown an elevation [50,51].

It has been reported that fat transport mechanisms, oxidation and synthesis are present in some regions of the brain [52,53]. In our findings, we verified an increase in the rate of brain lipogenesis from ¹⁴C-glucose in the H30-90 and CH30-90 groups in media with or without insulin.

Bukato et al. [54] have shown that the activity of malic enzyme and the citrate synthase enzyme in rat brains

Table 6 Brain glucose oxidation (μmol/g tissue/h)

		C	H	CH
Glucose (μmol/g tissue/h)	30-60	0.66 ± 0.07 ^a	0.71 ± 0.06 ^a	0.71 ± 0.04 ^a
	60-90	0.63 ± 0.07 ^a	0.60 ± 0.10 ^a	0.71 ± 0.07 ^a
	30-90	0.63 ± 0.07 ^a	0.70 ± 0.11 ^a	0.77 ± 0.12 ^a
Glucose + Insulin (μmol/g tissue/h)	30-60	0.68 ± 0.06 ^a	0.56 ± 0.06 ^a	0.52 ± 0.04 ^a
	60-90	0.57 ± 0.06 ^a	0.55 ± 0.11 ^a	0.57 ± 0.06 ^a
	30-90	0.57 ± 0.06 ^a	0.81 ± 0.09 ^a	0.61 ± 0.11 ^a

Control rats (C), rats treated with palatable hyperlipidic diet (H) and rats treated with food cycles alternating chow and palatable hyperlipidic diets (CH). The animals received these diets from 30th to 60th days of life (30-60), from 30th to 90th days of life (30-90), or from 60th to 90th days of life (60-90). Values are means ± standard error of the mean of 10 animals. Values in the same row with different superscript letters are significantly different from one another at p < 0.05 (Tukey's test)




Figure 1 Insulin receptor protein (IR) (A, B, and C) and leptin receptor protein (OB-R) (D, E, and F) expression (A.U. - Arbitrary Units) from hypothalamus whole tissue extract. (A, D) groups C30-60, H30-60 and CH30-60; (B, E) groups C60-90, H60-90 and CH60-90 and (C, F) groups C30-90, H30-90 and CH30-90. Values are means \pm standard error of the mean of 6 to 7 animals. Values with different superscript letters are significantly different from one another at $p < 0.05$ (Tukey's test).

increased gradually up to 60 days of life and was maintained at similar and stable levels until at least 160 days of life.

In summary, according to these findings, both the continuous intake of a palatable hyperlipidic diet and alternation of a high-fat intake with periods of chow intake increased the body adiposity in rats, which was independent from the duration of their administration and the age at which the altered diet was begun. In

contrast, these diets caused hyperglycemia and affected brain metabolism when they were begun at an early age.

Acknowledgements

The present work was supported by a grant from the State of São Paulo Research Foundation (FAPESP 2001/07187-1).

Author details

¹Disciplina de Fisiologia da Nutrição, Departamento de Fisiologia, Universidade Federal de São Paulo, São Paulo, SP, Brazil. ²Institute of Brain

Chemistry and Human Nutrition, London Metropolitan University, London, UK. ³Instituto de Ciências Biológicas e da Saúde, Campus Universitário do Araguaia, Universidade Federal do Mato Grosso, MT, Brazil.

Authors' contributions

DE has made substantial contributions to conception and design, carried out all the experiments, performed the statistical analysis, interpretation of data and drafted the manuscript. LMO has made substantial contributions to conception and design, analysis and interpretation of data and coordination to draft the manuscript. AAB helped to carried out the experiments and acquisition of data. CAH helped to carried out the experiments and acquisition of data. GIS carried out the fat composition of the diet and revised and helped to draft the manuscript. CSMM helped to carried out the experiments and acquisition of data. EBR participated in its design, coordination and helped to draft the manuscript. CMON has made substantial contributions to conception and design, analysis and interpretation of data, coordination and drafted the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 20 June 2011 Accepted: 23 September 2011

Published: 23 September 2011

References

- Dhopeswarkar GA, Subramanian C: Biosynthesis of polyunsaturated fatty acids in the developing brain: I. Metabolic transformations of intracranially administered 1-14C linolenic acid. *Lipids* 1976, **11**(1):67-71.
- Owen L, Finnegan Y, Hu H, Scholey AB, Sünram-Lea SI: Glucose effects on long-term memory performance: duration and domain specificity. *Psychopharmacology Berl* 2010, **211**(2):131-40.
- Greenwood CE, Winocur G: Glucose treatment reduces memory deficits in young adult rats fed high-fat diets. *Neurobiol Learn Mem* 2001, **75**: 179-189.
- Greenwood CE, Winocur G: High-fat diets, insulin resistance and declining cognitive function. *Neurobiol Aging* 2005, **26**:42-45.
- Das UN: Folic acid and polyunsaturated fatty acids improve cognitive function and prevent depression, dementia, and Alzheimer's disease—but how and why? *Prostaglandins Leukot Essent Fatty Acids* 2008, **78**(1):11-9.
- Kaiyala KJ, Prigeon RL, Kahn SE, Woods SC, Schwartz MW: Obesity induced by a high-fat diet is associated with reduced brain insulin transport in dogs. *Diabetes* 2000, **49**:1525-1533.
- Baskin DG, Figlewicz DP, Woods SC, Porte D, Dorsa DM: Insulin in the brain. *Annu Rev Physiol* 1987, **49**:335-347.
- Havrankova J, Roth J, Brownstein M: Insulin receptors are widely distributed in the central nervous system of the rat. *Nature* 1978, **272**:827-829.
- Schwartz MW, Figlewicz DP, Baskin DG, Woods SC, Porte D Jr: Insulin in the brain: a hormonal regulator of energy balance. *Endocr Rev* 1992, **13**:387-414.
- Sipols AJ, Baskin DG, Schwartz MW: Effect of intracerebroventricular insulin infusion on diabetic hyperphagia and hypothalamic neuropeptide gene expression. *Diabetes* 1995, **44**:147-151.
- Bruning JC, Gautam D, Burks DJ, Gillette J, Schubert M, Orban PC, Klein R, Krone W, Muller-Wieland D, Kahn CR: Role of brain insulin receptor in control of body weight and reproduction. *Science* 2000, **289**:2122-2125.
- Heidenreich KA: Insulin and IGF-I receptor signaling in cultured neurons. *Ann N Y Acad Sci* 1993, **692**:72-88.
- Robinson LJ, Leitner W, Draznin B, Heidenreich KA: Evidence that p21ras mediates the neurotrophic effects of insulin and insulin-like growth factor I in chick forebrain neurons. *Endocrinology* 1994, **135**:2568-2573.
- Carvalho JB, Siloto RM, Ignacchitti I, Brenelli SL, Carvalho CR, Leite A, Velloso LA, Gontijo JA, Saad MJ: Insulin modulates leptin-induced STAT3 activation in rat hypothalamus. *FEBS Lett* 2001, **500**:119-124.
- De Schepper J, Zhou X, De Bock S, Smits J, Louis O, Hooghe Peters E, Vandenplas Y: Study of serum leptin in cafeteria-diet-overfed rats. Influence of diet, insulin and corticosterone. *Horm Res* 1998, **50**:271-275.
- Gaíva MH, Couto RC, Oyama LM, Couto GE, Silveira VL, Riberio EB, Nascimento CM: Polyunsaturated fatty acid-rich diets: effect on adipose tissue metabolism in rats. *Br J Nutr* 2001, **86**:371-377.
- Estadella D, Oyama LM, Dâmaso AR, Ribeiro EB, Oller do Nascimento CM: Effect of palatable hyperlipidic diet on lipid metabolism of sedentary and exercised rats. *Nutrition* 2004, **20**:218-224.
- Esteve M, Rafecas I, Fernandez Lopes JA, Remesar X, Alemany M: Effect of cafeteria diet on energy-intake and balance in wistar rats. *Physiol Behav* 1994, **56**:65-71.
- Brooks SP, Lampi BJ: Effect of dietary fat on whole body fatty acid synthesis in weanling rats. *J Nutr Biochem* 1999, **10**:291-8.
- Leonhardt M, Langhans W: Fatty acid oxidation and control of food intake. *Physiol Behav* 2004, **83**:645-651.
- Madihe AM, Schaffhauser AO, Braymer DH, Bray GA, York DA: Differential expression of leptin receptor in high- and low-fat-fed Osborne-Mendel and S5B/Pl rats. *Obes Res* 2000, **8**:467-474.
- Lin S, Storlien LH, Huang X: Leptin receptor, NPY, POMC mRNA expression in the diet-induced obese mouse brain. *Brain Res* 2000, **875**:89-95.
- Boado RJ, Golden PL, Levin N, Pardridge WM: Up-regulation of blood-brain barrier short-form leptin receptor gene products in rats fed a high-fat diet. *J Neurochem* 1998, **71**:1761-1764.
- Ferré P, Leturque A, Burnol AF, Penicaud L, Girard J: A method to quantify glucose utilization in vivo in skeletal muscle and white adipose tissue of the anaesthetized rat. *Biochem J* 1985, **228**:103-110.
- Girard J, Ferré P, Pegorier JP, Duee PH: Adaptations of glucose and fatty acid metabolism during perinatal period and suckling-weaning transition. *Physiol Rev* 1992, **72**:507-562.
- Hansen PA, Han DH, Marshall BA, Nolte LA, Chen MM, Mueckler M, Holloszy JO: A high-fat diet impairs stimulation of glucose transport in muscle. Functional evaluation of potential mechanisms. *J Biol Chem* 1998, **273**:26157-26163.
- Liu R, Pei JJ, Wang XC, Zhou XW, Tian Q, Winblad B, Wang JZ: Acute anoxia induces tau dephosphorylation in rat brain slices and its possible underlying mechanisms. *J Neurochem* 2005, **94**(5):1225-34.
- Stansbie D, Brownsey RW, Crettaz M, Denton RM: Acute eVects in vivo of anti-insulin serum on rates of fatty acid synthesis and activities of acetyl-coenzyme A carboxylase and pyruvate dehydrogenase in liver and epididymal adipose issue of fed rats. *Biochem J* 1976, **160**:413-416.
- Oller do Nascimento CM, Williamson DH: Evidence for conservation of dietary lipid in the rat during lactation and the immediate period after removal of the litter. *Biochem J* 1986, **239**:233-236.
- Doyle P, Cusin I, Rohner-Jeanrenaud F, Jeanrenaud B: Four-day hyperinsulinemia in euglycemic conditions alters local cerebral glucose utilization in specific brain nuclei of freely moving rats. *Brain Res* 1995, **684**:47-55.
- Oyama LM, Oller do Nascimento CM: Effect of ethanol intake during lactation on male and female pups' liver and brain metabolism during the suckling-weaning transition period. *Nutr Neurosci* 2003, **3**:183-188.
- Lowry OH, Rosebrough NJ, Farr A, Randall RJ: Protein measurement with the Folin phenol reagent. *J Biol Chem* 1951, **193**:265-275.
- Gaíva MH, Couto RC, Oyama LM, Couto GE, Silveira VL, Riberio EB, Nascimento CM: Polyunsaturated fatty acid-rich diets: effect on adipose tissue metabolism in rats. *Br J Nutr* 2001, **86**(3):371-7.
- Schrauwen P, Westerterp KR: The role of high-fat diets and physical activity in the regulation of body weight. *Br J Nutr* 2000, **84**(4):417-27, Review.
- Roberts CK, Vaziri ND, Barnard RJ: Effect of diet and exercise intervention on blood pressure, insulin, oxidative stress, and nitric oxide availability. *Circulation* 2002, **106**(20):2530-2.
- Rozen R, Brigant L, Apfelbaum M: Effects of cycles of food restriction followed by ad-libitum refeeding on body composition and energy-expenditure in obese rats. *Am J Clin Nutr* 1994, **59**:560-565.
- Huang BW, Chiang MT, Yao HT, Chiang W: The effect of high-fat and high-fructose diets on glucose tolerance and plasma lipid and leptin levels in rats. *Diabetes Obes Metab* 2004, **6**(2):120-6.
- Svensson AM, Hellerström C, Jansson L: Diet-induced obesity and pancreatic islet blood flow in the rat: a preferential increase in islet blood perfusion persists after withdrawal of the diet and normalization of body weight. *J Endocrinol* 1996, **151**(3):507-11.
- Prats E, Monfar M, Castellà J, Iglesias R, Alemany M: Energy intake of rats fed a cafeteria diet. *Physiol Behav* 1989, **45**(2):263-72.
- Dawson R, Pellemounter MA, Millard WJ, Liu S, Eppler B: Attenuation of leptin-mediated effects by monosodium glutamate-induced arcuate nucleus damage. *Am J Physiol* 1997, **36**:E202-E206.

41. Storlien LH, Jenkins AB, Chisholm DJ, Pascoe WS, Khouri S, Kraegen EW: **Influence of dietary fat composition on development of insulin resistance in rats. Relationship to muscle triglyceride and omega-3 fatty acids in muscle phospholipid.** *Diabetes* 1991, **40**:280-289.
42. Yaspelkis BB, Davis JR, Saberi M, Smith TL, Jazayeri R, Singh M, Fernandez V, Trevino B, Chinookoswong N, Wang J, Shi ZQ, Levin N: **Leptin administration improves skeletal muscle insulin responsiveness in diet-induced insulin-resistant rats.** *Am J Physiol* 2001, **280**:E130-E142.
43. Sahu A, Nguyen L, O'doherty RM: **Nutritional regulation of hypothalamic leptin receptor gene expression is defective in diet-induced obesity.** *J Neuroendocrinol* 2002, **14**:887-893.
44. Peiser C, McGregor GP, Lang RE: **Leptin receptor expression and suppressor of cytokine signaling transcript levels in high-fat-fed rats.** *Life Sci* 2000, **67**:2971-2981.
45. Munzberg H, Flier JS, Bjorbaek C: **Region-specific leptin resistance within the hypothalamus of diet-induced obese mice.** *Endocrinology* 2004, **145**:4880-4889.
46. Das UN: **Long-chain polyunsaturated fatty acids in the growth and development of the brain and memory.** *Nutrition* 2003, **19**(1):62-65.
47. Yu N, Martin JL, Stella N, Magistretti PJ: **Arachidonic acid stimulates glucose uptake in cerebral cortical astrocytes.** *Proc Natl Acad Sci USA* 1993, **90**(9):4042-4046.
48. Duelli R, Maurer MH, Staudt R, Heiland S, Duembgen L, Kuschinsky W: **Increased chronic cerebral glucose utilization and decreased glucose transporter Glut1 during hyperglycemia in rat brain.** *Brain Res* 2000, **858**:338-347.
49. Duckrow RB, Bryan RM: **Regional cerebral glucose utilization during hyperglycemia.** *J Neurochem* 1987, **48**:989-993.
50. Mans AM, DeJoseph MR, Davis DW, Hawkins RA: **Brain energy metabolism in streptozotocin-diabetes.** *Biochem J* 1988, **249**:57-62.
51. Jakobsen J, Nedergaard M, Aarslew-Jensen M, Diemer NH: **Regional brain glucose metabolism and blood flow in streptozocin-induced diabetic rats.** *Diabetes* 1990, **39**:437-440.
52. Havel PJ: **Peripheral signals conveying metabolic information to the brain: short-term and long-term regulation of food intake and energy homeostasis.** *Exp Biol Med* 2001, **226**:963-977.
53. Kusakabe T, Maeda M, Hoshi N, Sugino T, Watanabe K, Fukuda T, Suzuki T: **Fatty acid synthase is expressed mainly in adult hormone-sensitive cells or cells with high lipid metabolism and in proliferating fetal cells.** *J Histochem Cytochem* 2000, **48**:613-622.
54. Bukato G, Kochan Z, Swierczynski J: **Changes of malic enzyme activity in the developing rat brain are due to both the increase of mitochondrial protein content and the increase of specific activity.** *Int J Biochem* 1992, **24**:267-273.

doi:10.1186/1476-511X-10-168

Cite this article as: Estadella et al: A palatable hyperlipidic diet causes obesity and affects brain glucose metabolism in rats. *Lipids in Health and Disease* 2011 **10**:168.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

