

## Editorial - Wound recognition across the tree of life

Martin Heil<sup>1\*</sup>, Walter G. Land<sup>2</sup>, Mahmut Tör<sup>3</sup>

<sup>1</sup>Departamento de Ingeniería Genética, CINVESTAV - Irapuato, Mexico, <sup>2</sup>Faculty of Medicine, University of Strasbourg, France, <sup>3</sup>Institute of Science and the Environment, University of Worcester, United Kingdom

*Submitted to Journal:*  
Frontiers in Plant Science

*Specialty Section:*  
Plant Biotic Interactions

*ISSN:*  
1664-462X

*Article type:*  
Editorial Article

*Received on:*  
27 Jul 2016

*Accepted on:*  
17 Aug 2016

*Provisional PDF published on:*  
17 Aug 2016

*Frontiers website link:*  
[www.frontiersin.org](http://www.frontiersin.org)

*Citation:*  
Heil M, Land WG and Tör M(2016) Editorial - Wound recognition across the tree of life. *Front. Plant Sci.* 7:1319. doi:10.3389/fpls.2016.01319

*Copyright statement:*  
© 2016 Heil, Land and Tör. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

This Provisional PDF corresponds to the article as it appeared upon acceptance, after peer-review. Fully formatted PDF and full text (HTML) versions will be made available soon.

## **Editorial – Wound recognition across the tree of life**

**Martin Heil<sup>1\*</sup>, Walter G. Land<sup>2</sup> & Mahmut Tör<sup>3</sup>**

<sup>1</sup>Departamento de Ingeniería Genética, CINVESTAV-Irapuato, Irapuato, Guanajuato, México

<sup>2</sup>Molecular ImmunoRheumatology, INSERM UMR S1109, Laboratory of Excellence

Transplantex, Faculty of Medicine, University of Strasbourg, France

<sup>3</sup>Institute of Science and the Environment, University of Worcester, Worcester WR2 6AJ, UK.

**\*Correspondence:** Martin Heil; Fax: +51 (462) 623 9650; Phone: +52 (462) 623 9657;

[mheil@ira.cinvestav.mx](mailto:mheil@ira.cinvestav.mx)

10

All multi-cellular organisms share the necessity to perceive damage and to employ an adequate immune response to withstand injury and infection. The role of damage-associated molecular patterns (DAMPs) in the mammalian adaptive immune system and in allograft rejection was discovered by Polly Matzinger and Walter Land (Land et al., 1994; Matzinger, 1994). These discoveries revolutionised the research into transplantation and immunity (Land et al., 2016) and improved the understanding of chronic and inflammation-related diseases such as Alzheimer's disease, Diabetes, Lupus, Rheuma (Land, 2015a,b), and many forms of cancer (Land, 2015c; Candeias and GaipI, 2016). Unfortunately, the tendency towards specialisation in contemporary science, albeit allowing for an incredible increase in the efficiency at which knowledge is being generated, enhances the risk to lose the communication across disciplines. A prime example of this situation is the research into injury perception and immunity, which developed in distinct disciplines for mammals and plants. In consequence, the first application of the DAMPs concept to plants appeared 13 years after their first description for mammals (Lotze et al., 2007). Two years later, four review papers discussed the role of DAMPs and 'damaged-self recognition' in plants (Boller and Felix, 2009; Heil, 2009; Metraux et al., 2009; Tör et al., 2009).

In an attempt to close this gap, 'DAMPs 2016' the first international and trans-disciplinary congress on injury perception and immunity, aims at promoting the trans-

disciplinary research into wound recognition in organisms across the tree of life. A central  
30 step towards a better cross-disciplinary communication in this field was the Research Topic  
"Wound recognition across the tree of life". Eleven articles co-authored by 43 researchers  
were published between July and November 2014 and attracted over 55,000 views by now  
(<http://journal.frontiersin.org/researchtopic/2173/wound-recognition-across-the-tree-of-life>).  
Reviews summarised the functions of DAMPs in insects (Krautz et al., 2014) and plants  
(Savatin et al., 2014), applied the 'danger model' to mosquitoes (Moreno-García et al., 2014),  
and discussed the role of extracellular ATP (eATP) as a DAMP in plants (Tanaka et al.,  
2014). It was known before that eATP induces plant defence (Roux and Steinebrunner,  
2007; Chivasa et al., 2009; Heil et al., 2012), but only the discovery of its specific receptor  
(Choi et al., 2014) provided unambiguous support for a role of eATP as a DAMP (Tanaka et  
40 al., 2014). Interestingly, eATP also acts as DAMP in the fungus, *Trichoderma viride* (Medina-  
Castellanos et al., 2014).

Three papers reported how *Arabidopsis* responds to enemies with different degrees  
of specialisation and combined transcriptional with metabolomic data to distinguish  
responses to a chewing insect versus bacterial infection (Appel et al., 2014a,b; Rehrig et al.,  
2014). Responses to an aphid and a caterpillar shared only a surprising 10% of the up-  
regulated and 8% of the down-regulated genes, and even responses to caterpillars from  
different species (*Spodoptera exigua* and *Pieris rapae*) shared only 21% of the up-regulated  
and 12% of the down-regulated genes (Appel et al., 2014a). Transcriptional changes were  
frequently weaker or absent in response to the specialist (*P. rapae*) (Rehrig et al., 2014). The  
50 degree to which DAMPs contribute to this specificity remains subject of speculation (Duran-  
Flores and Heil, 2016), although responses of bean to leaf homogenates from various  
species demonstrated specificity when plants only perceive endogenous 'danger signals'  
(Duran-Flores and Heil, 2014). A fine-tuning of plant defences was also reported for  
methanol, a wound-generated molecule that functions in within- and between-plant signalling  
(Dorokhov et al., 2012; Komarova et al., 2014) and strongly modulated the response of

Arabidopsis and Tomato to pathogen-associated molecular patterns (PAMPs) (Hann et al., 2014).

Recent studies, mostly published after the Research Tropic, revealed multiple similarities between 'trained immunity' in mammals (Crisan et al., 2016) and resistance  
60 'priming' in plants (Martinez-Medina et al., 2016). Firstly, parasites of mammals, plants and insects frequently target the same processes to manipulate host immunity (Guiguet et al., 2016;Heil, 2016), and mammals, insects, plants and fungi respond to damage employing similar mechanisms (Hernández-Oñate and Herrera-Estrella, 2015). Secondly, preparing the immune system for more efficient responses ('priming') appears to be a general feature of DAMPs (Crisan et al., 2016;Martinez-Medina et al., 2016). The perception of DAMPs initiates the maturation of dendritic cells to antigen-presenting cells (Matzinger, 2002) and gene expression for the *NOD-Like Receptor family Protein 3* (NLRP3)-inflammasome in macrophages. Consecutive sensing of DAMPs or PAMPs by NLRP3 activates the inflammasome (Figure 4 in Heil and Land, 2014). Thus, DAMPs prime the immune system  
70 for more directed and sensitive responses to future problems. At least in plants, this effect depends on epigenetic alterations and can last into the next generation (Rasmann et al., 2012). Thirdly, many DAMPs exert a double-function as direct anti-microbial compound and signal. For example, mammalian type-I Interferons have antiviral effects, and plant secondary compounds such as DIMBOA, various glucosinolate breakdown products and herbivore-induced plant volatiles are quickly synthesised – or released from stored precursors – when plants, are damaged and have both, biocidal and signalling (immunity enhancing) activity (Gallucci and Matzinger, 2001;Ahmad et al., 2011;Andersson et al., 2015;Veyrat et al., 2016).

Finally, the development of ROS and the involvement of NADPH oxidase, Ca<sup>2+</sup>  
80 influxes and downstream MAPKinase signalling cascades are common features of DAMP-induced immune responses in organisms across the tree of life (Duran-Flores and Heil, 2014a,b;Medina-Castellanos et al., 2014;Crisan et al., 2016;Segal, 2016). An inhibitor of the mammalian NADPH oxidase inhibited ROS development in plants (Dwyer et al.,

1995;Tenhaken et al., 1995), anti-sera to key mammalian proteins cross-reacted with the respective plant proteins (Dwyer et al., 1995; Tenhaken et al., 1995), and the human DAMP, high mobility group box (HMGB) protein 3, activated immunity in plants (Choi et al., 2016). Immunological and pharmacological cross-reactions make homology likely in these cases. By contrast, the function of eATP as a DAMP in plants, fungi and mammals appears to be the result of independent evolution, because eATP receptors in mammals and plants belong  
90 to different families (Choi et al., 2014). In summary, we conclude that damaged-self recognition and the involved perception mechanisms and signalling pathways contain both homologous and analogous elements among plants and mammals (Heil and Land, 2014).

### **Author Contributions**

MH prepared a first draft of the manuscript and all authors listed have made substantial, direct and intellectual contribution to the work, and approved it for publication.

### **Conflict of Interest Statement**

100 The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

### **Acknowledgments**

MH is supported by a grant from Consejo Nacional de Ciencia y Tecnología de México (CONACyT, grant 212715), MT is supported by a grant from the UK Biotechnology and Biological Sciences Research Council (BBSRC grant BB/E02484X/1). We thank all contributors to this Research Topic and apologize to those whose work could not be discussed in great detail due to space limitations.

## References

- 110 Ahmad, S., Veyrat, N., Gordon-Weeks, R., Zhang, Y.H., Martin, J., Smart, L., Glauser, G., Erb, M., Flors, V., Frey, M., and Ton, J. (2011). Benzoxazinoid metabolites regulate innate immunity against aphids and fungi in maize. *Plant Physiol.* 157, 317-327. doi: 10.1104/pp.111.180224
- Andersson, M.X., Nilsson, A.K., Johansson, O.N., Boztas, G., Adolfsson, L.E., Pinosa, F., Petit, C.G., Aronsson, H., Mackey, D., Tor, M., Hamberg, M., and Ellerstrom, M. (2015). Involvement of the electrophilic isothiocyanate sulforaphane in *Arabidopsis* local defense responses. *Plant Physiol.* 167, 251-261. doi: 10.1104/pp.114.251892
- Appel, H.M., Fescemyer, H., Ehltng, J., Weston, D., Rehrig, E., Joshi, T., Xu, D., Bohlmann, J., and Schultz, J. (2014a). Transcriptional responses of *Arabidopsis thaliana* to chewing and sucking insect herbivores. *Front. Plant Sci.* 5, art. no. 565. doi: 120 10.3389/fpls.2014.00565
- Appel, H.M., Maqbool, S.B., Raina, S., Jagadeeswaran, G., Acharya, B.R., Hanley, J., Pickering, K.K., Hearne, L., Jones, A.D., Raina, R., and Schultz, J.C. (2014b). Transcriptional and metabolic signatures of *Arabidopsis* responses to chewing damage by an insect herbivore and bacterial infection and the consequences of their interaction. *Front. Plant Sci.* 5, art.no 441. doi: 10.3389/fpls.2014.00441
- Boller, T., and Felix, G. (2009). A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu. Rev. Plant Biol.* 60, 379-406. doi: doi:10.1146/annurev.arplant.57.032905.105346
- 130 Candeias, S., and Gaipf, U. (2016). The immune system in cancer prevention, development and therapy *Anti-Cancer Agents in Medical Chemistry* 16, 101-107 doi: 10.2174/1871520615666150824153523
- Chivasa, S., Murphy, A.M., Hamilton, J.M., Lindsey, K., Carr, J.P., and Slabas, A.R. (2009). Extracellular ATP is a regulator of pathogen defence in plants. *Plant J.* 60, 436-448. doi: 10.1111/j.1365-313X.2009.03968.x

- Choi, H.W., Manohar, M., Manosalva, P., Tian, M.Y., Moreau, M., and Klessig, D.F. (2016). Activation of plant innate immunity by extracellular high mobility group box 3 and its inhibition by salicylic acid. *PLoS Pathog.* 12. doi: 10.1371/journal.ppat.1005518
- Choi, J., Tanaka, K., Cao, Y., Qi, Y., Qiu, J., Liang, Y., Lee, S.Y., and Stacey, G. (2014). Identification of a plant receptor for extracellular ATP. *Science* 343, 290-294. doi: 140 10.1126/science.343.6168.290
- Crisan, T.O., Netea, M.G., and Joosten, L.a.B. (2016). Innate immune memory: Implications for host responses to damage-associated molecular patterns. *Eur. J. Immunol.* 46, 817-828. doi: 10.1002/eji.201545497
- DAMPs 2016. First international congress for the trans-disciplinary research into damage recognition from plants to humans. *Guanajuato, GTO, México*, Nov 6<sup>th</sup>-11<sup>th</sup> 2016.
- Dorokhov, Y.L., Komarova, T.V., Petrunia, I.V., Frolova, O.Y., Pozdyshev, D.V., and Gleba, Y.Y. (2012). Airborne signals from a wounded leaf facilitate viral spreading and induce antibacterial resistance in neighboring plants. *PLoS Pathog.* 8, e1002640. doi: 150 10.1371/journal.ppat.1002640
- Duran-Flores, D., and Heil, M. (2014). Damaged-self recognition in common bean (*Phaseolus vulgaris*) shows taxonomic specificity and depends on reactive oxygen species (ROS) signalling. *Front. Plant Sci.* 5, art. 585. doi: 10.3389/fpls.2014.00585
- Duran-Flores, D., and Heil, M. (2016). Sources of specificity in plant damaged-self recognition. *Curr. Opin. Plant Biol.* 32, 77-87. doi: 10.1016/j.pbi.2016.06.019
- Dwyer, S.C., Legendre, L., Low, P.S., and Leto, T.L. (1995). Plant and human neutrophil oxidative burst complexes contain immunologically related proteins. *Biochemical and Biophysical Acta - General Aspects* 1289, 231-237. doi: 10.1016/0304-4165(95)00156-5
- 160 Gallucci, S., and Matzinger, P. (2001). Danger signals: SOS to the immune system. *Curr. Opin. Immunol.* 13, 114-119. doi: 10.1016/S0952-7915(00)00191-6
- Guiguet, A., Dubreuil, G., Harris, M.O., Appel, H.M., Schultz, J.C., Pereira, M.H., and Giron, D. (2016). Shared weapons of blood- and plant-feeding insects: Surprising

- commonalities for manipulating hosts. *J. Insect Physiol.* 84, 4-21. doi:  
10.1016/j.jinsphys.2015.12.006
- Hann, C.T., Bequette, C.J., Dombrowski, J.E., and Stratmann, J.W. (2014). Methanol and ethanol modulate responses to danger- and microbe-associated molecular patterns. *Front. Plant Sci.* 5, art. no 550. doi: 10.3389/fpls.2014.00550
- 170 Heil, M. (2009). Damaged-self recognition in plant herbivore defence. *Trends Plant Sci.* 14, 356-363. doi: 10.1016/j.tplants.2009.04.002
- Heil, M. (2016). Host manipulation by parasites: cases, patterns and remaining doubts. *Frontiers Ecol. Evol.*, art. no 80. doi: 10.3389/fevo.2016.00080
- Heil, M., Ibarra-Laclette, E., Adame-Álvarez, R.M., Martínez, O., Ramirez-Chávez, E., Molina-Torres, J., and Herrera-Estrella, L. (2012). How plants sense wounds: damaged-self recognition is based on plant-derived elicitors and induces octadecanoid signaling. *PLoS ONE* 7, e30537. doi: 10.1371/journal.pone.003053
- Heil, M., and Land, W.G. (2014). Danger signals - damaged-self recognition across the tree of life. *Front. Plant Sci.* 5, art. 579. doi: 10.3389/fpls.2014.00578
- 180 Hernández-Oñate, M.A., and Herrera-Estrella, A. (2015). Damage response involves mechanisms conserved across plants, animals and fungi. *Curr. Genet.* 61, 359-372 doi: 10.1007/s00294-014-0467-5
- Komarova, T.V., Sheshukova, E.V., and Dorokhov, Y.L. (2014). Cell wall methanol as a signal in plant immunity. *Front. Plant Sci.* 5, art. 101. doi: 10.3389/fpls.2014.00101
- Krautz, R., Arefin, B., and Theopold, U. (2014). Damage signals in the insect immune response. *Front. Plant Sci.* 5, art. no 342. doi: 10.3389/fpls.2014.00342
- Land, W., Schneeberger, H., Schleibner, S., Illner, W.D., Abendroth, D., Rutili, G., and Al, E. (1994). The beneficial effect of human recombinant superoxide dismutase on acute and chronic rejection events in recipients of cadaveric renal transplants. *Transplantation* 57, 211-217.
- 190 Land, W.G. (2015a). How evolution tells us to induce allotolerance. *Exp. Clin. Transplant.* 13, 46-54. doi: 10.6002/ect.mesot2014.L69



- Land, W.G. (2015b). The role of damage-associated molecular patterns in human diseases: part I - promoting inflammation and immunity. *SQU Med. J.* 15, e9-e21.
- Land, W.G. (2015c). The role of damage-associated molecular patterns (DAMPs) in human diseases: Part II: DAMPs as diagnostics, prognostics and therapeutics in clinical medicine. *SQU Med. J.* 15, e157-e170.
- Land, W.G., Agostinis, P., Gasser, S., Garg, A.D., and Linkermann, A. (2016). Transplantation and damage associated molecular patterns (DAMPs). *Am. J. Transplant.*, n/a-n/a. doi: 10.1111/ajt.13963
- 200 Lotze, M.T., Zeh, H.J., Rubartelli, A., Sparvero, L.J., Amoscato, A.A., Washburn, N.R., Devera, M.E., Liang, X., Tor, M., and Billiar, T. (2007). The grateful dead: damage-associated molecular pattern molecules and reduction/oxidation regulate immunity. *Immunol. Rev.* 220, 60-81. doi: 10.1111/j.1600-065X.2007.00579.x
- Martinez-Medina, A., Flors, V., Heil, M., Mauch-Mani, B., Pieterse, C.M.J., Pozo, M.J., Ton, J., Van Dam, N.M., and Conrath, U. (2016). Recognizing plant defense priming. *Trends Plant Sci.* publ. online. doi: 10.1016/j.tplants.2016.07.009
- Matzinger, P. (1994). Tolerance, danger, and the extended family. *Annu. Rev. Immunol.* 12, 991-1045. doi: doi:10.1146/annurev.iy.12.040194.005015
- Matzinger, P. (2002). The danger model: A renewed sense of self. *Science* 296, 301-305. doi: 10.1126/science.1071059
- 210 Medina-Castellanos, E., Esquivel-Naranjo, E.U., Heil, M., and Herrera-Estrella, A. (2014). Extracellular ATP activates MAPK and ROS signaling during injury response in the fungus *Trichoderma atroviride*. *Front. Plant Sci.* 5, art.no. 659. doi: 10.3389/fpls.2014.00659
- Metraux, J.P., Jackson, R.W., Schnettler, E., and Goldbach, R.W. (2009). "Plant pathogens as suppressors of host defense," in *Plant Innate Immunity*, ed. L.C. Van Loon. (London: Academic Press Ltd-Elsevier Science Ltd), 39-89.

- Moreno-García, M., Recio-Tótoro, B., Claudio-Piedras, F., and Lanz-Mendoza, H. (2014). Injury and immune response: applying the danger theory to mosquitoes. *Front. Plant Sci.* 5, art.no. 451. doi: 10.3389/fpls.2014.00451
- 220
- Rasmann, S., De Vos, M., Casteel, C.L., Tian, D., Halitschke, R., Sun, J.Y., Agrawal, A.A., Felton, G.W., and Jander, G. (2012). Herbivory in the previous generation primes plants for enhanced insect resistance. *Plant Physiol.* 158, 854-863. doi: 10.1104/pp.111.187831
- Rehrig, E.M., Appel, H.M., Jones, A.D., and Schultz, J.C. (2014). Roles for jasmonate- and ethylene- induced transcription factors in the ability of Arabidopsis to respond differentially to damage caused by two insect herbivores. *Front. Plant Sci.* 5, art. no 407. doi: 10.3389/fpls.2014.00407
- Roux, S.J., and Steinebrunner, I. (2007). Extracellular ATP: an unexpected role as a signaler in plants. *Trends Plant Sci.* 12, 522-527. doi: 10.1016/j.tplants.2007.09.003
- 230
- Savatin, D.V., Gramegna, G., Modesti, V., and Cervone, F. (2014). Wounding in the plant tissue: the defense of a dangerous passage. *Front. Plant Sci.* 5, art. no 470. doi: 10.3389/fpls.2014.00470
- Segal, A.W. (2016). NADPH oxidases as electrochemical generators to produce ion fluxes and turgor in fungi, plants and humans. *Open Biol.* 6, art.no. 160028. doi: 10.1098/rsob.160028
- Tanaka, K., Choi, J., Cao, Y., and Stacey, G. (2014). Extracellular ATP acts as a damage-associated molecular pattern (DAMP) signal in plants. *Front. Plant Sci.* 5, art. no. 446. doi: 10.3389/fpls.2014.00446
- 240
- Tenhaken, R., Levine, A., Brisson, L.F., Dixon, R.A., and Lamb, C. (1995). Function of the oxidative burst in hypersensitive disease resistance. *Proc. Natl. Acad. Sci. U.S.A.* 92, 4158-4163. doi:
- Tör, M., Lotze, M.T., and Holton, N. (2009). Receptor-mediated signalling in plants: molecular patterns and programmes. *J. Exp. Bot.* 60, 3645-3654. doi: 10.1093/jxb/erp233

Veyrat, N., Robert, C.a.M., Turlings, T.C.J., and Erb, M. (2016). Herbivore intoxication as a potential primary function of an inducible volatile plant signal. *J. Ecol.* 104, 591-600.  
doi: 10.1111/1365-2745.12526

250

Provisional