



This is , post-print (final draft) version of the following published document and is licensed under All Rights Reserved license:

**Fryer, Simon M and Giles, David and Garrido, Inmaculada and Stone, Keeron J and de la Puerta, Alex and España-Romero, Vanesa (2016) Cardiorespiratory and hemodynamic predictors of sport rock climbing performance (Abstract Only). In: British Association of Sport and Exercise Sciences, 29 - 30 November 2016, Nottingham.**

Official URL: <https://www.nutrition-society.org/events/british-association-sport-and-exercise-sciences-conference-2016>

EPrint URI: <http://eprints.glos.ac.uk/id/eprint/4245>

#### **Disclaimer**

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

## **Title: Cardio-respiratory and Hemodynamic Predictors of Sport Rock Climbing Performance**

**Simon Fryer<sup>1</sup>; David Giles<sup>2</sup>; Inmaculada Garrido<sup>3</sup>; Keeron Stone<sup>1</sup>; Alex de la Puerta<sup>4</sup>; Vanesa España Romero<sup>3</sup>**

Rock climbing performance has been suggested to involve a notable contribution from the aerobic metabolism. Previously forearm oxygenation kinetics has been used to distinguish between ability groups and predict the performance of sport rock climbers. Currently it is not known if forearm oxygenation kinetics, or a sport specific assessment of cardiorespiratory fitness best predicts sport rock climbing performance. Therefore the aim of this study was to determine whether forearm oxidative capacity index, maximal de-oxygenation of the flexor digitorum profundus (FDP), treadmill  $\dot{V}O_{2peak}$ , or running  $\dot{V}O_{2peak}$  best predicts red-point climbing performance. **Methods:** Following institutional ethical approval, twenty-nine sport rock climbers (age  $34.5 \pm 6.6$  yrs; height  $1.71 \pm 0.07$  m; mass  $64 \pm 8.2$  kg and body fat  $17.2 \pm 7.6$ %) visited the laboratory on two separate occasions within 7 days. During visit one participants self-reported their indoor sport rock climbing ability, previously shown to be a valid measure, and completed a treadmill  $\dot{V}O_{2peak}$  test during which maximal de-oxygenation of the FDP was assessed using near infrared spectroscopy (NIRS). During the second visit oxidative capacity index of the FDP was determined using NIRS, and this was followed by a running  $\dot{V}O_{2peak}$  test. **Results:** Linear regression, adjusted for age, sex and body fat percentage revealed that forearm oxidative capacity index, treadmill maximal de-oxygenation and treadmill  $\dot{V}O_{2peak}$  all significantly predicted self-reported red-point sport climbing ability ( $R^2 = 0.766; 0.538; 0.495$  respectively), whereas running  $\dot{V}O_{2max}$  did not ( $R^2 = 0.215$ ). **Conclusions:** Sport rock climbers should consider additional sport specific aerobic training rather than traditional aerobic exercise such as running to improve climbing performance.