

O'Donnell, C., Workman, H., Hosking, S. L. & Huntjens, B. (2007). Diurnal Refractive Error Fluctuations in Diabetic and Control Subjects. Paper presented at the Association for Research in Vision and Ophthalmology (ARVO) 2007 Annual Meeting, 06-05-2007 - 11-05-2007, Fort Lauderdale, USA.



**CITY UNIVERSITY
LONDON**

[City Research Online](#)

Original citation: O'Donnell, C., Workman, H., Hosking, S. L. & Huntjens, B. (2007). Diurnal Refractive Error Fluctuations in Diabetic and Control Subjects. Paper presented at the Association for Research in Vision and Ophthalmology (ARVO) 2007 Annual Meeting, 06-05-2007 - 11-05-2007, Fort Lauderdale, USA.

Permanent City Research Online URL: <http://openaccess.city.ac.uk/5603/>

Copyright & reuse

City University London has developed City Research Online so that its users may access the research outputs of City University London's staff. Copyright © and Moral Rights for this paper are retained by the individual author(s) and/ or other copyright holders. All material in City Research Online is checked for eligibility for copyright before being made available in the live archive. URLs from City Research Online may be freely distributed and linked to from other web pages.

Versions of research

The version in City Research Online may differ from the final published version. Users are advised to check the Permanent City Research Online URL above for the status of the paper.

Enquiries

If you have any enquiries about any aspect of City Research Online, or if you wish to make contact with the author(s) of this paper, please email the team at publications@city.ac.uk.

ABSTRACT ARVO

AUTHORS

Clare O'Donnell
Helena Workman
Sarah Hosking
Byki Huntjens

TITLE

Diurnal refractive error fluctuations in diabetic and control subjects

Purpose:

Refractive error has been shown to fluctuate in poorly controlled diabetic patients. The purpose of this study was to measure acute diurnal fluctuations in refractive error and blood glucose levels (BGLs) in diabetic and control subjects.

Methods:

Twenty-one type 2 diabetic subjects (age 56 ± 11 years), 20 type 1 diabetic subjects (age 38 ± 15 years) and 20 non-diabetic controls (age 49 ± 23 years) took part in the study. Refractive error was measured with an OPD ARK-10000 autorefractometer (Nidek) and BGLs were measured using a finger stick test (Hemocue). All measurements were taken between 8AM and 8PM at approximately two-hourly intervals. Using power vector analysis the variability in refractive error was mapped against time of day and related to BGLs, HbA1c, diabetic status and duration of disease.

Results:

Refractive error did not fluctuate significantly during the day in any of the three groups (ANOVA $p > 0.05$). The mean \pm SD values for BGLs during the day were 10.4 ± 4.40 mM/l in DM type 2, 10.3 ± 5.30 mM/l in DM type 1, and 5.4 ± 1.04 mM/l in control subjects. BGLs changed significantly during the day and between groups (ANOVA $p < 0.0005$). The correlations between refractive error components (mean sphere, J180 and J45) and BGLs did not reach statistical significance (ANOVA $p > 0.05$). Multiple regression analysis showed that neither BGLs, HbA1c, diabetic status, nor duration of disease had a significant effect on diurnal refractive error measurements. ($p > 0.05$).

Conclusions:

Diurnal changes in BGLs do not result in significant acute refractive error fluctuations in diabetic patients, as measured with an autorefractometer.

Acknowledgements: Supported by a PhD studentship from Lein Applied Diagnostics Ltd.