

Henry, K. M., Reyes-Aldasoro, C. C. & Renshaw, S. A. (2012). Exploring neutrophil behaviour in a zebrafish model of inflammation through the generation of novel parameters using MatLab algorithms. Paper presented at the European Congress of Immunology, 05-09-2012 - 08-09-2012, Glasgow, UK.



**CITY UNIVERSITY  
LONDON**

[City Research Online](#)

**Original citation:** Henry, K. M., Reyes-Aldasoro, C. C. & Renshaw, S. A. (2012). Exploring neutrophil behaviour in a zebrafish model of inflammation through the generation of novel parameters using MatLab algorithms. Paper presented at the European Congress of Immunology, 05-09-2012 - 08-09-2012, Glasgow, UK.

**Permanent City Research Online URL:** <http://openaccess.city.ac.uk/5763/>

### **Copyright & reuse**

City University London has developed City Research Online so that its users may access the research outputs of City University London's staff. Copyright © and Moral Rights for this paper are retained by the individual author(s) and/ or other copyright holders. All material in City Research Online is checked for eligibility for copyright before being made available in the live archive. URLs from City Research Online may be freely distributed and linked to from other web pages.

### **Versions of research**

The version in City Research Online may differ from the final published version. Users are advised to check the Permanent City Research Online URL above for the status of the paper.

### **Enquiries**

If you have any enquiries about any aspect of City Research Online, or if you wish to make contact with the author(s) of this paper, please email the team at [publications@city.ac.uk](mailto:publications@city.ac.uk).

Exploring neutrophil behaviour in a zebrafish model of inflammation through the generation of novel parameters using MatLab algorithms

K. M. Henry,\* S. A. Renshaw\* & C. C. Reyes-Aldasoro

\*Centre for Developmental and Biomedical Genetics, University of Sheffield, Sheffield, UK, Biomedical Engineering School of Engineering and Informatics, University of Sussex, Sussex, UK

**Purpose/Objective:** Tracking of immune cells is key to understanding their behaviour during inflammation. Current software available for tracking of immune cells is limited. The aim of this study was to develop a MatLab package of segmentation and tracking algorithms to apply in the tracking of neutrophils, fluorescently labelled in our zebrafish model [1]. The development of algorithms in MatLab allows us to explore parameters not available in other software packages such as directionality of neutrophil movement and neutrophil behaviour inside and outside of a wound region.

**Materials and methods:** Tail fin transection was performed on Tg(mpx:GFP) zebrafish (3 dpf) which were imaged on a spinning disk confocal from 1 h post injury (hpi) to 7 hpi. Images were exported from Velocity<sup>TM</sup> and analysed using MatLab m-files written for the tracking of immune cells. This is a fully automated analysis, after the user defines the initial thresholds based on fluorescent intensity of the images.

**Results:** Neutrophils from injured embryos had a lower meandering ratio and a greater speed than neutrophils tracked in uninjured embryos (meandering ratio  $0.24 \pm 0.03$  versus  $0.42 \pm 0.05$ ,  $P = 0.003$ ; speed  $4.03 \pm 0.32$  versus  $1.31 \pm 0.21$  pixels/frame,  $P < 0.0001$ ,  $n = 3$ ), however there was no significant difference in total distance travelled ( $166.9 \pm 17.4$  versus  $210.1 \pm 30.28$  pixels,  $P > 0.05$ ). In injured embryos with a defined wound region, the oriented velocity towards the wound was  $0.31 \pm 0.24$  pixels/frame. Once within the wound region, the oriented velocity of neutrophil tracks was  $-0.39 \pm 0.32$  pixels/frame; indicating that while the neutrophils travel at a similar speed, they are now travelling away from the wound. The 'in wound ratio' was  $0.91 \pm 0.04$ , indicating that once neutrophils enter the wound region they tend to stay, in the timeframe studied. The 'leave wound ratio' was  $0.37 \pm 0.03$ , a measure of the rate at which neutrophils move away from the site of injury once they have entered the wound region.

**Conclusions:** Using these algorithms, we can analyse the behaviour of immune cells in a more detailed way. In addition to previously available parameters such as meandering ratio and speed, more complex parameters such as velocity towards or away from a wound region and a measurement of how neutrophils behave while in a wound region are available. Combining this novel tracking technology with established assays in our laboratory will enable the further dissection of neutrophil fate following an inflammatory stimulus.

[1] K. M. Henry, L. Pase, C. F. Ramos-Lopez, G. J. Lieschke, S. A. Renshaw, and C. C. Reyes-Aldasoro, 'PhagoSight: an open-source MATLAB® package for the analysis of fluorescent neutrophil and macrophage migration in a zebrafish model', *PloS One*, vol. 8, no. 8, p. e72636, 2013.