

Functionalised Organocuprates: The Structures of Lithium and Magnesium-Grignard 2-Methoxyphenylcuprates

Roberta Bomparola, Robert P. Davies*, Steven Lal, Andrew J. P. White

*Department of Chemistry, Imperial College London, South Kensington,
London, UK SW7 2AZ.*

E-mail: r.davies@imperial.ac.uk;

Tel: +44 207 5945754

Abstract

Lithium and magnesium-Grignard diorganocuprates incorporating the functionalised aryl group 2-methoxyphenyl have been prepared and structurally characterised in the solid state. $[\text{Cu}_4\text{Li}_2(\text{C}_6\text{H}_4\text{OMe-2})_6(\text{THF})_2]$, **2**, and $[\text{Cu}(\text{C}_6\text{H}_4\text{OCH}_3\text{-2})_2\text{Mg}(\text{THF})_2\text{X}]$ (X = Cl, Br), **3-X**, all exhibit coordination of the s-block metal centre by the methoxy oxygen, resulting in the formation of novel aggregates and favouring contact ion pair structures. In contrast separate ion pair structures had previously been observed under similar conditions for non-functionalised arylcuprates. The magnesium organocuprates **3-Cl** and **3-Br** are of particular interest being rare examples of structurally characterised Grignard-derived organocuprates, and the first examples of functionalised Grignard organocuprates. All reported organocuprates undergo oxidative aryl coupling in the presence of O_2 or PhNO_2 to give 2,2'-dimethoxybiphenyl.

Introduction

Organocuprates are excellent reagents for the formation of carbon-carbon bonds and have been extensively employed in synthetic methodology since the 1960s for a number of important reactions including conjugate addition, substitution reactions, coupling reactions, and carbocupration.^{1,2} However, it is only relatively recently that significant insights have been obtained into the mechanism of operation of these reagents and the origin of their unique reactivity,³⁻⁵ including recent evidence of the important role played by Cu(III) intermediates.⁶




Figure 1. CIP (**I**) and SSIP (**II**) structures of lithium homocuprates

One of the first key steps in building current understanding of organocuprates was identification of the structural forms adopted by these complex species.³ Thus the resting state and also reactive form of lithium organocuprates (“R₂CuLi”, R = organo group) in non-polar or weakly coordinating solvents such as diethyl ether is now commonly accepted to be a dimeric contact ion pair (CIP) species **I** (Figure 1). However in more strongly coordinating solvents, such as THF, a less reactive solvent separated ion pair (SSIP) species **II** is predominant. Studies on the structures of analogous magnesium-Grignard derived organocuprates (“R₂CuMgX”, X = halide) are far less developed, despite the fact the heritage

and synthetic usage of these reagents rivals that of their lithium based analogs.¹ We have recently reported the first structural characterizations of magnesium-Grignard organocuprates, revealing $[\text{Ph}_4\text{Cu}_2(\text{Mg}(\text{OEt}_2)\text{I})_2]$ to adopt a dimeric CIP structure in diethyl ether which is analogous to lithium diorganocuprate CIPs (**I**) but with MgI formally replacing Li in the ring.⁷ Larger “inverse crown” type aggregates⁸ were observed when employing the bulkier mesityl ($\text{C}_6\text{H}_2\text{Me}_3$ -2,4,6) aryl group, and similar to lithium organocuprates solvent separate species were discerned when more strongly coordinating THF solvent molecules were present.⁷

Most structural studies on organocuprates to date have employed simple non-functionalised aryls (such as phenyl⁹⁻¹⁴ or mesityl¹⁵⁻¹⁷) as the organo R group. However, many organocuprate synthetic protocols employ aryl or alkyl groups with additional functionalization such as amine, ether, alkenyl or alkynyl groups¹⁸ and it is still unclear how closely the structures and behavior of these species match those of their unfunctionalised cousins. A small number of lithium organocuprates containing dimethylamino functionalized aryl groups have been studied, including the lithium homocuprate $[\text{Cu}_2\text{Li}_2(\text{C}_6\text{H}_4\text{CH}_2\text{NMe}_2)_4]$ **III** which was shown to adopt a CIP structure with dimethylamino substituents coordinating to the lithium atoms via their nitrogen lone pairs (Figure 2).¹⁹ An additional advantage to studying such species is the demonstrated ability of intramolecular Lewis donor groups to stabilize reagents or aggregates which are otherwise too reactive or unstable to be isolated. Hence the aforementioned dimethylamino substituted aryl was also employed by van Koten and co-workers to prepare $[(\text{C}_6\text{H}_4(\text{CH}_2\text{NMe}_2)_2)_2\text{CuLi}_2(\text{CN})(\text{THF})_4]_\infty$, **IV**, which proved key in understanding the chemistry of cyano-Gilman (Lipshultz) cuprates at a time when there was much controversy regarding the nature of these species.²⁰ In addition, a diamine functionalized aryl was thought to play a large role in contributing to the

thermodynamic stability of the lithium diorganocuprate-lithium halide complex

$[\text{Cu}(\text{C}_6\text{H}_4(\text{CH}_2\text{N}(\text{Et})\text{CH}_2\text{CH}_2\text{NEt}_2)-2)_2\text{Li}_2\text{Br}]$, **V**.²¹ Using similar principles Ribas *et al.*

employed a triaza-macrocyclic ligand to yield the first example of an isolable Cu(III) mono-aryl species.²²




Figure 2. Organocuprates containing amine functionalised aryl groups

In this work we investigate the role of ether functionalized aryl groups in lithium and Grignard organocuprates. There are currently no structurally characterized examples of Grignard-derived organocuprates with functionalized aryls of any sort, and as discussed above studies on functionalized lithium diarylcuprates are currently limited to just those with one or more amine donor groups. The 2-methoxyphenyl group has been selected as the organo group since organocuprates containing this ligand are readily accessible and have previously been employed in a number of synthetic protocols including the syntheses of dihydromultifidene,²³ hallucinogenic amphetamine derivatives,²⁴ and the anti-cancer natural product yomogin.²⁵

Experimental Section

General Considerations

All experimental work was carried out under an inert atmosphere of nitrogen using standard Schlenk double manifold and glovebox techniques. Purification and drying of the solvents was carried out following standard methods or using an Innovative Technologies PureSolv Solvent Purification System with purification grade solvents. NMR spectra were recorded on a Bruker DPX400 spectrometer with internal standards. Melting points were measured in capillaries sealed under nitrogen and microanalytical data were obtained from the Science Technical Support Unit, London Metropolitan University.

Synthesis of $[\text{Cu}(\text{C}_6\text{H}_4\text{OMe-2})]_8 \cdot 8\text{THF}$ (1)

40 mmol of 2-bromoanisole (5.0 ml) were added drop-wise to a suspension of magnesium metal (44 mmol, 1.07 g) in THF (40 ml) at 0 °C. After complete addition the reaction was brought to room temperature and stirred for 1 h. The resultant Grignard reagent was then added to a suspension of copper(I) chloride (40 mmol, 3.94 g) in THF (40 ml) at 0 °C and the mixture was stirred overnight. 20 ml of 1,4-dioxane were added to aid the precipitation of MgBrCl, which was removed by filtration and the filtrate volume was concentrated under vacuum. Storage at -35 °C for 3 days yielded orange crystals (2.58 g, 38% yield relative to 2-bromoanisole); m.p. 128 °C (decomp.). $^1\text{H-NMR}$ (400 MHz, 25 °C, C_6D_6): δ = 9.2 - 6.1 (m, 32 H, Ar-H), 3.70 (m, 32 H, $\text{CH}_2\text{O-THF}$), 3.5 - 2.1 (s, 24 H, $-\text{OCH}_3$), 1.52 (m, 32 H, $\text{CH}_2\text{-THF}$). Elemental analysis for $\text{C}_{56}\text{H}_{56}\text{Cu}_8\text{O}_8$ (FW = 1365.3): Calculated, %: C = 49.26, H = 4.13. Found, %: C = 49.15; H = 4.19 (note that THF solvent of crystallisation was lost during the isolation procedure required to prepare the sample for analysis).

Synthesis of $[\text{Cu}_4\text{Li}_2(\text{C}_6\text{H}_4\text{OMe-2})_6(\text{THF})_2]$ (**2**)

A solution of *n*BuLi in hexane/cyclohexane (2.5 M, 1.06 mmol, 0.43 ml) was added dropwise to a solution of 2-bromoanisole (1.06 mmol, 0.13 ml) in hexane (5 ml) at 0 °C. The reaction mixture was allowed to warm to room temperature and stirred for 1 h. A solution of 2-methoxyphenylcopper (**1**) (0.182 g, 1.06 mmol) in toluene (4.6 ml) was added to give a yellow precipitate, which was dissolved with the addition of THF (0.5 ml) and then filtered over celite. Storage at room temperature for 5 days yielded yellow blocks of **2** suitable for X-ray crystallography (85 mg, 24% yield based on 2-bromoanisole). $^1\text{H-NMR}$ (400 MHz, 25 °C, C_6D_6): δ = 8.02 (m, 6H, H6), 7.23 (m, 6H, H4), 7.05 (m, 6H, H5), 6.69 (m, 6H, H3), 3.68 (m, 8 H, $\text{CH}_2\text{O-THF}$), 3.40 (s, 18H, $-\text{OCH}_3$), 1.49 (m, 8 H, $\text{CH}_2\text{-THF}$). $^7\text{Li NMR}$ (155.6 MHz, 25 °C, C_6D_6) δ = 1.46.

Exposure of **2** to atmospheric oxygen over several days or treatment with PhNO_2 at room temperature produced 2,2'-dimethoxybiphenyl. $^1\text{H NMR}$ (400 MHz, 25 °C, C_6D_6) δ = 7.49 (d, 2H, J = 7.4 Hz, H6); 7.28 (m, 2H, H4); 7.06 (m, 2H, H5); 6.78 (d, 2H, J = 8 Hz, H3); 3.39 (s, 6H, $-\text{OCH}_3$).

Synthesis of $[\text{Cu}(\text{C}_6\text{H}_4\text{OCH}_3\text{-2})_2\text{Mg}(\text{THF})_2\text{Br}]$ (**3-Br**)

A solution of 2-methoxyphenylcopper (**1**) (182 mg, 1.10 mmol) in toluene (2.6 ml) was added to a solution of (2-methoxyphenyl)magnesium bromide in THF (1 M, 1.10 ml, 1.10 mmol) - previously prepared from the addition of 2-bromoanisole (20 mmol, 2.5 ml) to a suspension of magnesium metal (22 mmol, 0.48 g) in THF (20 ml). In order to dissolve the yellow precipitate formed additional toluene (2 ml) and THF (2 ml) were added; the solution was stirred at room temperature for 15 min before filtration through celite. Storage at -35 °C for 19 days yielded colourless crystals suitable for X-ray analysis (142 mg, 25 %); m.p. 282 °C (decomp.). $^1\text{H NMR}$ (400 MHz, C_6D_6 , 25 °C): δ = 7.48 (d, 2H, J = 8.0 Hz, H6), 7.22 (m,

2H, H4), 7.05 (m, 2H, H5), 6.93 (m, 2H, H3), 3.82 (m, 8H, CH₂O-THF), 3.39 (s, 6H, -OCH₃), 1.45 (m, 8H, CH₂-THF).

Synthesis of [Cu(C₆H₄OCH₃-2)₂Mg(THF)₂Cl] (3-Cl)

A solution of 2-methoxyphenylcopper (**1**) (182 mg, 1.10 mmol) in toluene (2.6 ml) was added to a solution of (2-methoxyphenyl)magnesium chloride in THF (1 M, 1.10 ml, 1.10 mmol) - previously prepared from the addition of 2-chloroanisole (20 mmol, 2.5 ml) to a suspension of magnesium metal (22 mmol, 0.48 g) in THF (20 ml). Toluene (2 ml) and THF (2 ml) were added to dissolve the solid and the solution was stirred at room temperature for 15 min before filtration through celite. Storage at room temperature for 14 days yielded colourless crystals suitable for X-ray analysis (98 mg, 19 %); ¹H NMR (400 MHz, C₆D₆, 25 °C): δ = 7.48 (d, 2H, J = 7.2 Hz, H6), 7.22 (m, 2H, H4), 7.05 (m, 2H, H5), 6.93 (m, 2H, H3), 3.69 (m, 8H, CH₂O-THF), 3.39 (s, 6H, -OCH₃), 1.52 (m, 8H, CH₂-THF).

Synthesis of [Cu(C₆H₄OCH₃-2)₂Mg(THF)₂I] (3-I)

A solution of 2-methoxyphenylcopper (182 mg, 1.10 mmol) in toluene (2.6 ml) was added to a solution of (2-methoxyphenyl)magnesium iodide in THF (0.5 M, 2.20 ml, 1.10 mmol) - previously prepared from the addition of 2-iodoanisole (20 mmol, 2.6 ml) to a suspension of magnesium metal (22 mmol, 0.48 g) in THF (20 ml). Toluene (2 ml) and THF (2 ml) were added and the mixture was heated to 50 °C before filtration through celite. Storage of the filtrate at room temperature for 12 h yielded a yellow solid, which was dried *in vacuo* (95 mg, yield = 16%). ¹H-NMR (400 MHz, C₆D₆, 25 °C): δ = 7.48 (d, 2H, J = 7.6 Hz, H6), 7.22 (m, 2H, H4); 7.05 (m, 2H, H5), 6.93 (m, 2H, H4), 3.71 (m, 8H, CH₂O-THF), 3.39 (s, 6H, -OCH₃), 1.48 (m, 8H, CH₂-THF).

X-ray Structure Determinations of **1**, **2**, **3-Cl** and **3-Br**

The crystals were all taken directly from the mother liquor, covered with a perfluorinated ether, and mounted on the top of a glass capillary under a flow of cold gaseous nitrogen. The data were collected using a Oxford Diffraction Xcalibur PX Ultra diffractometer fitted with an Oxford Cryostream low-temperature device, and the structures were refined based on F^2 using the SHLEXTL and SHELX-97 program systems. Table 2 provides a summary of the crystallographic data for all compounds. Full details of the X-ray structure solutions, including the handling of any disorder present in the structures, is given in the Supporting Information. The absolute structure of **3-Br** was determined by a combination of R -factor tests [$R_1^+ = 0.0163$, $R_1^- = 0.0294$] and by use of the Flack parameter [$\chi^+ = 0.000(10)$, $\chi^- = 1.018(10)$]. Similarly for **3-Cl** $R_1^+ = 0.0610$, $R_1^- = 0.0687$ and $\chi^+ = 0.00(4)$, $\chi^- = 1.00(4)$. The crystal structure data have been deposited with the Cambridge Crystallographic Data Center under deposition numbers CCDC 882792 (**1**), 882793 (**2**), 882794 (**3-Br**) and 882795 (**3-Cl**). This material can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, by emailing data_request@ccdc.cam.ac.uk, or by contacting the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB2 1EZ, UK.

Table 1. Summary of Crystal Structure Data for **1**, **2**, **3-Br** and **3-Cl**

	1	2	3-Br	3-Cl
empirical formula	C ₅₆ H ₅₆ Cu ₈ O ₈ · 8(C ₄ H ₈ O)	C ₅₀ H ₅₈ Cu ₄ Li ₂ O ₈	C ₂₂ H ₃₀ BrCu MgO ₄	C ₂₂ H ₃₀ ClCu MgO ₄
fw	1942.16	1055.00	526.22	481.76
temperature (K)	173	173	173	173
wavelength	Cu K α 1.54184 Å	Mo K α 0.71073 Å	Cu K α 1.54184 Å	Cu K α 1.54184 Å
crystal system	tetragonal	orthorhombic	orthorhombic	orthorhombic
space group	<i>P4/n</i> (no. 85)	<i>Pbca</i> (no. 61)	<i>P2₁2₁2₁</i> (no. 19)	<i>P2₁2₁2₁</i> (no. 19)
<i>a</i> (Å)	18.15205(9)	17.96778(17)	8.18331(6)	7.9461(15)
<i>b</i> (Å)	-	17.60950(18)	8.43100(6)	8.5104(7)
<i>c</i> (Å)	13.48120(9)	30.1255(4)	33.1182(2)	33.255(3)
<i>V</i> (Å ³)	4442.01(4)	9531.82(18)	2284.94(3)	2248.9(5)
<i>Z</i>	2	8	4	4
ρ_{calc} (gcm ⁻³)	1.452	1.470	1.530	1.423
θ range (deg)	3.44-72.46	3.72-32.46	2.67-71.32	2.66-72.30
μ (mm ⁻¹)	2.549	1.813	3.868	2.941
reflns collected	40837	119452	31950	5720
reflns unique (<i>R</i> _{int})	4398 (0.0345)	15909 (0.0494)	4480 (0.0217)	3779 (0.0391)
parameters	165	617	282	264
goodness-of-fit on <i>F</i> ²	1.148	1.131	1.101	1.179
<i>R</i> ₁ [<i>F</i> >4 σ (<i>F</i>)]	0.0577	0.0563	0.0163	0.0610
<i>wR</i> ₂	0.2240	0.1410	0.0434	0.1467

Results and Discussion


2-Methoxyphenylcopper(I)

The synthesis of 2-methoxyphenylcopper was first reported by Camus and Marsich in 1968, where it was observed that compared to phenylcopper and tolylcopper complexes it was less air and temperature sensitive and more soluble in organic solvents.²⁶ The reported synthesis of this complex was from the reaction of 2-methoxyphenyllithium with copper(I) bromide, however reproducibility of the reaction was poor in part due to contamination from lithium (most likely in the form of co-complexed 2-methoxyphenyllithium). We therefore adopted an

alternative synthesis route via reaction of the Grignard $(\text{C}_6\text{H}_4\text{OMe-2})\text{MgCl}$ with copper(I) chloride in THF to give a reproducible and high purity yield of the orange crystalline product $[\text{Cu}(\text{C}_6\text{H}_4\text{OMe-2})]_8 \cdot 8\text{THF}$, **1** (see Experimental Section). Though somewhat complicated due to the highly aggregated nature of the complex, the ^1H NMR spectrum of **1** is fully consistent with Camus's previously reported spectroscopic analysis of this compound.²⁷

A solid-state structure of 2-methoxyphenylcopper crystallised from toluene was previously reported in 1971, however the data were reported to be of poor quality due to "decomposition of the crystal" during the data collection and is consequently not published on the Cambridge Structural Database.²⁸ Hence, in order to allow comparisons to be drawn with the novel lithium and Grignard-derived 2-methoxyphenylcuprates discussed below, a new structural analysis of **1** is briefly reported herein. Note that the crystals of **1** obtained in this work were from a THF solution and contain non-coordinating THF within the crystal lattice, whereas the 1971 structural analysis by Camus contained toluene molecules with the lattice thus giving rise to very different crystallographic parameters. However in both cases octomeric $[\text{Cu}(\text{C}_6\text{H}_4\text{OMe-2})]_8$ aggregates are observed as shown in Figure 3 for **1**.

a)



b)





Figure 3. a) Molecular structure of $[\text{Cu}(\text{C}_6\text{H}_4\text{OMe-2})]_8 \cdot 8\text{THF}$, **1**. Hydrogen atoms and solvent THF molecules are omitted for clarity. b) Simplified view of **1** showing the aryl *ipso* carbon atoms and just one of the methoxyphenyl groups. Thermal ellipsoids are displayed at 30% probability level. Symmetry transformations used to generate equivalent atoms: $y, -x+3/2, z$; $-y+3/2, x, z$

The structure of **1** is best considered as consisting of two tetrameric rings which lie on top of one another in a staggered conformation, so that the aryl group in one ring lies above a copper atom in the ring below. The rings are held together via coordination from anisyl methoxy oxygens in one ring to the copper centres in the adjacent ring (Cu(1)-O(7), 2.350(3) Å; Cu(2)-O(17), 2.387(3) Å), and there is no evidence of any intra-ring Cu-O interactions. Cu-C_{ipso} bond distances range from 2.013(4) to 2.051(4) Å, and are therefore comparable to analogous Cu-C_{ipso} distances in tetrameric arylcuprates such as [Cu₄Mes₄] (Mes = C₆H₂Me₃-2,4,6), 1.986(10)-1.999(9) Å,^{29,30} and [Cu₄(C₆H₄CH₂NMe₂-2)₄], 1.994(2)-2.102(2) Å,³¹ and are indicative of Cu-C-Cu two-electron-three-centre (2e-3c) bonding. The Cu-C-Cu angles within the rings are acute (Cu(1)-C(12)-Cu(1A), 74.66(12)°; Cu(2)-C(2)-Cu(2A), 75.00(13)°) and are also typical for 2e-3c bonding of an aryl group to two copper atoms.²⁹⁻³¹ There is no evidence for Cu-C interactions between the tetrameric rings in **1**, the shortest Cu...C inter-ring distance being 3.130 Å. The shortest inter-ring Cu...Cu distance of 2.690 Å is shorter than the combined van der Waals radii of two copper atoms, but is not thought to represent any significant bonding interaction: computational studies have shown that solid-state Cu(d¹⁰)-Cu(d¹⁰) interactions are likely to be very weak in nature with Cu(I)...Cu(I) distances primarily governed by the presence of other supramolecular interactions within the crystal lattice.³²

Lithium 2-methoxyphenylcuprate **2**


Reaction of 2-methoxyphenylcopper (**1**) with 2-methoxyphenyllithium in toluene / THF gave a yellow solution from which crystals of [Cu₄Li₂(C₆H₄OMe-2)₆(THF)₂] **2** were grown,

Scheme 1. The solid-state structure for complex **2**, as determined using single crystal X-ray diffraction, is shown in Figure 4 with key bond lengths and angles in Table 2.



Scheme 1. Synthesis of lithium diarylcuprate **2**

a)



b)




Figure 4. a) Molecular structure of $[\text{Cu}_4\text{Li}_2(\text{C}_6\text{H}_4\text{OMe-2})_6(\text{THF})_2]$, **2** with hydrogen atoms and disorder in the THF molecules omitted for clarity; b) Picture of the core connectivity in **2** omitting all other atoms. Thermal ellipsoids are displayed at 40% probability level.

Table 2. Selected bond lengths (Å) and angles (°) in [Cu₄Li₂(C₆H₄OMe-2)₆(THF)₂], **2**

Cu(1)-C(11)	1.924(3)	Li(1)-O(7)	1.970(6)
Cu(1)-C(1)	1.992(3)	Li(1)-O(17)	1.953(6)
Cu(2)-C(1)	1.998(3)	Li(1)-O(27)	1.921(7)
Cu(2)-C(41)	1.922(3)	Li(1)-O(60)	1.964(6)
Cu(3)-C(21)	1.925(3)	Li(2)-O(37)	1.951(7)
Cu(3)-C(31)	1.980(3)	Li(2)-O(47)	1.940(7)
Cu(4)-C(31)	1.991(3)	Li(2)-O(57)	1.924(7)
Cu(4)-C(51)	1.911(3)	Li(2)-O(70)	1.959(7)
C(1)-Cu(1)-C(11)	141.39(13)	C(31)-Cu(4)-C(51)	139.57(14)
C(1)-Cu(2)-C(41)	144.30(13)	Cu(1)-C(1)-Cu(2)	76.71(12)
C(21)-Cu(3)-C(31)	145.05(13)	Cu(3)-C(31)-Cu(4)	77.26(12)

Lithium organocuprate **2** possesses a copper to lithium ratio of two to one, despite the fact that it was formed from an equimolar mixture of its organocopper and organolithium precursors. Similar copper-rich lithium homocuprates with Cu:Li ratios greater than one are well known in the literature and include the anionic clusters [Cu₄LiPh₆]⁻¹² and [Cu₃Li₂Ph₆]⁻¹³ and the neutral complex [Cu₃LiMes₄]¹⁵. These species have been speculated to arise due to interaggregate exchange between Cu₂Li₂R₄ and Cu₄R₄ molecules in solution.^{33,34}


Closer analysis of the structure of **2** reveals the aggregate to be constructed from two [Cu₂(C₆H₄OMe-2)₃]⁻ anionic organocuprate units joined together by two THF-solvated lithium cations. Within each of the [Cu₂R₃]⁻ units there are two different types of aryl groups: the first bridges the two copper(I) atoms symmetrically to give a 3c-2e bond with Cu-C distances in the range 1.980(3) to 1.998(3) Å (mean 1.990 Å). The second type of aryl group is terminal to just one Cu(I) atom, resulting in the formation of shorter Cu-C distances in the range 1.911(3) to 1.925(3) Å (mean 1.920 Å) and indicative of 2c-2e bonds. A similar but ion separate [Cu₂Mes₃]⁻ anion has previously been reported which adopts an equivalent motif in

which the two terminal mesityl groups form 2c-2e bonds with the Cu atoms (Cu-C range 1.923(8) to 1.925(8) Å) and the central bridging mesityl group forms a longer 3c-2e bond (Cu-C range 2.003(8) to 2.020(7) Å).⁷ However, unique to **2** the two [Cu₂R₃]⁻ cuprate units are held together via coordination of lithium cations by the methoxy oxygen atoms with Li-O(Me) distances in the range 1.921(7) to 1.970(6) Å (mean 1.943 Å). The coordination sphere of the lithium is completed by a single THF molecule (Li-O 1.959(7), 1.964(6) Å).

In contrast to previously reported amino substituted homocuprates such as **III** (Figure 2), there are no Li-C_{ipso} interactions evident within the solid-state structure of **2**. The bonding behaviour of the methoxyphenyl groups therefore also differs significantly from that observed for the parent homometallic 2-methoxyphenyllithium complex, which adopts a dimeric structure in THF solution exhibiting both Li-C_{ipso} and Li-OMe coordination.³⁵ In the case of **2** this can be rationalised by the strong carbophilicity of copper(I) when compared to lithium. However, despite the absence of any Li-C_{ipso} bonding complex **2** still forms a CIP structure as opposed to a SSIP structure (Figure 1), which is rare for a lithium organocuprate in the presence of THF^{17,36} and can be attributed to the strong propensity of the anisyl oxygen atoms towards lithium cation coordination.

At first glance the four copper atoms in **2** appear to form a central tetrahedron with Cu...Cu distances within the range 2.476(5) to 2.824(6) Å and C-Cu-C bond angles distorted from linearity and pointing towards the centre of the tetrahedron (mean C-Cu-C, 142.74 °). However, as discussed above for **1** and based on previous bonding studies³² this should not be automatically interpreted to support the existence of any significant copper(I)-copper(I) bonding. Nevertheless the Cu₄ cluster does differ from previously reported Cu(I)₄ clusters -

for example $[\text{Cu}_4\text{I}_6]^{2-}$,³⁷ $[\text{Cu}_4(\text{SPh})_6]^{2-}$,³⁸ and $[\text{Cu}_4(\text{SePh})_6]^{2-}$ ³⁹ - in that in **2** only two of the six anionic ligands directly bridge copper atoms (these being the central aryl groups in the $[\text{Cu}_2\text{R}_3]^-$ units), and instead it is the peripheral Li-O(anisyl) bonding which must therefore play a large part in the supramolecular assembly of the structure.




Scheme 2. Oxidative homocoupling of 2-methoxyphenyl

On exposure to atmospheric oxygen, lithium organocuprate **2** was observed to decompose via oxidative coupling of the aryl groups to give small quantities of 2,2'-dimethoxybiphenyl (Scheme 2). By using the controlled introduction of nitrobenzene as the oxidising agent in place of atmospheric dioxygen we have been able to improve the yield of this coupling reaction to close to quantitative (>95%). In addition and as noted in the introduction above the application of lithium 2-methoxyphenyl cuprate **2** in conjugate addition reactions is already well established in the organic chemistry literature.²³⁻²⁵

Magnesium-Grignard 2-methoxyphenylcuprates **3-X** (X = Cl, Br, I)

Despite their widespread application in synthesis,¹ studies on the structures of Grignard organocuprates (commonly prepared from the reaction of a copper(I) salt with two

equivalents of Grignard reagent) are sparse in the literature, even when compared to lithium organocuprates. We recently reported upon the first structurally characterised examples of Grignard organocuprates: an iodo-Grignard CIP organocuprate $[\text{Ph}_4\text{Cu}_2(\text{Mg}(\text{OEt}_2)\text{I})_2]$ and bromo-Grignard CIP and SSIP organocuprates $[\text{Cu}_4\text{Mg}_2\text{Mes}_6\text{Br}_2]$ and $[\text{Cu}_2\text{Mes}_3]^- [\text{MgBr}(\text{THF})_5]^+$ respectively.⁷ Two related structures derived from diorgano-magnesium reagents have also been reported: $[\text{Cu}_4\text{MgPh}_6(\text{OEt}_2)]^{12}$ and $[\text{Cu}_4\text{Mes}_4][\mu\text{-SAr}]_2[\text{MgSAr}]_2$ (SAr = $\text{SC}_6\text{H}_4\text{CH}(\text{Me})\text{NMe}_2$).^{40,41} Expanding upon these previous studies, employment of the 2-methoxyphenyl ligand has led to the first structural characterization of a functionalised Grignard organocuprate. Thus reaction of copper 2-methoxyphenyl with the corresponding aryl Grignard (itself prepared from the reaction of the aryl halide with magnesium metal) in THF solution gave the magnesium Grignard organocuprate complexes **3-X** (X = Cl, Br, I) as shown in Scheme 3.



Scheme 3. Synthesis of Magnesium-Grignard diarylcuprates **3-X** (X = Cl, Br, I)

The preparation of **3-Br** and **3-Cl** gave crystalline products which were shown by X-ray diffraction to exist as isostructural complexes with similar crystallographic parameters (see

Table 1). The molecular structure of **3-Br** is shown in Figure 5 and the selected bond lengths and angles for **3-Br** and **3-Cl** are compared in Table 3.

Both **3-Br** and **3-Cl** are monomeric but dinuclear R_2CuMgX complexes with a 1:1 stoichiometric ratio of copper to magnesium halide. This ratio is therefore as expected from the original reaction stoichiometry, but differs from that observed in the homologous copper-rich organocuprate **2**, Cu:Li = 2:1 (*vide supra*). In addition, the structure of **3-X** contrasts with previously reported Grignard organocuprates, being the first example of a monomeric CIP complex. Although the crystals of **3-Br** and **3-Cl** were obtained from a THF solution, the organocuprates do not form THF solvated SSIP structures as might be initially expected (Figure 1), but rather give CIP structures, albeit with no apparent C_{ipso} -Mg bonding. Similar to lithium organocuprate **2**, the formation of a CIP structure can be attributed to the strong coordination of the s-block metal by the anisole methoxy group.




Figure 5. Molecular structure of $[\text{Cu}(\text{C}_6\text{H}_4\text{OCH}_3\text{-}2)_2\text{Mg}(\text{THF})_2\text{Br}]$ (**3-Br**). Hydrogen atoms are omitted for clarity and thermal ellipsoids are displayed at 40% probability level.

Table 3. Selected bond lengths (Å) and angles (°) in **3-X** (X = Cl, Br)

	X = Br	X = Cl
Cu-C(1)	1.9176(15)	1.923(6)
Cu-C(11)	1.9190(16)	1.919(6)
Mg-O(7)	2.0756(13)	2.075(5)
Mg-O(17)	2.0973(12)	2.091(5)
Mg-O(20)	2.0771(12)	2.071(5)
Mg-O(25)	2.0559(13)	2.061(5)
Mg-X	2.5183(6)	2.352(3)
C(1)-Cu-C(11)	172.57(7)	172.6(3)
O(7)-Mg-O(17)	155.37(6)	154.1(2)
O(7)-Mg-X	101.90(4)	101.94(16)
O(17)-Mg-X	102.54(4)	103.77(16)

Closer inspection of the make-up of **3-X** reveals two key structural components: a RCuR anionic fragment and a $\text{MgX}(\text{THF})_2$ cationic fragment. The anionic RCuR organocuprate unit is close to linear at the copper centre (C-Cu-C, $172.57(7)^\circ$ (**3-Br**), $172.6(3)^\circ$ (**3-Cl**)) with mean Cu-C distances of 1.918 \AA (**3-Br**) and 1.921 \AA (**3-Cl**). These copper angles and distances are consistent with the presence of 2c-2e copper(I)-carbon bonds and are directly comparable in length to similar 2c-2e Cu-C bonds in **2** (mean 1.920 \AA) as well as being significantly shorter than the 3c-2e bond lengths observed in **1** (mean 2.147 \AA) and **2** (mean 1.990 \AA). The diarylcuprate unit is connected to the Mg centre via coordination of the two methoxy groups, with mean Mg-O distances of 2.086 \AA (**3-Br**) and 2.083 \AA (**3-Cl**). The magnesium metal centres are additionally complexed by two THF oxygens and a halide anion. The geometry at the magnesium is close to square based pyramid with the halide in the

apical position: the sum of *cis* O-Mg-O bond angles is 358.57° in **3-Br** and 358.4° in **3-Cl**.

The position *trans* to the halide is effectively shielded by the copper atom thus preventing any additional donor coordination at this site (Cu...Mg distances are 2.7409(6) Å in **3-Br** and 2.765(2) Å in **3-Cl**). The positioning of the methyl groups as well as twisting in the THF molecules results in a chiral configuration for the molecule in the solid state, although the bulk product is racemic.

Although it was not possible to prepare crystals of the iodo-Grignard derivative **3-I** suitable for X-ray diffraction analysis, the NMR data of this complex is almost identical to that observed for both **3-Br** and **3-Cl** suggesting a similar Grignard diarylcuprate structure (see Experimental Section). In addition all **3-X** complexes exhibit aryl-aryl oxidative coupling, either slowly over several days in the presence of atmospheric oxygen or quantitatively over 1 hour in the presence of nitrobenzene, to give 2,2'-dimethoxybiphenyl in an analogous reaction to that reported for **2** (Scheme 2).

Summary and Conclusions

Despite recent advances in the understanding of organocuprate chemistry, studies concerning the structures of organocuprates containing functionalised organo groups remain limited to a few examples of amino-functionalised diaryl lithium cuprates. Given the important role functionalised organocuprates can play in natural product and other organic synthesis protocols, we set out to explore the coordination chemistry of the 2-methoxyphenyl group in lithium and Grignard organocuprates.

In a re-visitation of early work by Camus,^{27,28} 2-methoxyphenylcopper (**1**) was prepared and shown to exist in the solid state as an octameric aggregate formed from two tetrameric copper(I)-aryl rings conjoined via copper-oxygen(OMe) bonding interactions. This organocopper species was subsequently used in the preparation of the novel lithium organocuprate **2**, [Cu₄Li₂(C₆H₄OMe-2)₆(THF)₂], and the Grignard organocuprates **3-X**, [Cu(C₆H₄OCH₃-2)₂Mg(THF)₂X] X = Cl, Br, I. Solid-state structural studies show the donor methoxy group to play a large role in determining the structures adopted by these organocuprate species, coordinating in each case to the s-block metal centre with no copper-oxygen bonding present. Most notably this intramolecular methoxy coordination to lithium or magnesium is preferred over coordination of solvent THF, thus giving contact ion pair (CIP) structures rather than THF-solvated separated ion pairs (SSIPs) as observed for non-functionalised arylcuprates. Previous studies have shown how organocuprates structure can be key to understanding the reactivity and mechanism of reaction of these species,³ with the balance between CIP and SSIP species particularly important in this respect³⁶. In the past such structural studies have predominately concentrated upon the role of the solvent or the steric size of the organo group. However, from this work it is apparent how the presence of additional functional groups on the organo group can also play a key role in determining the overall structure and aggregation of these reagents. It is through such studies that we are able to build a more thorough understanding of these important and widely-used reagents.

References

- (1) Krause, N. *Modern organocopper chemistry*; Wiley-VCH: Weinheim, 2002.
- (2) Taylor, R. J. K. *Organocopper reagents : a practical approach*; Oxford University Press: Oxford, 1994.
- (3) Davies, R. P. *Coord. Chem. Rev.* **2011**, 255, 1226-1251.

- (4) Yoshikai, N.; Nakamura, E. *Chem. Rev.* **2012**, *112*, 2339-2372.
- (5) Gschwind, R. M. *Chem. Rev.* **2008**, *108*, 3029-3053.
- (6) Hickman, A. J.; Sanford, M. S. *Nature* **2012**, *484*, 177-185.
- (7) Bomparola, R.; Davies, R. P.; Hornauer, S.; White, A. J. *Angew. Chem. Int. Ed.* **2008**, *47*, 5812-5815.
- (8) Mulvey, R. E. *Organometallics* **2006**, *25*, 1060-1075.
- (9) Lorenzen, N. P.; Weiss, E. *Angew. Chem. Int. Ed. Eng.* **1990**, *29*, 300-302.
- (10) Olmstead, M. M.; Power, P. P. *J. Am. Chem. Soc.* **1990**, *112*, 8008-8014.
- (11) Edwards, P. G.; Gellert, R. W.; Marks, M. W.; Bau, R. *J. Am. Chem. Soc.* **1982**, *104*, 2072-2073.
- (12) Khan, S. I.; Edwards, P. G.; Yuan, H. S. H.; Bau, R. *J. Am. Chem. Soc.* **1985**, *107*, 1682-1684.
- (13) Hope, H.; Oram, D.; Power, P. P. *J. Am. Chem. Soc.* **1984**, *106*, 1149-1150.
- (14) Olmstead, M. M.; Power, P. P. *J. Am. Chem. Soc.* **1989**, *111*, 4135-4136.
- (15) Davies, R. P.; Hornauer, S. *Chem. Commun.* **2007**, 304-306.
- (16) Davies, R. P.; Hornauer, S.; Hitchcock, P. B. *Angew. Chem. Int. Ed.* **2007**, *46*, 5191-5194.
- (17) Bomparola, R.; Davies, R. P.; Hornauer, S.; White, A. J. P. *Dalton Trans.* **2009**, 1104-1106.
- (18) Perlmutter, P. *Conjugate addition reactions in organic synthesis*; Pergamon, 1992.
- (19) van Koten, G.; Jastrzebski, J. T. B. H.; Muller, F.; Stam, C. H. *J. Am. Chem. Soc.* **1985**, *107*, 697-698.
- (20) Kronenburg, C. M. P.; Jastrzebski, J. T. B. H.; Spek, A. L.; van Koten, G. *J. Am. Chem. Soc.* **1998**, *120*, 9688-9689.

- (21) Kronenburg, C. M. P.; Jastrzebski, J. T. B. H.; Boersma, J.; Lutz, M.; Spek, A. L.; van Koten, G. *J. Am. Chem. Soc.* **2002**, *124*, 11675-11683.
- (22) Ribas, X.; Jackson, D. A.; Donnadiou, B.; Mahia, J.; Parella, T.; Xifra, R.; Hedman, B.; Hodgson, K. O.; Llobet, A.; Stack, T. D. P. *Angew. Chem. Int. Ed.* **2002**, *41*, 2991-2994.
- (23) Ainai, T.; Ito, M.; Kobayashi, Y. *Tetrahedron Lett.* **2003**, *44*, 3983-3986.
- (24) Oberlender, R.; Ramachandran, P. V.; Johnson, M. P.; Huang, X. M.; Nichols, D. E. *J. Med. Chem.* **1995**, *38*, 3593-3601.
- (25) Kitson, R. R. A.; McAllister, G. D.; Taylor, R. J. K. *Tetrahedron Lett.* **2011**, *52*, 561-564.
- (26) Camus, A.; Marsich, N. *J. Organomet. Chem.* **1968**, *14*, 441-446.
- (27) Baici, A.; Camus, A.; Pellizer, G. *J. Organomet. Chem.* **1971**, *26*, 431-437.
- (28) Camus, A.; Marsich, N.; Nardin, G.; Randaccio, L. *J. Organomet. Chem.* **1979**, *174*, 121-128.
- (29) Meyer, E. M.; Gambarotta, S.; Floriani, C.; Chiesivilla, A.; Guastini, C. *Organometallics* **1989**, *8*, 1067-1079.
- (30) Eriksson, H.; Hakansson, M. *Organometallics* **1997**, *16*, 4243-4244.
- (31) Janssen, M. D.; Corsten, M. A.; Spek, A. L.; Grove, D. M.; van Koten, G. *Organometallics* **1996**, *15*, 2810-2820.
- (32) Carvajal, M. A.; Alvarez, S.; Novoa, J. J. *Chem. Eur. J.* **2004**, *10*, 2117-2132.
- (33) van Koten, G.; Noltes, J. G. *J. Am. Chem. Soc.* **1979**, *101*, 6593-6599.
- (34) van Koten, G.; Noltes, J. G. *J. Organomet. Chem.* **1979**, *174*, 367-387.
- (35) Harder, S.; Boersma, J.; Brandsma, L.; van Mier, G. P. M.; Kanters, J. A. *J. Organomet. Chem.* **1989**, *364*, 1-15.

- (36) John, M.; Auel, C.; Behrens, C.; Marsch, M.; Harms, K.; Bosold, F.; Gschwind, R. M.; Rajamohanan, P. R.; Boche, G. *Chem. Eur. J.* **2000**, *6*, 3060-3068.
- (37) Bowmaker, G. A.; Clark, G. R.; Yuen, D. K. P. *J. Chem. Soc., Dalton Trans.* **1976**, 2329-2334.
- (38) Maiti, B. K.; Pal, K.; Sarkar, S. *Eur. J. Inorg. Chem.* **2007**, *2007*, 5548-5555.
- (39) Jin, X.; Tang, K.; Long, Y.; Tang, Y. *Acta Crystallographica Section C* **1999**, *55*, 1799-1800.
- (40) Knotter, D. M.; Grove, D. M.; Smeets, W. J. J.; Spek, A. L.; van Koten, G. *J. Am. Chem. Soc.* **1992**, *114*, 3400-3410.
- (41) Knotter, D. M.; Smeets, W. J. J.; Spek, A. L.; van Koten, G. *J. Am. Chem. Soc.* **1990**, *112*, 5895-5896.