

Interleukin-6 mediated upregulation of CYP1B1 and CYP2E1 in colorectal cancer involves DNA methylation, miR27b and STAT3.

Saroor A. A. Patel¹, Upinder Bhambra¹, Michalis P. Charalambous¹, Rhiannon M. David¹, Robert J. Edwards², Tracy Lightfoot³, Alan R. Boobis² and Nigel J. Gooderham¹

¹ Surgery and Cancer, Imperial College London, London, SW7 2AZ, UK,

² Experimental Medicine and Toxicology, Imperial College London, W12 0NN,

³ Department of Health Sciences, University of York, YO10 5DD, UK.

Correspondence: Professor Nigel J. Gooderham
Computational and Systems Medicine,
Surgery and Cancer,
Imperial College London,
Sir Alexander Fleming Building,
London SW7 2AZ, UK
E-mail: n.gooderham@imperial.ac.uk
Tel: (+44) 020 7594 3188

Running title: IL6 upregulates CYP1B1 and CYP2E1 in colorectal cancer.

Competing interests: The authors declare that they have no competing interests.

Word count: 3932 (excluding abstract, references and figure legends)

Number of figures: 6

Number of tables: 1

Key words: Inflammation, microRNA, cytochrome P450, colorectal cancer, immunohistochemistry.

Conflict of Interest: None declared

Abstract

Background: The pro-inflammatory cytokine interleukin-6 (IL6) promotes colorectal cancer (CRC) development. IL6 is also known to regulate cytochrome P450 (CYP450) enzymes, which are involved in CRC tumour initiation and promotion via activation of chemical carcinogens. Here, IL6 regulation of CYP450 expression was investigated in CRC.

Methods: IL6 effect on CYP 1A1, 1B1 and 2E1 expression was determined *in vitro* using CRC cell lines HCT116 and SW480, and CYP450 expression was determined by immunohistochemistry in CRC tissues previously shown to have increased levels of IL6.

Results: In mechanistic studies, IL6 treatment significantly induced *CYP1B1* and *CYP2E1*, but not *CYP1A1*, gene expression in HCT116 and SW480 cells. *CYP2E1* expression regulation occurred *via* a transcriptional mechanism involving STAT3. For *CYP1B1* regulation, IL6 downregulated the *CYP1B1*-targeting microRNA miR27b through a mechanism involving DNA methylation. In clinical samples, expression of *CYP1B1* and *CYP2E1*, but not *CYP1A1*, were significantly increased in malignant tissue compared to matched adjacent normal tissue.

Conclusions: Colonic inflammation with the presence of IL6 associated with neoplastic tissue can alter metabolic competency of epithelial cells by manipulating *CYP2E1* and *CYP1B1* expression through transcriptional and epigenetic mechanisms. This can lead to increased activation of dietary carcinogens and DNA damage, thus promoting colorectal carcinogenesis. (199 words)

Introduction

Colorectal cancer (CRC) is one of the most common malignancies in the Western world where it is the second highest cause of cancer-related mortality (Tenesa and Dunlop, 2009). Despite multiple advances in treating the disease, 5-year survival rates have not significantly improved in the last 20 years highlighting the need for new diagnosis strategies.

Local inflammation in the colon is a known risk for CRC development and progression (Feagins *et al*, 2009). We have previously reported increased expression of a number of inflammatory markers including pro-inflammatory cytokine interleukin-6 (IL6), a key regulator of inflammation, in colon tumour tissue and stroma (Fig1) (Maihofner *et al*, 2003; Charalambous *et al*, 2003; Charalambous *et al*, 2009). Indeed, stromal cells including tumour-associated immune cells and fibroblasts release high levels of IL6 (up to 6000pg/ml) in the colorectal tumour microenvironment (Nagasaki *et al*, 2014). Furthermore, elevated serum levels of IL6 are associated with poor disease outcome and non-steroidal anti-inflammatory drugs are known to reduce CRC risk by lowering levels of inflammation (Williams *et al*, 2007; Smalley *et al*, 1999).

IL6 is also known to regulate CYP enzyme expression. Studies in hepatocytes have generally shown that IL6 inhibits CYP450 expression including CYP 1A1, 1A2, 3A4 and 2E1 (Abdel-Razzak *et al*, 1993; Jover *et al*, 2002; Hakkola *et al*, 2003). However conflicting results have been reported; IL6 has also been shown to induce CYP2E1 in astrocytes (Tindberg *et al*, 1996), suggesting that the effects of IL6 on CYP450 expression vary depending on tissue type and the relation between IL6 and CYP450 expression has not yet been defined in the colon. CYP1B1 is known to be overexpressed in a number of tumour types including lung, colon and breast (Murray *et al*, 1997). A recent study in human

hepatocellular carcinoma cells has reported that IL6 treatment induced CYP1B1 gene and protein expression (Kurzawski *et al*, 2012), suggesting that IL6 is able to regulate CYP1B1. However, there have been very few other studies to confirm this effect of IL6 on CYP1B1. Furthermore, despite these numerous reports of IL6 regulating CYP450s, the specific molecular mechanisms underlying the effects are not yet well understood.

MicroRNAs (miRNA) are small, approximately 22 nucleotide long non-coding gene-regulating RNAs; they silence gene expression by binding to target mRNAs.

MiRNAs only require partial base complementarity to bind to their targets and can thus target a number of different mRNAs.

Recent studies have found that miRNAs can modulate CYP450 expression. For instance, miR27b and miR378 have been reported to directly regulate CYP1B1 and CYP2E1 expression respectively (Tsuchiya *et al*, 2006; Mohri *et al*, 2010). Furthermore, emerging evidence suggests that epigenetic events including DNA methylation and miRNA expression are involved in CRC initiation and progression (Suzuki *et al*, 2012). Interestingly, IL6 has been shown to alter miRNA expression profiles (Yang *et al*, 2010) and to induce genome wide methylation of promoter regions (Webbe *et al*, 2006) resulting in gene silencing, thus epigenetic mechanisms may be involved in IL6-mediated regulation of CYP450s.

To investigate the relationship between IL6 and CYP450 expression in the neoplastic colon, we first determined the expression of the CYP450 enzymes CYP 1A1, 1B1 and 2E1 in the human CRC cell lines HCT116 and SW480 with subsequent mechanistic studies to confirm increased levels of IL6 could alter CYP450 expression and propose underlying mechanisms. Here to extend the *in vitro* results, we measured the expression of the same CYP450s in malignant tissues resected from CRC patients that have increased expression of IL6 in the

epithelium and stroma (Fig1; Maihofner *et al*, 2003), and compared expression to that in adjacent normal tissues.

Materials and methods

Cell culture

The human colorectal adenocarcinoma cell lines HCT116 and SW480 were obtained from ATCC (LGC Prochem, Middlesex, UK) and were routinely cultured in RPMI1640 medium (GIBCO, Life technologies, Paisley, UK) supplemented with 10% foetal bovine serum (FBS), 100units/ml penicillin, 100µg/ml streptomycin and 2mM *L*-glutamine (GIBCO, Life technologies). All cells were incubated at 37°C in a humidified incubator (5% CO₂). Cells between passages 3-7 were used in the experiments.

IL6 treatment and demethylation/STAT3 inhibition

Prior to treatment, HCT116 and SW480 were maintained in culture medium supplemented with 5% dextran-coated charcoal-stripped FBS for at least 72 hours. Cells were seeded at a density of 1x10⁵ cells per well of a 6-well plate. HCT116 and SW480 do not express IL6; cells were treated for 24 and 48 hours with recombinant human IL6 (HumanKine, Sigma-Aldrich) dissolved in PBS containing 0.1% human serum albumin (Sigma-Aldrich) at doses of 0, 100 and 1000pg/ml (chosen within the range secreted by stromal cells in colon tumour microenvironment). For demethylation/STAT3 inhibition, cells were co-treated for 24 and/or 48 hours with 1000pg/ml IL6 and either 4µM 5-aza-2'-deoxycytidine (Sigma-Aldrich) or 25µM STAT3 inhibitor (STAT3 inhibitor VIII 5,15-diphenylporphyrin, Millipore, Feltham, UK), respectively. STAT3 inhibitor and 5-aza-2'-deoxycytidine were dissolved in DMSO (final vehicle control concentration of 0.1%).

RNA extraction

Total RNA was isolated from cells using TRIzol reagent (Invitrogen, Life technologies) according to the manufacturer's protocol. RNA extracts were quantified by UV spectroscopy (UV-VIS Spectrophotometer, Implen, Essex, UK) with purity assessed from 260/280nm and 260/230nm ratios. Extracts were stored at -80°C until used.

Reverse transcription and quantitative PCR

For mRNA reverse transcription, RNA extracts (100-500ng) from each sample were added to 300ng of random primers, heated for 5min at 65°C and then immediately placed on ice. Each sample was incubated with 0.5mM dNTPs, 1x first strand buffer, 8µM dithiothreitol and 100units of Superscript II reverse transcriptase (Invitrogen, Paisley, UK) for 10min at 25°C, 90min at 42°C and 15min at 70°C on a thermocycler (Peltier Thermal Cycler PTC-200, MJ Research, Waltham, Massachusetts, USA). A miRNA reverse transcription kit was used for miRNA expression according to the manufacturer's protocol (Taqman, Applied Biosystems, Life technologies). Quantitative PCR was performed using pre-designed gene expression assays and FAST PCR master mix (Taqman, Applied Biosystems, Life technologies), and measured in a StepOnePlus fast real-time PCR system (Applied Biosystems, Life technologies) according to the manufacturer's protocol. Gene expression was quantified using the delta-C_t method.

Chromatin immunoprecipitation-quantitative PCR (ChIP-qPCR)

SW480 cells were treated with 1000pg/ml IL6 and 25µM STAT3 inhibitor for 60min prior to collecting the cells. ChIP assay was performed using a magnetic ChIP kit (Thermo Fisher Scientific, Cramlington, UK) as per manufacturer's

protocol with an anti-STAT3 antibody (sc-482X, Santa Cruz Biotechnology, Heidelberg, Germany). Rabbit IgG was used as a negative control for non-specific binding. Binding was calculated as a percentage of the total input chromatin.

Sequences of the primers used were as follows:

CYP2E1 site 1 forward: 5'-TGAATTTTCCTTCTGGCCCCAT-3',

reverse 5'- TGATGAGGAGGTTTGTCTGAGC-3';

CYP2E1 site 2 forward: 5'- CTCCATCCTCACCAGGTCAC-3',

reverse: 5'- CCAACCAATGCCCTCTTGCT-3'.

Patients

The study adhered to the tenets of the Declaration of Helsinki. Surgical specimens of primary tumours and adjacent normal colon were obtained with informed consent from 40 patients (28 men and 12 women; aged 49-80 years, mean age 64.7 years \pm 8.1 SEM), with histologically verified colorectal cancer, treated at the Department of Surgery, York District Hospital, York, UK. Ethical approval for the study was obtained from the Human Research Ethics Committee at York District Hospital. Nineteen patients had colon cancer and 21 rectal cancers. Tumours were classified according to Duke's classification (Table1). The entire study was carried out blind using coded tissue sections. Eligibility criteria for patients recruited to the study included Caucasian origin, 45-80 years of age and no history of previous gastrointestinal disease or any form of cancer, including familial adenomatous polyposis.

Tissue specimens

Tissue samples taken at operation for histopathological confirmation of disease were fixed in 4% buffered formaldehyde and embedded in paraffin wax; sections surplus to pathology requirements were made available for the study. For all 40

patients, tissue sections of both malignant and normal (taken outside of the tumour margin) colon or rectum were provided.

Antibodies and reagents

Specific anti-peptide antibodies raised against individual CYP450 enzymes and the corresponding pre-immune sera have been previously described (Edwards *et al*, 1998; Kapucuoglu *et al*, 2003). These antibodies have previously been used to detect human CYP450 protein expression (Piipari *et al*, 2000, Rodriguez-Antona *et al*, 2001). Biotin labeled secondary antibodies and streptavidin/HRP complex were obtained from DAKO Ltd. All other chemicals were supplied by either Sigma-Aldrich (Dorset, UK) or Merck (Nottingham, UK) unless otherwise specified.

Immunohistochemistry

The levels of expression and localization of CYPs 1A1, 1B1 and 2E1 in malignant and adjacent normal colorectal tissue was determined using a modified avidin/biotin immunohistochemistry procedure (Goggi *et al*, 1986). In preliminary experiments, each of the immunohistochemistry assays was optimized using a range of antisera dilutions (1/200 – 1/8000). For each assay, the negative control antisera (pre-immune sera) were confirmed negative for staining at the dilution optimized for the primary antibody and blocking peptides confirmed specificity. The dilutions used were 1/5000 for the anti-CYP1A1 antibody, and 1/1000 for the anti-CYP1B1 and anti-CYP2E1 antibodies. The sections were deparaffinised and rehydrated through xylene and a series of graded alcohol solutions. Endogenous peroxidase activity was blocked by immersing the sections into a solution of 3% hydrogen peroxide in distilled water for 30 minutes at room temperature, and then rinsed in cold running tap water for 10 minutes. Incubating the sections with 5% normal swine serum for 30 minutes at room temperature reduced non-specific background staining. Sections were then washed twice with phosphate-buffered

saline (PBS) and either the primary antibody or the normal goat or rabbit IgGs (negative control) was applied to each section and left at 4°C overnight. The next day, the slides were washed twice with PBS, and then incubated with the secondary antibody solution (Biotinylated swine anti-goat, mouse, rabbit immunoglobulin; 1/150 dilution), for one hour at room temperature. After being washed twice with PBS, they were incubated with the StrepABCComplex solution for one hour at room temperature, washed twice with PBS and immersed into the substrate (300ml PBS, 90µl hydrogen peroxide and 2.5ml 3,3-diaminobenzidine) for 3 minutes, and then rinsed with PBS and cold running tap water. Sections were then successively immersed into haematoxylin, acid alcohol and Scott's tap water to counterstain. Finally, the sections were dehydrated by successive immersion into 70% ethanol, 100% ethanol twice and xylene twice and mounted.

Immunohistochemical evaluation

Processed specimens were scored under the light microscope and the intensity and localization of staining with CYPs 1A1, 1B1 and 2E1 antibodies graded blind using coded slides. In order to assess and grade intensity and distribution of immunoreactivity in the epithelial cells, a previously described method of scoring the sections was used (Maihofner *et al*, 2003, Charalambous *et al.*, 2003, 2009). The distribution was scored according to the number of positive cells; none (not stained), 0; focal (<1/3 of cells stained), 1; multi-focal (1/3 – 2/3 of cells stained), 2; and diffuse (>2/3 stained), 3; staining intensity was scored as: none (not stained), 0; lightly stained, 1; and strong staining, 2. The distribution and intensity scores were added to produce the grade of staining. Sections treated with the normal goat or rabbit IgGs (negative controls) or omitting the primary antibody were devoid of staining. Positive staining controls included sections of kidney, duodenum and liver.

Statistical analysis

The Wilcoxon's signed rank test was used to compare the scoring of the respective immunoreactivity for CYP 1A1, 1B1 and 2E1 between tumour and adjacent normal tissues (Stata Statistical Software 9). Gene expression data were obtained from measurements made in at least three biological replicates and presented as a mean \pm standard error. Significant differences ($p < 0.05$) were determined using Student's t-test and one-way analysis of variance (ANOVA) followed by a Dunnett post-test or a linear trend analysis (GraphPad Prism 5). Pearson's product-moment correlation coefficient test was used for correlation analysis (GraphPad Prism 5).

Results

Can IL6 regulate CYP450 expression in CRC cells?

To determine whether IL6 can regulate CYP450s in colon cancer cells, we performed mechanistic studies in *in vitro* models (human CRC cell lines HCT116 and SW480) to examine the effect of IL6 treatment on *CYP 1A1*, *1B1* and *2E1* gene expression at various time points using quantitative PCR.

CYP1A1 gene expression was detected but not significantly changed in either cell line following 24- and 48-hour IL6 treatment (Fig2. A). However, *CYP1B1* and *CYP2E1* mRNA expression was regulated dose dependently by IL6 as determined by positive trend analyses and was significantly increased at the highest dose of 1000pg/ml IL6 in both cell lines (Fig2. B and C). To the best of our knowledge, this is the first account of *CYP1B1* and *CYP2E1* being upregulated by IL6 in colon tumour-derived epithelial cells.

What are the mechanisms involved in IL6-mediated upregulation of CYP2E1 and CYP1B1?

IL6 regulates CYP2E1 expression through STAT3 transcription factor.

To understand the mechanism underlying IL6 induction of *CYP2E1* expression, we examined the different pathways involved in *CYP2E1* regulation. *CYP2E1* is regulated at various stages of its synthesis and includes transcriptional and post-transcriptional mechanisms. We looked at miRNA-mediated regulation of *CYP2E1* mRNA by determining miR378 expression, a miRNA reported to target *CYP2E1* (Mohri *et al*, 2010). However no change in miR378 expression by IL6 was observed (Fig3. A) and we found no correlation between *CYP2E1* and miR378 expression in our model (Fig3. B).

IL6 is a potent inducer of the JAK/STAT3 pathway. An analysis of the *CYP2E1* promoter region revealed multiple potential STAT binding sites (Fig4. A; TFSEARCH ver1.3; Heinemeyer *et al*, 1998). We thus studied involvement of STAT3 in regulating IL6-mediated *CYP2E1* induction using a STAT3 inhibitor, STAT3 inhibitor VIII 5,15-diphenylporphyrin. Treatment with the inhibitor prevented IL6-mediated *CYP2E1* induction after 24 hours in both HCT116 and SW480 cell lines (Fig4. B). Furthermore, a ChIP analysis in SW480 cells revealed that STAT3 does bind to the *CYP2E1* promoter region following IL-6 treatment (Fig4. D), compatible with a STAT3-mediated mechanism for induction of *CYP2E1* expression by IL6.

IL6 regulates CYP1B1 expression through repression of miR27b.

We next examined the mechanisms underlying IL6-mediated induction of *CYP1B1* expression. The aryl hydrocarbon receptor (AhR) pathway is a well-known transcriptional regulator of *CYP1B1* and *CYP1A1* expression. However, *CYP1A1* mRNA expression was not induced upon IL6 treatment (Fig2. A), thus the AhR pathway is unlikely to be involved in IL6-mediated induction of *CYP1B1*. Furthermore, we determined that STAT3 was not involved in regulating IL6-

mediated *CYP1B1* induction, as treatment with the STAT3 inhibitor did not affect IL6-mediated induction of *CYP1B1* (Fig4. C).

MiR27b has been reported to directly target *CYP1B1* mRNA by binding to its 3'UTR to regulate its expression (Tsuchiya *et al*, 2006). IL6 significantly downregulated miR27b expression in both cell lines in the current study (Fig3. C). Furthermore, there was a significant inverse correlation between miR27b expression and *CYP1B1* expression (Fig3. D), suggesting that downregulation of miR27b could be responsible for the increase in *CYP1B1* mRNA observed. To our knowledge, this is the first account of IL6 modulating miR27b expression, thus providing a potential post-transcriptional mechanism by which *CYP1B1* is regulated by IL6.

How does IL6 cause miR27b downregulation?

We next determined the mechanism underlying miR27b regulation by IL6. MiR27b is an intragenic miRNA located within the *C9orf3* gene on chromosome 9. Previous reports have shown that an intragenic CpG island (chr9:96887100-96887300) located close to miR27b can be methylated in colon cancer cells and regulate miR27b expression (Yan *et al*, 2011). IL6 is known to induce genome-wide DNA methylation through phosphorylation of DNMT1 by activated AKT, which increases DNMT1 nuclear translocation and activity (Hodge *et al*, 2007); therefore we examined the role of DNA methylation in IL6 regulation of miR27b expression by inhibiting DNA methylation using 5-aza-2'-deoxycytidine treatment. We observed that by inhibiting DNA methylation, we were able to prevent miR27b downregulation by IL6 in both cell lines (Fig5. A), implying that DNA methylation plays a role in regulating miR27b expression. Again, these changes inversely correlated with *CYP1B1* expression (Fig5. B and C). These mechanistic data

suggest that IL6-mediated upregulation of CYP1B1 involves downregulation of miR27b via DNA methylation.

Is CYP450 expression altered in malignant colorectal tissues that overexpress IL6?

Having shown that IL-6 regulates expression of CYP1B1 and 2E1, but did not alter expression of CYP1A1 in CRC cells, to confirm whether this may be relevant *in vivo*, we measured the expression of these CYP450s by immunohistochemistry (Fig6) in tissue sections of malignant and normal bowel from colorectal cancer patients, in which IL6 expression has previously been determined (Fig1; Maihofner *et al*, 2003).

In most cases, we observed immunostaining only in epithelial cells. CYP expression was generally greatest in luminal enterocytes with involvement of the upper third of the crypts in some individuals. We did not observe any expression in the base of crypts. Within the cells, we noted the expression was cytoplasmic with the intensity of expression greatest in perinuclear regions, suggesting localization within the endoplasmic reticulum. No expression was observed in the mucous inclusions of goblet cells. In tumour tissue, staining was confined to epithelial cells and there was no significant intratumour heterogeneity in enzyme expression.

CYP1A1 expression was observed in epithelial cells of normal colorectal tissue in 7/40 patients, while 11/40 patients expressed CYP1A1 in malignant epithelial cells (Fig6. C and D). Using our previously described immunohistochemistry scoring procedures (Mainhofner *et al.*, 2003, Charalambous *et al.*, 2003, 2009), we found that this did not translate into a significant difference (Fig6. K). Expression of CYP1B1 in normal colorectal tissue was detected in 18/40 patients (Fig6. E), while the majority of patients expressed CYP1B1 in malignant tissue (35/40 patients; Fig6. F). CYP1B1 was also expressed relatively intensely in histologically normal

crypts that were immediately adjacent to tumour cells (Fig6. I). CYP2E1 was expressed in both normal and tumour tissue, although expression was more intense in malignant tissue (Fig6. G and H). In addition to epithelial cells, plasma cells (found in the lamina propria) also expressed CYP2E1 intensely (Fig6. J inset).

We compared these CYP450 expression levels with patient demographics and found no significant correlation between alcohol, age, sex or tumour stage. (data not shown).

More importantly, in line with our *in vitro* data, CYP1B1 and CYP2E1 expression were significantly increased in tumour samples compared to adjacent normal tissue while the CYP1A1 expression levels were not significantly changed (Fig6. K). We have previously reported increased expression of inflammatory cytokine IL6 in these same tumour tissue samples (Fig1; Maihofner *et al*, 2003). Furthermore, CYP1B1 and CYP2E1 gene expression were also the only CYP450s to be induced by IL6 in our *in vitro* mechanistic models and as CYP450 mRNA levels in human cell culture have been reported to correlate well with protein expression (Rodriguez-Antona *et al*, 2002), we therefore propose that IL6 may be involved in the increased expression of these proteins in tumour tissue.

Discussion

The effect of IL6 on CYP450 expression in colorectal cancer has not been widely investigated. Furthermore, only a few studies have attempted to characterize the expression of individual CYP isozymes using immunohistochemistry (Kumarakulasingham *et al*, 2005; Androutsopoulos *et al*, 2013). Certain members of the CYP450 family, including CYP1A1 and 1B1, can activate dietary pro-carcinogens such as heterocyclic amines and polycyclic aromatic hydrocarbons, which have been associated with colorectal cancer (Ito *et al*, 1991; Barrett *et al*,

2003; Sinha *et al*, 2005; Nothlings *et al*, 2009; Crofts *et al*, 1997; Boyce *et al*, 2004), and the expression of CYP450 enzymes in tissues that are targets for xenobiotic genotoxicity implies a potential for *in situ* activation. CYP450s, in particular CYP2E1, can also metabolise several drugs resulting in either inhibition or activation, and thus play an important role in drug bioavailability (Koop, 1992). CYP450 enzymes play significant roles in xenobiotic activation, tumour initiation and promotion (Murray *et al*, 1997; Rodriguez *et al*, 2013; Kasai *et al*, 2013; Braeuning *et al*, 2011, Gooderham *et al*, 2007), hence, characterization of CYP450 enzyme profiles in colon tissue and understanding the mechanisms involved in their regulation can be important for diagnosis, prognosis and treatment of disease.

IL6 has been reported to regulate CYP enzyme expression (Abdel-Razzak *et al*, 1993; Joyer *et al*, 2002; Hokkola *et al*, 2003; Tindberg *et al*, 1996; Kurzawski *et al*, 2012) and is also thought to be involved in colon cancer initiation and progression (Feagins *et al*, 2009). We have previously shown IL6 expression is present at higher levels in the neoplastic colon (Maihofner *et al*, 2003) and the current study extends these observations and provides evidence for the localization of several CYP450 enzymes in colon tissue from the same patient cohort and we demonstrate that CYP1B1 and CYP2E1 expression are significantly upregulated in these tumour tissue samples. Moreover, we have conducted mechanistic studies that give an insight into some of the regulatory mechanisms involving IL6 that underlie CYP1B1 and 2E1 enzyme expression in the colon.

Regarding CYP2E1, we found that IL6 induced *CYP2E1* expression in human colon tumour-derived cell lines and in support, we showed CYP2E1 levels were increased in tumour tissue expressing higher levels of IL6 than matched normal tissue. It is commonly thought that IL6 as well as other pro-inflammatory cytokines

repress CYP450 expression (Abdel-Razzak *et al*, 1993; Joyer *et al*, 2002) and *CYP2E1* gene expression was previously shown to be downregulated by IL6 in hepatic tissue (Hakkola *et al*, 2003), which is in contrast to our data. However, a study published by Tindberg *et al*. (Tindberg *et al*, 1996) reported that inflammation induced *CYP2E1* gene expression in astrocytes. Taken together, these data suggest that the effect of inflammation on *CYP2E1* expression is dependent on tissue type. We identified that IL6-mediated induction of *CYP2E1* *in vitro* was through a mechanism involving direct binding of STAT3 to the *CYP2E1* promoter. This is the first account of a direct involvement of IL6 and STAT3 in *CYP2E1* expression in colonic cells. It has previously been shown that both STAT3 activation and *CYP2E1* overexpression are linked to alcohol consumption (Norkina *et al*, 2007; Roberts *et al*, 1995). Epidemiological and animal studies have identified an association between excessive alcohol intake and increased risk of colorectal cancer (Pollack *et al*, 1984; Seitz *et al*, 1984; Kune *et al*, 1987; Kato *et al*, 1990). More recently, a study published by Morita *et al*. (2009) demonstrated a correlation between *CYP2E1* activity, alcohol intake, meat consumption and the development of colorectal cancer. Our findings provide a potential inflammatory STAT3-mediated mechanism by which *CYP2E1* expression could be regulated by alcohol in the colon.

CYP1B1 has been suggested to be a tumour specific enzyme as it is rarely found in healthy tissue (Murray *et al*, 1997). As expected, there was a significant increase in *CYP1B1* expression in tumour tissue when compared with matched normal tissue in the current study. These results are in agreement with previous studies, which have shown increased *CYP1B1* expression in colon cancer (Murray *et al*, 1997; Kumarakulasingham *et al*, 2005; Androutsopoulos *et al*, 2013; Chang *et al*, 2005; Gibson *et al*, 2003). Moreover, in mechanistic studies we found that

IL6 induced *CYP1B1* gene expression in colon tumour-derived epithelial cells. The relationship between inflammation and CYP1B1 is not as well studied as other CYP450 enzymes, despite both of these factors being of significant interest in tumour development. However, inflammation was previously shown to increase *CYP1B1* expression in brain cells (Malaplate-Armand *et al*, 2003). In a recent study, IL6 was shown to induce *CYP1B1* but not *CYP1A1* gene expression in human hepatocellular carcinoma cell line HepG2 in an AhR-independent manner (Kurzawski *et al*, 2012). This is in line with our results, where IL6 induced CYP1B1 but not CYP1A1 in CRC cells and only CYP1B1 levels were significantly elevated in tumour tissue where IL6 expression was also increased. While Kurzawski *et al*. (2012) did not establish a specific mechanism for the induction of CYP1B1 by IL6, here we show that *CYP1B1* is post-transcriptionally regulated by IL6 through miR27b, a miRNA reported to directly target *CYP1B1* mRNA (Tsuchiya *et al*, 2006). Furthermore, our data suggest that DNA methylation plays a role in IL6-mediated downregulation of miR27b. Yan *et al*. reported a CpG island located near miR27b that is methylated in CRC cells and can regulate miR27b expression (Yan *et al*, 2011), thus supporting our findings. MiR27b has been shown to play various roles in the inflammatory process. In neuroblastoma cells, inhibition of miR27b activated NFκB signaling leading to increased expression of IL6 (Lee *et al*, 2011), whereas the opposite has been found in breast cancer cells (Jin *et al*, 2013). Despite these conflicting reports, miR27b and IL6 appear to be involved in the same pathways and this is the first account of IL6 directly regulating miR27b expression.

In summary, this study has shown that CYP450 enzymes expressed in colon tissue are significantly influenced by disease as evidenced by the increased expression of CYP2E1 and CYP1B1 in tumour tissue samples. **However, it is**

important to note that a sample size of 40 patients was used in this study, and while our data are in agreement with previous reports of increased CYP1B1 expression, studies with larger patient numbers are needed in order to validate these findings, in particular for CYP2E1 expression. Furthermore, we noted the changes in CYP450 expression observed in malignant tissue did not correlate to sex or age. Interestingly, CYP450 expression was also not associated to tumour grade, suggesting these changes occur at an early disease stage. We have previously shown these same colorectal tumours (from the same patient cohort) have increased levels of IL6 compared to adjacent normal tissues (Maihofner *et al*, 2003) and here we demonstrate for the first time that addition of IL6 can regulate CYP1B1 and CYP2E1 in colon tumour-derived cell lines. We describe a transcriptional mechanism for IL6-mediated CYP2E1 induction *via* STAT3 (Fig7). We also demonstrate an epigenetic mechanism by which IL6 represses miR27b expression involving DNA methylation resulting in CYP1B1 upregulation (Fig7). Taken together, our study provides further insight into the mechanisms by which IL6 promotes tumour development and progression. While altering CYP450 expression is only one of the mechanisms by which IL6 exerts its pro-tumourigenic effects, this pathway is of particular importance when it comes to designing therapies. New CRC treatment strategies could exploit this novel pathway potentially through the use of drugs that can only be activated by CYP2E1 or CYP1B1, and avoiding treatments that may be deactivated by these CYP450s. These findings also suggest that preventive dietary measures are of particular importance for patients with inflammatory bowel conditions to reduce their risk of CRC. By manipulating CYP450 enzymes, IL6 can induce phenotypic changes in colon tumour cells possibly rendering them drug resistant, or encouraging the *in*

situ metabolism of carcinogens, thus resulting in DNA damage and potentially tumour promotion.

Acknowledgements

The work was supported by funding from the United Kingdom Food Standards Agency and Medical Research Council.

References

Abdel-Razzak Z, Loyer P, Fautrel A, Gautier JC, Corcos L, Turlin B, Beaune P, Guillouzo A (1993) Cytokines down-regulate expression of major cytochrome P-450 enzymes in adult human hepatocytes in primary culture. *Mol Pharmacol* **44**: 707–715.

Androutsopoulos VP, Spyrou I, Ploumidis A, Papalampros AE, Kyriakakis M, Delakas D, Spandidos DA, Tsatsakis AM (2013) Expression profile of CYP1A1 and CYP1B1 enzymes in colon and bladder tumours. *PLoS One* **8**(12): e82487. doi:10.1371/journal.pone.0082487.

Barrett JH, Smith G, Waxman R, Gooderham N, Lightfoot T, Garner RC, Augustsson K, Wolf CR, Bishop DT, Forman D, Colorectal Cancer Study Group (2003) Investigation of interaction between N-acetyltransferase 2 and heterocyclic amines as potential risk factors for colorectal cancer. *Carcinogenesis* **24**: 275-282.

Boyce A, Doehmer J, Gooderham NJ (2004) Phytoalexin resveratrol attenuates the mutagenicity of the heterocyclic amines 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine and 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline. *J Chromatogr B Analyt Technol Biomed Life Sci* **802**(1): 217-223.

Braeuning A, Kohle C, Buchmann A, Schwarz M (2011) Coordinate regulation of cytochrome P450 1a1 expression in mouse liver by the aryl hydrocarbon receptor and the beta-catenin pathway. *Toxicol Sci* **122**: 16-25.

Chang H, Su JM, Huang CC, Liu LC, Tsai CH, Chou MC, Lin P (2005) Using a combination of cytochrome P450 1B1 and beta-catenin for early diagnosis and prevention of colorectal cancer. *Cancer Detect Prev* **29**(6): 562-569.

Charalambous MP, Lightfoot T, Speirs V, Horgan K, Gooderham NJ (2009) Expression of COX-2, NF-kappaB-p65, NF-kappa-p50 and IKKalpha in malignant and adjacent normal human colorectal tissue. *Br J Cancer* **101**(1): 106-115.

Charalambous MP, Maihofner C, Bhambra U, Lightfoot T, Gooderham NJ, and the colorectal cancer study group (2003) Upregulation of cyclooxygenase-2 is accompanied by increased expression of nuclear factor-kappa B and I kappa B kinase-alpha in human colorectal cancer epithelial cells. *Br J Cancer* **88**(10): 1598-1604.

Crofts FG, Strickland PT, Hayes CL, Sutter TR (1997) Metabolism of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) by human cytochrome P450 1B1. *Carcinogenesis* **18**: 1793-1798.

Edwards RJ, Adams DA, Watts PS, Davies DS, Boobis AR (1998) Development of a comprehensive panel of antibodies against the major xenobiotic metabolising forms of cytochrome P450 in humans. *Biochem Pharmacol* **56**: 377-387.

Feagins LA, Souza RF, Spechler SJ (2009) Carcinogenesis in IBD: potential targets for the prevention of colorectal cancer. *Nat Rev Gastroenterol Hepatol* **6**: 297-305.

Gibson P, Gill JH, Khan PA, Seargent JM, Martin SW, Batman PA, Griffith J, Bradley C, Double JA, Bibby MC, Loadman PM (2003) Cytochrome P450 1B1 (CYP1B1) is overexpressed in human colon adenocarcinomas relative to normal colon: implications for drug development. *Mol Cancer Ther* **2**(6): 527-534.

Goggi G, Dell'Orto P, Viale G (1986) Immunohistochemistry, Modern Methods and Applications. pp 54-70. Butterworth-Heinemann: London.

Gooderham NJ, Creton S, Lauber SN, Zhu H (2007) Mechanisms of action of the carcinogenic heterocyclic amine PhIP. *Toxicol Lett* **168**(3): 269-277.

Hakkola J, Hu Y, Ingelman-Sundberg M (2003) Mechanisms of down-regulation of CYP2E1 expression by inflammatory cytokines in rat hepatoma cells. *J Pharm Exp Ther* **304**(3): 1048-1054.

Heinemeyer T, Wingender E, Reuter I, Hermjakob H, Kel AE, Kel OV, Ignatieva EV, Ananko EA, Podkolodnaya OA, Kolpakov FA, Podkolodny NL, Kolchanov NA (1998) Databases on Transcriptional Regulation: TRANSFAC, TRRD, and COMPEL. *Nucleic Acids Res* **26**: 364-370.

Hodge DR, Cho E, Copeland TR, Guszczynski T, Yang E, Seth AK, Farrar WL (2007) IL-6 enhances the nuclear translocation of DNA cytosine-5-methyltransferase (DNMT1) via phosphorylation of the nuclear localization sequence by AKT kinase. *Cancer Genomics Proteomics* **4**: 387-398.

Ito N, Hasegawa R, Sano M, Tamano S, Esumi H, Takayama S, Sugimura T (1991) A new colon and mammary carcinogen in cooked food, 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP). *Carcinogenesis* **12**: 1503-1506.

Jin L, Wessely O, Marcusson EG, Ivan C, Calin GA, Alahari SK (2013) Prooncogenic factors miR-23b and miR-27b are regulated by Her2/Neu, EGF, and TNF α in breast cancer. *Cancer Res* **73**(9): 2884-2896.

Jover R, Bort R, Gomez-Lechon MJ, Castell JV (2002) Down-regulation of human CYP3A4 by the inflammatory signal interleukin-6: molecular mechanism and transcription factors involved. *FASEB J* **16**(13): 1799-1801.

Kapucuoglu N, Coban T, Raunio H, Pelkonen O, Edwards RJ, Boobis AR, Iscan M (2003) Expression of CYP3A4 in human breast tumour and non-tumour tissues. *Cancer Lett* **202**; 17-23.

Kasai S, Ishigaki T, Takumi R, Kamimura T, Kikuchi H (2013) Beta-catenin signaling induces CYP1A1 expression by disrupting adherent junctions in Caco-2 human colon carcinoma cells. *Biochim Biophys Acta* **1830**: 2509-2516.

Kato S, Kawase T, Alderman J, Inatomi N, Lieber CS (1990) Role of xanthine oxidase in ethanol-induced lipid peroxidation in rats. *Gastroenterology* **98**: 203-210.

Koop DR (1992) Oxidative and reductive metabolism of cytochrome P450 2E1. *FASEB J* **6**(2): 724-730.

Kumarakulasingham M, Rooney PH, Dundas SR, Telfer C, Melvin WT, Curran S, Murray GI. Cytochrome p450 profile of colorectal cancer: identification of markers of prognosis. *Clin Cancer Res* **11**(10): 3758-3765.

Kune S, Kune GA, Watson LF (1987) Case-control study of alcoholic beverages as etiological factors: the Melbourne Colorectal Cancer Study. *Nutr Cancer* **9**: 43-56.

Kurzawski M, Dzieziejko V, Post M, Wojcicki M, Urasinska E, Mietkiewski J, Drozdik M (2012) Expression of genes involved in xenobiotic metabolism and transport in end-stage liver disease: up-regulation of ABCC4 and CYP1B1. *Pharmacol Rep* **64**(4): 927-939.

Lee J-J, Drakaki A, Iliopoulos D, Struhl K (2011) MiR-27b targets PPAR γ to inhibit growth, tumour progression, and the inflammatory response in neuroblastoma cells. *Oncogene* **31**(33): 3818-3825.

Maihofner C, Charalambous MP, Gooderham NJ, and the colorectal cancer group (2003) Expression of cyclooxygenase-2 parallels expression of interleukin-1 β , interleukin-6 and NF-kappaB in human colorectal cancer. *Carcinogenesis* **24**(4): 665-671.

Malaplate-Armand C, Ferrari L, Masson C, Siest G, Batt AM (2003) Astroglial CYP1B1 up-regulation in inflammatory/oxidative toxic conditions: IL-1 β effect and protection by N-acetylcysteine. *Toxicol Lett* **138**(3): 243-251.

Mohri T, Nakajima M, Fukami T, Takamiya M, Aoki Y, Yokoi T (2010) Human CYP2E1 is regulated by miR-378. *Biochem Pharmacol* **79**(7): 1045-1052.

Morita M, Le Marchand L, Kono S, Yin G, Toyomura K, Nagano J, Mizoue T, Mibu R, Tanaka M, Kakeji Y, Maehara Y, Okamura T, Ikejiri K, Futami K, Maekawa T, Yasunami Y, Takenaka K, Ichimiya H, Imaizumi N (2009) Genetic polymorphisms of CYP2E1 and risk of colorectal cancer: the Fukuoka Colorectal Cancer Study. *Cancer Epidemiol Biomarkers Prev* **18**: 235-241.

Murray GI, Taylor, MC, McFadyen MCE, McKJay JA, Greenlee WF, Burke MD, Melvin WT (1997) Tumour-specific expression of cytochrome P450 CYP1B1. *Cancer Res* **57**: 3026-3031.

Nagasaki T, Hara M, Nakanishi H, Takahashi H, Sato M and Takeyama H (2014) Interleukin-6 released by colon cancer-associated fibroblasts is critical for tumour angiogenesis: anti-interleukin-6 receptor antibody suppressed angiogenesis and inhibited tumour-stroma interaction. *Br J Cancer* **110**: 469-478.

Norkina O, Dolganiuc A, Shapiro T, Kodys K, Mandrekar P, Szabo G (2007) Acute alcohol activates STAT3, AP-1 and Sp-1 transcription factors via the family of Src kinases to promote IL-10 production in human monocytes. *J Leukoc Biol* **82**(3): 752-762.

Nothlings U, Yamamoto JF, Wilkens LR, Murphy SP, Park SY, Henderson BE, Kolonel LN, Le Marchand L (2009) Meat and heterocyclic amine intake, smoking, NAT1 and NAT2 polymorphisms, and colorectal cancer risk in the multiethnic cohort study. *Cancer Epidemiol Biomarkers Prev* **18**: 2098-2106.

Pollack ES, Nomura AM, Heilbrun LK, Stemmermann GN, Green SB (1984) Prospective study of alcohol consumption and cancer. *N Engl J Med* **310**: 617-621.

Piipari R, Savela K, Nurminen T, Hukkanen J, Raunio H, Hakkola J, Mantyla T, Beaune P, Edwards RJ, Boobis AR, Anttila S (2000) Expression of CYP1A1, CYP1B1 and CYP3A, and polycyclic aromatic hydrocarbon-DNA adduct formation in bronchoalveolar macrophages of smokers and non-smokers. *Int J Cancer* **86**(5): 610-616.

Roberts BJ, Song B-J, Soh Y, Park SS, Shoaf SE (1995) Ethanol induces CYP2E1 by protein stabilization. *J Biol Chem* **270**: 29632-29635.

Rodriguez-Antona C, Donato MT, Boobis AR, Edwards RJ, Watts PS, Vicente Castell J, Gomez-Lechon MJ (2002) Cytochrome P450 expression in human hepatocytes and hepatoma cell lines: molecular mechanisms that determine lower expression in cultured cells. *Xenobiotica* **32**(6): 505-520.

Rodriguez M, Potter DA (2013) CYP1A1 regulates breast cancer proliferation and survival. *Mol Cancer Res* **11**(7): 780-792.

Seitz HK, Czygan P, Waldherr R, Veith S, Raedsch R, Kassmodel H, Kommerell B (1984) Enhancement of 1,2-dimethylhydrazine-induced rectal carcinogenesis following chronic ethanol consumption in the rat. *Gastroenterology* **86**: 886-891.

Sinha R, Peters U, Cross AJ, Kulldorf M, Weissfeld JL, Pinsky PF, Rothman N, Hayes RB (2005) Meat, meat cooking methods and preservation, and risk for colorectal adenoma. *Cancer Res* **65**: 8034-8041.

Smalley,W., Ray,WA, Daugherty,J, Griffin,MR (1999) Use of nonsteroidal anti-inflammatory drugs and incidence of colorectal cancer: a population-based study. *Arch Int Med* **159**: 161-166.

Suzuki H, Maruyama R, Yamamoto E and Kai M (2012) DNA methylation and microRNA dysregulation in cancer. *Mol oncol* **6**(6): 567-578.

Tenesa A, Dunlop MG (2009) New insights into the aetiology of colorectal cancer from genome-wide association studies. *Nat Rev Genet* **10**: 353-358.

Tindberg N, Baldwin HA, Cross AJ, Ingelman-Sundberg M (1996) Induction of cytochrome P450 2E1 expression in rat and gerbil astrocytes by inflammatory factors and ischemic injury. *Mol Pharmacol* **50**(5): 1065-1072.

Tsuchiya Y, Nakajima M, Takagi S, Taniya T, Yokoi T (2006) MicroRNA regulates the expression of human cytochrome P450 1B1. *Cancer Res* **66**(18): 9090-9098.

Wehbe H, Henson R, Meng F, Mize-Berge J, Patel T (2006) Experimental Therapeutics, Molecular Targets, and Chemical Biology: Interleukin-6 Contributes to Growth in Cholangiocarcinoma Cells by Aberrant Promoter Methylation and Gene Expression. *Cancer Res* **66**: 10517-10524.

Williams CS, Smalley W, DuBois RN (1997) Aspirin use and potential mechanisms for colorectal cancer prevention. *J Clin Invest* **100**: 1325-1329.

Yan H, Choi A, Lee BH, Ting AH (2011) Identification and functional analysis of epigenetically silenced in colorectal cancer cells. *PLoS One* **6**(6): e20628. doi: 10.1371/journal.pone.0020628.

Yang CH, Yue J, Fan M, Pfeffer LM (2010) IFN induces miR-21 through a signal transducer and activator of transcription 3-dependent pathway as a suppressive negative feedback on IFN-induced apoptosis. *Cancer Res* **70**(20): 8108-8116.

Figure titles and legends

Fig1. IL6 expression is increased in epithelial cells and stroma of colorectal tumour samples. Data are adapted from our previous publication (Maihofer *et al*,

2003) and are shown as fold change of immunohistochemistry score relative to control. Significant differences from normal tissue were calculated using Wilcoxon's signed-ranked test (* $p < 0.05$, *** $p < 0.001$).

Fig2. IL6 effect on CYP450 gene expression. HCT116 and SW480 cells were treated with 0, 100 and 1000pg/ml IL6 for 24 and 48 hours. *CYP1A1* (A), *CYP1B1* (B) and *CYP2E1* (C) expression were measured by RT-qPCR. Data were normalised to expression of *GAPDH* housekeeping gene and are shown relative to control. Significance was calculated using one-way ANOVA with a Dunnett post-test comparing treated groups to vehicle control and linear trend analysis (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). Data are presented as a mean of at least three biological replicates. Error bars represent the SEM.

Fig3. MiRNA involvement in IL6-mediated regulation of CYP1B1 and CYP2E1 gene expression. (A-D) HCT116 and SW480 cells were treated with 0 and 1000pg/ml IL6 for 24 and 48 hours. MiR378 (A) and miR27b (C) expression were measured by RT-qPCR. Fold change expression of miR378 was correlated with fold change expression of *CYP2E1* (B) and fold change expression of miR27b with fold change *CYP1B1* expression (D). Data were normalised to expression of U6 RNA and are shown relative to control. Significance was calculated using Student's t-test; Pearson's product-moment correlation coefficient test was used for correlation analysis (* $p < 0.05$, ** $p < 0.01$). Data are presented as a mean of at least three biological replicates. Error bars represent the SEM.

Fig4. STAT3 involvement in IL6-mediated regulation of CYP1B1 and CYP2E1 gene expression. (A) Potential STAT3 binding sites in the *CYP2E1* promoter region (1000bp upstream of the *CYP2E1* start site) predicted using TFSEARCH ver1.3 (Heinemeyer *et al*, 1998) (B, C) HCT116 and SW480 were treated with 1000pg/ml IL6 or a combination of IL6 and 25 μ M STAT3 inhibitor VII 5,15-

diphenylporphyrin for 24 hours (B) and 48 hours (C). *CYP2E1* (B) and *CYP1B1* (C) expression was measured by RT-qPCR. Data were normalised to expression of *GAPDH* housekeeping gene and are shown relative to control. (D) SW480 cells were treated with 1000pg/ml IL6 or a combination of IL6 and 25 μ M STAT3 inhibitor VII 5,15-diphenylporphyrin for 60min prior to being fixed and STAT3 binding to predicted sites in the *CYP2E1* promoter region were measured by ChIP-qPCR using an anti-STAT3 antibody. Data are presented as a mean of at least three biological replicates for STAT3 ChIP experiments and error bars represent the SEM. Significance was calculated using two-way ANOVA with a Bonferroni post-test comparing treated groups to vehicle control (** $p < 0.01$). ChIP experiments were repeated using IgG antibody to control for non-specific binding and performed in duplicate.

Fig5. IL6 induced downregulation of miR27b is mediated by DNA methylation. HCT116 and SW480 cells were treated with 1000pg/ml IL6, 4 μ M 5-aza-2'-deoxycytidine or a combination of IL6 and 5-aza-2'-deoxycytidine for 24 and 48 hours. (A) MiR27b expression was measured by RT-qPCR. Data were normalised to expression of U6 RNA and are shown relative to control. (B) *CYP1B1* expression was measured by RT-qPCR. Data were normalised to expression of *GAPDH* housekeeping gene and are shown relative to control. (C) Fold change expression of miR27b was correlated with fold change expression of *CYP1B1*. Significance was calculated using Student's t-test and one-way ANOVA a Dunnett post-test comparing treated groups to vehicle control; Pearson's product-moment correlation coefficient test was used for correlation analysis (* $p < 0.05$, ** $p < 0.01$). Data are presented as a mean of at least three biological replicates. Error bars represent the SEM.

Fig6. Immunohistochemical localisation of CYP450s in normal and adjacent malignant colonic epithelia. The presence of the immunoreactive protein is indicated by brown staining. Normal (A) and tumour (B) tissue treated with preimmune serum as primary antibody (negative control); normal (C) and tumour (D) tissue treated with anti-CYP1A1 primary antibody; normal (E) and tumour (F) tissue treated with anti-CYP1B1 primary antibody; normal (G) and tumour (H) tissue treated with anti-CYP2E1 primary antibody. (I) Expression of CYP1B1 in non-malignant colonic epithelial cells lying adjacent to tumour cells. (J) CYP2E1 expression in plasma cells. All sections are shown at x200 or x400 (inset magnification). (K) Expression of CYP450s in matched normal and adjacent malignant colonic epithelia from 40 patients. Significant differences from normal tissue were calculated using Wilcoxon's signed-ranked test (** $p < 0.01$, *** $p < 0.001$).

Fig7. Potential mechanism of IL6-mediated induction of CYP1B1 and CYP2E1 mRNA expression in colorectal cancer cells. IL6 secreted by stromal cells binds to the soluble IL6 receptor (sIL6R) in the stroma, which interacts with transmembrane protein gp130 at the cellular surface resulting in activation of JAK/STAT3 and PI3K/AKT pathways. Activated STAT3 forms a homodimer, translocates to the nucleus and binds to the CYP2E1 promoter region, thus inducing its transcription. Activated AKT phosphorylates DNMT1 leading to its nuclear translocation. Once in the nucleus, DNMT1 induces DNA methylation at a CpG island located near miR27b, thus preventing its expression. Lower expression of miR27b results in less binding to the CYP1B1 mRNA leading to less degradation and increased CYP1B1 mRNA expression.

Figure 1








Figure 2

A



B



C







Figure 3


A



B



C



D




Figure 4








Figure 5

A



B



C





Figure 6



K




Figure 7




Table1. Patient demographic information.

Patient	Age ¹	Sex	Tumor Site	Dukes' Stage	Drug History	Tobacco use	Alcohol ²
1	59	M	Rectum	A	None	No	1-7
2	58	F	Colon	B	Azathioprine, Insulin, Prednisolone, Lisinopril, Frusemide, Ferrous sulphate	No	1-7
3	79	M	Rectum	B	Atenolol, Bendrofluazide	No	8+
4	72	M	Colon	B	None	No	8+
5	60	F	Rectum	B	Lipostat	No	8+
6	66	F	Colon	C	5-Fluorouracil, Enalapril	No	0
7	69	M	Rectum	C	Co-codamol	No	1-7
8	52	M	Rectum	C	Adalat	No	1-7
9	68	M	Colon	B	Atenolol, Prednisolone, Warfarin, Ferrous sulphate, Diltiazem, Isosorbite mononitrate, Gliclazide, Co-danthramer	No	1-7
10	69	M	Rectum	C	Atenolol	No	8+
11	70	F	Rectum	B	Lithium, Levothyroxine	No	8+
12	72	M	Rectum	B	Captopril, Naproxen, Allopurinol, Isosorbite mononitrate, Frusemide, Atenolol, Prochlorperazine	No	1-7
13	56	M	Colon	A	None	No	8+
14	76	M	Colon	A	None	No	1-7
15	58	F	Colon	N/K	None	No	0
16	66	F	Colon	B	None	No	0
17	54	M	Rectum	C	None	No	1-7
18	49	M	Colon	B	None	No	8+
19	52	M	Rectum	B	None	No	8+
20	68	M	Colon	B	Salbutamol, Ferrous sulphate	No	0
21	63	F	Rectum	A	Salbutamol, Beclomethasone, Bendrofluazide	No	1-7
22	56	M	Colon	C	Losec	No	8+
23	68	M	Rectum	B	Sotalol, Aspirin	No	8+
24	80	F	Rectum	B	None	No	0
25	59	M	Rectum	C	None	No	8+
26	66	M	Rectum	N/K	None	Current	8+
27	59	F	Colon	N/K	Fibrogel	No	0
28	74	M	Colon	N/K	None	No	1-7
29	64	M	Rectum	B	Aspirin, Omeprazole, Simvastatin	No	0
30	72	F	Rectum	C	None	No	8+
31	50	M	Rectum	B	Betagan	Current	1-7
32	72	M	Rectum	B	Diclofenac, Propanolol, Adalat, GTN, Isosorbide mononitrate	No	8+
33	55	M	Colon	B	None	No	1-7
34	67	F	Rectum	C	Ibuprofen, Aspirin, Bendrofluazide	No	1-7
35	74	M	Colon	C	5-Fluorouracil	No	8+
36	73	M	Colon	A	None	No	8+
37	60	M	Colon	C	None	No	1-7
38	59	F	Rectum	B	Voltarol	No	0
39	73	M	Colon	C	5-Fluorouracil, Co-danthrusate, Manevac	No	1-7
40	70	M	Colon	C	Ferrous sulphate	No	0

N/K - Not known

¹ Age in years² Alcohol consumption in units per week (1 unit = half a pint of beer or one glass of wine or one shot of spirits)