

First evidence of a functional interaction between DNA quadruplexes and Poly(ADP-ribose) polymerase-1

V.A. Soldatenkov^{2*}, A.A. Vetcher^{3†}, T. Duka², S. Ladame^{1*}

¹*Institut de Science et Ingénierie Supramoléculaires, Université Louis Pasteur - CNRS, 8 allée Gaspard Monge, 67083 Strasbourg, France*

²*Department of Radiation Medicine, Lombardi Comprehensive Cancer Center, Georgetown University Medical Center, 3970 Reservoir Rd, NW, Washington, DC 20007, USA*

³*National Center for Biodefense and Infectious Diseases, George Mason University, 10900 University Blvd, Manassas, VA 20110, USA*

[†]*Present address: AlfaGene BioSciences, Inc., Gaithersburg, MD 20877*

(*)These authors contributed equally to this work and shall be regarded as the senior authors.

Correspondence to: Sylvain Ladame and Viatcheslav Soldatenkov

Emails : s.ladame@isis.u-strasbg.fr, soldates@georgetown.edu

Abstract: We discovered that the abundant human nuclear protein poly(ADP-ribose) polymerase (hPARP-1) binds to intramolecular DNA quadruplexes *in vitro* with high affinity and with a stoichiometry of two proteins for one quadruplex. Using an enzymatic assay, we have shown that hPARP-1 gets catalytically activated upon binding to G-quadruplexes localized at the c-kit promoter and human telomere regions. This is the first example of a truly functional quadruplex-protein interaction which has possible implications in understanding hPARP-1 mediated mechanisms of transcription regulation and telomere end protection.

It has been known for several decades that DNA sequences containing a high density of contiguous guanines grouped as clusters are able to adopt four-stranded secondary structures named guanine (G)-quadruplexes or tetraplexes.¹ Converging *in silico* and *in vitro* data have recently revealed high prevalence of such G rich DNA sequences throughout the human genome.²⁻⁴ The identification of prokaryotic and eukaryotic proteins that interact with these motifs also reinforces the hypothesis that quadruplexes do form *in vivo* and that their formation is biologically relevant.⁵⁻⁷ Although several lines of evidence link G-quadruplexes with DNA recombination,⁸ telomere maintenance⁹ and more recently regulation of gene transcription,¹⁰⁻¹² there is noticeable absence of examples for DNA quadruplex induced modulation of nuclear protein(s) function. When exploring protein-quadruplex interactions, attention has been almost exclusively focused on the effects of the natural DNA binding protein on the quadruplex structure and proteins have been classified according to their ability to stabilise, destabilise or promote the formation of DNA quadruplexes.⁶⁻⁷ Unnatural Cys₂His₂ zinc finger proteins have also been engineered that bind to quadruplex DNA with high affinity¹³⁻¹⁴ and block the biochemical functions of DNA polymerase and telomerase *via* quadruplex stabilization.¹⁵ However, to our knowledge, there is no example of a quadruplex - induced protein catalytic activation which would provide an additional proof of quadruplex existence *in vivo* and contribute to our understanding of quadruplex functions in nuclei. Human poly(ADP-ribose) polymerase (hPARP-1), a protein abundant in chromatin of eukaryotic cells, has been regarded as an intracellular sensor of DNA strand breaks.¹⁶ Upon binding to DNA breaks *via* its two zinc finger DNA binding domain (DBD), activated hPARP-1 cleaves NAD⁺ into nicotinamide and ADP-ribose and polymerases the latter onto nuclear acceptor proteins and hPARP-1 itself. It has also been implicated in the transcription of eukaryotic genes possibly by perturbing chromatin structure after being recruited to non-B DNA structures in gene regulating sequences.¹⁷⁻¹⁹

Although its importance, only limited data is available regarding interactions between hPARP-1 and undamaged genomic DNA.

An increasing number of oncogenes have been reported that possess a quadruplex forming sequence in their promoter.²⁰ An attractive hypothesis suggests that formation of these promoter quadruplexes *in vivo* could interfere with the transcription machinery, thus leading to a change of the level of expression of the corresponding gene. Although a significant body of *in vitro* and indirect evidences have recently been gathered that all support this idea of a quadruplex mediated regulatory mechanism¹⁰⁻¹², the exact role quadruplexes may play remains unknown. Herein, we discovered that the abundant human nuclear protein poly(ADP-ribose) polymerase (hPARP-1) binds to intramolecular DNA quadruplexes *in vitro* with high affinity and gets catalytically activated upon binding to G-quadruplexes. This is the first example of a truly functional quadruplex-protein interaction which has possible implications in understanding hPARP-1 mediated mechanisms of transcription regulation and telomere end protection.

Our first goal was to demonstrate the interaction between hPARP-1 and an intramolecular promoter DNA quadruplex. Two G-rich sequences have been identified within the nuclease hypersensitive region of the human proto-oncogene c-kit that encodes for a tyrosine kinase receptor. Both sequences are 31 nucleotides apart from each other and were shown to form G-quadruplex structures *in vitro*.²¹⁻²² As a model system we have chosen to study in detail the interaction between hPARP-1 and the quadruplex from the c-kit promoter which proved the most stable (c-kit-1).²¹ However we also investigated the interaction with other intramolecular quadruplexes, including the second c-kit quadruplex (c-kit-2)²² or the human telomeric quadruplex (Htelo). Surface Plasmon Resonance (SPR) was first used to simultaneously evaluate the affinity of purified hPARP-1 for biotinylated quadruplex, duplex and single strand DNAs immobilized on a streptavidin coated sensor chip (**Fig. 1a**).¹⁴ The sequence of the single-stranded oligonucleotide control was analogous to that of the c-kit-1 quadruplex but carrying five G→T

mutations to prevent quadruplex formation. Whilst no detectable binding to single stranded DNA was observed up to high protein concentrations, responses of comparable amplitude and proportional to protein concentration were observed for both c-kit-1 quadruplex and double-stranded DNA, indicative of a specific interaction with either structural transitions (quadruplex) or structural discontinuities (free ends) in DNA. Corrected sensorgrams were analyzed using a langmuirian global fit model and hPARP-1 was found to bind to double-stranded DNA break and to the c-kit-1 quadruplex with apparent Kd values of 18 and 65 nM respectively. We also found that hPARP-1 could bind to the human telomeric quadruplex, although with a significantly weaker affinity than to the c-kit-1 quadruplex (apparent Kd value of 5.8 μ M). hPARP-1 specific binding to the DNA quadruplexes was confirmed by a pull-down assay using the same single-stranded and c-kit-1 quadruplex biotinylated oligonucleotides as for the SPR experiments and streptavidin coated magnetic beads. Data indicate that only folded quadruplex sequences provide a binding site for hPARP-1, thus further supporting the notion that hPARP-1 binding to c-kit-1 sequence requires DNA structure specific recognition (**Fig. 1b**).

Insert Figure 1 here

We explored further the nature of the protein-DNA interaction using Atomic Force Microscopy (AFM) and a 200 bases long oligonucleotide from the c-kit promoter gene and carrying both natural quadruplex-forming sequences (bases 76 to 94 and 125 to 146). Quadruplex free single-stranded oligonucleotides are invisible or indistinguishable from the background under the commonly employed conditions of AFM observation in tapping mode and unmodified tip.²³ Therefore neither single-stranded nor quadruplex arms appear on corresponding panels. In the absence of DNA or in the presence of unfolded single-stranded DNA, hPARP-1 molecular volume has the typical unimodal distribution²⁴ with the maximum at 200-225 nm³. When hPARP-1 was incubated in the presence of the quadruplex containing oligonucleotide the distribution of frequencies has quasi-trimodal character, because of the

appearance of 1:1 hPARP-1-quadruplex complexes (maximum at 275-300 nm³) and hPARP-1 dimerization on DNA leading to formation of 2:1 complexes with the maximum of volume at 475-500 nm³ (**Fig. 2**). In addition to confirming hPARP-1 affinity for DNA quadruplexes, this AFM experiment also suggests that the protein binds to DNA as a dimer. However, because of the presence of two quadruplex forming sequences within the fragment of the c-kit promoter, one cannot completely rule out that the 2:1 protein/DNA complexes observed by AFM correspond to the co-existence of both quadruplexes, each bound to one hPARP-1 protein.

Insert Figure 2 here

In order to unambiguously clarify the stoichiometry of the complexes observed on a surface, we subsequently employed a spectroscopic method that could allow us to further characterize the nature of individual complexes in solution. hPARP-1 contains multiple tryptophan residues, and in particular four in its DBD that are highly sensitive to DNA binding.²⁵⁻²⁶ We observed the formation of the protein: DNA complexes by monitoring the tryptophan fluorescence quenching upon addition of different DNA sequences. Titrations were carried out using both c-kit quadruplexes, the human telomeric quadruplex, the c-myc quadruplex and a single-stranded mutated DNA fragment that cannot adopt a quadruplex conformation and which was used as a negative control. Whilst no significant fluorescence changes were observed with single-stranded DNA, even at high DNA concentration, a DNA concentration-dependent fluorescence quenching was obtained for all four quadruplexes tested which all showed similar profiles. A first fast fluorescence decrease was observed at low quadruplex concentration until a plateau was slowly reached at higher quadruplex concentration (**Fig. 1c** and **Supplementary Fig. 1**). The titration experiment was repeated with two initial protein concentrations (40 and 60 nM) and the binding stoichiometry was determined by the “tangent method”. In both cases, quenching saturation was obtained with half equivalent quadruplex, indicating the implication of two proteins for one quadruplex per complex, which is

in agreement with our AFM experiments. hPARP-1 was previously demonstrated to dimerize in solution only at high (submicromolar) protein concentration suggesting that the observed protein dimerization happens upon binding to quadruplex only. The observed level of fluorescence quenching at saturating quadruplex concentration (40-50%) is consistent with that observed by others when investigating hPARP-1 DBD binding to double-stranded DNA breaks. This result in addition to our previous studies on engineered zinc fingers binding to quadruplexes¹³⁻¹⁵ suggests that hPARP-1 could possibly bind to DNA quadruplexes via its zinc finger DBD. Protein dimerization has been previously demonstrated for hPARP-1 binding to 5'-recessed DNA ends, although considering the DBD of hPARP-1 only.²⁵⁻²⁶ Our observation that native human hPARP-1 protein dimerizes upon interaction with DNA quadruplexes is in line with previously reported suggestion on hPARP-1 dimerization derived from the enzyme kinetics study.²⁷ This supports the notion that protein dimer formation is required for hPARP-1 catalytic activation, and further substantiate the recent reports that hPARP-1 function is not limited to DNA damage repair, but that this protein also has a prominent role in normal cell physiology.

Cellular proteins have been identified that bind to quadruplexes and exhibit either a stabilising or destabilising effect on the DNA structure. In order to assess the effect of hPARP-1 on quadruplex stability and conformation we have carried out fluorescence resonance energy transfer (FRET) experiment using a c-kit-1 quadruplex dually labelled with two fluorophores, fluorescein (FAM) and tetramethyl-rhodamine (TAMRA) at the 5' and 3' termini respectively. When quadruplex is formed, fluorescence emission of FAM is efficiently quenched by the neighbouring TAMRA. When the quadruplex unfolds the distance between the fluorophore and the quencher increases and the emissions of both FAM and TAMRA become independent of each other.²⁸ Fluorescence experiments were carried out by titrating in a solution of dual-labelled c-kit-1 quadruplex with increasing amounts of hPARP-1 and monitoring the fluorescence emission of both fluorophores whilst exciting fluorescein at 470 nm. No significant change in the

fluorescence spectra was observed even at high protein concentration (up to 4 equivalents) suggesting that hPARP-1 does not unwind quadruplexes upon binding (data not shown).

Insert Figure 3 here

We finally investigated whether hPARP-1: quadruplex interaction was functional by measuring the effect of quadruplex DNA on hPARP-1 catalytic activity in the reaction of auto-poly(ADP-ribosylation). The formation of poly(ADP-ribose) induced by the interaction of hPARP-1 with the c-kit-1 and Htelo quadruplexes was compared to that obtained in the presence of a nuclease-treated salmon testes DNA acting as a hPARP-1 activator and used as a reference (**Fig. 3**). We show that DNA quadruplex induced the *in vitro* hPARP-1 catalyzed poly(ADP-ribose) synthesis while single-stranded oligonucleotides containing mutated sequences that cannot form a quadruplex exhibited no potential to stimulate hPARP-1 enzymatic activity. These results therefore establish DNA quadruplex as an effective co-enzymatic activator of hPARP-1 and indicate that poly(ADP-ribose) synthesis is affected by quadruplex structure. Although the requirement of DNA co-factor for hPARP-1 activity is well established in many studies²⁹ the precise mechanisms for enzymatic transition from inactive to active state are yet to be investigated. Our data, together with previous observations indicate that hPARP-1 can utilize different types of DNA co-factors (DNA strand interruptions, non B-DNA and DNA quadruplexes) to catalyse poly(ADP-ribosylation).^{27, 29-30} The structural diversity of enzyme co-factors that can be formed in the genome in several physiological contexts and following various pathophysiological stimuli provides a rationale for pleiotropic functions of hPARP-1 in eukaryotic cells.

Our data provide the first reported example of a functional interaction linking a promoter DNA quadruplex and an abundant natural protein. This has implications in the yet to be demonstrated existence of promoter quadruplexes *in vivo* but also in understanding the role hPARP-1 may play in undamaged cells, and in regulation of gene transcription in particular. It

has been proposed that hPARP-1 may exert its function in transcription regulation through direct binding to the gene-regulating sequences.¹⁸⁻¹⁹ However, the mechanism by which hPARP-1 is recruited to gene promoters in the absence of DNA damage remains unknown. We propose that promoter quadruplexes may act as hPARP-1 recruiting elements as part of its mechanism of transcription regulation. Our finding that at least two genes (pS2 and iNOS linked to breast cancer and oxidative stress respectively) for which transcription is known to be regulated by hPARP-1³¹⁻³² possess in their promoter a sequence that forms extremely stable quadruplexes *in vitro* (**Supplementary Fig. 2**) supports this novel hypothesis that we are now in the process of exploring. This hPARP-1: quadruplex interaction may also have implications in the mechanisms of telomere maintenance. Using SPR, we have demonstrated that hPARP-1 could also bind the intramolecular quadruplex formed by the single-stranded 3'-end of the human telomeric DNA although with a significantly weaker (90-fold) affinity (**Supplementary Fig. 3**). This would support the recent suggestion that hPARP-1 may be activated by and directly bind eroded telomeres as part of a DNA damage repair process.³³⁻³⁴

METHODS

DNA oligonucleotide preparation. The dual labelled oligonucleotide used for the FRET experiment and the 200 bases long oligonucleotide used for AFM were purchased from IBA (Germany). All other oligonucleotides were purchased from Sigma Genosis. All concentrations were expressed in strand molarity with a nearest-neighbour approximation for the absorption concentrations of the unfolded species.

The DNA quadruplexes were prepared in buffer containing 50 mM Tris.HCl at pH 7.4 and 100 mM KCl by heating to 95 °C for 5 min. After slow cooling for 6 h to room temperature, the oligonucleotide was stored at 4 °C. Biotinylated oligonucleotides of identical sequences,

quadruplex or single-stranded oligonucleotides, were used for the SPR, “pull-down” and hPARP-1 activity assays.

Surface Plasmon Resonance. All experiments were carried out on a Biacore 2000 Biosensor using freshly filtered and degassed buffers. The experiments were performed using four different immobilized DNA targets: two quadruplexes (**c-kit-1**) of sequence d(biotin-[C₃G₃CG₃CGCGAG₃AG₄AG₂]) and (**Htelo**) of sequence d(biotin-[GT₂A(G₃T₂A)₄G₂]), one double-stranded DNA (**ds DNA**) comprising the oligonucleotide d(biotin-[G₂CATAGTGCGTG₃CGT₂AGC]) hybridized with its complementary sequence and one unstructured single-stranded DNA of sequence comparable to that of **c-kit-1** but carrying G→T mutations to prevent quadruplex formation d(biotin-[C₃GTGCGTGCGCGAGTGAGTTG]) (**ss DNA**). Biotinylated oligonucleotides were loaded onto four separate lanes of a SA chip (Biacore) using the MANUAL INJECT command (Biacore 2000 control software) at a flow rate of 10 µl/min in loading buffer (50 mM Tris pH 7.4, 100 mM KCl). Running buffer (50 mM Tris pH 7.4, 100 mM KCl, 1 mM DTT, 50 µM zinc acetate) was then run through the four channels at a flow rate of 10µl/min for one hour before any protein was injected. hPARP-1 proteins were diluted in running buffer to concentrations of 0.98, 1.95, 3.9, 7.8, 15.6, 31.2, 62.5 and 125 nM. The KINJECT command was used to inject 60µL protein samples for a period of 3 min followed by a dissociation phase of 180 s. In between successive protein injections, the chip was regenerated by injecting running buffer containing 1 M KCl for 1 min.

Oligonucleotide “pull-down” assay. Streptavidin coated polystyrene beads (Dynabeads M-280, DYNALAB) were incubated with a 2.5 µM solution of either c-kit quadruplex (**c-kit-1**) or mutated single-stranded oligonucleotide (**ss ODN**) at room temperature for 30 min in accordance with manufacturer’s instructions. Protein binding assays were carried out in the presence of 12 nM recombinant full-length human PARP-1 protein (specific activity 10 U/µg, Trevigen) for 30 min with gentle agitation at room temperature in the binding buffer: 50 mM

Tris-HCl (pH 7.4), 50 mM KCl, 2 mM MgCl₂, 1 mM DTT, 50 μM Zn(OAc)₂. The protein-bound beads were separated using a magnetic separator (Dynal) and washed 5 times with 50 μl of low-salt buffer (50 mM Tris-HCl, pH 7.4, 100 mM KCl). Bound hPARP-1 proteins were eluted with 20 μl of NaCl (0.5-2.0 M) and subsequently analyzed by Western blotting using mouse anti-PARP-1 antibody (1:500, clone 42, BD Transduction Labs). Signals were detected using an enhanced chemoluminescence system (Amersham Biosciences).

Atomic Force Microscopy. 200 bases long oligonucleotide carrying c-kit quadruplex forming sequence (0.5 pmol) was incubated with hPARP-1 (0.1-10 pmol) in 10 μl of binding buffer (50 mM KOAc, 20 mM Tris-AcOH pH 7.9, 10 mM Mg(OAc)₂, 1 mM DTT) for ten minutes at 25°C and the complex was cross-linked with 0.8% glutaraldehyde for 15 minutes. The reaction was terminated by the addition of 5 μl of 2 M Tris-HCl (pH 7.7). High molecular weight components were transferred in the binding buffer using G-25 spin columns, samples were diluted 5-fold with this buffer placed onto freshly cleaved mica (Ted Pella), rinsed with deionized water and dried with argon. In some experiments oligonucleotides were heat-denatured at 70 °C for 10 minutes and subsequently annealed in the K⁺ and Na⁺ - free deposition buffer (10 mM Bis-Tris-Propane-HCl, 10 mM MgCl₂, 1 mM DTT, pH 7.0) to achieve DNA quadruplexes unfolding. AFM images were acquired in dry air with a Nanoscope III AFM Instrument (Veeco/Digital Instruments) in close contact (tapping) mode using tapping mode etched OMCL-AC160TS silicon AFM probes (Olympus Optical). The images were processed, and the measurements were performed with version 2.2.85(5.1) Femtoscan software (Advanced Technologies Center). The histograms were created from the measurements of at least 80 unobstructed objects.

Fluorescence Spectroscopy. Fluorescence emission spectra were recorded in quartz cells at 20 °C on a Jobin Yvon Fluorolog 3.22 instrument. The excitation and emission bandwidths were fixed to 8 and 5 nm respectively. FRET experiments were carried out using a 500 μL quartz

cuvette containing a solution of 10 nM dual-labelled annealed c-kit quadruplex in 50 mM Tris pH 7.4, 100 mM KCl, 50 μ M Zn(OAc)₂ and 1 mM DTT. Concentrated protein aliquots (from a 1 μ M stock solution in buffer) were directly added to the quadruplex solution. The spectra were recorded between 485 and 680 nm while exciting at 470 nm.

Fluorescence titration experiments were carried out in a 500 μ L quartz cuvette containing a solution of 500 μ L hPARP-1 (40 nM) in 50 mM Tris pH 7.4, 100 mM KCl, 50 μ M Zinc acetate and 1 mM DTT. Concentrated DNA quadruplex aliquots (from 500 nM, 5 μ M and 50 μ M stock solutions in buffer) were directly added to the protein solution. The spectra were recorded between 300 and 550 nm while exciting at 295 nm. Prior to experiment, the protein at the desired concentration was kept in a low-binding Eppendorf tube at 0 °C for 1 hour. It was then transferred into a quartz cell for the time of the experiment. Fluorescence of the free protein was shown to be stable for the average duration of an experiment, proving that photobleaching and protein adsorption on cell walls previously observed by others²⁴⁻²⁵ were negligible within our experimental conditions.

hPARP-1 Activity Assays. Reactions of auto-poly(ADP-ribosylation) were carried out in a buffer containing 50 mM Tris-HCl (pH 7.4), 50 mM KCl, 2 mM MgCl₂, 1 mM DTT, 50 μ M Zn(OAc)₂ in the presence of 133 μ M NAD⁺ (Trevigen) and structurally different types of DNA (see figure legend) as a co-factor. ADP-ribosylation was induced by the addition of recombinant full-length hPARP-1 (specific activity 10 U/ μ g, Trevigen) and the reaction mixture (15 μ l volume) was incubated at 25 °C for 30 min. In some reactions, the hPARP-1 inhibitor EB-47 (Axxora) was added in final concentration of 400 nM. The reactions were terminated by adding 2X SDS-PAGE loading buffer, samples were heated at 100 °C for 5 min and resolved by SDS-PAGE in a 4-12% gradient bis-Tris gel (NuPAGE, Invitrogen). In vitro auto-modified hPARP-1 (Biomol) was loaded onto the gel (50 ng/lane) alongside the reaction samples as an immunoblotting standard. The extent of hPARP-1 auto-poly(ADP-ribosylation) was determined

by Western blot analysis with a mouse monoclonal anti-poly(ADP-ribose) antibody (1:500, Biomol). For immunodetection of hPARP-1 the blots were re-probed with mouse anti-PARP antibody (1:500, clone 42, BD Transduction Labs). Signals were detected using an enhanced chemiluminescence system (Amersham Biosciences).

ACKNOWLEDGEMENTS

This work was supported in part by National Institutes of Health grant CA074175 (VAS) and by the CNRS (SL). The authors thank Pr. S. Balasubramanian for critically reading this manuscript.

AUTHOR CONTRIBUTIONS

S.L. and V.A.S. designed the project and wrote the manuscript. S.L. designed and performed SPR and fluorescence experiments. A.A.V. designed and performed AFM experiment. T.D. performed pull-down and enzyme activation assays.

REFERENCES

1. Parkinson, G.N. (2006) Fundamentals of quadruplex structures. In *Quadruplex Nucleic Acids*; Neidle, S., Balasubramanian, S., Eds; RSC Biomolecular Sciences Publishing, Cambridge.
2. Huppert, J.L., and Balasubramanian, S. (2005) Prevalence of quadruplexes in the human genome, *Nucleic Acids Res.* *33*, 2908-2916.
3. Todd, A.K., Johnston, M., and Neidle, S. (2005) Highly prevalent putative quadruplex sequence motifs in human DNA, *Nucleic Acids Res.* *33*, 2901-2907.
4. Eddy, J., and Maizels, N. (2006) Gene function correlates with potential for G4 DNA formation in the human genome, *Nucleic Acids Res.* *34*, 3887-3896.
5. Simonsson, T. (2001) G-quadruplex DNA structures: variations on a theme, *Biol. Chem.* *382*, 621-628.

6. Maizels, N. (2006) Dynamic roles for G4 DNA in the biology of eukaryotic cells, *Nat. Struct. Mol. Biol.* *13*, 1055-1059.
7. Fry, M. (2007) Tetraplex DNA and its interacting proteins, *Frontiers in Biosciences* *12*, 4336-4351.
8. Shukla, A.K., and Roy, K.B. (2006) Rec A-independent homologous recombination induced by a putative fold-back tetraplex DNA, *Biol. Chem.* *387*, 251-256.
9. Paeschke, K., Simonsson, T., Postberg, J., Rhodes, D., and Lipps, H.J. (2005) Telomere end-binding proteins control the formation of G-quadruplex DNA structures *in vivo*, *Nat. Struct. Mol. Biol.* *12*, 847-854.
10. Siddiqui-Jain, A., Grand, C.L., Bearss, D.J., and Hurley, L.H. (2002) Direct evidence for a G-quadruplex in a promoter region and its targeting with a small molecule to repress c-MYC transcription, *Proc. Natl. Acad. Sci. U.S.A.* *99*, 11593-11598.
11. Bejugam, M., Sewitz, S., Shirude, P.S., Rodriguez, R., Shahid, R., and Balasubramanian, S. (2007) Trisubstituted isoalloxazines as a new class of G-quadruplex binding ligands: small molecule regulation of c-kit oncogene expression, *J. Am. Chem. Soc.* *129*, 12926-12927.
12. Dexheimer, T.S., Fry, M., and Hurley, L.H. (2006) DNA quadruplexes and gene regulation. In *Quadruplex Nucleic Acids*; Neidle, S., Balasubramanian, S., Eds; RSC Biomolecular Sciences Publishing, Cambridge.
13. Isalan, M., Patel, S.D., Balasubramanian, S., and Choo, Y. (2001) Selection of zinc fingers that bind single-stranded telomeric DNA in the G-quadruplex conformation, *Biochemistry* *40*, 830-836.
14. Ladame, S., Schouten, J.A., Roldan, J., Redman, J.E., Neidle, S., and Balasubramanian, S. (2006) Exploring the recognition of quadruplex DNA by an engineered Cys2-His2 zinc finger protein, *Biochemistry* *45*, 1393-1399.

15. Patel, S.D., Isalan, M., Gavory, G., Ladame, S., Choo, Y., and Balasubramanian, S. (2004) Inhibition of human telomerase activity by an engineered zinc finger protein that binds G-quadruplexes, *Biochemistry* 43, 13452-13458.
16. Dantzer, F., Ame, J.C., Schreiber, V., Nakamura, J., Menissier-de Murcia, J., and de Murcia, G. (2006) Poly(ADP-ribose) polymerase-1 activation during DNA damage and repair, *Methods Enzymol.* 409, 493-510.
17. Kraus, W.L., and Lis, J.T. (2003) Parp goes transcription, *Cell* 113, 677-683.
18. Soldatenkov, V.A., Chasovskikh, S., Potaman, V.N., Trofimova, I., Smulson, M.E., and Dritschilo, A. (2002) Transcriptional repression by binding of poly(ADP-ribose) polymerase to promoter sequences, *J. Biol. Chem.* 277, 665-670.
19. Potaman, V.N., Shlyakhtenko, L.S., Oussatcheva, E.A., Lyubchenko, Y.L., and Soldatenkov, V.A. (2005) Specific binding of poly(ADP-ribose) polymerase-1 to cruciform hairpins, *J. Mol. Biol.* 348, 609-615.
20. Huppert, J.L., and Balasubramanian S. (2007) G-quadruplexes in promoters throughout the genome, *Nucleic Acids Res.* 35, 406-413.
21. Fernando, H., Reszka, A., Huppert, J.L., Ladame, S., Rankin, S.R., Neidle, S., and Balasubramanian, S. (2006) A conserved quadruplex motif located in a transcription activation site of the human c-kit oncogene, *Biochemistry* 45, 7854-7860.
22. Rankin, S., Reszka, A.P., Huppert, J., Zloh, M., Parkinson, G.N., Todd, A.K., Ladame, S., Balasubramanian, S., and Neidle, S. (2005) Putative DNA quadruplex formation within the human c-kit oncogene, *J. Am. Chem. Soc.* 127, 10584-10589.
23. Potaman, V.N., Bissler, J.J., Hashem, V.I., Oussatcheva, E.A., Lu, L., Shlyakhtenko, L.S., Lyubchenko, Y.L., Matsuura, T., Ashizawa, T., Leffak, M., Benham, C.J., and Sinden, R.R. (2003) Unpaired structures in SCA10 (ATTCT)_n(AGAAT)_n repeats, *J. Mol. Biol.* 326, 1095-1111.

24. Bergkvist, M., Carlsson, J., and Oscarsson, S. (2001) A method for studying protein orientation with atomic force microscopy using relative protein volumes, *J. Phys. Chem. B* 105, 2062-2069.
25. Pion, E., Bombarda, E., Stiegler, P., Ullmann, G.M., Mely, Y., de Murcia, G., and Gerard, D. (2003) Poly(ADP-ribose) polymerase-1 dimerizes at a 5' recessed DNA end in vitro: a fluorescence study, *Biochemistry* 42, 12409-12417.
26. Pion, E., Ullmann, G.M., Ame, J.C., Gerard, D., de Murcia, G., and Bombarda E. (2005) DNA-induced dimerization of poly(ADP-ribose) polymerase-1 triggers its activation, *Biochemistry* 44, 14670-14681.
27. Lonskaya, I., Potaman, V.N., Shlyakhtenko, L.S., Oussatcheva, E.A., Lyubchenko, Y.L., and Soldatenkov, V.A. (2005) Regulation of poly(ADP-ribose) polymerase-1 by DNA structure-specific binding, *J. Biol. Chem.* 280, 17076-17083.
28. Salas, T.R., Petruseva, I., Lavrik, O., Bourdoncle, A., Mergny, J.L., Favre, A., and Saintomé, C. (2006) Human replication protein A unfolds telomeric G-quadruplexes, *Nucleic Acids Res.* 34, 4857-4865.
29. Schreiber, V., Dantzer, F., Ame, J.C., and de Murcia, G. (2006) Poly(ADP-ribose): novel functions for an old molecule, *Nat. Rev. Mol. Cell Biol.* 7, 517-528.
30. Kun, E., Kirsten, E., Mendeleyev, J., and Ordahl, C.P. (2004) Regulation of the enzymatic catalysis of poly(ADP-ribose) polymerase by dsDNA, polyamines, Mg²⁺, Ca²⁺, histones H1 and H3, and ATP, *Biochemistry* 43, 210-216.
31. Yu, Z., Kunczewicz, T., Dubinsky, W.P., and Kone, B.C. (2006) Nitric oxide-dependent negative feedback of PARP-1 trans-activation of the inducible nitric-oxide synthase gene, *J. Biol. Chem.* 281, 9101-9109.

32. Ju, B.G., Lunyak, V.V., Perissi, V., Garcia-Bassets, I., Rose, D.W., Glass, C.K., and Rosenfeld, M.G. (2006) A topoisomerase IIbeta-mediated dsDNA break required for regulated transcription, *Science* 312, 1798-1802.
33. d'Adda di Fagagna, F., Hande, M.P., Tong, W.M., Lansdorp, P.M., Wang, Z.Q., and Jackson, S.P. (1999) Functions of poly(ADP-ribose) polymerase in controlling telomere length and chromosomal stability, *Nat Genet.* 23, 76-80.
34. Gomez, M., Wu, J., Schreiber, V., Dunlap, J., Dantzer, F., Wang, Y., and Liu, Y. (2006) PARP1 is a TRF2-associated poly(ADP-ribose)polymerase and protects eroded telomeres, *Mol. Biol. Cell* 17, 1686-1696.

FIGURE LEGENDS

Figure 1 *hPARP-1 binds to the c-kit-1 quadruplex in vitro.* (a) *left:* sensorgram overlay for hPARP-1 binding to immobilized c-kit quadruplex at 6 different hPARP-1 concentrations (0.98 to 31.25 nM bottom to top); *right:* sensorgram overlay for hPARP-1 (3.9 nM) binding to single-stranded DNA (blue), double-stranded DNA (red) and c-kit-1 quadruplex (green). (b) Purified recombinant hPARP-1 was bound to streptavidin coated magnetic beads coupled with either 5'-biotinylated c-kit-1 quadruplex oligonucleotides (**Q-ODN**) or control single-stranded quadruplex-free oligonucleotide (**ssODN**). Bound proteins were eluted with increasing concentrations of NaCl (0.5-2.0M) and analyzed by Western blotting using monoclonal anti-hPARP-1 antibodies. (c) Titration curve for binding of hPARP-1 to the c-kit-1 quadruplex. The hPARP-1 concentration was 40 nM. The linear parts of the binding curve are fitted and extrapolated separately with a linear fitting (---). The intersection at a quadruplex concentration of 20 nM indicates a protein/DNA stoichiometry of 2:1.

Figure 2 *AFM analysis of hPARP-1 binding to DNA quadruplexes.* 200bp oligonucleotides carrying c-kit quadruplex-forming sequences and purified human hPARP-1 protein were used in the binding assays. **A:** mica before deposition. **B:** heat-denatured (unfolded) oligonucleotides in the K^+ , Na^+ - free buffer. **C:** folded DNA quadruplexes **D:** representative AFM images of free hPARP-1 proteins (top panel) and hPARP-1-quadruplex binding reactions (middle and bottom panels). The length of the scale bars is 50 nm. **E:** histograms representing the distribution of molecular volume frequencies in samples shown in **D**. The molar ratios of DNA to protein are indicated on the linkers between AFM image and corresponding histogram. Pictograms on the top indicate hPARP-1 protein in its monomeric form (red), hPARP-1 in complex with DNA quadruplex (blue), and hPARP-1 dimers (green) formed upon binding to DNA.

Figure 3 *hPARP-1 enzymatic activity is stimulated by DNA G-quadruplexes.* Immunoblot analyses of the *in vitro* poly(ADP-ribose) [**PAR**] synthesis by human PARP-1 stimulated by different types of DNA substrates in the presence of 133 μM βNAD^+ . *In vitro* poly(ADP-ribosyl)ated hPARP-1 [**AM-hPARP**] and PAR synthesis stimulated by 16.6 ng/ μl nuclease-treated salmon testes DNA [**NT-DNA**] are shown as a reference (lines 1-3). Lane 4, the reaction mixture contained no hPARP-1 and DNA [**Blank**]. The formation of PAR induced by interaction of hPARP-1 with c-kit-1 promoter [**c-kit-1**, lines 5-10] and human telomeric [**Htelo**, lines 11-13] quadruplexes (8.3 μM) was evaluated in the presence of increasing hPARP-1 content. c-kit-1 and Htelo DNA quadruplexes both induced the *in vitro* hPARP-1 catalyzed PAR synthesis while corresponding single-stranded oligonucleotides (lines 5 and 11) containing mutated sequences that cannot form a quadruplex exhibited no potential to stimulate hPARP-1 enzymatic activity. The inhibitor of hPARP-1 enzyme, EB-47 (400 nM), prevents hPARP-1 activation by both NT-DNA and c-kit-1 quadruplex (lines 3 and 10). Immunodetection of hPARP-1 in the corresponding samples is shown in the bottom panel.

Figure 1




Figure 2





Figure 3



Graphical abstract

