



University of Dundee

The timing of mandibular tooth formation in two African groups

Elamin, Fadil; Hector, Mark; Liversidge, Helen M.

Published in:
Annals of Human Biology

DOI:
[10.1080/03014460.2016.1213313](https://doi.org/10.1080/03014460.2016.1213313)

Publication date:
2017

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Elamin, F., Hector, M. P., & Liversidge, H. M. (2017). The timing of mandibular tooth formation in two African groups. *Annals of Human Biology*, 44(3), 261-272. DOI: 10.1080/03014460.2016.1213313

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

"This is an Accepted Manuscript of an article published by Taylor & Francis in Annals of Human Biology on 13th July 2016, available at <http://www.tandfonline.com/10.1080/03014460.2016.1213313>."

ORIGINAL ARTICLE

The timing of mandibular tooth formation in two African groups

Fadil Elamin^{a,b,c}, Mark P. Hector^d and Helen M. Liversidge^a

^aQueen Mary University of London, Institute of Dentistry, Bart's and The London School of Medicine and Dentistry, London, UK; ^bKhartoum Centre for Research and Medical Training, Khartoum, Sudan; ^cEl Razi Dental School, Khartoum, Sudan; ^dDental School, University of Dundee, Dundee, UK

ABSTRACT

Background: Ethnic differences in the timing of human tooth development are unclear.

Aim: To describe similarities and differences in the timing of tooth formation in two groups of Sudanese children and young adults.

Subjects and methods: The sample consisted of healthy individuals from Khartoum, Sudan, aged 2–23 years. The Northern group was of Arab origin (848 males, 802 females) and the Western group was of African origin (846 males, 402 females). Each mandibular left permanent tooth from first incisor to third molar was assessed from dental radiographs into one of 15 development stages. Mean ages at entry for 306 tooth stages were calculated using probit regression in males/females in each group and compared using a *t*-test.

Results: Mean ages were not significantly different in most tooth stage comparisons *between* ethnic groups for both males (61/75) and females (56/76), despite a tendency of earlier mean ages in the Western group. Mean ages for most tooth stage comparisons between males and females (137/155) were not significantly different *within* ethnic groups suggesting low sexual dimorphism.

Conclusion: The mean ages of most mandibular tooth formation stages were generally not significantly different between ethnic groups or between males and females in this study.

Abbreviations: I1: mandibular central incisor; I2: mandibular lateral incisor; C: mandibular canine; P1: mandibular first premolar; P2: mandibular second premolar; M1: mandibular first molar; M2: mandibular second molar; M3: mandibular third molar; Cr: crypt; Ci: initial cusp formation; Cco: coalescence on cusps; Coc: crown outline complete; C1/2: crown half complete; C3/4: crown three quarters complete; Cc: crown complete; Ri: initial root formation; Rcl: root cleft formation; R1/4: root length one quarter; R1/2: root length one half; R3/4: root length three quarters; Rc: root length complete; A1/2: apex half closed; Ac: apex closed

ARTICLE HISTORY

Received 19 August 2014

Revised 3 June 2016

Accepted 14 June 2016

Published online 22 June 2016

KEYWORDS

Ethnic origin; tooth formation; sexual dimorphism; individual variation; group differences

Introduction

Human teeth grow in a highly organised sequence in relation to each other prior to their eruption into the oral cavity. Teeth appear in maxillary and mandibular bones as individual units and follow a sequence of development until all teeth are fully formed. Tooth formation from crown tip to root apex follows a predictable morphological path. How this relates to the timing of consecutive stage formation between human groups is less clear. Understanding the effects of biological factors on the timing of tooth formation is important in human biology, forensic anthropology, evolutionary, medical and anthropological fields.

Maturing teeth are regulated by complex biological clocks in their trajectory to full maturity (Papagerakis et al., 2014). This is particularly evident in crown formation (Antoine et al., 2009). Children vary considerably with respect to maturing teeth in relation to chronological age by a number of years, but are still considered biologically normal (Demirjian & Levesque, 1980; Levesque et al., 1981; Liversidge, 2011; Liversidge et al., 2006; Nystrom et al., 2007).

In light of the large age variation between individuals, the precise effect of biological factors such as sex and ethnicity on the timing of dental maturity events needs to be explored in large groups or populations across world regions. The effect of biological factors on the complex and lengthy process of forming teeth requires further attention.

Methods of sampling and analysis have been called into question (Bennett et al., 1999). In addition, the overall timing of tooth formation is relatively unaffected by severe biological insults such as malnutrition, despite affecting skeletal and somatic growth (Elamin & Liversidge, 2013; Garn et al., 1965). The mechanisms that insulate the forming teeth from severe environmental pressures such as malnutrition are not clear. Other evidence suggests that biological factors such as sex and ethnicity influence the timing of tooth formation than previously (Liversidge, 2011).

Our understanding of the timing of tooth formation

hdw046

2016-08-02 11:06:07

Delete Bennett et al. insert Smith

hdw046

2016-08-02 10:49:25

Nyström

h.m.liversidge@qmul.ac.uk insert
f.elamin@qmul.ac.uk
Khartoum, Sudan

Table 1. Radiographic studies that describe timing of individual permanent tooth maturation.

Country	Authors	n male, female	Age range	Teeth*	Statistics	Dispersion
Canada	Anderson et al. (1976)	121, 111	3–20	1	mean age L	SD
	Thompson et al. (1975)	121, 111	3–20	2	mean age L	—
	Demirjian & Levesque (1980)	2705, 2732	2–19	3	% smoothed curves	—
China	Levesque et al. (1981)	2278, 2367	7–25	8	% smoothed curves	—
	Zhao et al. (1990)	465, 438	3–16	3	50th percentile	—
Finland	Haavikko (1970)	615, 547	2–21	2	50th percentile	10th, 90th percentile
	Nyström et al. (2007)	966, 1004	0–25	2	50th percentile	—
South Africa, UK	Liversidge (2008)	390, 335	5–23	8	mean, logistic regression	SD †, SE
UK	Liversidge & Speechly (2001)	263, 258	4–9	3	mean, probit regression	SD
	Liversidge (2011)	529, 521	2–22	2	mean, logistic regression	SD †, SE
USA	Garn et al. (1958)	255	0–15	6	mean age L	—
	Garn et al. (1959)	285	0–15	5	50th percentile	—
	Fanning (1961)	48, 51	3–12	3, 7	mean age L	—
	Fanning & Brown (1971)	151, 139	0–22	4	mean age L	—
	Moorrees et al. (1963)	184, 161	0–22	2, 7	mean age L	—
Various	Liversidge et al. (2006)	4522, 4480	2–16	3	mean, logistic regression	—

Age range in years.


*1, all maxillary and mandibular; 2, all mandibular; 3, all mandibular except M₃; 4, mandibular C, P₁, P₂, M₁, M₂, M₃; 5, maxillary P₁, P₂, M₁, M₂; 7, maxillary incisors; 8, M₃; L, longitudinal; SD, standard deviation; SE, standard error.

† needs correcting factor, ‡ illustrated as error bar.

number of methodological factors. Most studies on the timing of tooth formation have samples of limited age ranges, small numbers, and used non-cumulative statistical approaches. Samples that contain limited age ranges allow only a few tooth stages to be investigated and statistically significant differences may not necessarily amount to biological importance in light of the inherent variation seen between individuals (Bennett et al., 1991). Part of the difficulty is due to maturation of the tooth being divided into many stages (usually 8–15). This reduces the number of children per stage, despite a large sample size and wide age range. Another challenge is that teeth mature over a long period of time in relation to other body systems, developing from prior to birth into early adulthood when individuals undergo marked morphologic and skeletal change with genetic and other environmental factors playing a role.

Only a handful of studies describe the timing of individual tooth stages using appropriate statistical methods and dispersion parameters across age and these are listed in Table 1 (Anderson et al., 1976; Demirjian & Levesque, 1980; Fanning & Brown, 1971; Garn et al., 1959; Haavikko, 1970; Levesque et al., 1981; Liversidge, 2008, 2011; Liversidge & Speechly, 2001; Liversidge et al., 2006; Moorrees et al., 1963; Nyström et al., 2007; Thompson et al., 1975; Zhao et al., 1990). Some directly compare ethnic/regional groups (Liversidge, 2008, 2011; Liversidge et al., 2006), one notes differences for the third molar (Liversidge, 2008), whilst others showed similarities (Liversidge, 2011; Liversidge et al., 2006) in formation times. Most of these studies describe tooth formation in children of European origin and there are gaps in the literature describing the timing of tooth formation in other world groups, particularly those in Africa. The timing of formation of the mandibular third molar in South African Blacks (Liversidge, 2008) is the only tooth documented in this way from an African population.

The aim of our study was to investigate the effect of ethnic group and sex on the timing of mandibular tooth formation in two groups of dental patients in Sudan, with a wide age range using well-defined ethnic grouping, age structure and statistical analysis.

**Figure 1.** Map of Sudan, pre-cessation into two countries in 2011, where the study was conducted.

Materials and methods

The subjects (2.83–23.96 years, $n = 2898$) for this investigation were part of a population survey that considered tooth formation, anomalies and disease in Khartoum, Sudan (Elamin, 2011). Sample selection followed the design from the Strengthening and Reporting of Observational Studies in Epidemiology guidelines (STROBE) (von Elm et al., 2007). Arab tribes from the North of Sudan (Jaali, Mahasi, Shaigi, Bedairi, Halfawi and Dongalawi groups) and Western Sudanese tribes of African origin, namely Fur and Zagawa, were studied (Figure 1).

Table 2. Age and sex distribution of study sample. The Northern group was of Arab origin and the Western group was of African origin.

Age	North group			West group		
	F	M	Total	F	M	Total
2p	1	4	5	—	—	—
3p	31	37	68	4	4	8
4p	28	32	60	4	6	10
5p	31	47	78	30	21	51
6p	71	68	139	23	54	77
7p	62	56	118	22	61	83
8p	60	32	92	28	35	63
9p	45	35	80	52	45	97
10p	30	30	60	25	49	74
11p	20	29	49	57	150	207
12p	28	19	47	52	130	182
13p	18	41	59	26	113	139
14p	30	50	80	14	86	100
15p	31	34	65	16	43	59
16p	37	36	73	12	21	33
17p	61	52	113	6	7	13
18p	55	65	120	4	3	7
19p	34	48	82	9	11	20
20p	35	41	76	5	4	9
21p	34	39	73	9	2	11
22p	41	31	72	3	1	4
23p	19	22	41	—	—	—
Total	802	848	1650	402	846	1248

M, male; F, female; 2p, includes individual aged from 2.00–2.99 years, etc.

Subjects from the Northern group were randomly selected from pre-schools, Khalwas (religious schools), mainstream schools and universities between January 2007 and May 2012 and stratified by school, using a cluster sampling method (Bennett et al., 1991). Schools, pre-schools, Khalwas and universities in the three localities (Bahri, Umdurman and Khartoum City) in Khartoum were chosen from a list of schools in these localities obtained from the Ministry of Education and sampled where safety permitted. The Western group were drawn from schools and Khalwas located in and around camps for the internally displaced in Khartoum. Children were excluded if the date of birth was unknown or if they had craniofacial anomalies. The age and sex distribution of the group is shown in Table 2. For practical reasons, radiographs were limited to individuals ≥ 3 years and older. Ethical approval was granted by the Ethics committee at El Razi Dental School prior to the study (01/11/2006). Verbal and written consent were obtained from individuals and from parents of minors. A dental examination was carried out and, where appropriate, a dental panoramic radiograph taken (Orthophos Model: D3200, Siemens, Germany). The same machine was used in this study but was moved between two locations, a dental school in southern Khartoum and a dental hospital in the northern part. The radiograph involved taking a standard view with the head in the Frankfort plane. The film was manually developed by experienced dental staff in preparation for clinical use.

The radiographs were digitised, decoded and randomised for blind scoring by a person other than the investigator. Age of each subject was converted to decimal age. Permanent mandibular left teeth were staged by the first author (FE) from the radiographs using the 14 crown and




Figure 2. Mandibular tooth formation stages after Moorrees et al. (1963), with the addition of crypt stage for molars.

root stages after Moorrees et al. (1963) in addition to staging the crypt of the third molar, i.e. 120 stages (Figure 2). We only assessed mandibular teeth as maxillary teeth are less clearly visualised on panoramic radiographs due to superimposition of the hard palate. Intra-examiner reliability of stage assessment from 90 panoramic radiographs was assessed by Cohen's Kappa, showing excellent agreement ($\kappa = 0.91$). SPSS, Release Version 17.0 (#SPSS, Inc., 2009, Chicago IL) was used to analyse data. Probit regression was used to calculate the cumulative mean age at entry or transition into tooth stages, where at least 10 individuals per year of age were available.

A total of 309 mean ages at entry were available to compare from a total of 420 possible comparisons in four groups of different ethnic origin and sex (male–female groups from North and West; male–male and female–female groups from both regions). Missing data in Tables 3–9 are the tooth stages that develop before the third birthday, prior to the minimum age in our sample. Most tooth stages were observed across a wide age range and for most calculations at least 10 individuals were present in consecutive year age groups.

The number of available comparisons is shown in summary in Tables 3 and 4: 75 and 76 comparisons of Northern–Western males and females, respectively, 80 comparisons of Northern male–female and 75 Western male–female groups. The mean age of stage attainment for each tooth stage was compared between male (North/West) and female (North/West) groups of different ethnicity. Sex comparisons were carried out within ethnic groups using a *t*-test, with a significance level of $p < 0.05$.

Table 3. Ethnic group comparisons. Summary of 75 comparisons of mean age entering mandibular tooth stages in North-West males and 76 in North-West females showing predominant similarity between ethnic groups. Missing tooth stages occurred at a younger age than our sample and could not be calculated.

Comparisons*	I1	I2	C	P1	P2	M1	M2	M3
<i>Male</i>								
Cr								NS
Ci								"
Cco								NS
Coc					NS		NS	NS
C1/2					NS		NS	NS
C3/4				NS	NS		NS	#
Cc			NS	NS	"		NS	NS
Ri			NS	"	#	NS	NS	NS
Rcl			—	—	—	NS	NS	NS
R1/4	NS	NS	NS	"	NS	NS	"	NS
R1/2	NS	NS	NS	NS	#	NS	NS	NS
R3/4	NS	NS	NS	#	NS	NS	NS	NS
Rc	NS	"	NS	NS	NS	NS	NS	NS
A1/2	NS	"	#	#	NS	NS	NS	NS
Ac	NS	"	NS	NS	NS	NS	NS	NS
<i>Female</i>								
Cr								NS
Ci								NS
Cco							NS	NS
Coc					NS		NS	NS
C1/2					NS		NS	NS
C3/4				NS	NS		NS	NS
Cc			NS	NS	NS		NS	NS
Ri		NS	NS	NS	NS		"	NS
Rcl		—	—	—	—	NS	"	NS
R1/4	NS	NS	NS	#	#	NS	NS	NS
R1/2	NS	NS	NS	#	#	NS	"	NS
R3/4	NS	NS	NS	#	#	NS	"	NS
Rc	NS	NS	#	NS	NS	NS	NS	"
A1/2	NS	NS	NS	#	#	NS	NS	"
Ac	NS	NS	#	#	#	#	NS	"

*Significant difference in mean age entering tooth stage ($p < 0.05$).
 " Significant advance in mean age for Western group (Northern group is reference).
 # Significant delay in mean age for Western group (Northern group is reference).
 NS, Non-significant differences ($p > 0.05$); —, No cleft stage for anterior teeth.

Results

A summary of significant/non-significant comparisons for all available mandibular tooth stages between ethnic groups and male/female groups is shown in Table 3 and 4, respectively. Results for M3 are shown in Table 5, M2 in Table 6, M1 in Table 7, P2 in Table 8, P1 in Table 9, canine in Table 10, I2 in Table 11 and I1 in Table 12. Figures 3–8 show mean age at entry with 95% confidence interval of mean for stages of M3, M2 and C, comparing means in Northern and Western male groups and Northern male and female groups.

Other results of individual tooth stages including average age of individuals within a tooth stage and minimum and maximum age for tooth stages are available in Elamin (2011), although some root stages were re-assessed and ages checked subsequently for this paper. The reported standard deviation values for mean age at entry in this paper require a multiplication correction factor of p/\sqrt{n} , as mean ages were calculated using logistic regression (Greene & Hensher, 2010).

Ethnic comparisons

A high percentage of tooth stage comparisons of mandibular teeth were not significantly different. In males, 61 out of 75

Table 4. Male–female comparisons. Summary of 80 comparisons of mean age entering mandibular tooth stages in North male–female and 75 West male–female groups showing predominant similarity between males and females. Missing tooth stages occurred at a younger age than our sample and could not be calculated.

Comparisons*	I1	I2	C	P1	P2	M1	M2	M3
<i>North</i>								
Cr								NS
Ci								"
Cco								NS
Coc					NS		NS	NS
C1/2					NS		NS	NS
C3/4				NS	NS		NS	#
Cc			NS	NS	"		NS	NS
Ri			NS	"	#	NS	NS	NS
Rcl			—	—	—	NS	NS	NS
R1/4	NS	NS	NS	"	NS	NS	NS	NS
R1/2	NS	NS	"	NS	NS	NS	NS	NS
R3/4	NS	NS	"	NS	NS	NS	NS	NS
Rc	NS	NS	NS	NS	NS	NS	NS	NS
A1/2	NS	"	"	"	"	NS	NS	NS
Ac	NS	"	NS	NS	NS	NS	NS	#
<i>West</i>								
Cr								NS
Ci								NS
Cco							NS	NS
Coc					NS		NS	NS
C1/2					NS		NS	"
C3/4				NS	NS		NS	NS
Cc			NS	NS	NS		NS	NS
Ri			NS	NS	NS		"	NS
Rcl			—	—	—	NS	"	NS
R1/4	NS	NS	NS	NS	NS	NS	NS	NS
R1/2	NS	NS	NS	NS	NS	NS	NS	NS
R3/4	NS	NS	"	NS	NS	NS	NS	NS
Rc	NS	NS	NS	NS	NS	NS	NS	NS
A1/2	NS	NS	NS	NS	NS	NS	NS	NS
Ac	NS	NS	"	NS	NS	NS	NS	NS

*Significant difference in mean age entering tooth stage ($p < 0.05$).
 " Significant advance in mean age for females (Males are reference group).
 # Significant delay in mean age for females (Males are reference group).
 NS, Non-significant differences ($p > 0.05$); —, No cleft stage for anterior teeth.

tooth stages (81%) were not significantly different. Of the 14 stages that were significant, mean age was earlier for eight stages in the Western males compared to the Northern males. In females, 56 out of 76 tooth stages (74%) were not significantly different. Of the 20 stages that were significant, mean age was earlier for seven stages in the Western group. The tooth stages involved were the lateral incisor apical root stages in males and premolars in both sexes. Mean ages of most root stages of both premolars in the Western females were later compared to Northern females. The reason for this is unclear. The Western female group was considerably smaller than other groups and late root stages of M3 are based on small numbers. These results are summarised in Table 3 and illustrated in Figures 3, 5 and 7 for M3, M2 and C, respectively.

Sex comparisons

Most tooth stage comparisons of eight mandibular teeth were not significantly different. In the Northern groups, 69 out of 80 tooth stages (86%) were not significantly different. Of the 11 stages that were significant, mean age was earlier for eight stages in females compared to males. In the Western group, 68 out of 75 tooth stages (91%) were not significantly different. Mean age was earlier in the females for

Table 5. Comparison of mean age in years, standard deviation (SD), 95% confidence interval (95% CI) for entering mandibular M3 stages in Northern and Western Sudanese groups.

Stage	North females (n¼ 802)		West females (n¼ 401)		North males (n¼ 848)		West males (n¼ 846)	
	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI
Cr	8.19, 1.41	7.95–8.45	8.23, 1.68	7.28–9.18	8.49, 2.05	7.75–9.23	8.35, 2.66	8.01–8.68
Ci	8.88, 1.18	8.65–9.14	8.75, 2.06	8.34–9.16	9.54, 3.76	9.08–10.01	8.45, 1.45	8.02–8.89
Cco	9.28, 1.19	9.04–9.59	9.50, 1.32	9.13–9.87	9.72, 2.06	9.21–10.23	9.45, 1.36	9.17–9.73
Coc	9.77, 1.26	9.50–10.10	10.11, 1.54	9.79–10.44	10.17, 2.12	9.72–10.62	10.29, 1.50	10.05–10.53
C1/2	10.71, 1.57	10.11–11.49	11.23, 1.48	10.95–11.51	11.14, 1.74	10.80–11.47	11.55, 1.63	11.36–11.74
C3/4	12.27, 1.81	11.79–12.71	12.26, 1.18	12.02–12.51	12.44, 1.86	12.09–12.79	13.30, 1.41	13.12–13.48
Cc	13.56, 1.96	13.11–13.97	13.04, 1.45	12.71–13.36	13.72, 1.72	13.40–14.05	14.04, 1.25	13.85–14.23
Ri	14.47, 1.87	14.06–14.85	14.23, 1.23	13.86–14.59	14.76, 1.29	14.48–15.03	14.58, 1.18	14.37–14.80
Rcl	14.95, 1.77	14.56–15.31	15.25, 1.47	14.80–15.71	15.10, 1.36	14.83–15.37	14.91, 1.12	14.71–15.12
R1/4	15.70, 2.14	15.34–16.06	15.49, 1.21	15.07–15.92	15.66, 1.38	15.39–15.93	15.49, 1.36	15.19–15.79
R1/2	16.69, 2.50	16.34–17.05	16.27, 1.12	15.82–16.72	16.62, 1.77	16.32–16.91	17.26, 1.63	16.64–17.87
R3/4	18.06, 2.73	17.68–18.45	17.28, 0.85	16.70–17.86	17.64, 2.03	17.33–17.95	18.01, 1.36	17.26–18.74
Rc	19.16, 2.39	18.78–19.54	18.11, 1.00	17.55–18.66	18.51, 1.86	18.21–18.81	18.63, 1.19	18.01–19.25
A1/2	19.57, 2.28	19.21–19.93	18.44, 0.80	17.95–18.93	18.89, 1.70	18.55–19.22	19.05, 0.98	18.44–19.66
Ac	20.61, 1.94	20.24–20.99	19.43, 1.34	18.72–20.14	19.70, 1.54	19.41–20.00	20.17, 1.47	19.33–21.00

n, Number of individuals in the group.

Table 6. Comparison of mean age in years for entering mandibular M2 stages in Northern and Western Sudanese groups.

Stage	North females (n¼ 802)		West females (n¼ 402)		North males (n¼ 848)		West males (n¼ 846)	
	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI
Ci	3.48, 0.22	3.25–3.71			3.03, 0.94	2.59–3.47		
Cco	3.78, 0.56	3.56–3.99	3.90, 0.52	3.36–4.44	3.44, 0.94	3.12–3.76		
Coc	3.98, 0.69	3.76–4.21	4.19, 0.33	3.81–4.57	3.86, 0.76	3.62–4.10	4.13, 0.54	3.67–4.60
C1/2	4.34, 0.98	4.07–4.61	4.66, 0.96	4.15–5.17	4.23, 0.96	3.97–4.49	4.79, 0.89	4.29–5.28
C3/4	5.63, 0.89	5.41–5.86	5.58, 1.14	5.39–5.76	5.40, 1.18	5.15–5.65	5.88, 1.07	5.52–6.23
Cc	7.61, 1.50	7.32–7.90	7.52, 1.14	7.14–7.90	7.33, 1.83	7.00–7.65	7.59, 1.19	7.33–7.85
Ri	9.71, 1.38	9.44–9.98	9.00, 0.81	8.76–9.25	9.76, 1.52	9.47–10.04	9.44, 0.63	9.29–9.58
Rcl	10.49, 1.30	10.24–10.74	9.84, 0.65	9.65–10.02	10.29, 1.25	10.04–10.54	10.14, 0.67	9.98–10.29
R1/4	11.24, 1.61	10.95–11.53	10.79, 0.72	10.62–10.95	11.23, 1.21	10.99–11.47	10.83, 0.62	10.72–10.93
R1/2	12.07, 1.19	11.81–12.33	11.60, 0.49	11.45–11.75	11.79, 1.12	11.53–12.05	11.82, 0.65	11.73–11.92
R3/4	12.87, 1.23	12.60–13.14	12.33, 0.63	12.17–12.49	12.54, 1.18	12.27–12.80	12.49, 0.60	12.39–12.59
Rc	13.44, 1.18	13.18–13.69	13.09, 0.56	12.89–13.28	13.50, 1.09	13.27–13.74	13.29, 0.76	13.19–13.39
A1/2	14.01, 1.12	13.74–14.29	14.30, 1.12	13.93–14.67	14.29, 1.03	14.04–14.55	14.08, 1.01	13.92–14.23
Ac	15.55, 1.67	15.24–15.85	15.62, 1.09	15.19–16.06	15.32, 1.00	15.10–15.53	15.57, 1.27	15.28–15.86

Abbreviations, see Table 5.

Table 7. Comparison of mean age in years entering mandibular M1 stages in Northern and Western Sudanese groups.

Stage	North females (n¼ 802)		West females (n¼ 402)		North males (n¼ 848)		West males (n¼ 846)	
	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI
Cc	2.97, 0.57	2.30–3.65						
Ri	4.07, 0.23	3.90–4.25			4.18, 0.42	3.95–4.41	4.10, 0.18	3.71–4.32
Rcl	4.20, 0.21	4.04–4.37	4.33, 0.50	3.71–4.96	4.51, 0.34	4.31–4.71	4.58, 0.41	4.10–5.07
R1/4	5.22, 0.31	5.03–5.41	4.86, 0.32	4.53–5.19	5.23, 0.32	5.06–5.41	5.39, 0.13	5.20–5.57
R1/2	6.00, 0.41	5.80–6.20	6.09, 0.15	5.92–6.27	6.08, 0.47	5.88–6.27	6.12, 0.23	5.94–6.30
R3/4	6.47, 0.97	5.76–6.91	6.80, 0.31	6.51–7.09	7.06, 0.47	6.84–7.28	6.81, 0.57	6.55–7.08
Rc	7.34, 0.88	6.92–7.77	7.18, 0.81	6.89–7.45	7.27, 0.89	6.27–8.26	7.03, 1.31	5.84–7.86
A1/2	7.60, 0.71	6.45–9.02	8.31, 1.23	7.36–8.88	7.89, 1.07	7.29–8.69	8.29, 1.27	8.04–8.52
Ac	8.08, 0.58	7.94–8.24	9.71, 1.09	9.45–9.96	8.99, 1.30	8.18–10.14	9.50, 1.07	8.95–9.95

Abbreviations, see Table 5.

all seven tooth stages that were significant in this group. These results are summarised in Table 4 and illustrated in Figures 4, 6 and 8 for M3, M2 and C, respectively.

Discussion

Our results show that the timing of tooth formation is not significantly different in the vast majority of the 306 tooth stage comparisons. There was no clear pattern, i.e. the stages that were significant were not specific to crown stage, root stage or tooth type, although mean ages in a number of

root stages of successional teeth (canine and premolars) were delayed in the Western group compared to the Northern group. Mean ages were earlier in the Western group compared to the Northern group in around one third of comparisons.

Comparison between Northern and Western groups

Our main finding was that most mandibular tooth stages were not significantly different in timing in the two ethnic groups in Khartoum. Around 83% of comparisons in our

532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590

Table 8. Comparison of mean age in years entering mandibular P2 stages in Northern and Western Sudanese groups.

Stage	North females (n¼ 802)		West females (n¼ 402)		North males (n¼ 848)		West males (n¼ 846)	
	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI
Cco	3.40, 0.70	2.85–3.96			3.66, 0.50	2.99–3.72		
Coc	3.98, 0.52	3.66–4.30	4.02, 0.79	2.61–4.55	3.57, 0.40	3.30–3.82	4.15, 0.39	3.69–4.60
C1/2	4.26, 0.56	3.95–4.57	4.43, 0.66	3.34–4.85	4.15, 0.45	3.90–4.40	4.20, 0.47	3.51–4.89
C3/4	5.25, 0.59	4.99–5.52	5.18, 0.61	4.52–5.44	5.21, 0.94	4.97–5.42	5.16, 0.65	4.64–5.68
Cc	6.50, 0.71	6.24–6.76	6.48, 1.10	6.12–6.80	5.95, 1.18	5.66–6.12	6.85, 0.61	6.58–7.12
Ri	8.73, 0.97	8.42–9.04	8.16, 0.67	7.80–8.53	7.91, 1.56	7.63–8.22	8.45, 0.50	8.24–8.66
R1/4	9.00, 1.20	8.77–9.25	9.54, 0.54	9.32–9.96	9.33, 1.26	9.05–9.64	9.78, 0.54	9.61–9.96
R1/2	9.83, 1.41	9.55–10.14	10.58, 0.58	10.37–10.80	10.18, 0.63	9.97–10.40	10.79, 0.53	10.64–10.94
R3/4	11.02, 1.33	10.71–11.37	11.80, 0.43	11.61–11.99	11.43, 1.64	11.07–11.77	11.82, 0.64	11.68–11.96
Rc	12.18, 1.45	11.44–12.90	12.70, 0.46	12.48–12.92	12.66, 1.33	12.33–12.96	12.87, 0.60	12.73–13.02
A1/2	12.96, 1.38	12.60–13.31	14.27, 0.78	13.95–14.61	13.77, 1.13	13.50–14.03	14.50, 0.60	13.90–14.20
Ac	14.19, 1.54	13.82–14.56	15.30, 1.32	14.88–15.83	14.29, 0.78	12.75–15.39	14.82, 1.03	14.40–15.38

Abbreviations, see Table 5.

Table 9. Comparison of mean ages in years for entering mandibular P1 stages in Northern and Western Sudanese groups.

Stage	North females (n¼ 802)		West females (n¼ 402)		North males (n¼ 848)		West males (n¼ 846)	
	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI
C1/2	2.99, 1.18	2.31–3.68						
C3/4	4.20, 0.91	3.93–4.46	3.66, 1.21	2.51–4.80	4.14–0.86	3.58–4.5		
Cc	5.47, 0.80	5.26–5.69	4.97, 1.41	4.41–5.52	5.23–0.90	5.00–5.4		
Ri	6.54, 1.16	6.30–6.78	6.82, 1.18	6.43–7.20	6.04–1.47	5.32–6.5		
R1/4	7.76, 0.92	7.58–7.95	8.32, 1.16	7.97–8.66	8.03–1.30	7.78–8.3		
R1/2	8.69, 1.07	8.19–9.30	9.70, 0.98	9.49–9.91	9.68–1.23	9.26–10		
R3/4	9.63, 1.65	9.14–10.28	10.82, 1.00	10.63–11.01	10.53–0.87	10.26–10		
Rc	10.92, 1.46	10.12–11.89	11.97, 0.72	11.77–12.16	11.61–0.41	10.69–12		
A1/2	11.88, 1.35	11.54–12.24	12.95, 0.92	12.72–13.17	12.62–1.37	12.30–12		
Ac	12.76, 1.31	12.41–13.14	13.94, 0.96	13.61–14.26	13.46–1.36	12.52–14		

Abbreviations, see Table 5.

Table 10. Comparison of mean ages in years for entering mandibular C stages in Northern Sudanese and Western Sudanese groups.

Stage	North females (n¼ 802)		West females (n¼ 402)		North males (n¼ 848)		West males (n¼ 846)	
	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI
C3/4					2.97–0.89	2.52–3.4		
Cc	3.93, 1.00	3.62–4.24	4.19, 0.74	3.10–4.67	4.05, 1.21	3.73–4.3		
Ri	5.18, 0.87	4.95–5.41	5.08, 1.34	4.57–5.58	5.21, 1.10	4.96–5.4		
R1/4	6.69, 1.43	6.42–6.96	6.90, 0.92	6.54–7.25	7.33, 1.52	7.04–7.6		
R1/2	8.04, 1.13	7.67–8.44	8.19, 1.09	7.39–8.89	8.77, 0.24	8.52–9.0		
R3/4	8.91, 1.03	8.70–9.14	8.89, 1.18	8.54–9.24	9.76, 1.47	9.25–10		
Rc	10.24, 1.25	9.96–10.54	10.83, 1.00	10.64–11.02	10.98, 0.98	10.33–11		
A1/2	11.13, 1.30	10.83–11.47	12.47, 0.63	12.30–12.66	11.96, 1.27	11.64–12		
Ac	12.11, 1.24	11.78–12.46	12.97, 0.49	12.81–13.17	12.65, 0.97	11.36–13		

Abbreviations, see Table 5.

study were not significantly different and there was no clear pattern in the tooth stages that were different. This finding is in agreement with two previous reports comparing the timing of age at entry of individual mandibular teeth between groups. The first study compared mean age entering Demirjian stages in children from Australia, Belgium, Canada, England, Finland, France, South Korea and Sweden (Liversidge et al., 2006). About 13% of comparisons were significantly different between groups with no clear pattern. The second (Liversidge et al. 2006) noted few differences between Whites and Bangladeshi groups in London (Liversidge, 2011). Comparing our results with the London groups show some stages in the Sudanese groups to be earlier (particularly incisors) and some later. Incisor stages assessment in some of the radiographs of the younger children in our study was difficult due to occasional poor image quality and this may have influenced our results.

Mean ages of M3 tooth stages were slightly later than one previous study of third molars in South African Blacks (Liversidge, 2008). Sub-Saharan African groups are known to have extensive genetic diversity (Tishkoff et al., 2009) and future studies on tooth formation in Africa are needed.

Several studies use non-cumulative methods of analyses and report ethnic differences. Two studies compare timing of tooth formation directly. Harris and McKee (1990) describe mean age using Moorrees stages aged 3–13 years, reporting earlier mean ages in middle southern US black children compared to Whites. Another study in South African children found Blacks earlier than Whites (Blankenship et al., 2007; Harris, 2007; Mincer et al., 2007; Olze et al., 2004). Other studies reported in Black individuals mean age earlier in Blacks than other groups (Olze et al., 2007) and first molars in Germany and Japan

650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708

Table 11. Comparison of mean ages in years for entering mandibular I2 stages in Northern and Western Sudanese groups.

Stage	North females (n = 802)		West females (n = 402)		North males (n = 848)		West males (n = 846)	
	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI
Ri	3.61, 0.83	3.29–3.93	3.83, 0.62	3.19–4.47	3.27, 1.18	2.84–3.70		
R1/4	4.80, 0.89	4.56–5.04	4.63, 0.72	4.17–5.09	4.78, 1.07	4.52–5.03	4.77, 0.91	4.27–5.28
R1/2	5.97, 0.92	5.75–6.19	5.88, 0.83	5.58–6.18	5.90, 1.14	5.63–6.16	5.40, 1.25	4.95–5.86
R3/4	7.16, 1.29	6.90–7.42	7.09, 0.74	6.76–7.43	7.00, 0.95	6.80–7.20	6.70, 1.10	6.42–6.98
Rc	7.37, 0.84	7.20–7.54	8.25, 1.05	6.98–7.59	7.54, 1.00	7.33–7.76	7.12, 0.96	6.90–7.32
A1/2	7.68, 0.84	7.21–8.18	7.94, 1.25	7.60–8.25	8.52, 1.42	8.21–8.84	7.81, 1.19	7.57–8.05
Ac	8.07, 0.86	7.89–8.26	8.31, 1.27	7.98–8.60	9.12, 1.37	8.82–9.48	8.31, 1.08	8.08–8.54

Abbreviations, see Table 5.

Table 12. Comparison of mean age in years entering mandibular I1 stages in Northern and Western Sudanese groups.

Stage	North females (n = 802)		West females (n = 402)		North males (n = 848)		West males (n = 846)	
	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI	Mean, SD	95% CI
R1/4	4.00, 0.43	3.92–4.24	3.83, 0.38	3.19–4.47	3.58, 0.57	3.19–3.99	3.46, 0.75	2.24–4.68
R3/4	6.17, 0.61	5.93–6.41	5.88, 0.46	5.58–6.18	6.16, 0.64	5.92–6.39	5.40, 0.69	4.95–5.86
Rc	6.75, 0.91	6.56–6.94	6.59, 0.77	6.32–6.86	6.54, 0.43	6.42–6.67	6.18, 1.15	4.38–6.74
A1/2	7.97, 0.86	6.62–7.48	7.92, 0.94	6.73–7.32	9.20, 0.88	6.52–9.40	8.85, 0.99	6.83–7.08

Abbreviations, see Table 5.

nation people in Canada (Olze et al., 2010). Forensic age estimation from third molars in the living is reviewed by Olze et al. (2004), who compare descriptive data in German, Japanese, and South African groups. They note that mean ages of some root stages in South Africans were earlier than Germans and mean ages in Japanese were later than Germans and they recommend population-specific reference data. The use of mean/median ages to estimate age has been questioned and age of transition into maturity stages (including probit regression analysis) is now seen as more appropriate to estimate age and to compare groups (Boldsen et al., 2002; Konigsberg & Frankenberg, 2002; Konigsberg et al., 2008).

Comparison between male–female within groups

Our study found few statistical differences between mean ages of mandibular tooth stages in the Northern male–female group and Western male–female group. The results from this study differ from previous findings of an advancement in root formation in females compared to males (Anderson et al., 1976; Garn et al., 1958; Haavikko, 1970; Liversidge, 2012; Liversidge et al., 2006; Moorrees et al., 1963; Nyström et al., 2007; Thompson et al., 1975). Numerous previous studies report sexual dimorphism in timing between males and females in some teeth, particularly canines and third molar root stages (Anderson et al., 1976; Demirjian & Levesque, 1980; Garn et al., 1959; Haavikko, 1970; Levesque et al., 1981; Liversidge, 2008; Liversidge et al., 2006; Moorrees et al., 1963; Nyström et al., 2007; Thompson et al., 1975).

The reason for the lack of significant difference in the timing of tooth formation in males and females in our study is unclear. The size of the Western female group was considerably smaller than other groups. Sampling is possibly a factor when numbers are small; however, the cumulative statistical approach partly overcomes this difficulty.

This lack of significant difference in timing of tooth formation supports a recent histological study reconstructing longitudinal growth in enamel and dentine that reports no clear difference in modern male and female canine formation rates (Dean et al., 2014).

Strengths and limitations

A major strength of our study, which addresses some of the limitations of previous studies, was the structured design and statistical analysis to compare mandibular tooth formation at the population level in the Northern group. The stratified structured sampling strategy enabled comparisons to be representative. The study design included a large number of children per age group (3–23 years) selected to represent the Northern group. The age distribution of children in our sample was spread across a wide age range from as young as was practical up to dental maturity of the third molar. In order to assess the entire dentition, large sample sizes from very young to fully mature are required to validate the conclusions of effects of biological factors on the timing of tooth formation. This was accomplished for the Northern group.

Most tooth stages are observed in individuals in a given population. It has to ensure sufficient children in the age range and a cumulative approach. Nyström (2016) reported differences that may have been influenced by the large variation in age that is seen between individuals or small groups.

The mean age entering a tooth stage is calculated as the age at which 50% of children have reached that stage. This cumulative statistical calculation of mean age of entry into a tooth stage uses all individuals who have reached or passed each specific tooth stage in consecutive age categories. For example, to calculate mean age entering stage X, the proportion of individuals who have




Figure 3. North-West male mean age comparisons for mandibular third molar (M3) stages across the age span of the developing tooth showing similarity between the groups. black diamond, north; unfilled diamond, west.




Figure 4. North male-female mean age comparisons for mandibular third molar (M3) stages across the age span of the developing tooth showing similarity between the groups. unfilled diamond, female; black diamond, male.

entered stage X (or later stages) is noted for consecutive age categories from the youngest to the age category where 99% of individuals have reached/passed stage X. This means that sufficient individuals are included in each analysis as the

age range from 1–99% can include up to nine consecutive age categories (for late root stages of M3).

A limitation of the study was that a convenience sample of the Western group was necessary due to the ongoing civil

886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944




Figure 5. North-West male mean age comparisons for mandibular second molar (M2) stages across the age span of the developing tooth showing similarity between the groups. black diamond, north; unfilled diamond, west.




Figure 6. North male-female mean age comparisons for mandibular second molar (M2) stages across the age span of the developing tooth showing similarity between the groups. unfilled diamond, female; black diamond, male.

war, resulting in a different sample structure of considerably smaller size. Despite a reasonable number of Western males and females for ages 5–16 (see Table 2), the younger and older age categories are not well represented. This resulted

in some mandibular tooth stages containing small numbers in the unrepresented age categories. The number of individuals in the Western group over the age of 16 is small and similarity in the timing of later root stages of M3 between

1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062




Figure 7. North-West male mean age comparisons for mandibular canine (C) stages across the age span of the developing tooth showing similarity between the groups. black diamonds, north; unfilled diamonds, west.




Figure 8. North male-female mean age comparisons for mandibular canine (C) stages across the age span of the developing tooth showing similarity between the groups. unfilled diamond, female; black diamond, male.

the sexes should be further investigated with larger numbers.

Another limitation, in common with many other studies using dental panoramic radiographs of living children, is the lack of very young children. The mean ages of most crown

stages of permanent incisors, canines and first molars occur during the first 3 years.

A limitation of any maturity study is that maturation, a continuous process, is divided into arbitrarily selected, discrete stages that are not equally spaced in time. A tooth

must be allocated to a crown or root stage and some tooth stages rely on subjective estimation of final crown and root lengths, for example Cr³⁼⁴ and R³⁼⁴ and these are more difficult to assign. Other difficulties include superimposition of the vertebral column with anterior teeth and the mandible not being correctly positioned in very young children. These influence the clarity of the radiographic image. Other factors such as individual variation in crown height and root length, as well as the duration of development of both crown and roots, may explain some inconsistencies in the pattern of differences in our results.

Conclusions

Mean ages of 23% of 151 tooth stage comparisons of mandibular teeth between 2–23 years of age were significantly different between Northern and Western males and Northern and Western females. The age variation within each group was considerably greater than the age variation between groups for all mandibular tooth stages we could assess in this study.

Mean ages of 12% of 155 tooth stage comparisons of mandibular teeth between 2–23 years of age were significantly different between males and females in both Northern and Western groups. A pattern of sexual dimorphism in tooth formation was not apparent in our results. This is in contrast to most previous radiographic studies that report a clear pattern with earlier mean ages of canine root stages and late M3 root stages in females compared to males.

This means that healthcare providers in the region can use these results of timing of mandibular teeth from the Northern group as a reference to assess dental maturity for dental treatment or to plan the timing of orthodontic treatment. More research is needed on the rate of tooth development, root proportions of mature teeth, as well as timing and root stage at tooth eruption in these and other African groups.

Acknowledgements

We are grateful to Wagner Marcenes (Dental Institute, Queen Mary University of London) and Mudather Widaa, Nihal Abdelazeem, Israa Salah, Ahmed Yahia and Motaman Elhadi (Khartoum Centre for Research and Medical Training) for help during the planning and data collection of this study. We are grateful to the two anonymous reviewers whose suggestions improved the manuscript.

Disclosure statement

The authors report no conflict of interest, financial or personal relations with other organisations or people who may influence the study findings inappropriately.

References

Anderson DL, Thompson GW, Popovich F. 1976. Age of attainment of mineralization stages of the permanent dentition. *J Forensic Sci* 21:191–200.

Antoine D, Hillson S, Dean MC. 2009. The developmental clock of dental enamel: a test for the periodicity of prism cross-striations in modern

humans and an evaluation of the most likely sources of error in histological studies of this kind. *J Anat* 214:45–55.

Bennett S, Woods T, Liyanage WM, Smith DL. 1991. A simplified general method for cluster-sample surveys of health in developing countries. *World Health Stat Q* 44:98–106.

Blankenship JA, Mincer HH, Anderson KM, Woods MA, Burton EL. 2007. Third molar development in the estimation of chronologic age in American blacks as compared with whites. *J Forensic Sci* 52:428–433.

Boldsen JL, Milner GR, Konigsberg LW, Wood JW. 2002. Transition analysis: a new method for estimating age from skeletons. In: Hoppa RD, Vaupel JW, editors. *Paleodemography: age distributions from skeletal samples*. Cambridge Studies in Biological Anthropology and Evolutionary Anthropology. Cambridge: Cambridge University Press. p 36.

Dean MC, Liversidge HM, Elamin F. 2014. Combining radiographic and histological data for dental development to compare growth in the past and the present. *Ann Hum Biol* 41:336–347.

Demirjian A, Levesque GY. 1980. Sexual differences in dental development and prediction of emergence. *J Dent Res* 59:1110–1122.

Elamin F. 2011. Tooth formation in Sudanese children. Ph.D. Dissertation, University of London. Available online at: <https://qmro.qmul.ac.uk/jspui/bitstream/123456789/2339/1/ELAMINToothFormation2011.pdf>, accessed

Elamin F, Liversidge HM. 2013. Malnutrition has no effect on the timing of human tooth formation. *PLoS One* 8:e72274.

Fanning EA, Brown T. 1971. Primary and permanent tooth development. *Aust Dent J* 16:41–43.

Garn SM, Lewis AB, Kerewsky RS. 1965. Genetic, nutritional, and maturational correlates of dental development. *J Dent Res* 44:(Suppl):228–242.

Garn SM, Lewis AB, Koski K, Polacheck DL. 1958. The sex difference in tooth calcification. *J Dent Res* 37:561–567.

Garn SM, Lewis AB, Polacheck DL. 1959. Variability of tooth formation. *J Dent Res* 38:135–148.

Greene WH, Hensher DA. 2010. *Modeling ordered choices. A primer*. Cambridge: Cambridge University Press.

Haavikko K. 1970. The formation and the alveolar and clinical eruption of the permanent teeth. An orthopantomographic study. *Proc Finn Dent Soc* 66:103–170.

Harris EF. 2007. Mineralization of the mandibular third molar: a study of American blacks and whites. *Am J Phys Anthropol* 132:98–109.

Harris EF, McKee JH. 1990. Tooth mineralization standards for blacks and whites from the middle southern United States. *J Forensic Sci* 35:859–872.

Konigsberg LW, Frankenberg SR. 2002. Deconstructing death in paleodemography. *Am J Phys Anthropol* 117:297–309.

Konigsberg LW, Herrmann NP, Wescott DJ, Kimmerle EH. 2008. Estimation and evidence in forensic anthropology: age-at-death. *J Forensic Sci* 53:541–557.

Levesque GY, Demirjian A, Tanguay R. 1981. Sexual dimorphism in the development, emergence, and agenesis of the mandibular third molar. *J Dent Res* 60:1735–1741.

Liversidge HM. 2008. Timing of human mandibular third molar formation. *Ann Hum Biol* 35:294–321.

Liversidge HM. 2011. Similarity in dental maturation in two ethnic groups of London children. *Ann Hum Biol* 38:702–715.

Liversidge HM. 2012. The assessment and interpretation of Demirjian, Goldstein and Tanner's dental maturity. *Ann Hum Biol* 39:412–431.

Liversidge HM, Chaillet N, Mørnstad H, Nyström M, Rowlings K, Taylor J, Willems G. 2006. Timing of Demirjian's tooth formation stages. *Ann Hum Biol* 33:454–470.

Liversidge HM, Speechley T. 2001. Growth of permanent mandibular teeth of British children aged 4 to 9 years. *Ann Hum Biol* 28:256–262.

Liversidge HM, Speechley T, Hector MP. 1999. Dental maturation in British children: are Demirjian's standards applicable? *Int J Paediatr Dent* 9:263–269.

Mincer HH, Harris EF, Berryman HE. 1993. The A.B.F.O. study of third molar development and its use as an estimator of chronological age. *J Forensic Sci* 38:379–390.

1299	Moorrees CF, Fanning EA, Hunt EE, Jr. 1963. Age Variation of formation stages for ten permanent teeth. <i>J Dent Res</i> 42:1490–1502.	1358
1300	Nyström M, Ranta HM, Peltola JS, Kataja JM. 2007. Timing of developmental stages in permanent mandibular teeth of Finns from birth to age 25. <i>Acta Odontol Scand</i> 65:36–43.	1359
1301		1360
1302		1361
1303	Olze A, Pynn BR, Kraul V, Schulz R, Heinecke H, Pfeiffer H, Schmeling A. 2010. Studies on the chronology of third molar mineralization in First Nations people of Canada. <i>Int J Legal Med</i> 124:433–437.	1362
1304		1363
1305	Olze A, Schmeling A, Taniguchi M, Maeda H, Van Niekerk P, Wernecke KD, Geserick G. 2004. Forensic age estimation in living subjects: the ethnic factor in wisdom tooth mineralization. <i>Int J Legal Med</i> 118:170–173.	1364
1306		1365
1307		1366
1308		1367
1309	Olze A, Van Niekerk P, Ishikawa T, Zhu BL, Schulz R, Maeda H, Schmeling A. 2007. Comparative study on the effect of ethnicity on wisdom tooth eruption. <i>Int J Legal Med</i> 121:445–448.	1368
1310		1369
1311	Papagerakis S, Zheng L, Schnell S, Sartor MA, Somers E, Marder W, et al. 2014. The circadian clock in oral health and diseases. <i>J Dent Res</i> 93:27–35.	1370
1312		1371
1313		1372
1314		1373
1315		1374
1316		1375
1317		1376
1318		1377
1319		1378
1320		1379
1321		1380
1322		1381
1323		1382
1324		1383
1325		1384
1326		1385
1327		1386
1328		1387
1329		1388
1330		1389
1331		1390
1332		1391
1333		1392
1334		1393
1335		1394
1336		1395
1337		1396
1338		1397
1339		1398
1340		1399
1341		1400
1342		1401
1343		1402
1344		1403
1345		1404
1346		1405
1347		1406
1348		1407
1349		1408
1350		1409
1351		1410
1352		1411
1353		1412
1354		1413
1355		1414
1356		1415
1357		1416