



**University of Dundee**

### **Identification of RNF168 as a PML nuclear body regulator**

Shire, Kathy; Wong, Andrew I.; Tatham, Michael H.; Anderson, Oliver F.; Ripsman, David; Gulstene, Stephanie; Moffat, Jason; Hay, Ronald T.; Frappier, Lori

*Published in:*  
Journal of Cell Science

*DOI:*  
[10.1242/jcs.176446](https://doi.org/10.1242/jcs.176446)

*Publication date:*  
2016

*Document Version*  
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

*Citation for published version (APA):*

Shire, K., Wong, A. I., Tatham, M. H., Anderson, O. F., Ripsman, D., Gulstene, S., ... Frappier, L. (2016). Identification of RNF168 as a PML nuclear body regulator. *Journal of Cell Science*, 129(3), 580-591. DOI: 10.1242/jcs.176446

#### **General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## **Identification of RNF168 as a PML Nuclear Body Regulator**

Kathy Shire<sup>1</sup>, Andrew I. Wong<sup>1</sup>, Michael H. Tatham<sup>2</sup>, Oliver F. Anderson<sup>2</sup>, David Ripsman<sup>1</sup>, Stephanie Gulstene<sup>1</sup>, Jason Moffat<sup>1</sup>, Ronald T. Hay<sup>2</sup> and Lori Frappier<sup>1\*</sup>

<sup>1</sup>Department of Molecular Genetics, University of Toronto

<sup>2</sup>Centre for Gene Regulation and Expression, College of Life Sciences, University of Dundee

\* Author for Correspondence ([lori.frappier@utoronto.ca](mailto:lori.frappier@utoronto.ca))

Running title: RNF168 as a PML Regulator

Word count: 6,503

## 1 **Abstract**

2 Promyelocytic leukemia (PML) proteins form the basis of PML nuclear bodies (NBs), which control  
3 many important processes. We screened an shRNA library targeting ubiquitin pathway proteins for  
4 effects on PML NBs and identified RNF8 and RNF168 DNA damage response proteins as negative  
5 regulators of PML NBs. Additional studies confirmed that depletion of either RNF8 or RNF168  
6 increased the level of PML NBs and proteins, while overexpression induced loss of PML NBs.  
7 RNF168 partially localized to PML NBs through its UMI/MIU1 ubiquitin interacting region and  
8 associated with NBs formed by any PML isoform. The association of RNF168 with PML NBs resulted  
9 in increased ubiquitylation and SUMO2 modification of PML proteins. In addition, RNF168 was  
10 found to associate with SUMO2/3 modified proteins in a manner dependent on its ubiquitin binding  
11 sequences, suggesting that hybrid SUMO-ubiquitin chains might be bound. *In vitro* assays confirmed  
12 that RNF168 preferentially binds hybrid SUMO2-K63 ubiquitin chains over individual SUMO2 or  
13 K63 ubiquitin chains. Our study identified previously unrecognized roles for RNF8 and RNF168 in  
14 PML regulation and a previously unknown preference of RNF168 for hybrid SUMO-ubiquitin chains.

15

16

## 17 **INTRODUCTION**

18 Promyelocytic leukemia (PML) nuclear bodies (NBs) are discreet foci in mammalian cells  
19 formed on the basis of PML proteins. PML proteins were so named because a translocation resulting  
20 in a PML-retinoic acid receptor alpha (RAR $\alpha$ ) fusion protein results in the development of acute  
21 promyelocytic leukemia (APL) (Salomoni et al. 2008). Considerable evidence now indicates that PML  
22 proteins and NBs are tumour suppressors in general and that increasing PML levels is an effective way  
23 of inhibiting the growth of tumour cells (Scaglioni et al. 2006; Wu et al. 2014; Salomoni et al. 2008).  
24 The tumour-suppressing properties of PML NBs stems from their important contributions to multiple  
25 cellular processes including apoptosis, p53 activation, DNA repair and senescence (Salomoni et al.  
26 2008; Lallemand-Breitenbach and de The 2010; Nisole et al. 2013).

27 PML NBs contain six different nuclear PML isoforms (PML I to VI) that are derived from  
28 alternative splicing events in the C-terminal portion of the protein, and there is evidence that specific  
29 PML isoforms mediate distinct functions associated with the NBs (Bernardi and Pandolfi 2007;  
30 Condemine et al. 2006; Nisole et al. 2013). PML proteins contain a SUMO interacting motif (SIM)  
31 and multiple SUMO-modified sites that mediate interactions between the PML proteins allowing for

32 the formation of the NB (Bernardi and Pandolfi 2007; Shen et al. 2006b). In addition to PML proteins,  
33 the NBs contain many other proteins that vary in their dynamics of association and mediate the various  
34 NB functions. Many proteins (eg. Daxx) associate with PML NBs through SUMO-SIM interactions  
35 with PML core sequences, while others (eg. p53) associate with the C-terminal tail sequences specific  
36 to a particular isoform (Bernardi and Pandolfi 2007; Nisole et al. 2013).

37 In addition to their cellular roles, PML NBs are part of the innate immune response that  
38 suppresses infection by several viruses (Geoffroy and Chelbi-Alix 2011; Everett and Chelbi-Alix  
39 2007; Nisole et al. 2013; Tavalai and Stamminger 2011; Sivachandran et al. 2012b). To overcome  
40 PML suppression, these viruses encode proteins that disrupt PML NBs by a variety of mechanisms  
41 that include inducing degradation of PML proteins, interfering with PML protein interactions by  
42 inhibiting SUMOylation, and restructuring of PML NBs into tracts (Tavalai and Stamminger 2009;  
43 Nisole et al. 2013; Geoffroy and Chelbi-Alix 2011; Everett and Chelbi-Alix 2007; Sivachandran et al.  
44 2008). These studies showed that PML NB functions can be regulated by manipulating PML proteins  
45 and provided a framework for studies on cellular regulation of PML NBs. In addition, considerable  
46 information on the regulation of PML NBs has come from studying how they are affected by arsenic  
47 trioxide, a treatment for APL that induces the loss of NBs formed by PML-RAR $\alpha$  fusion proteins,  
48 allowing restoration of NBs formed by the nonmutated copy of PML (Ablain et al. 2014; de The et al.  
49 2012). Arsenic induces hyper-SUMOylation of PML proteins, resulting in recruitment of the SUMO-  
50 targeted E3 ligases RNF4 and Arkadia (Erker et al. 2013; Lallemand-Breitenbach et al. 2008; Tatham  
51 et al. 2008). RNF4 in particular has been shown to be important for the arsenic-induced PML  
52 degradation (Lallemand-Breitenbach et al. 2008; Tatham et al. 2008).

53 Under normal cell growth conditions, the level of PML proteins also appears to be largely  
54 regulated by ubiquitylation and proteasomal-mediated degradation (Rabellino and Scaglioni 2013).  
55 PML proteins contain a phosphodegron that, upon phosphorylation by casein kinase 2 (CK2), triggers  
56 polyubiquitylation (Scaglioni et al. 2006; Scaglioni et al. 2008). However the E3 ligase responsible  
57 for this ubiquitylation has not been identified. A few ubiquitin E3 ligases have been reported to  
58 promote PML loss under specific circumstances. For example, the KLHL20-Cul3-ROC1 complex  
59 down-regulates PML in response to hypoxia (Yuan et al. 2011) and SIAH1 and SIAH2 can induce  
60 PML degradation upon overexpression (Fanelli et al. 2004). In addition, E6AP has been identified as a  
61 PML regulator in Burkitt's lymphoma (Wolyniec et al. 2012). The ubiquitin specific protease USP7  
62 has also been shown to negatively regulate PML NBs in nasopharyngeal carcinoma cells, by

63 promoting the degradation of PML proteins in a manner independent of its catalytic activity (Sarkari et  
64 al. 2011; Sivachandran et al. 2008). Despite these specific studies, there is still much to be learned  
65 about which ubiquitin pathway proteins regulate the levels of PML proteins and NBs.

66 To gain a more comprehensive understanding of cellular ubiquitin pathway proteins that  
67 regulate PML NBs, we screened an shRNA library targeting ~500 ubiquitin pathway proteins for  
68 effects on PML NBs. The screen identified RNF8 and RNF168, known to work together in DNA  
69 repair, as E3 ubiquitin ligases whose depletion increases the number and intensity of PML NBs.  
70 Additional studies confirmed that silencing of RNF8 or 168 increased PML NBs and protein levels,  
71 while overexpression of either protein induced loss of PML NBs. RNF168 was further shown to  
72 localize to PML NBs through the UMI/MIU1 sequence in a PML isoform-independent manner, and to  
73 bind to hybrid SUMO2-ubiquitin chains. The results suggest additional roles for RNF8 and 168 in  
74 PML regulation.

75

## 76 **RESULTS**

### 77 **Identification of RNF8 and RNF168 as PML regulators**

78 We have previously identified a mechanism by which an Epstein-Barr virus (EBV) protein,  
79 EBNA1, induces loss of PML NBs and degradation proteins in nasopharyngeal carcinoma (NPC) and  
80 gastric carcinoma, two EBV-associated cancers (Sivachandran et al. 2012a; Sivachandran et al. 2008;  
81 Sivachandran et al. 2010). However, little is known about the cellular ubiquitin pathway proteins that  
82 regulate PML proteins in these cells. To identify cellular ubiquitin pathway proteins that regulate  
83 PML NBs in NPC cells, we conducted a high-content screen in which ~2500 lentiviruses expressing  
84 shRNAs designed to silence ~500 ubiquitin pathway proteins (E1s, E2s, E3s and deubiquitylating  
85 proteins) were used to infect the NPC cell line CNE2Z in triplicate. Several days later, cells were  
86 stained for PML and imaged by confocal microscopy and the number and intensity of PML NBs was  
87 determined. Target proteins in which two or more of the five shRNAs increased the average PML NB  
88 number 1.5 fold or more or increased average PML NB intensity 2-fold or more were considered for  
89 further study. The E3 ligases RNF8 and RNF168, which are known to work together to repair double  
90 stranded DNA breaks (Al-Hakim et al. 2010), met these criteria. As shown in Figure 1, multiple  
91 shRNAs targeting RNF8 or RNF168 increased the number and/or intensity of PML NBs relative to  
92 uninfected cells and also to a negative control lentivirus expressing shRNA targeting RNF111. The

93 latter lentivirus serves as a negative control for lentivirus infection, because RNF111 expression levels  
94 are already undetectable in these cells (data not shown).

95         The effect of RNF8 and RNF168 on PML NBs was further examined, first by depleting these  
96 proteins with siRNA. siRNA-mediated depletion of either protein resulted in an obvious increase in  
97 the number and intensity of the PML NBs (Fig. 2A and B). In keeping with these results, Western  
98 blots showed that the total level of PML proteins was increased upon silencing RNF8 or RNF168 (Fig.  
99 2C). siRNA-mediated depletion of RNF8 or RNF168 also increased the levels of PML NBs and  
100 proteins in U2OS cells, showing that the effects are not cell line specific (Fig. 2D to F). Conversely,  
101 overexpression of FLAG-tagged RNF8 or RNF168 decreased the number of PML NBs (Fig. 3).

102

### 103 **RNF168 can modulate PML NBs independently from RNF8**

104         In response to DNA damage, RNF8 is recruited to sites of double-stranded DNA breaks  
105 (DSBs) where it ubiquitylates histones H2A/H2AX, which results in recruitment of RNF168.  
106 Therefore the role of RNF168 in the DNA damage response (DDR) depends on RNF8. We asked  
107 whether a similar relationship exists for the roles of RNF8 and RNF168 in regulating PML NBs. To  
108 this end, we treated CNE2Z cells with siRNA targeting RNF8 or negative control siRNA, then  
109 transfected them with a plasmid expressing FLAG-RNF168 (to overexpress RNF168) or an empty  
110 plasmid (pcDNA3). Cells were then stained for FLAG and PML and the number of PML NBs in  
111 FLAG-positive cells were compared to the empty plasmid control (Fig.4). As expected, RNF8  
112 silencing (which was confirmed by Western blotting; Fig 4B) in the presence of pcDNA3 resulted in a  
113 notable increase in PML NBs (from 11 to 16 on average; Fig. 4A and C). Also as expected, in control  
114 siRNA samples, RNF168 overexpression notably decreased the number of PML NBs relative to the  
115 pcDNA3 control (from 11 to 6 on average; Fig. 4A and C). However, RNF168 overexpression was  
116 also found to induce the loss of PML NBs in RNF8-silenced samples (from 16 to 9 on average; Fig.  
117 4A and C). The results indicate that RNF168 is able to induce PML loss in the absence of RNF8, at  
118 least under conditions of RNF168 overexpression. For completeness, we also did the opposite  
119 experiment, in which the ability of overexpressed RNF8 to induce PML NB loss was tested with and  
120 without RNF168 silencing (Fig. S1). RNF8 retained the ability to induce loss of PML NBs after  
121 RNF168 depletion, indicating the RNF8 and RNF168 can work independently of each other. We  
122 focused our further studies on RNF168.

123

## 124 **Localization of RNF168 to PML NBs**

125 To gain insight into whether RNF168 was acting directly at PML NBs, we asked whether  
126 RNF168 localized to PML NBs. To this end, FLAG-tagged RNF168 with and without deletions of  
127 previously characterized functional elements were expressed in CNE2Z cells, followed by staining  
128 for FLAG and PML (Fig. 5A and B). As previously reported, overexpression of WT RNF168 results  
129 in the formation of aggregates interfering with assessment of nuclear localization (Pinato et al.  
130 2009). This aggregation is not seen with RNF168 mutants lacking the RING domain ( $\Delta$ RING) or  
131 MIU2 ubiquitin-binding sequence ( $\Delta$ MIU2). Both of these proteins showed nuclear diffuse staining  
132 with some foci that localize to the PML NBs (Fig. 5B). The localization to PML NBs was not an  
133 artifact of overexpression, as  $\Delta$ RING localized to PML NBs even at very early post-transfection  
134 times, when  $\Delta$ RING expression levels were close to those of endogenous RNF168 (Fig. S2A). The  
135 localization to PML NBs was not particular to CNE2Z cells, as  $\Delta$ MIU2 was also found to form foci  
136 in U2OS cells, some of which coincide with PML NBs (Fig. S2B). The localization to PML NBs  
137 was abrogated when the UMI and MIU1 ubiquitin-binding sequences were deleted individually or  
138 together within the context of  $\Delta$ MIU2 (compare  $\Delta$ MIU2 to  $\Delta$ 100-201 $\Delta$ MIU2,  $\Delta$ MIU1 $\Delta$ MIU2 or  
139  $\Delta$ UMI $\Delta$ MIU2 in Fig 5B), implicating the UMI/MIU1 region as a PML-targeting sequence. While  
140 the expression level of the different mutants examined for PML localization varied considerably  
141 (Fig. 5C), there was no correlation between PML NB localization and expression level. The ability  
142 of the UMI/MIU1 region to interact with PML NBs was further examined by expressing an RNF168  
143 fragment containing amino acids 100-201 fused to GFP (Fig. 5C and D). GFP-100-201 formed  
144 prominent foci that localized to PML NBs, whereas no such foci were seen with the expression of  
145 GFP alone (Fig. 5D). In addition, removal of MIU1 from this construct (GFP-100-166) resulted in  
146 the loss of prominent PML-localizing foci, confirming that MIU1 plays a role in recruitment to PML  
147 NBs.

148 Since RNF168 is known to have a role in the DNA damage response (DDR) involving its  
149 recruitment to DNA repair sites, we examined whether RNF168 can also associate with PML NBs  
150 during the DNA damage response. To this end, FLAG- $\Delta$ RING was expressed in CNE2Z cells,  
151 which were then treated with etoposide to induce the DDR (as evidenced by 53BP1 foci formation;  
152 Fig. 5E top panels), followed by staining for FLAG and PML. As shown in Fig. 5E,  $\Delta$ RING was still  
153 detected at PML NBs, indicating that RNF168 retains its ability to associate with PML NBs during  
154 the DDR. This result is similar to studies on the localization of the Bloom (BLM) DDR protein,

155 which localizes with PML NBs both before and after induction of DNA damage (Bischof et al.  
156 2001).

157 PML NBs contain 6 different PML isoforms that vary in the length and sequence of their C-  
158 terminal regions. Some proteins associate with PML NBs through isoform-specific sequences while  
159 others associate with the conserved PML core sequence. To determine if a specific PML isoform  
160 was responsible for RNF168 recruitment to PML NBs, we examined the localization of RNF168  
161  $\Delta$ MIU2 in CNE2Z cells that express single PML isoforms. These cells were previously generated by  
162 silencing total PML with lentivirus-delivered shRNA, then adding back a silencing-resistant PML  
163 isoform with a second lentivirus (Sarkari et al. 2011; Cuchet et al. 2011). Each PML isoform forms  
164 NBs and this system has been previously used to reveal isoform specificities of some viral and  
165 cellular proteins (Sarkari et al. 2011; Sivachandran et al. 2012b; Cuchet-Lourenco et al. 2012).  
166 Examination of the localization of RNF168  $\Delta$ MIU2 in cells with single PML isoforms, revealed that  
167 this protein can localize to NBs formed by any PML isoform, indicating that the interaction is not  
168 through the PML C-terminal tails (Fig 6).

169

### 170 **RNF168 increases PML ubiquitylation and SUMOylation**

171 The loss of PML NBs can result from either polyubiquitylation and degradation of PML  
172 proteins, or from interference with SUMOylation of PML proteins, which interferes with the ability  
173 of the PML proteins to interact to form NBs. Therefore we examined the effect of RNF168  
174 overexpression on the ubiquitylation and SUMOylation of PML proteins (Fig. 7). To this end,  
175 CNE2Z cells were co-transfected with a plasmid expressing HA-tagged ubiquitin and a second  
176 plasmid expressing FLAG-tagged RNF168, an RNF168 mutant or empty plasmid, then treated with  
177 proteasomal inhibitor (MG132) to inhibit degradation of the ubiquitylated proteins. PML proteins  
178 were immunoprecipitated (IP) with antibody recognizing all PML isoforms under denaturing  
179 conditions, then immuno-blotted for PML and HA (Fig. 7A). Both full-length and  $\Delta$ MIU2 versions  
180 of RNF168 were found to induce the accumulation of polyubiquitylated forms of PML, whereas an  
181 RNF168 mutant lacking all 3 ubiquitin interacting motifs ( $\Delta$ UMI $\Delta$ MIU1 $\Delta$ MIU2 called  $\Delta$ Ub)  
182 resembled the empty plasmid negative control. Therefore the sequences that are necessary for  
183 RNF168 to associate with PML NBs are also needed for RNF168 to induce PML ubiquitylation. An  
184 RNF168 mutant lacking the catalytic RING domain ( $\Delta$ RING) was also tested for the induction of



185 PML ubiquitylation and found to be impaired relative to RNF168 or  $\Delta$ MIU2. However the very low  
186 expression level of this mutant relative to other RNF168 proteins makes the result inconclusive.

187 To further examine the nature of the ubiquitin modifications of PML that are induced by  
188 RNF168, we repeated the above experiment in the presence and absence of MG132 and, in addition  
189 to detecting total ubiquitin (HA blot), we also blotted with antibodies specific for K48- and K63-  
190 linked ubiquitin chains (Fig. 7B). As expected, MG132 treatment resulted in increased levels of total  
191 ubiquitylated products (HA blot; input lanes) and K48-linked ubiquitin products (K48 blot; input  
192 lanes). In addition, a smaller increase in K63-linked ubiquitin products (K63 blot; input lanes) was  
193 detected in the presence of MG132, consistent with reports that hybrid chains containing SUMO2  
194 and K63 ubiquitin can trigger proteasomal degradation (Tatham et al. 2011). Consistent with the  
195 known role of RNF168 in synthesizing K63 ubiquitin chains, total levels of these chains (in input  
196 lanes) were increased when RNF168 was overexpressed (relative to the pcDNA3 control lane). PML  
197 IP experiments showed induction of PML ubiquitylation (detected in HA blot) by RNF168 and  
198  $\Delta$ MIU2 (relative to the control lane) only when proteasomal degradation was blocked, consistent  
199 with a role of this ubiquitylation in PML degradation. Immunoprecipitated PML showed some  
200 reactivity with both K48 and K63 ubiquitin antibodies under all conditions, but RNF168 and  $\Delta$ MIU2  
201 were only found to increase levels of K48 ubiquitin (and to a lesser degree, K63 ubiquitin) on PML  
202 (compared to control lanes) when the proteasome was blocked. Together the results suggest that  
203 RNF168 can increase the levels of both K48 and K63 ubiquitin chains on PML.

204 PML proteins are modified by the addition of SUMO1, SUMO2 and SUMO3. These  
205 modifications are needed for nuclear body formation, while hyper-SUMOylation can be a step in  
206 PML degradation (as induced by arsenic trioxide). We examined the effect of RNF168  
207 overexpression on PML SUMOylation by co-transfecting CNE2Z cells with a plasmid expressing  
208 myc-tagged SUMO2 and the indicated FLAG-tagged RNF168 protein or empty plasmid, followed  
209 by IP of total PML (Fig. 7C). Relative to the empty plasmid negative control, RNF168 was  
210 consistently found to increase the levels of SUMO2-modified PML proteins, whereas  $\Delta$ Ub  
211 ( $\Delta$ UMI $\Delta$ MIU1 $\Delta$ MIU2) did not exhibit this effect and  $\Delta$ MIU2 and  $\Delta$ RING had an intermediate  
212 effect. The results suggest that RNF168-induced loss of PML NBs is not due to interference with  
213 PML SUMOylation, but rather that RNF168 can increase PML SUMOylation. In addition, the fact  
214 that the low-expressing  $\Delta$ RING still induced PML SUMOylation to some degree, suggests that the  
215 increase in PML SUMOylation is independent of the ubiquitylation activity of RNF168.

216

## 217 **RNF168 associates with hybrid SUMO2-ubiquitin chains**

218         The findings that RNF168 associated with PML core sequences, which are known to be  
219 SUMOylated, and could increase SUMOylation of PML proteins prompted us to further examine the  
220 ability of RNF168 to bind SUMO. To this end, CNE2Z cells were co-transfected with a plasmid  
221 expressing myc-tagged SUMO1, SUMO2 or SUMO3 and a plasmid expressing FLAG-tagged  
222 RNF168 or  $\Delta$ Ub ( $\Delta$ UMI $\Delta$ MIU1 $\Delta$ MIU2) or empty plasmid. FLAG-RNF168 was recovered by IP and  
223 recovery of Myc-SUMO examined by Western blots (Fig. 8A). Consistent with previous reports,  
224 input lanes showed that SUMO2 and SUMO3 formed high molecular weight chains more efficiently  
225 than SUMO1. In addition, an increase in the total level of SUMO2 or SUMO3 chains was  
226 consistently seen in the presence of RNF168 relative to the empty plasmid control (compare lane 2  
227 to lane 5 and lane 3 to lane 6). FLAG IPs showed that full length RNF168 associates with the  
228 SUMO2 and SUMO3 chains (compare lanes 11 and 12 to lanes 14 and 15), and that this interaction  
229 is greatly decreased by deletion of the three ubiquitin interacting regions in the  $\Delta$ Ub mutant (lanes 17  
230 and 18). We did not detect any shifted forms of FLAG-RNF168 in either the input or FLAG-IP  
231 samples (see FLAG blots), suggesting that RNF168 itself is not SUMO-modified, but rather can bind  
232 to SUMO2/3 chains. However, a direct interaction between purified RNF168 and free SUMO2 or  
233 SUMO3 chains was not detected in *in vitro* binding assays (data not shown). This suggests that  
234 RNF168 is not simply recognizing SUMO2/3 but rather SUMO2/3 in a particular context.

235         Hybrid SUMO-ubiquitin chains have been described consisting of a SUMO2 chain joined  
236 to a K63-linked ubiquitin chain (Tatham et al. 2013), and these can be specifically recognized by  
237 some proteins (Geoffroy and Hay 2009; Guzzo et al. 2012; Hu et al. 2012). The UMI/MUI1  
238 sequences of RNF168 that we found to be important for recovery of SUMO2 chains have been  
239 previously shown to recognize K63-linked ubiquitin (Panier et al. 2012), raising the possibility  
240 that the SUMO2-containing material that we recovered are hybrid SUMO-ubiquitin chains. To  
241 investigate this possibility, affinity resins were generated that contain one of the following: K63-  
242 ubiquitin polymers (K63-Ub), linear 4xSUMO2 chains (4xSUMO-2), N-terminally  
243 monoubiquitinated 4xSUMO2 chains (Ub-4xSUMO2) or hybrid chains containing K63-ubiquitin  
244 polymers linked to 4xSUMO2 (K63Ub-4xSUMO2) (see Fig. S3A). Nuclear extracts from 293  
245 cells were incubated with each of these resins as well as with empty resin (blank), and proteins  
246 retained on each resin were identified and quantified by mass spectrometry-based label-free

247 quantitative proteomics (Fig. S3B). Ratios of the recovery of each protein on a particular  
248 SUMO/ubiquitin resin relative to the blank resin were then determined (Fig. S3C and D).  
249 RNF168 was one of the highest scoring proteins with much higher recovery on resins containing  
250 K63Ub-SUMO2 hybrid chains than any of the other resins (Figs. 8B, S3C and S3D).

251 To determine whether RNF168 binds directly to the hybrid chains, assays with the above  
252 affinity resins were repeated using purified RNF168 generated in *E.coli*, and eluted proteins were  
253 analysed by SDS-PAGE and Coomassie staining. As shown in Fig. 8C, RNF168 was recovered  
254 most efficiently on the resin containing the hybrid K63Ub-4xSUMO2 chains. Consistent with  
255 previous reports (Panier et al. 2012), some binding was also detected to resin containing K63-  
256 linked polyubiquitin. The results indicate that the association of RNF168 with SUMO2 chains  
257 involves direct binding to hybrid SUMO2-ubiquitin chains.

258

## 259 **DISCUSSION**

260 Our screen for proteins that regulate the level of PML NBs identified RNF8 and RNF168 as  
261 proteins whose depletion results in increased number and intensity of PML NBs and increased level  
262 of PML proteins. Conversely overexpression of RNF8 or RNF168 can lead to loss of PML NBs. To  
263 date, RNF8 and RNF168 are known for their important roles in the DDR to double stranded DNA  
264 breaks (Bartocci and Denchi 2013; Al-Hakim et al. 2010). K63-linked ubiquitylation of histones  
265 H2A/H2AX and/or other proteins by RNF8 at DSBs results in the recruitment of RNF168, which  
266 interacts with the ubiquitylated histones through its three ubiquitin-binding motifs (UMI, MIU1, and  
267 MIU2) (Stewart et al. 2009; Bartocci and Denchi 2013; Panier and Durocher 2009; Panier et al.  
268 2012; Mailand et al. 2007; Doil et al. 2009). Further ubiquitylation of H2A/H2AX by RNF168 and  
269 RNF8 then leads to recruitment of additional DNA repair proteins including 53BP1 and BRCA1.  
270 Our current studies indicate additional roles for RNF8 and RNF168 as negative regulators of PML  
271 NBs.

272 Our studies show that RNF8 and RNF168 can down-regulate PML NBs and proteins. We  
273 focused subsequent studies on RNF168 due to its partial co-localization with PML NBs, which  
274 suggests it is acting directly on the PML NBs. Overexpression of full length RNF168 is known to  
275 aggregate in cells and hence is not typically used for functional studies (Pinato et al. 2009). However  
276 deletion of MIU2 or the RING domain abrogates this aggregation and reveals prominent foci that  
277 correspond to PML NBs. Analyses of additional deletion mutants and GFP-fusion proteins

278 identified the UMI/MIU1 region as important for PML targeting. Since all of our UMI/ MIU1  
279 deletion constructs lacked MIU2, we do not know whether PML targeting is unique to UMI/MIU1 or  
280 can also be mediated by MIU2.

281 The mechanism by which RNF168 induces PML NB loss appears to involve ubiquitylation  
282 of PML proteins, since overexpression of RNF168 or  $\Delta$ MIU2 resulted in increased levels of  
283 ubiquitylated PML proteins. The effect on PML ubiquitylation also fits with the increase in PML  
284 protein levels that were observed upon RNF168 depletion. Induction of PML ubiquitylation requires  
285 UMI/MIU1 PML-targeting sequences in RNF168 (that were deleted in the  $\Delta$ Ub mutant in Fig. 7),  
286 consistent with a requirement to associate with the PML NB to induce PML degradation. It remains  
287 unclear whether PML is ubiquitylated directly by RNF168, although this is the simplest  
288 interpretation. Interestingly, Tikoo et al (Tikoo et al. 2013) previously found that RNF8 could  
289 conjugate K63-linked ubiquitin to PML proteins *in vitro*, although this led to an increase rather than  
290 a loss of PML NBs in their system. We also observed that RNF168 but not the  $\Delta$ Ub mutant  
291 increased the level of SUMO2-modified PML. SUMO modifications of PML are also factors in the  
292 stability of PML proteins. For example, arsenic trioxide triggers the loss of PML proteins by first  
293 inducing their hyperSUMOylation (Lallemand-Breitenbach et al. 2008; Lallemand-Breitenbach et al.  
294 2001; Tatham et al. 2008). Therefore the effect of RNF168 on PML SUMOylation may also be a  
295 factor in its ability to regulate the level of PML NBs.

296 RNF168 $\Delta$ MIU2 was found to localize to NBs formed by any single PML isoform. The PML  
297 isoforms are identical except for their C-terminal tails, which differ in length and sequence, and have  
298 been shown to mediate some isoform-specific interactions (Cuchet-Lourenco et al. 2012;  
299 Sivachandran et al. 2012b; Nisole et al. 2013). The fact that RNF168 localization to PML NBs was  
300 not PML isoform-dependent indicates that recruitment of RNF168 to PML NBs involves the PML  
301 core sequences. These core sequences contain multiple SUMO interacting motifs (SIM) and SUMO  
302 modifications, and proteins that interact with PML core sequences typically do so through SUMO-  
303 SIM interactions. This prompted us to examine whether RNF168 is SUMOylated or can bind  
304 SUMO-modified proteins (Fig. 8). While RNF168 can be modified by SUMO1 in response to DNA  
305 damage (Danielsen et al. 2012), we did not detect any SUMO modifications of RNF168 under our  
306 conditions. Rather, RNF168 was found to bind high molecular weight complexes containing  
307 SUMO2 and SUMO3 chains. These interactions were dependent on the presence of the UMI /MIU1/  
308 MIU2 ubiquitin interacting sequences suggesting, either that these sequences can also interact with

309 SUMO2/3 chains, or that the SUMO2/3 chains that were bound were hybrid chains that also  
310 contained ubiquitin.

311 To clarify whether or not RNF168 can bind hybrid SUMO-ubiquitin chains, we conducted *in*  
312 *vitro* assays comparing recovery of RNF168 on resin containing chains of SUMO2, K63 ubiquitin or  
313 K63Ub-SUMO2 hybrids. We found that the hybrid chains were most efficiently bound by RNF168,  
314 whereas binding of purified RNF168 to SUMO2 chains was not detected (Fig. 8B, 8C and additional  
315 unpublished data). This suggests that the SUMO2/3-modified proteins that we recovered from cells  
316 contain hybrid SUMO-ubiquitin chains. This would explain the requirement for the ubiquitin-  
317 binding sequences of RNF168 and suggests that RNF168 also contains a weak SIM.

318 Taken together, our data suggest a model in which RNF168 is recruited to PML NBs through  
319 direct interaction with hybrid SUMO-ubiquitin chains which could be on PML proteins themselves  
320 or other PML-NB constituents, such as Daxx, Sp100 and BLM, which are also SUMO-modified  
321 (Seeler and Dejean 2001). In this respect, RNF168 may act downstream of RNF4, a SUMO-targeted  
322 ubiquitin ligase that generates hybrid chains and is known to negatively regulate PML NBs  
323 (Geoffroy and Hay 2009). Once recruited to PML NBs, RNF168 causes increased ubiquitylation and  
324 SUMOylation of PML proteins, which stimulates their proteasomal degradation. It remains to be  
325 determined whether RNF 168 is recruited to PML NBs through interactions with hybrid SUMO-  
326 ubiquitin chains on PML. Conducting experiments in cells expressing PML mutants lacking SUMO-  
327 modified sites may be one way of clarifying whether SUMO or SUMO-ubiquitin chains on PML are  
328 required for RNF168 recruitment and induced ubiquitylation of PML. However, since the lack of  
329 SUMOylation of PML affects the structure and composition of the NBs (Lallemand-Breitenbach et  
330 al. 2001), any changes in RNF168 behaviour could not be conclusively attributed to a direct  
331 requirement for SUMO chains on PML.

332 Our finding that RNF168 binds hybrid SUMO-ubiquitin chains is also likely to be relevant  
333 for RNF168 recruitment to sites of DNA damage. It has been previous reported that SUMO1,  
334 SUMO2 and SUMO3 accumulate at double stranded DNA breaks due to the action of PIAS4 and  
335 PIAS1, and that PIAS1/PIAS4 are needed for productive recruitment of RNF168 (Galanty et al.  
336 2009). We now suggest that this requirement may be due to the generation of hybrid SUMO-  
337 ubiquitin chains, which are then recognized by RNF168. This would be similar to the recruitment of  
338 RAP80 to double-stranded DNA breaks, which has been shown to involve binding to SUMO-  
339 ubiquitin hybrid chains generated by RNF4 (Hu et al. 2012; Guzzo et al. 2012).

340 Our studies have uncovered previously unrecognized roles for the RNF8 and RNF168 DNA  
341 damage response proteins in PML regulation. A relationship between DNA repair and PML NBs is  
342 well established, in that PML NBs are known to contribute to efficient DNA repair and have been  
343 shown to be associated with several DDR proteins (Bischof et al. 2001; Tikoo et al. 2013; Boichuk  
344 et al. 2011; Boe et al. 2006; Yeung et al. 2012; Zhong et al. 1999; Dellaire et al. 2006; Dellaire and  
345 Bazett-Jones 2004). For example BLM localizes to PML NBs in cells with and without induced  
346 DNA damage, and PML is required for the formation of BLM-containing repair foci as well as for  
347 BLM function in DNA repair (Zhong et al. 1999; Bischof et al. 2001). However, others report a lack  
348 of association of PML with sites of active DNA repair (Dellaire et al. 2006), suggesting that the role  
349 of PML in DNA repair may be indirect, for example in facilitating modifications of DNA repair  
350 proteins (Lallemant-Breitenbach and de The 2010). Like BLM, we have shown that RNF168  
351 associates with PML NBs and that this interaction also occurs during the DDR. Interestingly,  
352 RNF168 is modified by SUMO1 in response to DNA damage (Danielsen et al. 2012), and since  
353 PML NBs can promote SUMOylation, this might be one reason for the association of RNF168 with  
354 PML NBs. Whether or not the interaction of RNF168 with PML NBs is necessary for its function in  
355 the DDR or represents a distinct role of RNF168 in PML regulation remains to be determined.

356

## 357 **MATERIALS AND METHODS**

### 358 **Cell lines**

359 The EBV-negative NPC cell line CNE2Z (Sun et al. 1992) was maintained in alpha-MEM (Gibco)  
360 supplemented with 10% fetal bovine serum (FBS). CNE2Z cell lines expressing single PML  
361 isoforms are described in Sarkari et al (Sarkari et al. 2011). U2OS cells were grown in Dulbecco's  
362 modified essential medium (DMEM) supplemented with 10% FBS.

363

### 364 **shRNA screen**

365 CNE2Z cells were seeded in 384-well glass bottom plates (Perkin Elmer 6007550) in 50  $\mu$ l media  
366 containing 8  $\mu$ g/mL polybrene at a density of 500 cells/well. 1 hr later, 5  $\mu$ l of shRNA-expressing  
367 lentivirus was added, corresponding to culture supernatant from a 96-well plate lentivirus infection.  
368 The lentivirus library targeted approximately 500 ubiquitin pathway proteins with 3 to 5 shRNA  
369 constructs per gene and is a subset of the library described in Moffat et al (Moffat et al. 2006).  
370 Infections with each lentivirus were set up in triplicate. 24 hrs later, 2  $\mu$ g/ml puromycin was added to

371 select for the virus. The puromycin was removed 24 hours later and the cells were grown for 5 days  
372 before fixation and fluorescent staining for image acquisition. Cells were washed with PBS, fixed  
373 with 3.7% paraformaldehyde for 25 min, washed with PBS, permeabilized with PBS containing 1%  
374 Triton X-100 for 5 min, blocked with 4% BSA in PBS for 2 hours and incubated with primary  
375 antibody solution (1:100 in 4% BSA in PBS) recognizing PML (Santa Cruz Biotechnology; sc-966)  
376 for 18 hours at 4°C. The cells were washed with PBS and incubated with blocking buffer containing  
377 1:700 dilution of goat anti-mouse conjugated Alexa Fluor 488 (Life Technologies; A11029) and  
378 1:1000 dilution of 10 mg/mL Hoechst (Molecular Probes). After 1½ hours at room temperature in  
379 the dark, the cells were rinsed 3 times with PBS and stored in PBS. Images from ten areas per well  
380 in the green (488) and blue (Hoechst) channels were obtained using a 40X objective on an Opera  
381 confocal microscope (PerkinElmer). The number and intensity of PML NBs was determined using  
382 Acapella software.

383

#### 384 **Plasmids and siRNA**

385 Plasmids expressing FLAG-tagged RNF8 (Genecopoeia), RNF168 (pcDNA3:FLAG-RNF168wt)  
386 or the RNF168 mutants  $\Delta$ MIU1 (lacking amino acids 168-191),  $\Delta$ MIU1  $\Delta$ MIU2 (lacking amino  
387 acids 168-191 and 439-462),  $\Delta$ RING (lacking amino acids 1-58),  $\Delta$ MIU2 (lacking amino acids 439-  
388 462) were kindly supplied by Dan Durocher and are described in Stewart et al 2009.  
389 pcDNA3::FLAG  $\Delta$ UMI  $\Delta$ MIU1  $\Delta$ MIU2 (lacking amino acids 134-191 and 439-462) was generated  
390 by PCR amplification of the RNF168 sequences in pGFP-RNF168  $\Delta$ UMI  $\Delta$ MIU1  $\Delta$ MIU2 (Panier et  
391 al. 2012) and insertion into AscI and XbaI of pcDNA3:FLAG.  $\Delta$ 100-201 $\Delta$ MIU2 was generated  
392 from pcDNA3::FLAG  $\Delta$ MIU2 by Quik Change site directed mutagenesis. pEGFP:NLS-  
393 RNF168aa100-201 was obtained from Dan Durocher and is described in Panier et al (2012).  
394 pEGFP:NLS-RNF168 aa100-166 was generated by PCR amplification of the relevant RNF168  
395 sequences and ligation into AscI and XbaI sites of pEGFP:NLS. Stealth siRNA of RNF168  
396 (CCAUCCAGCCUCAUCUGGACCAGUU) and RNF8  
397 (GGGUUUGGAGAUAGCCCAAGGAGAA) were from Invitrogen. AllStar negative-control  
398 siRNA was obtained from Qiagen.

399

#### 400 **Antibodies**

401 PML antibodies used for microscopy were from Santa Cruz Biotechnology (sc-966: 1:80 dilution)  
402 and those used for Western blotting were from Bethyl (A301-167A; 1:2000 dilution). Other  
403 antibodies used were mouse anti-FLAG M2 (Sigma; 1:5000 dilution), rabbit anti-FLAG (Bethyl  
404 A190-102A; 1:5000 dilution), RNF 8 (Santa Cruz 271462; 1:200 dilution), RNF168 (Millipore 06-  
405 1130; 1:1000 dilution), HA (Santa Cruz 7392; 1:500 dilution), K63-linked ubiquitin (Millipore 05-  
406 1308; 1:1000 dilution), K48-linked ubiquitin (Millipore 05-1307; 1:10,000 dilution), myc (Santa  
407 Cruz 40; 1:500 dilution) and actin (Santa Cruz 1616; 1:2000 dilution). Secondary antibodies used for  
408 Western Blots (GAR/HRP -2055, GAR/HRP-2004; 1:5000 dilutions) were from Santa Cruz  
409 Biotechnology. Secondary antibodies for microscopy were from Invitrogen (GAR488-A11029,  
410 GAR555, GAR488, GAR555; all at 1:700 dilution).

411

### 412 **Transfections**

413 Approximately  $6 \times 10^5$  CNE2Z cells were plated in 5 mls of medium in 10 cm dishes or on  
414 coverslips (for microscopy). They were immediately transfected with 100 pmol of siRNA targeted  
415 against either RNF8 or RNF168 or AllStars negative control siRNA (Qiagen) using 2  $\mu$ L of  
416 lipofectamine 2000 (Invitrogen). The cells were subject to second and third rounds of the same  
417 transfection after 24 and 48 hours. 48 hours after the third round of transfections cells were  
418 harvested and either analyzed by Western blotting or processed for immunofluorescence  
419 microscopy. For overexpression experiments, cells plated as above were transfected with 2  $\mu$ g of the  
420 indicated plasmid using PolyJet (FroggaBio) then harvested 24 hrs (for PML NB localization  
421 experiments) to 48 hrs (for PML loss experiments) post transfection.

422

### 423 **Immunofluorescence microscopy**

424 Cells grown on coverslips were fixed with 3.7% formaldehyde in PBS for 20 min, rinsed twice in  
425 PBS, and permeabilized with 1% Triton X-100 in PBS for 5 min. Samples were blocked with 4%  
426 bovine serum albumin (BSA) in PBS followed by incubation with primary antibodies as indicated,  
427 followed by secondary antibodies. Coverslips were mounted on slides using ProLong Gold antifade  
428 medium containing 4',6-diamidino-2-phenylindole (DAPI) (Invitrogen). Images were obtained using  
429 the 40 $\times$  oil objective on a Leica inverted fluorescence microscope and processed using the  
430 OpenLAB (ver.X.0) software program. PML was quantified by counting 100 cells per sample, and



431 experiments were performed in triplicate. Averages and standard deviations were calculated in  
432 Excel. P values were determined using two tailed t-tests in Excel.

433

#### 434 **Western blotting**

435 Cells were lysed in 9 M urea, 10 mM Tris pH 6.8 followed by sonication. 50 µg of clarified lysates  
436 were loaded onto 10% SDS-PAGE and transferred onto nitrocellulose. Membranes were blocked in  
437 5% non-fat dry milk in PBS-T (PBS with 0.1% Tween) for 1 hour, followed by incubation with  
438 primary antibody in blocking buffer overnight at room temperature. Membranes were washed three  
439 times with PBS-T and then incubated with secondary antibodies conjugated to horseradish  
440 peroxidase (Santa Cruz Biotechnology) for 1 hour. Membranes were washed three times with PBS-  
441 T and signals were detected by enhanced chemiluminescence (ECL) (Perkin Elmer Life and  
442 Analytical Sciences).

443

#### 444 **PML ubiquitylation and SUMOylation assays**

445 For ubiquitylation assays, CNE2Z cells in a 15 cm dish were transfected with 5 µg of plasmid  
446 expressing HA-tagged ubiquitin and 5 µg of FLAG-RNF168 expression plasmid or empty control  
447 plasmid (pcDNA3) using Polyjet (FroggaBio Scientific Solutions). 40 hours post-transfection, cells  
448 were treated with 10 µM MG132 (Sigma) for 10 hours (except where indicated). Harvested cell  
449 pellets were frozen then thawed and boiled in 200 µl of SDS lysis buffer (62.5 mM Tris pH 6.8, 2%  
450 SDS, 10% glycerol, 1 mM *N*-ethyl maleimide). Clarified lysates were diluted with 1 ml IP buffer (50  
451 mM Tris pH 8.0, 150 mM NaCl, 1% NP-40). Lysates were incubated overnight with PML antibody  
452 (Bethyl, A301-167A) and Protein A/G agarose (SantaCruz sc-2003). After washing in IP buffer,  
453 immunoprecipitates were eluted in loading buffer (60 mM Tris.HCl pH 6.8, 1% SDS, 100 mM DTT,  
454 5% glycerol) prior to western blotting. SUMOylation assays were performed as above except that  
455 the plasmid expressing HA-ubiquitin was replaced by myc-tagged SUMO2 (MacPherson et al. 2009)  
456 and no MG132 treatment was used. A positive control for SUMOylation was also performed by  
457 treating pcDNA3/myc-SUMO2 transfected cells with 4 µM arsenic trioxide for 4 hrs.

458

#### 459 **Cell-based assay for interaction of RNF168 with SUMO**

460 293T cells were transfected with 10 µg of plasmid expressing myc-tagged SUMO1, SUMO2 or  
461 SUMO3 (MacPherson et al. 2009) (kindly supplied by Dr. Paul Sadowski) and 10 µg of FLAG-

462 RNF168 expression plasmid or empty control plasmid (pcDNA3). 48 hrs later, cells were lysed  
463 in 50 mM Tris pH 7.4, 150 mM NaCl, 1% TritonX-100, 1 mM N-ethyl maleimide, P8340  
464 Protease Inhibitor (Sigma). 1.7 mg of lysate was incubated with 20 µl of M2 FLAG affinity  
465 resin (Sigma A2220) for 3 hours and then washed 4 times with 1 ml of lysis buffer. Proteins  
466 were eluted in 2X SDS loading buffer (4% (w/v) SDS, 0.2% (w/v) bromophenol blue, 20% (v/v)  
467 glycerol, 200 mM DTT) then analysed by Western blotting.

468

#### 469 **Generation of SUMO2/ubiquitin affinity resins**

470 4xSUMO2 (SUMO2 residues 2-92 linked to three copies of SUMO2 residues 12-92) and Ub-  
471 4xSUMO2 were expressed and purified as described previously (Tatham et al. 2013).  
472 Unanchored K63-linked polyubiquitin chains (K63Ub), and K63-linked polyubiquitin chains  
473 conjugated to the N-terminus of 4xSUMO2 (K63Ub-4xSUMO2) were synthesized using  
474 recombinant Ubc13 and Ube2v2 as described previously (Branigan et al. 2015). These proteins  
475 were coupled to N-hydroxysuccinimide (NHS) resin in 0.2M NaHCO<sub>3</sub>, 0.5M NaCl as per  
476 manufacturer's instructions (GE Healthcare). The remaining active groups were blocked with  
477 ethanolamine, and resins were stored at 4°C.

478

#### 479 **Identification of cellular proteins with affinity for SUMO2 and ubiquitin polymers.**

480 Nuclear extracts were generated from 293 N3S cells as previously described for U2OS cells  
481 (Seifert et al. 2015). Nuclear lysate (10 mls at 5.5 mg/ml) was incubated with 50 µl of the affinity  
482 resin overnight at 4°C with rotation, then resins were washed three times with RIPA buffer.  
483 Bound proteins were eluted by sequential incubation of the resins with 5 µM USP2 and 5 µM  
484 SENP1 for 2 hours at 22°C, followed by a 5 minute incubation at 70°C in denaturing buffer  
485 (Novex NuPAGE LDS sample buffer; Life technologies). SENP1 (415-644) was prepared as  
486 described previously (Shen et al. 2006a). 6His-USP2(259-605) was purified as described for  
487 RNF168 below. Since RNF168 was most abundant in the denaturing elutions, only these elutions  
488 were used in further analyses. Eluates were fractionated on a NuPAGE Novex 10% Bis-Tris gel  
489 run in MOPS buffer, followed by Coomassie staining. Each lane was cut into upper and lower  
490 portions, followed by peptide extraction and in gel tryptic digestion (Shevchenko et al. 2006).  
491 Extracted peptides were resuspended in 35µl 0.5% acetic acid, 0.1% TFA and 8 µl was analysed  
492 by LC-MS/MS on Q Exactive Orbitrap mass spectrometer (Thermo Fisher Scientific). Data were

493 analysed with MaxQuant incorporating the Andromeda search engine (version 1.5.2.8) as in  
494 Tatham et al (Tatham et al. 2011). Protein intensity values based on extracted ion chromatograms  
495 (XICs) were reported for each purification. Protein intensities were arithmetically converted to  
496 ratio values and ratios were normalized to the median ratio of the entire group to allow for  
497 sample loading errors. Ratio scores were calculated for each protein by comparing data from  
498 each affinity resin to the blank resin. Only 69 proteins of the 1827 proteins processed had a ratio  
499 score >12 in at least one of the four comparisons. Hierarchical clustering was used to combine  
500 the four ratio scores of these proteins into a single heatmap using Perseus.

501

### 502 **Assay of purified RNF168 binding to SUMO2/ubiquitin polymers**

503 6His-RNF168 (full length) was expressed in *E.coli* and purified by nickel affinity  
504 chromatography, followed by removal of the 6His-tag by TEV protease digestion and  
505 repurification of the untagged product as previously described (Branigan et al. 2015). 10 µM  
506 purified RNF168 was incubated with 20 µl of each affinity resin in 100 µl of binding buffer (50  
507 mM Tris, 150 mM NaCl, 0.5 mM TCEP, 0.05% NP-40) overnight at 4°C. Resins were collected  
508 by centrifugation at 500 x g for 1 minute, then washed twice with binding buffer. Bound proteins  
509 were eluted by sequential incubation with USP2, SENP1 and denaturing buffer as described  
510 above. Proteins from the denaturing elution were analysed by SDS-PAGE and Coomassie  
511 staining.

512

### 513 **Acknowledgements**

514 We are grateful to Daniel Durocher for providing several constructs for RNF8 and RNF168  
515 expression and for helpful advice throughout the course of this work. We thank Patricia Mero and  
516 Cameron Landry for assistance in setting up the shRNA screen. We also thank Paul Sadowski and  
517 Melissa MacPherson for the plasmids expressing myc-tagged SUMO.

518

### 519 **Competing interests**

520 The authors declare no competing or financial interests.

521

### 522 **Author Contributions**

523 L.F., K.S. and R.T.H. conceived of and designed the experiments. K.S., A.I.W., M.H.T., O.F.A.,  
524 D.R. and S.G. performed the experiments. J.M. provided the lentivirus shRNA library and  
525 microscope for the initial screen. L.F. wrote the manuscript.

526

527

## 528 **Funding**

529 This work was supported by a Canadian Institutes of Health Research (CIHR) operating grant to LF  
530 and by a Programme grant from Cancer Research UK to RTH. OFA was supported by a studentship  
531 from the Biotechnology and Biological Sciences Research Council. LF is a tier one Canada Research  
532 Chair. RTH is a Senior Investigator of the Wellcome Trust.

533

534

535 **REFERENCES**

- 536 **Ablain J, Rice K, Soilihi H, de Reynies A, Minucci S, De The H** (2014). Activation of a  
 537 promyelocytic leukemia–tumor protein 53 axis underlies acute promyelocytic leukemia  
 538 cure. *Nature Medicine* **20**,167-174
- 539 **Al-Hakim A, Escribano-Diaz C, Landry MC, O'Donnell L, Panier S, Szilard RK, Durocher**  
 540 **D** (2010). The ubiquitous role of ubiquitin in the DNA damage response. *DNA Repair*  
 541 (*Amst*) **9** (12),1229-1240. doi,10.1016/j.dnarep.2010.09.011  
 542 S1568-7864(10)00316-2 [pii]
- 543 **Bartocci C, Denchi EL** (2013). Put a RING on it: regulation and inhibition of RNF8 and  
 544 RNF168 RING finger E3 ligases at DNA damage sites. *Front Genet* **4**,128.  
 545 doi,10.3389/fgene.2013.00128
- 546 **Bernardi R, Pandolfi PP** (2007). Structure, dynamics and functions of promyelocytic leukaemia  
 547 nuclear bodies. *Nat Rev Mol Cell Biol* **8**,1006-1016
- 548 **Bischof O, Kim SH, Irving J, Beresten S, Ellis NA, Campisi J** (2001). Regulation and  
 549 localization of the Bloom syndrome protein in response to DNA damage. *J Cell Biol* **153**  
 550 (2),367-380
- 551 **Boe SO, Haave M, Jul-Larsen A, Grudic A, Bjerkvig R, Lonning PE** (2006). Promyelocytic  
 552 leukemia nuclear bodies are predetermined processing sites for damaged DNA. *J Cell Sci*  
 553 **119** (Pt 16),3284-3295
- 554 **Boichuk S, Hu L, Makielski K, Pandolfi PP, Gjoerup OV** (2011). Functional connection  
 555 between Rad51 and PML in homology-directed repair. *PLoS ONE* **6** (10),e25814.  
 556 doi,10.1371/journal.pone.0025814  
 557 PONE-D-11-11910 [pii]
- 558 **Branigan E, Plechanovova A, Jaffray EG, Naismith JH, Hay RT** (2015). Structural basis for  
 559 the RING-catalyzed synthesis of K63-linked ubiquitin chains. *Nat Struct Mol Biol* **22**  
 560 (8),597-602. doi,10.1038/nsmb.3052  
 561 nsmb.3052 [pii]
- 562 **Condemine W, Takahashi Y, Zhu J, Puvion-Dutilleul F, Guegan S, Janin A, de The H**  
 563 (2006). Characterization of endogenous human promyelocytic leukemia isoforms. *Cancer*  
 564 *Res* **66** (12),6192-6198
- 565 **Cuchet-Lourenco D, Vanni E, Glass M, Orr A, Everett RD** (2012). Herpes simplex virus 1  
 566 ubiquitin ligase ICP0 interacts with PML isoform I and induces its SUMO-independent  
 567 degradation. *J Virol* **86** (20),11209-11222. doi,JVI.01145-12 [pii]  
 568 10.1128/JVI.01145-12
- 569 **Cuchet D, Sykes A, Nicolas A, Orr A, Murray J, Sirma H, Heeren J, Bartelt A, Everett RD**  
 570 (2011). PML isoforms I and II participate in PML-dependent restriction of HSV-1  
 571 replication. *J Cell Sci* **124** (Pt 2),280-291. doi,jcs.075390 [pii]  
 572 10.1242/jcs.075390
- 573 **Danielsen JR, Povlsen LK, Villumsen BH, Streicher W, Nilsson J, Wikstrom M, Bekker-**  
 574 **Jensen S, Mailand N** (2012). DNA damage-inducible SUMOylation of HERC2  
 575 promotes RNF8 binding via a novel SUMO-binding Zinc finger. *J Cell Biol* **197** (2),179-  
 576 187. doi,10.1083/jcb.201106152  
 577 jcb.201106152 [pii]
- 578 **de The H, Le Bras M, Lallemand-Breitenbach V** (2012). The cell biology of disease: Acute  
 579 promyelocytic leukemia, arsenic, and PML bodies. *J Cell Biol* **198** (1),11-21.  
 580 doi,10.1083/jcb.201112044

581 jcb.201112044 [pii]  
582 **Dellaire G, Bazett-Jones DP** (2004). PML nuclear bodies: dynamic sensors of DNA damage  
583 and cellular stress. *Bioessays* **26** (9),963-977  
584 **Dellaire G, Ching RW, Ahmed K, Jalali F, Tse KC, Bristow RG, Bazett-Jones DP** (2006).  
585 Promyelocytic leukemia nuclear bodies behave as DNA damage sensors whose response  
586 to DNA double-strand breaks is regulated by NBS1 and the kinases ATM, Chk2, and  
587 ATR. *J Cell Biol* **175** (1),55-66  
588 **Doil C, Mailand N, Bekker-Jensen S, Menard P, Larsen DH, Pepperkok R, Ellenberg J,**  
589 **Panier S, Durocher D, Bartek J, Lukas J, Lukas C** (2009). RNF168 binds and  
590 amplifies ubiquitin conjugates on damaged chromosomes to allow accumulation of repair  
591 proteins. *Cell* **136** (3),435-446. doi,10.1016/j.cell.2008.12.041  
592 S0092-8674(09)00004-X [pii]  
593 **Erker Y, Neyret-Kahn H, Seeler JS, Dejean A, Atfi A, Levy L** (2013). Arkadia, a novel  
594 SUMO-targeted ubiquitin ligase involved in PML degradation. *Mol Cell Biol* **33**  
595 (11),2163-2177. doi,10.1128/MCB.01019-12  
596 MCB.01019-12 [pii]  
597 **Everett RD, Chelbi-Alix MK** (2007). PML and PML nuclear bodies: implications in antiviral  
598 defence. *Biochimie* **89** (6-7),819-830  
599 **Fanelli M, Fantozzi A, De Luca P, Caprodossi S, Matsuzawa S, Lazar MA, Pelicci PG,**  
600 **Minucci S** (2004). The coiled-coil domain is the structural determinant for mammalian  
601 homologues of Drosophila Sina-mediated degradation of promyelocytic leukemia protein  
602 and other tripartite motif proteins by the proteasome. *J Biol Chem* **279** (7),5374-5379.  
603 doi,10.1074/jbc.M306407200  
604 M306407200 [pii]  
605 **Galanty Y, Belotserkovskaya R, Coates J, Polo S, Miller KM, Jackson SP** (2009).  
606 Mammalian SUMO E3-ligases PIAS1 and PIAS4 promote responses to DNA double-  
607 strand breaks. *Nature* **462** (7275),935-939. doi,10.1038/nature08657  
608 nature08657 [pii]  
609 **Geoffroy MC, Chelbi-Alix MK** (2011). Role of promyelocytic leukemia protein in host  
610 antiviral defense. *J Interferon Cytokine Res* **31** (1),145-158. doi,10.1089/jir.2010.0111  
611 **Geoffroy MC, Hay RT** (2009). An additional role for SUMO in ubiquitin-mediated proteolysis.  
612 *Nat Rev Mol Cell Biol* **10** (8),564-568. doi,10.1038/nrm2707  
613 nrm2707 [pii]  
614 **Guzzo CM, Berndsen CE, Zhu J, Gupta V, Datta A, Greenberg RA, Wolberger C, Matunis**  
615 **MJ** (2012). RNF4-dependent hybrid SUMO-ubiquitin chains are signals for RAP80 and  
616 thereby mediate the recruitment of BRCA1 to sites of DNA damage. *Sci Signal* **5**  
617 (253),ra88. doi,10.1126/scisignal.2003485  
618 5/253/ra88 [pii]  
619 **Hu X, Paul A, Wang B** (2012). Rap80 protein recruitment to DNA double-strand breaks  
620 requires binding to both small ubiquitin-like modifier (SUMO) and ubiquitin conjugates.  
621 *J Biol Chem* **287** (30),25510-25519. doi,10.1074/jbc.M112.374116  
622 M112.374116 [pii]  
623 **Lallemant-Breitenbach V, de The H** (2010). PML nuclear bodies. *Cold Spring Harb Perspect*  
624 *Biol* **2** (5),a000661

625 **Lallemand-Breitenbach V, Jeanne M, Benhenda S, Nasr R, Lei M, Peres L, Zhou J, Zhu J,**  
626 **Raught B, de The H** (2008). Arsenic degrades PML or PML-RARalpha through a  
627 SUMO-triggered RNF4/ubiquitin-mediated pathway. *Nat Cell Biol* **10** (5),547-555

628 **Lallemand-Breitenbach V, Zhu J, Puvion F, Koken M, Honore N, Doubeikovsky A, Duprez**  
629 **E, Pandolfi PP, Puvion E, Freemont P, de The H** (2001). Role of promyelocytic  
630 leukemia (PML) sumolation in nuclear body formation, 11S proteasome recruitment, and  
631 As2O3-induced PML or PML/retinoic acid receptor alpha degradation. *J Exp Med* **193**  
632 (12),1361-1371

633 **MacPherson MJ, Beatty LG, Zhou W, Du M, Sadowski PD** (2009). The CTCF insulator  
634 protein is posttranslationally modified by SUMO. *Mol Cell Biol* **29** (3),714-725.  
635 doi,10.1128/MCB.00825-08  
636 MCB.00825-08 [pii]

637 **Mailand N, Bekker-Jensen S, Faustrup H, Melander F, Bartek J, Lukas C, Lukas J** (2007).  
638 RNF8 ubiquitylates histones at DNA double-strand breaks and promotes assembly of  
639 repair proteins. *Cell* **131** (5),887-900. doi,S0092-8674(07)01271-8 [pii]  
640 10.1016/j.cell.2007.09.040

641 **Moffat J, Grueneberg DA, Yang X, Kim SY, Kloepfer AM, Hinkle G, Piqani B, Eisenhaure**  
642 **TM, Luo B, Grenier JK, Carpenter AE, Foo SY, Stewart SA, Stockwell BR,**  
643 **Hacohen N, Hahn WC, Lander ES, Sabatini DM, Root DE** (2006). A lentiviral RNAi  
644 library for human and mouse genes applied to an arrayed viral high-content screen. *Cell*  
645 **124** (6),1283-1298. doi,S0092-8674(06)00238-8 [pii]  
646 10.1016/j.cell.2006.01.040

647 **Nisole S, Maroui MA, Mascle XH, Aubry M, Chelbi-Alix MK** (2013). Differential Roles of  
648 PML Isoforms. *Front Oncol* **3**,125. doi,10.3389/fonc.2013.00125

649 **Panier S, Durocher D** (2009). Regulatory ubiquitylation in response to DNA double-strand  
650 breaks. *DNA Repair (Amst)* **8** (4),436-443. doi,S1568-7864(09)00008-1 [pii]  
651 10.1016/j.dnarep.2009.01.013

652 **Panier S, Ichijima Y, Fradet-Turcotte A, Leung CC, Kaustov L, Arrowsmith CH,**  
653 **Durocher D** (2012). Tandem protein interaction modules organize the ubiquitin-  
654 dependent response to DNA double-strand breaks. *Mol Cell* **47** (3),383-395.  
655 doi,10.1016/j.molcel.2012.05.045  
656 S1097-2765(12)00490-X [pii]

657 **Pinato S, Scandiuzzi C, Arnaudo N, Citterio E, Gaudino G, Penengo L** (2009). RNF168, a  
658 new RING finger, MIU-containing protein that modifies chromatin by ubiquitination of  
659 histones H2A and H2AX. *BMC Mol Biol* **10**,55. doi,10.1186/1471-2199-10-55  
660 1471-2199-10-55 [pii]

661 **Rabellino A, Scaglioni PP** (2013). PML Degradation: Multiple Ways to Eliminate PML. *Front*  
662 *Oncol* **3**,60. doi,10.3389/fonc.2013.00060

663 **Salomoni P, Ferguson BJ, Wyllie AH, Rich T** (2008). New insights into the role of PML in  
664 tumour suppression. *Cell Res* **18** (6),622-640

665 **Sarkari F, Wang X, Nguyen T, Frappier L** (2011). The herpesvirus associated ubiquitin  
666 specific protease, USP7, is a negative regulator of PML proteins and PML nuclear  
667 bodies. *PLoS ONE* **6** (1),e16598. doi,10.1371/journal.pone.0016598

668 **Scaglioni PP, Yung TM, Cai LF, Erdjument-Bromage H, Kaufman AJ, Singh B, Teruya-**  
669 **Feldstein J, Tempst P, Pandolfi PP** (2006). A CK2-dependent mechanism for  
670 degradation of the PML tumor suppressor. *Cell* **126** (2),269-283

671 **Scaglioni PP, Yung TM, Choi SC, Baldini C, Konstantinidou G, Pandolfi PP** (2008). CK2  
672 mediates phosphorylation and ubiquitin-mediated degradation of the PML tumor  
673 suppressor. *Mol Cell Biochem* **316**,149-154

674 **Seeler JS, Dejean A** (2001). SUMO: of branched proteins and nuclear bodies. *Oncogene* **20**  
675 (49),7243-7249. doi,10.1038/sj.onc.1204758

676 **Seifert A, Schofield P, Barton GJ, Hay RT** (2015). Proteotoxic stress reprograms the  
677 chromatin landscape of SUMO modification. *Sci Signal* **8** (384),rs7.  
678 doi,10.1126/scisignal.aaa2213  
679 8/384/rs7 [pii]

680 **Shen L, Tatham MH, Dong C, Zagorska A, Naismith JH, Hay RT** (2006a). SUMO protease  
681 SENP1 induces isomerization of the scissile peptide bond. *Nat Struct Mol Biol* **13**  
682 (12),1069-1077. doi,nsmb1172 [pii]  
683 10.1038/nsmb1172

684 **Shen TH, Lin HK, Scaglioni PP, Yung TM, Pandolfi PP** (2006b). The mechanisms of PML-  
685 nuclear body formation. *Mol Cell* **24** (3),331-339

686 **Shevchenko A, Tomas H, Havlis J, Olsen JV, Mann M** (2006). In-gel digestion for mass  
687 spectrometric characterization of proteins and proteomes. *Nat Protoc* **1** (6),2856-2860.  
688 doi,nprot.2006.468 [pii]  
689 10.1038/nprot.2006.468

690 **Sivachandran N, Cao JY, Frappier L** (2010). Epstein-Barr virus nuclear antigen 1 Hijacks the  
691 host kinase CK2 to disrupt PML nuclear bodies. *J Virol* **84** (21),11113-11123

692 **Sivachandran N, Dawson CW, Young LS, Liu FF, Middeldorp J, Frappier L** (2012a).  
693 Contributions of the Epstein-Barr virus EBNA1 protein to gastric carcinoma. *J Virol* **86**  
694 (1),60-68. doi,JVI.05623-11 [pii]  
695 10.1128/JVI.05623-11

696 **Sivachandran N, Sarkari F, Frappier L** (2008). Epstein-Barr nuclear antigen 1 contributes to  
697 nasopharyngeal carcinoma through disruption of PML nuclear bodies. *PLoS Pathog* **4**  
698 (10),e1000170. doi,10.1371/journal.ppat.1000170

699 **Sivachandran N, Wang X, Frappier L** (2012b). Functions of the Epstein-Barr Virus EBNA1  
700 Protein in Viral Reactivation and Lytic Infection. *J Virol* **86** (11),6146-6158.  
701 doi,JVI.00013-12 [pii]  
702 10.1128/JVI.00013-12

703 **Stewart GS, Panier S, Townsend K, Al-Hakim AK, Kolas NK, Miller ES, Nakada S,**  
704 **Ylanko J, Olivarius S, Mendez M, Oldreive C, Wildenhain J, Tagliaferro A,**  
705 **Pelletier L, Taubenheim N, Durandy A, Byrd PJ, Stankovic T, Taylor AM,**  
706 **Durocher D** (2009). The RIDDLE syndrome protein mediates a ubiquitin-dependent  
707 signaling cascade at sites of DNA damage. *Cell* **136** (3),420-434. doi,S0092-  
708 8674(09)00005-1 [pii]  
709 10.1016/j.cell.2008.12.042

710 **Sun Y, Hegamyer G, Cheng YJ, Hildesheim A, Chen JY, Chen IH, Cao Y, Yao KT,**  
711 **Colburn NH** (1992). An infrequent point mutation of the p53 gene in human  
712 nasopharyngeal carcinoma. *Proc Natl Acad Sci U S A* **89** (14),6516-6520

713 **Tatham MH, Geoffroy MC, Shen L, Plechanovova A, Hattersley N, Jaffray EG, Palvimo**  
714 **JJ, Hay RT** (2008). RNF4 is a poly-SUMO-specific E3 ubiquitin ligase required for  
715 arsenic-induced PML degradation. *Nat Cell Biol* **10** (5),538-546



716 **Tatham MH, Matic I, Mann M, Hay RT** (2011). Comparative proteomic analysis identifies a  
717 role for SUMO in protein quality control. *Sci Signal* **4** (178),rs4.  
718 doi,10.1126/scisignal.2001484  
719 4/178/rs4 [pii]

720 **Tatham MH, Plechanovova A, Jaffray EG, Salmen H, Hay RT** (2013). Ube2W conjugates  
721 ubiquitin to alpha-amino groups of protein N-termini. *Biochem J* **453** (1),137-145.  
722 doi,10.1042/BJ20130244  
723 BJ20130244 [pii]

724 **Tavalai N, Stamminger T** (2009). Interplay between Herpesvirus Infection and Host Defense  
725 by PML Nuclear Bodies. *Viruses* **1** (3),1240-1264. doi,10.3390/v1031240  
726 viruses-01-01240 [pii]

727 **Tavalai N, Stamminger T** (2011). Intrinsic cellular defense mechanisms targeting human  
728 cytomegalovirus. *Virus Res* **157** (2),128-133. doi,S0168-1702(10)00361-8 [pii]  
729 10.1016/j.virusres.2010.10.002

730 **Tikoo S, Madhavan V, Hussain M, Miller ES, Arora P, Zlatanou A, Modi P, Townsend K,  
731 Stewart GS, Sengupta S** (2013). Ubiquitin-dependent recruitment of the Bloom  
732 syndrome helicase upon replication stress is required to suppress homologous  
733 recombination. *Embo J* **32** (12),1778-1792. doi,10.1038/emboj.2013.117  
734 emboj2013117 [pii]

735 **Wolyniec K, Shortt J, de Stanchina E, Levav-Cohen Y, Alsheich-Bartok O, Louria-Hayon  
736 I, Corneille V, Kumar B, Woods SJ, Opat S, Johnstone RW, Scott CL, Segal D,  
737 Pandolfi PP, Fox S, Strasser A, Jiang YH, Lowe SW, Haupt S, Haupt Y** (2012).  
738 E6AP ubiquitin ligase regulates PML-induced senescence in Myc-driven  
739 lymphomagenesis. *Blood* **120** (4),822-832. doi,10.1182/blood-2011-10-387647  
740 blood-2011-10-387647 [pii]

741 **Wu HC, Lin YC, Liu CH, Chung HC, Wang YT, Lin YW, Ma HI, Tu PH, Lawler SE,  
742 Chen RH** (2014). USP11 regulates PML stability to control Notch-induced malignancy  
743 in brain tumours. *Nat Commun* **5**,3214. doi,10.1038/ncomms4214  
744 ncomms4214 [pii]

745 **Yeung PL, Denissova NG, Nasello C, Hakhverdyan Z, Chen JD, Brenneman MA** (2012).  
746 Promyelocytic leukemia nuclear bodies support a late step in DNA double-strand break  
747 repair by homologous recombination. *J Cell Biochem* **113** (5),1787-1799.  
748 doi,10.1002/jcb.24050

749 **Yuan WC, Lee YR, Huang SF, Lin YM, Chen TY, Chung HC, Tsai CH, Chen HY, Chiang  
750 CT, Lai CK, Lu LT, Chen CH, Gu DL, Pu YS, Jou YS, Lu KP, Hsiao PW, Shih  
751 HM, Chen RH** (2011). A Cullin3-KLHL20 Ubiquitin ligase-dependent pathway targets  
752 PML to potentiate HIF-1 signaling and prostate cancer progression. *Cancer Cell* **20**  
753 (2),214-228. doi,10.1016/j.ccr.2011.07.008  
754 S1535-6108(11)00263-7 [pii]

755 **Zhong S, Hu P, Ye TZ, Stan R, Ellis NA, Pandolfi PP** (1999). A role for PML and the nuclear  
756 body in genomic stability. *Oncogene* **18** (56),7941-7947  
757  
758  
759  
760

761 **FIGURE LEGENDS**

762 **Fig. 1.** An shRNA screen identifies RNF8 and RNF168 as regulators of PML NBs. CNE2Z cells  
763 were infected with a lentivirus expressing an shRNA targeted to RNF111 (negative control),  
764 RNF168 (shRNAs 1 to 5) or RNF8 (shRNAs 1 to 5) or left uninfected (none). Cells were then fixed,  
765 stained with Hoechst and with PML-specific antibody, then imaged by confocal microscopy. A. The  
766 number of PML NBs per cell and intensity of PML NBs was determined. Average values with  
767 standard deviation are shown. B. Sample images for the indicated shRNAs from A.

768  
769 **Fig. 2.** siRNA targeting RNF8 and RNF168 increases PML NBs and PML protein levels. A.  
770 CNE2Z cells were transfected with siRNA targeting RNF8 or RNF168 or negative control siRNA  
771 (control), then stained for PML. B. The number of PML NBs per cell was determined for 100 cells  
772 in 3 independent experiments and average values with standard deviation were plotted. *P* values  
773 (determined by t-tests) are indicated as follows: \* =  $0.01 < P < 0.05$ ; \*\* =  $0.001 < P < 0.01$ ; \*\*\* =  $P <$   
774  $0.001$ . C. CNE2Z cells treated with siRNA as in A were lysed and analysed by Western blotting  
775 using the indicated antibodies. PML bands were quantified by densitometry and normalized to actin,  
776 and values for each lane are shown relative to the Control lane under the gel. D to F. The same  
777 experiment as in A to C but performed in U2OS cells.

778  
779 **Fig. 3.** Overexpression of RNF8 or RNF 168 induces loss of PML NBs. A. CNE2Z cells were  
780 transfected with plasmids expressing FLAG-tagged RNF8 or RNF168 then stained for FLAG and  
781 PML. Sample images showing fewer PML NBs in FLAG-positive cells are shown. B. PML NBs  
782 were counted for cells expressing FLAG-RNF8 or FLAG-RNF168 and compared to cells transfected  
783 with empty plasmid (pcDNA3). Average values with standard deviation are shown from 3  
784 independent experiments, with *P* values defined as in Fig. 2.

785  
786 **Fig. 4.** RNF168 can induce loss of PML NBs independent of RNF8. CNE2Z cells were treated with  
787 siRNA targeting RNF8 (siRNF8) or negative control siRNA (control) then transfected with the  
788 FLAG-RNF168 expression plasmid (bottom two rows in A) or empty control plasmid pcDNA3 (top  
789 two rows in A). Cells were then either fixed and stained for PML and FLAG (A) or lysed and  
790 analysed by Western blotting using RNF8 specific antibody to confirm RNF8 silencing (B). C. PML  
791 NBs in A were counted for 100 cells in three independent experiments. For RNF168 samples, PML

792 NBs were counted only in FLAG-positive cells. Average values with standard deviations are plotted  
793 with *P* values (\*\*\*) = *P* < 0.001) for RNF168 overexpression shown relative to the pcDNA3 control  
794 in each condition.

795  
796 **Fig. 5.** RNF168 localization to PML NBs. A. Schematic representation of RNF168 showing the  
797 positions of the catalytic RING domain, ubiquitin binding sequences (UMI, MIU1, MIU2) and LR  
798 motifs (LRM) (Panier et al. 2012). B. CNE2Z cells were transiently transfected with plasmids  
799 expressing FLAG-tagged RNF168 with WT sequence or the indicated deletions or with pcDNA3  
800 empty plasmid. Cells were then stained for FLAG and PML and imaged by fluorescence  
801 microscopy. C. Western blots for FLAG-tagged and GFP-tagged proteins expressed in B and D. D.  
802 CNE2Z cells were transfected with plasmids expressing GFP fused to RNF168 amino acids 100-201  
803 (containing LRM1, UMI and MIU1) or 100-166 (containing LRM1 and UMI) or GFP alone, then  
804 stained for PML and imaged for GFP and PML. E. In the bottom panels, CNE2Z cells were  
805 transiently transfected with plasmids expressing FLAG- $\Delta$ RING, treated with etoposide (10  $\mu$ g/ml)  
806 for 5 hours then stained for FLAG and PML and imaged by fluorescence microscopy. In the top  
807 panels, CNE2Z cells were treated with etoposide as in bottom panels or left untreated as indicated,  
808 then stained for 53BP1 to confirm that DDR foci were induced by etoposide.

809  
810 **Fig. 6.** RNF168 localizes to NBs formed by any PML isoform. A plasmid expressing FLAG-tagged  
811 RNF168  $\Delta$ MIU2 was used to transfect CNE2Z cells or CNE2Z cells that lack WT PML but express  
812 a single recombinant PML isoform ( PML I to PML VI) (Sarkari et al. 2011; Sivachandran et al.  
813 2012b). Cells were then stained for FLAG and PML.

814  
815 **Fig. 7.** RNF168 induces PML ubiquitylation and SUMOylation. A. CNE2Z cells were transfected  
816 with a plasmid expressing HA-tagged ubiquitin and a second plasmid expressing RNF168 or the  
817 RNF168 mutants  $\Delta$ MIU2,  $\Delta$ Ub ( $\Delta$ UMI $\Delta$ MIU1 $\Delta$ MIU2) or  $\Delta$ RING or empty plasmid (pcDNA3) in  
818 the presence of proteasomal inhibition. Total PML was then immunoprecipitated from cell lysates  
819 and analysed by Western blotting for PML and HA (IP). A sample of the cell lysate prior to  
820 immunoprecipitation is also shown (INPUT). B. Experiments were performed as in A in either the  
821 presence (+) or absence (-) of MG132 proteasomal inhibitor. Ubiquitylated proteins in input and IP  
822 samples were detected by Western blotting using antibodies against HA, K48-linked ubiquitin or

823 K63-linked ubiquitin. C. The same experiments as in A except that the plasmid expressing HA-  
824 tagged ubiquitin was replaced by one expressing myc-tagged SUMO2. In addition, a positive control  
825 for PML SUMOylation is shown, in which cells transfected with pcDNA3 were treated with arsenic  
826 trioxide (pcDNA3+As). Two exposures of the Input FLAG blot are shown.

827


828 **Fig. 8.** RNF168 associates with SUMO2/3-modified proteins in cells and preferentially binds  
829 hybrid ubiquitin-SUMO chains *in vitro*. A. 293T cells were co-transfected with a plasmid  
830 expressing FLAG-tagged RNF168 (wt), RNF168 $\Delta$ Ub ( $\Delta$ UMI $\Delta$ MIU1 $\Delta$ MIU2) or empty plasmid  
831 (Control) and a second plasmid expressing myc-tagged SUMO1, SUMO2 or SUMO3. After cell  
832 lysis, a sample was analysed by Western blotting (Input), while the remaining sample was  
833 subjected to FLAG IP, followed by Western blotting for Myc and FLAG. The positions of  
834 molecular weight markers are indicated on the right. B. Comparison of retention of RNF168  
835 from 293 cell extracts on affinity resins containing K63-ubiquitin polymers (K63-Ub), a linear  
836 4xSUMO2 chain (4xSUMO-2), an N-terminally monoubiquitinated 4xSUMO2 chain (Ub-  
837 4xSUMO-2), hybrid chains containing K63-ubiquitin polymers linked to 4xSUMO-2 (K63Ub-  
838 4xSUMO-2) or no protein (blank). Total intensity of RNF168 peptides retained on each resin are  
839 shown. RNF168 identification was based on 26 unique peptides, 47% sequence coverage. C.  
840 Purified RNF168 was incubated with the affinity resins or blank resin from B and, after washing  
841 was eluted with denaturing buffer. A Coomassie stained gel is shown of the protein loaded onto  
842 the resins (load) and eluted from the indicated resins (Pulldown).







843


844

**A**

Lentivirus	PML bodies/cell	Intensity/NB
none	10.4 +/- 2.2	124.2 +/- 20
RNF111	11 +/- 3.0	122 +/- 19
RNF168-1	14.5 +/- 1.7	282 +/- 38
RNF168-2	11.8 +/- 1.5	180 +/- 10
RNF168-3	11.8 +/- 1.5	149 +/- 27
RNF168-4	19.5 +/- 1.3	417 +/- 39
RNF168-5	13.5 +/- 1.3	167 +/- 20
RNF8-1	18.5 +/- 2.1	218 +/- 9.9
RNF8-2	8.0 +/- 2.2	235 +/- 30
RNF8-3	10.3 +/- 3.1	154 +/- 18
RNF8-4	18.0 +/- 4.8	172 +/- 16
RNF8-5	13.0 +/- 3.8	196 +/- 55

**B**



**A****B**