



**University of Dundee**

**Stiffness at shear-wave elastography and patient presentation predicts upgrade at surgery following an ultrasound-guided core biopsy diagnosis of ductal carcinoma in situ**

Evans, A.; Purdie, C. A.; Jordan, L.; Macaskill, E. J.; Flynn, J.; Vinnicombe, S.

*Published in:*  
Clinical Radiology

*DOI:*  
[10.1016/j.crad.2016.07.004](https://doi.org/10.1016/j.crad.2016.07.004)

*Publication date:*  
2016

*Document Version*  
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

*Citation for published version (APA):*

Evans, A., Purdie, C. A., Jordan, L., Macaskill, E. J., Flynn, J., & Vinnicombe, S. (2016). Stiffness at shear-wave elastography and patient presentation predicts upgrade at surgery following an ultrasound-guided core biopsy diagnosis of ductal carcinoma in situ. *Clinical Radiology*, 71(11), 1156-1159. DOI: 10.1016/j.crad.2016.07.004

**General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

1     **Introduction**

2     Breast lesions yielding an image guided biopsy diagnosis of Ductal Carcinoma in Situ (DCIS) may contain  
3     invasive foci when the entire lesion is removed at surgery. Such an upgrade may lead to further surgery,  
4     especially to the axilla (1). Predicting the likelihood of upgrade to invasion is useful as it may change  
5     options for initial surgery, particularly with regard to sentinel node biopsy. Factors predicting upgrade  
6     of DCIS lesions diagnosed with stereotactic biopsy have been well established in the literature and  
7     include mammographic extent, palpability and grade of DCIS diagnosed on percutaneous biopsy (2-4).  
8     However, factors predicting upgrade following an US guided biopsy of DCIS have not been well  
9     established and only two previous **studies have** studied shear wave elastography (SWE) in this context  
10    (5-8). A number of reports of SWE of the breast have shown that DCIS lesions are less stiff than invasive  
11    cancers (9, 10). These findings suggest that SWE may be useful in predicting the presence of invasion.  
12    The aim of this study is to establish the role of mammographic, ultrasound, SWE and other pre-operative  
13    features in predicting invasion at surgical excision in lesions yielding an US guided 14g core biopsy  
14    diagnosis of DCIS.

15

16 **Methods**

17 SWE has been part of the routine breast ultrasound examination at our centre since November 2009.  
18 Quantitative data are routinely extracted prospectively within a few days of the examination and before  
19 biopsy results were known. In accordance with the applicable National Research Ethics Service guidance,  
20 ethical approval for this retrospective study of prospectively acquired anonymised data was not  
21 required (National Research Ethics Service, 2008(11)). However, written informed consent to use images  
22 was obtained, according to routine practice.

23 All patients with US visible lesions were scanned using the Aixplorer ultrasound system (SuperSonic  
24 Imagine™, Aix en Provence, France) between December 2010 and December 2015. Those patients with  
25 lesions visible on greyscale US and subjected to ultrasound guided 14g needle core biopsy yielding a  
26 biopsy diagnosis of DCIS were included in this study. Patients with biopsies from elsewhere in the  
27 ipsilateral breast showing invasive disease were excluded.

28 All patients were scanned and biopsied with ultrasound guidance by one of five breast radiologists or an  
29 advanced radiography practitioner trained to perform and interpret breast ultrasound. These  
30 practitioners had between 7 and 22 years of breast ultrasound experience and had at least 3 months  
31 experience of performing SWE of solid breast lesions prior to the commencement of the study.

32 Greyscale and SWE images were obtained during the standard ultrasound appointment. Four SWE  
33 images in two orthogonal planes were obtained. The region of interest (ROI) utilised in all cases was  
34 2mm in diameter. A threshold value of 50 kilopascals (kPa) was used for mean elasticity (E<sub>mean</sub>) to  
35 dichotomise the elasticity data. This threshold has been validated in previous studies using SWE to  
36 differentiate between benign and malignant masses (12, 13). The average of the 4 values from the 4  
37 images was used for analysis. Additionally the mean SWE value across the entire study group was also  
38 used as a possible cut off value.

39 The size and BIRADS classification of the greyscale US abnormalities were recorded as was the presence  
40 or absence of an US visible mass or calcification (some lesions were visible as textural change with or  
41 without US visible calcifications). The dominant mammographic feature was also recorded as was the  
42 BIRADS classification of any mammographic abnormality.

43 The source of the patient from either mammographic screening or the symptomatic referral service was  
44 noted as was the palpability of the lesion. The histological grade of DCIS present on the US guided 14g  
45 core biopsy was recorded. Regarding subsequent surgical excisions, micro-invasion was included in the  
46 DCIS group as such lesions do not routinely warrant an axillary nodal procedure. The size, grade,  
47 vascular invasion and nodal status of any invasive foci were collected from the surgical pathology report.

48 The significance of differences between groups was established using the chi-square test with  $p < 0.05$   
49 taken to indicate statistically significant differences.

50

51

52 **Results**

53 Analysis of 1954 US visible masses with histological correlation yielded 57(3%) patients with an US 14g  
54 core biopsy diagnosis of DCIS who had subsequent surgical excision. The mean age of the patients was  
55 60 yrs (range 23 -82 yrs). 33(58%) lesions were screen-detected while 24(42%) resulted from  
56 investigation of symptoms. At surgical excision 24(42%) patients had invasive foci while 33 had DCIS  
57 only. The features of the 24 invasive cancers are shown in Table 1. 12 of 24 (50%) invasive foci were  
58 less than 10mm in diameter and 2 of 24 (8%) were node positive. The mean histological whole tumour  
59 size (i.e. extent including DCIS) was 28mm for the pure DCIS lesions and 53 mm for those containing an  
60 invasive focus.

61 The relationship between pre-operative factors and presence of invasion are shown in Tables 2 to 4.  
62 Patients presenting with symptoms were more likely to have invasive foci than patients presenting  
63 through screening (58% [14 of 24] vs. 30% [10 of 33],  $p=0.03$ ). Patients whose lesions had a mean  
64 stiffness at SWE of  $> 50\text{kPa}$  were more likely to have an invasive focus than women with lesions whose  
65 mean stiffness was  $<50\text{kPa}$  (51% [20 of 39] vs. 22% [4 of 18],  $p=0.04$ ). The mean stiffness value in the  
66 entire study group was 84kPa. Using this as a cut off value did not aid the identification of those women  
67 with invasive foci. The mean stiffness values for women with and without invasion were 89 kPa and  
68 78kPa respectively, while the median values were 74 kPa and 56 kPa. Trends towards younger women  
69 and women with high greyscale US suspicion scores (BI-RADS) did not achieve statistical significance.

70 Patient age, core biopsy DCIS grade, US lesion size (mean size was 17mm), mammographic features and  
71 mammography BI-RADS score showed no significant relationship with the presence of invasion at  
72 excision. The presence of an ultrasound visible mass or calcification likewise did not have an association  
73 with invasion. Palpability of the lesion did not have a statistically significant relationship with invasion as  
74 10 of 18 (55%) of palpable lesions were invasive compared to 14 of 39(36%) impalpable lesions ( $p=0.16$ ).

75 Combining the two significant predictors of invasion further improves risk stratification (Table 5) with  
76 symptomatic and stiff lesions having a risk of invasion of 67% (12 of 18) while soft lesions presenting at  
77 screening had only a 17% (2 of 12) risk of invasion ( $p=0.02$ ).

78

79 **Discussion**

80 We have shown that the stiffness of lesions yielding a core biopsy diagnosis of DCIS has a significant  
81 relationship with the risk of occult invasion at excision. We have also found that occult invasion is  
82 commoner in women whose lesions are symptomatic compared to women presenting through  
83 mammographic screening. The combination of these two factors allows a high risk group of women  
84 with stiff lesions that are symptomatic to be identified. This group has a risk of invasion of 67% and it  
85 contains half the women in the study group with invasion. Thus, this is a group where sentinel lymph  
86 node biopsy (SNB) may be justified or, alternatively, another approach such as re-biopsing these women  
87 using a large volume vacuum-assisted biopsy (VAB) technique to try and confirm the presence of  
88 invasion could be considered. Women whose lesions are not symptomatic or stiff have a low risk of  
89 invasion (17%) and neither repeat biopsy nor SNB would seem justified. The two intermediate groups  
90 (screen detected and stiff, or symptomatic but soft) also have a relatively low risk of invasion and SNB or  
91 further biopsy is probably not required in these patients either. It should also be noted that the nodal  
92 positivity rate in those women with invasion in this study was very low (8%).

93 A number of previous studies have shown that invasive cancers tend to be stiffer than DCIS lesions (9,  
94 10) while two previous studies have shown that stiffness is associated with upgrade following an  
95 ultrasound guided biopsy with a DCIS diagnosis (5,8). Neither of these studies combined the significant  
96 factors to provide the risk stratification shown in table 5 which may aid clinical management.

97 What is unclear is whether the increased stiffness in the upgrade group is because the invasive focus  
98 itself was imaged but not biopsied or whether increased stiffness identifies DCIS lesions at particular risk  
99 of invasion. The stiffness of invasive cancers is usually most marked at the lesional/stromal boundary  
100 and is thought to be due in part, to the stiffness of the extracellular matrix (ECM) produced by cancer  
101 associated fibroblasts (CAFs). CAFs are important in tumour progression, invasion and metastasis (14-

102 16) and this is likely to be the reason why stiffness at SWE has been shown to be an independent  
103 predictor of nodal metastasis (17). Similarly the stiffness detected in DCIS lesions is likely to be due to  
104 the stiffness of the extracellular matrix produced by CAFs, which are also present within DCIS lesions.  
105 Stromal influences are known to be important in initiating invasion within DCIS lesions (14-16). It is  
106 therefore possible that the increased stiffness seen with DCIS lesions with occult invasion is due to stiff  
107 stroma produced by activated CAFs which also initiate invasion, in contrast to less stiff stroma in DCIS  
108 lesions with lower invasive potential.

109 We have also found that occult invasion is commoner in women whose lesions were symptomatic  
110 compared to women presenting through mammographic screening. It has been noted in previous  
111 studies that palpability is associated with upgrade following a biopsy diagnosis of DCIS (2-4). Not all  
112 patients with symptoms have a palpable lesion (some may have presented with nipple discharge) and  
113 some screen detected lesions are palpable. We found palpability less predictive than the source of  
114 referral.

115 A number of previous studies have studied the greyscale US features of lesions yielding an US guided  
116 core biopsy diagnosis of DCIS and upgrade to invasive cancer. The results of these studies have been  
117 inconsistent. Some have found abnormal axillary lymph nodes and large lesion size to be associated  
118 with higher rates of upgrade to invasive cancer (6, 7). In our study, we did not find US lesion size a  
119 useful predictor of invasion. We did not assess the value of abnormal axillary lymph nodes in our study  
120 as the node positivity rate was so low.

121 Studies looking at upgrade following stereotactic biopsy of malignant calcifications have found that DCIS  
122 grade of the biopsy and calcification extent are predictors of occult invasion (2-4). Neither of these  
123 factors was significant in the present study. The reason why grade of DCIS does not predict invasion in



124 the current study is not clear. There is not a hint of a difference so the small numbers in the current  
125 study does not appear to be the reason for this finding.

126 There are a number of limitations in the current study. The study was carried out in a single centre and  
127 has a relatively small numbers of patients. The centre has a special interest in SWE so whether the  
128 results are generalizable is open to question. However the results are consistent with the two other  
129 study examining the role of SWE in predicting invasion after an US guided core biopsy result of DCIS  
130 (5,8).

131 In conclusion, we have found that stiffness on SWE and the referral source of the patient are predictors  
132 of occult invasion in women with an US guided core biopsy diagnosis of DCIS and that the combination  
133 of these factors allows stratification of invasion risk which might be used to guide future patient  
134 management.

135

136 **References**

137

138 1. O'Flynn EA, Currie RJ, Mohammed K, Allen SD, Michell MJ. Pre-operative factors indicating risk of  
139 multiple operations versus a single operation in women undergoing surgery for screen detected breast  
140 cancer. *Breast*. 2013 Feb;22(1):78-82.

141 2. Lee SK, Yang JH, Woo SY, Lee JE, Nam SJ. Nomogram for predicting invasion in patients with a  
142 preoperative diagnosis of ductal carcinoma in situ of the breast. *Br J Surg*. 2013 Dec;100(13):1756-63.

143 3. Kim J, Han W, Lee JW, You JM, Shin HC, Ahn SK, Moon HG, Cho N, Moon WK, Park IA, Noh DY. Factors  
144 associated with upstaging from ductal carcinoma in situ following core needle biopsy to invasive cancer  
145 in subsequent surgical excision. *Breast*. 2012 Oct;21(5):641-5.

146 4. Brennan ME, Turner RM, Ciatto S, Marinovich ML, French JR, Macaskill P, Houssami N. Ductal  
147 carcinoma in situ at core-needle biopsy: meta-analysis of underestimation and predictors of invasive  
148 breast cancer. *Radiology*. 2011 Jul;260(1):119-28

149 5. Park AY, Son EJ, Kim JA, Han K, Youk JH. Lesion stiffness measured by shear-wave elastography:  
150 Preoperative predictor of the histologic underestimation of US-guided core needle breast biopsy. *Eur J*  
151 *Radiol*. 2015 Dec;84:2509-14.

152 6. Park SH, Kim MJ, Kim SJ, Kim EK. Ductal carcinoma in situ diagnosed using an ultrasound-guided 14-  
153 gauge core needle biopsy of breast masses: can underestimation be predicted preoperatively?  
154 *Ultrasonography*. 2014 Apr;33(2):128-35.

155 7. Park AY, Gweon HM, Son EJ, Yoo M, Kim JA, Youk JH. Ductal carcinoma in situ diagnosed at US-guided  
156 14-gauge core-needle biopsy for breast mass: preoperative predictors of invasive breast cancer. *Eur J*  
157 *Radiol*. 2014 Apr;83(4):654-9.

- 158 8. Bae JS, Chang JM<sup>2</sup>, Lee SH, Shin SU, Moon WK. Prediction of invasive breast cancer using shear-wave  
159 elastography in patients with biopsy-confirmed ductal carcinoma in situ. *Eur Radiol*. 2016 Apr 16. [Epub  
160 ahead of print]
- 161 9. Vinnicombe SJ, Whelehan P, Thomson K et al. What are the characteristics of breast cancers  
162 misclassified as benign by quantitative ultrasound shear wave elastography? *Eur Radiol*. 2014  
163 Apr;24(4):921-6.
- 164
- 165 10. Berg WA, Mendelson EB, Cosgrove DO, Doré CJ, Gay J, Henry JP, Cohen-Bacrie C. Quantitative  
166 Maximum Shear-Wave Stiffness of Breast Masses as a Predictor of Histopathologic Severity. *AJR Am J*  
167 *Roentgenol*. 2015 Aug;205(2):448-55
- 168 11. National Research Ethics Service (2008) Approval for medical devices research: guidance for  
169 researchers, manufacturers, research ethics committees and NHS R&D offices (Version 2). London:  
170 *National Patient Safety Agency*.
- 171 12. Evans A, Whelehan P, Thomson K et al Differentiating benign from malignant solid breast masses:  
172 value of shear wave elastography according to lesion stiff-ness combined with gray scale ultrasound  
173 according to BI-RADS classification. *Br J Cancer* 2012 107:224–229
- 174 13. Evans A, Whelehan P, Thomson K, et al. Quantitative shear wave ultrasound elastography: initial  
175 experience in solid breast masses. *Breast Cancer Research* 2010; 12:R1
- 176 14. Osuala KO, Sameni M, Shah S, Aggarwal N, Simonait ML, Franco OE, Hong Y, Hayward SW, Behbod F,  
177 Mattingly RR, Sloane BF. Il-6 signaling between ductal carcinoma in situ cells and carcinoma-associated  
178 fibroblasts mediates tumor cell growth and migration. *BMC Cancer*. 2015 Aug 13;15:584. doi:  
179 10.1186/s12885-015-1576-3.

- 180 15. Martins D1, Beça FF, Sousa B, Baltazar F, Paredes J, Schmitt F. Loss of caveolin-1 and gain of MCT4  
181 expression in the tumor stroma: key events in the progression from an in situ to an invasive breast  
182 carcinoma. *Cell Cycle*. 2013 Aug 15;12(16):2684-90.
- 183 16. Vargas AC, McCart Reed AE, Waddell N, Lane A, Reid LE, Smart CE, Cocciardi S, da Silva L, Song S,  
184 Chenevix-Trench G, Simpson PT, Lakhani SR. Gene expression profiling of tumour epithelial and stromal  
185 compartments during breast cancer progression. *Breast Cancer Res Treat*. 2012 Aug;135(1):153-65
- 186 17. Evans A, Rauchhaus P, Whelehan P, Thomson K, Purdie CA, Jordan LB, Michie CO, Thompson AM,  
187 Vinnicombe S. Does shear wave ultrasound independently predict axillary lymph node metastasis in  
188 women with invasive breast cancer? *Breast cancer research and treatment*. 2014;143:153-7

189

190