



University of Dundee

Multiscale modelling and analysis of signalling processes in tissues with non-periodic distribution of cells

Ptashnyk, Mariya

Published in:
Vietnam Journal of Mathematics

DOI:
[10.1007/s10013-016-0232-9](https://doi.org/10.1007/s10013-016-0232-9)

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Ptashnyk, M. (2016). Multiscale modelling and analysis of signalling processes in tissues with non-periodic distribution of cells. *Vietnam Journal of Mathematics*, 45(1), 295-316. DOI: 10.1007/s10013-016-0232-9

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Multiscale Modelling and Analysis of Signalling Processes in Tissues with Non-Periodic Distribution of Cells

Mariya Ptashnyk¹

Received: 7 July 2015 / Accepted: 2 September 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract In this paper, a microscopic model for a signalling process in the left ventricular wall of the heart, comprising a non-periodic fibrous microstructure, is considered. To derive the macroscopic equations, the non-periodic microstructure is approximated by the corresponding locally periodic microstructure. Then, applying the methods of locally periodic homogenization (the locally periodic (l-p) unfolding operator, locally periodic two-scale (l-t-s) convergence on oscillating surfaces and l-p boundary unfolding operator), we obtain the macroscopic model for a signalling process in the heart tissue.

Keywords Non-periodic microstructures · Plywood-like microstructures · Signalling processes · Domains with non-periodic perforations · Locally periodic homogenization · Unfolding operator

Mathematics Subject Classification (2010) 35Bxx · 35D30 · 35Kxx

1 Introduction

In this paper, we consider the multiscale analysis of microscopic problems posed in domains with non-periodic microstructures. We consider a model for a signalling process in the cardiac muscle tissue of the left ventricular wall, comprising plywood-like microstructure [25, 28]. The plywood-like structure is given by the superposition of planes of parallel aligned fibres, gradually rotated with a rotation angle γ , see Fig. 1. In the left ventricular wall, the orientation of the layers of muscle fibres changes from a negative angle at the epicardium

This article is dedicated to Professor Willi Jäger on his 75th birthday.

✉ Mariya Ptashnyk
mptashnyk@maths.dundee.ac.uk; m.ptashnyk@dundee.ac.uk

¹ Department of Mathematics, University of Dundee, Scotland, UK




Fig. 1 *Left*: Schematic representation of a plywood-like structure. *Right*: Cardiac muscle fiber orientations vary continuously through the left ventricular wall from a negative angle at the epicardium to positive values toward the endocardium. Republished from A.D. McCulloch, Cardiac biomechanics, in The Biomedical Engineering Handbook, 2nd edn., J.D. Bronzino (ed.), CRC Press, Boca Raton, FL, 2000 [25]

to a positive angle at the endocardium. In the microscopic model of a signalling process, we consider the diffusion of signalling molecules in the extracellular space and their interaction with receptors located on the surfaces of muscle cells. There are two main challenges in the multiscale analysis of microscopic problems posed in domains with non-periodic perforations: (i) the approximation of the non-periodic microstructure by a locally periodic one and (ii) derivation of limit equations for the non-linear equations defined on oscillating surfaces of the microstructure. First, assuming the C^2 -regularity for the rotation angle γ , we define the locally periodic microstructure which approximates the original non-periodic plywood-like structure. Similar approximation of non-periodic plywood-like microstructure by locally periodic one was considered in [7, 29]. Then, applying techniques of locally periodic homogenization (locally periodic two-scale convergence (l-t-s) and l-p unfolding operator), we derive macroscopic equations for the original microscopic model. The l-p two-scale convergence on oscillating surfaces and l-p boundary unfolding operator allow us to pass to the limit in the non-linear equations defined on surfaces of the locally periodic microstructure.

In this paper, we consider a simple model describing the interactions between processes defined in the perforated domain and on the surfaces of the microstructure. However, the techniques presented here can be also applied to more general microscopic models as well as to other non-periodic microstructures, provided the variations in the microscopic structure are sufficiently regular.

Previous results on homogenization in locally periodic media constitute the multiscale analysis of a heat-conductivity problem defined in domains with non-periodically distributed spherical balls [3, 8, 31], and elliptic and Stokes equations in non-periodic fibrous materials [4, 6, 7, 29]. Formal asymptotic expansion and two-scale convergence defined for periodic test functions, [27], were used to derive macroscopic equations for models posed in domains with locally periodic perforations, i.e., domains consisting of periodic cells with smoothly changing perforations [5, 10, 11, 22, 23, 33].

The paper is organized as follows. In Section 2, the microscopic model for a signalling process in a tissue with non-periodic plywood-like microstructure is formulated. In Section 3, we prove the existence and uniqueness results for the microscopic model and derive a priori estimates for a solution of the microscopic model. The approximation of the microscopic equations posed in the domain with non-periodic microstructure by the corresponding problem defined in a domain with locally periodic microstructure is given in Section 4. Then, applying the l-p unfolding operator, l-t-s convergence on oscillating surfaces, and l-p boundary unfolding operator we derive the macroscopic model for a signalling process in the heart muscle tissue. In Appendix, we summarize the definitions and main compactness results for the l-t-s convergence and l-p unfolding operator.

2 Microscopic Model for a Signalling Process in Heart Tissue

In this work, we consider a receptor-based microscopic model for a cellular signalling process in cardiac tissue. A signalling system is important for proper function of cells and appropriate respond to changes in the extracellular environment. Many regulatory events in cardiac tissue are mediated via surface receptors, located in cardiac cells membrane, that transmit signals through the activation of GTP binding proteins (G proteins) [34]. Cardiac cells (myocytes) in the heart wall tissue are joined in a linear arrangement to form muscle fibres. In the left ventricular wall of the heart the orientation of layers of muscle fibre changes with position through the wall. The layers of parallel aligned muscle fibres are rotated from -60° at epicardium to $+70^\circ$ at endocardium [25] and create a non-periodic plywood-like microstructure. For simplicity, we assume that the individual muscle fibres are not connected to each other. However, it is possible to consider a periodic distribution of connections between the fibres.

In the mathematical model for a signalling process in cardiac tissue, we consider the binding of signalling molecules to receptors located on the cell membrane, which through the activation of G proteins (not considered in our simple model) results in the activation of a cell signalling pathway. We consider the diffusion, production and decay of ligands (signalling molecules) c and binding of ligands to the membrane receptors. We shall distinguish between free receptors r_f and bound receptors r_b , which correspond to receptor-ligand complexes. We assume that the receptor-ligand complex can dissociate and result in a free receptor and ligand. We also consider the production of new free receptors and natural decay of free and bound receptors.

$$\begin{aligned}
 \partial_t c - \nabla \cdot (A(x)\nabla c) &= F(x, c) && \text{in the extracellular space,} \\
 A(x)\nabla c \cdot n &= -\alpha(x)c r_f + \beta(x)r_b && \text{on the cell membrane,} \\
 \partial_t r_f &= p(x, r_b) - \alpha(x)c r_f + \beta(x)r_b - d_f(x)r_f && \text{on the cell membrane,} \\
 \partial_t r_b &= \alpha(x)c r_f - \beta(x)r_b - d_b(x)r_b && \text{on the cell membrane,}
 \end{aligned}$$

where d_f and d_b are the decay rates, β denotes the dissociation rate for the receptor-ligand complex, α is the binding rate, the function p models the production of free receptors, and the function F describes the production and decay of ligands.

To define the plywood-like microstructure of the cardiac muscle tissue of the left ventricular wall, we consider a function $\gamma \in C^2(\mathbb{R})$, with $-\pi/2 \leq \gamma(x) \leq \pi/2$ for $x \in \mathbb{R}$ and define the rotation matrix around the x_3 -axis as

$$R(\gamma(x)) = \begin{pmatrix} \cos(\gamma(x)) & -\sin(\gamma(x)) & 0 \\ \sin(\gamma(x)) & \cos(\gamma(x)) & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

where $\gamma(x)$ denotes the rotation angle with the x_1 -axis. Denote $R_x := R(\gamma(x_3))$.

We consider an open, bounded subdomain $\Omega \subset \mathbb{R}^3$, with Lipschitz boundary, representing a part of the cardiac muscle tissue and the x_3 -axis to be orthogonal to the layers of parallel-aligned muscle fibres. We assume that the radius of the muscle fibres depends on the position in the tissue and define the characteristic function of a fibre by

$$\vartheta(x, y) = \begin{cases} 1, & |\hat{y}| \leq \rho(\hat{x}_R)a, \\ 0, & |\hat{y}| > \rho(\hat{x}_R)a, \end{cases}$$

where $\hat{y} = (y_2, y_3)$ and $\hat{x}_R = ((R_x^{-1}x)_2, (R_x^{-1}x)_3)$, $\rho \in C^1(\mathbb{R}^2)$, with $0 < \rho_0 \leq \rho(\hat{x}_R) \leq \rho_1 < \infty$ and $\rho(\hat{x}_R)a \leq 2/5$ for all $x \in \Omega$, i.e., $a = 2/(5\rho_1)$.

By small parameter ε we denote the characteristic size of the microstructure of cardiac tissue, given as a ratio between the characteristic diameter of muscle fibres and characteristic size of the cardiac tissue. Notice that for plywood-like microstructure the axis of each fibre can be defined by a rotated around the x_3 -axis line, parallel to the x_1 -axis and passing through a point of an ε -grid in the plane $x_1 = \text{const}$. Thus, for $j \in \mathbb{Z}^3$, we define $x_j^\varepsilon = R_{x_j^\varepsilon} \varepsilon j$ with $R_{x_j^\varepsilon} := R(\gamma(x_{j,3}^\varepsilon))$. Notice that $x_{j,3}^\varepsilon = \varepsilon j_3$ and the third variable is invariant under the rotation $R_{x_j^\varepsilon}$. This ensures that for each fixed εj_3 we obtain a layer of parallel aligned fibres.

Then the perforated domain Ω_ε^* , corresponding to the extracellular space of cardiac tissue, is defined as

$$\Omega_\varepsilon^* = \Omega \setminus \Omega_\varepsilon^0, \quad \text{with } \Omega_\varepsilon^0 = \bigcup_{j \in \Xi_\varepsilon} (\varepsilon R_{x_j^\varepsilon} K_{x_j^\varepsilon} Y_0 + x_j^\varepsilon) = \bigcup_{j \in \Xi_\varepsilon} \varepsilon R_{x_j^\varepsilon} (K_{x_j^\varepsilon} Y_0 + j),$$

where $\Xi_\varepsilon = \{j \in \mathbb{Z}^3 : \varepsilon R_{x_j^\varepsilon}(\bar{Y}_1 + j) \subset \Omega\}$, $Y_0 = \{y \in \bar{Y}_1 : |\hat{y}| \leq a\}$, with $a = 2/(5\rho_1)$ and $\rho_1 = \sup_{\bar{\Omega}} \rho(x)$, and

$$Y_1 = \left(-\frac{1}{2}, \frac{1}{2}\right)^3, \quad K_x = K(x), \quad K(x) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \rho(\hat{x}_R) & 0 \\ 0 & 0 & \rho(\hat{x}_R) \end{pmatrix}.$$

We also define $Y_{x_j^\varepsilon} = R_{x_j^\varepsilon} Y_1$, and $Y_{x_j^\varepsilon, K}^* = R_{x_j^\varepsilon} (Y_1 \setminus K_{x_j^\varepsilon} Y_0)$ for $j \in \Xi_\varepsilon$.

We denote $\Gamma_{0,1} = \{y \in \mathbb{R}^3 : |y_1| = \pm 1/2\}$. Then, assumptions on ρ and a ensure that $K_x(Y_0 \setminus \Gamma_{0,1}) \subset Y_1$ for all $x \in \bar{\Omega}$ and, since R is a rotation matrix, $(\varepsilon R_{x_n^\varepsilon} K_{x_n^\varepsilon} Y_0 + x_n^\varepsilon) \cap (\varepsilon R_{x_m^\varepsilon} K_{x_m^\varepsilon} Y_0 + x_m^\varepsilon) = \emptyset$ for any $m, n \in \Xi_\varepsilon$ with $n_2 \neq m_2$ or $n_3 \neq m_3$. Hence, Ω_ε^* is connected. This corresponds to our assumption that muscle fibres do not touch each other and are not directly connected, and the interactions between the muscle fibres are facilitated through the extracellular matrix.

Now, using the definition of ϑ , the characteristic function of muscle fibres in cardiac tissue reads

$$\chi_{\Omega_j^\varepsilon}(x) = \chi_\Omega(x) \sum_{j \in \Xi_\varepsilon} \vartheta \left(x_j^\varepsilon, R_{x_j^\varepsilon}^{-1}(x - x_j^\varepsilon) / \varepsilon \right)$$

and the extracellular space is characterised by

$$\chi_{\Omega_\varepsilon^*} = (1 - \chi_{\Omega_j^\varepsilon}) \chi_\Omega.$$

The surfaces of muscle cells, i.e., the boundaries of the microstructure, are denoted by

$$\Gamma^\varepsilon = \sum_{j \in \Xi_\varepsilon} \left(\varepsilon R_{x_j^\varepsilon} K_{x_j^\varepsilon} \Gamma + x_j^\varepsilon \right) = \sum_{j \in \Xi_\varepsilon} \varepsilon R_{x_j^\varepsilon} (K_{x_j^\varepsilon} \Gamma + j),$$

where $\Gamma = \overline{\partial Y_0 \setminus \Gamma_{0,1}}$.

Notice that the changes in the microstructure of Ω_ε^* are defined by changes in the periodicity given by the linear transformation (rotation) $R(x)$ and by changes in the shape of the microstructure (changes in the radius of muscle fibres) given by the linear transformation $K(x)$ for $x \in \Omega$.

To determine the non-constant reaction rates for binding and dissociation processes on cell membranes, we consider $\alpha, \beta \in C^1(\bar{\Omega}; C_0^1(Y_1))$, extended in y -variable by zero to \mathbb{R}^3 , and define

$$\alpha^\varepsilon(x) = \sum_{j \in \Xi_\varepsilon} \alpha(x, R_{x_j^\varepsilon}^{-1}(x - x_j^\varepsilon)/\varepsilon) \chi_{(\varepsilon Y_{x_j^\varepsilon} + x_j^\varepsilon)}(x),$$

$$\beta^\varepsilon(x) = \sum_{j \in \Xi_\varepsilon} \beta(x, R_{x_j^\varepsilon}^{-1}(x - x_j^\varepsilon)/\varepsilon) \chi_{(\varepsilon Y_{x_j^\varepsilon} + x_j^\varepsilon)}(x).$$

Then, the microscopic model for a signalling process in cardiac tissue reads

$$\begin{aligned} \partial_t c^\varepsilon - \operatorname{div}(A \nabla c^\varepsilon) &= F(c^\varepsilon) && \text{in } (0, T) \times \Omega_\varepsilon^*, \\ -A \nabla c^\varepsilon \cdot \mathbf{n} &= \varepsilon \left[\alpha^\varepsilon(x) c^\varepsilon r_f^\varepsilon - \beta^\varepsilon(x) r_b^\varepsilon \right] && \text{on } (0, T) \times \Gamma^\varepsilon, \\ A \nabla c^\varepsilon \cdot \mathbf{n} &= 0 && \text{on } (0, T) \times (\partial \Omega_\varepsilon^* \setminus \Gamma^\varepsilon), \\ c^\varepsilon(0, x) &= c_0(x) && \text{in } \Omega_\varepsilon^*, \end{aligned} \tag{1}$$

where the dynamics in the concentrations of free and bound receptors on cell surfaces is determined by two ordinary differential equations

$$\begin{aligned} \partial_t r_f^\varepsilon &= p(r_b^\varepsilon) - \alpha^\varepsilon(x) c^\varepsilon r_f^\varepsilon + \beta^\varepsilon(x) r_b^\varepsilon - d_f r_f^\varepsilon && \text{on } (0, T) \times \Gamma^\varepsilon, \\ \partial_t r_b^\varepsilon &= \alpha^\varepsilon(x) c^\varepsilon r_f^\varepsilon - \beta^\varepsilon(x) r_b^\varepsilon - d_b r_b^\varepsilon && \text{on } (0, T) \times \Gamma^\varepsilon, \\ r_f^\varepsilon(0, x) &= r_{f0}^\varepsilon(x), && r_b^\varepsilon(0, x) = r_{b0}^\varepsilon(x) && \text{on } \Gamma^\varepsilon, \end{aligned} \tag{2}$$

with initial conditions defined as

$$r_{l0}^\varepsilon(x) = r_{l0}^1(x) \sum_{j \in \Xi_\varepsilon} r_{l0}^2(R_{x_j^\varepsilon}^{-1}(x - x_j^\varepsilon)/\varepsilon) \chi_{(\varepsilon Y_{x_j^\varepsilon} + x_j^\varepsilon)}(x) \quad \text{for } l = f, b.$$

For simplicity of the presentation we shall assume that the diffusion coefficient A and the decay rates d_f, d_b are constant. We also assume that the functions F and p are independent of $x \in \Omega$. The dependence of A, d_f, d_b, F and p on the microscopic and macroscopic variables can be analyzed in the similar way as for α^ε and β^ε .

Assumption 1

- A is symmetric and $(A\xi, \xi) \geq a_0|\xi|^2$ for $\xi \in \mathbb{R}^3, a_0 > 0$.
- $\gamma \in C^2(\mathbb{R}), K \in C^1(\bar{\Omega})$ with $0 < \rho_0^2 \leq |\det K(x)| \leq \rho_1^2 < \infty$ and $K(x)(Y_0 \setminus \Gamma_{0,1}) \subset Y_1$ for all $x \in \bar{\Omega}$, and $d_l \geq 0$ for $l = f, b$.
- $F : \mathbb{R} \rightarrow \mathbb{R}$ is Lipschitz continuous, $F(\xi_-)\xi_- \leq \mu_F|\xi_-|^2$, with $\mu_F > 0$ and $\xi_- = \min\{0, \xi\}$.
- $p : \mathbb{R} \rightarrow \mathbb{R}$ is Lipschitz continuous and $p(\xi) \geq 0$ for $\xi \geq 0$.
- $\alpha, \beta \in C^1(\bar{\Omega}; C_0^1(Y_1))$ are nonnegative.
- $c_0 \in H^1(\Omega) \cap L^\infty(\Omega), r_{l0}^1 \in C^1(\bar{\Omega})$, and $r_{l0}^2 \in C_0^1(Y_1)$, extended by zero to \mathbb{R}^3 , and c_0, r_{l0}^j are nonnegative for $l = f, b$ and $j = 1, 2$.

Notice that the C^1 -regularity of α and β is required for the approximation of the integrals defined on the boundaries of the non-periodic microstructure by the integrals defined on the boundaries of the corresponding locally periodic microstructure.

We shall use the following notations $\Omega_{\varepsilon, T}^* = (0, T) \times \Omega_\varepsilon^*, \Gamma_T^\varepsilon = (0, T) \times \Gamma^\varepsilon, \Omega_T = (0, T) \times \Omega, \Gamma_T = (0, T) \times \Gamma$, and $\Gamma_{x, T} = (0, T) \times \Gamma_x$.

For $u \in L^q(0, \tau; L^p(G))$ and $v \in L^{q'}(0, \tau; L^{p'}(G))$, with $1/p + 1/p' = 1$ and $1/q + 1/q' = 1, \tau > 0$ and $G \subset \mathbb{R}^d$ for $d = 2$ or 3 , we denote $\langle u, v \rangle_{G_\tau} = \int_0^\tau \int_G uv dx dt$.

We shall consider a weak solution of the problem (1) and (2), defined in the following way.

Definition 1 A weak solution of the microscopic problem (1) and (2) are functions $c^\varepsilon, r_f^\varepsilon, r_b^\varepsilon$ such that

$$c^\varepsilon \in L^2(0, T; H^1(\Omega_\varepsilon^*)) \cap H^1(0, T; L^2(\Omega_\varepsilon^*)),$$

$$r_l^\varepsilon \in H^1(0, T; L^2(\Gamma^\varepsilon)), \quad r_l^\varepsilon \in L^\infty(0, T; L^\infty(\Gamma^\varepsilon)), \quad l = f, b,$$

satisfying (1) in the weak form

$$\langle \partial_t c^\varepsilon, \phi \rangle_{\Omega_\varepsilon^*, T} + \langle A \nabla c^\varepsilon, \nabla \phi \rangle_{\Omega_\varepsilon^*, T} = \langle F(c^\varepsilon), \phi \rangle_{\Omega_\varepsilon^*, T} + \varepsilon \langle \beta^\varepsilon r_b^\varepsilon - \alpha^\varepsilon c^\varepsilon r_f^\varepsilon, \phi \rangle_{\Gamma_T^\varepsilon} \quad (3)$$

for all $\phi \in L^2(0, T; H^1(\Omega_\varepsilon^*))$, (2) are satisfied a.e. on Γ_T^ε , and $c^\varepsilon \rightarrow c_0$ in $L^2(\Omega_\varepsilon^*)$, $r_l^\varepsilon \rightarrow r_{l0}^\varepsilon$ in $L^2(\Gamma^\varepsilon)$ for $l = f, b$, as $t \rightarrow 0$.

3 Existence, Uniqueness, and a Priori Estimates for a Weak Solution of the Microscopic Problem (1) and (2)

In a similar way as in [9, 21, 30], we can prove the existence, uniqueness, and a priori estimates for a weak solution of problem (1)–(2). Notice that for the derivation of a priori estimates a trace estimate, uniform in ε , for functions $\phi \in W^{1,p}(\Omega_\varepsilon^*)$ is required. The fact that $K_x(Y_0 \setminus \Gamma_{0,1}) \subset Y_1$ for all $x \in \bar{\Omega}$ and the uniform boundedness of $\det K$, i.e., $0 < \rho_0^2 \leq |\det K(x)| \leq \rho_1^2 < \infty$, ensure the trace estimate for $\phi \in H^1(Y_1 \setminus K_{x_j^\varepsilon} Y_0)$, i.e.,

$$\|\phi\|_{L^p(K_{x_j^\varepsilon} \Gamma)}^p \leq C \left[\|\phi\|_{L^p(Y_1 \setminus K_{x_j^\varepsilon} Y_0)}^p + \|\nabla_y \phi\|_{L^p(Y_1 \setminus K_{x_j^\varepsilon} Y_0)}^p \right],$$

where the constant C depends on Y_1, Y_0, K and is independent of ε and $j \in \Xi_\varepsilon$. Then, considering the change of variables $x = \varepsilon R_{x_j^\varepsilon} y + x_j^\varepsilon = \varepsilon R_{x_j^\varepsilon} (y + j)$ and summing up over $j \in \Xi_\varepsilon$, we obtain for $\phi \in W^{1,p}(\Omega_\varepsilon^*)$, with $p \in [1, \infty)$, that

$$\varepsilon \|\phi\|_{L^p(\Gamma^\varepsilon)}^p \leq \tilde{\mu} \left[\|\phi\|_{L^p(\Omega_\varepsilon^*)}^p + \varepsilon^p \|\nabla \phi\|_{L^p(\Omega_\varepsilon^*)}^p \right], \quad (4)$$

where the constant $\tilde{\mu}$ depends on Y_1, Y_0, R and K and is independent of ε .

Lemma 1 Under Assumption 1 there exists a unique non-negative weak solution of the microscopic problem (1) and (2) satisfying the following a priori estimates

$$\|c^\varepsilon\|_{L^\infty(0, T; L^2(\Omega_\varepsilon^*))} + \|\nabla c^\varepsilon\|_{L^2(\Omega_\varepsilon^*, T)} + \|\partial_t c^\varepsilon\|_{L^2(\Omega_\varepsilon^*, T)} + \varepsilon^{\frac{1}{2}} \|c^\varepsilon\|_{L^2(\Gamma_T^\varepsilon)} \leq \mu, \quad (5a)$$

$$\|r_f^\varepsilon\|_{L^\infty(\Gamma_T^\varepsilon)} + \|r_b^\varepsilon\|_{L^\infty(\Gamma_T^\varepsilon)} + \varepsilon^{\frac{1}{2}} \|\partial_t r_f^\varepsilon\|_{L^2(\Gamma_T^\varepsilon)} + \varepsilon^{\frac{1}{2}} \|\partial_t r_b^\varepsilon\|_{L^2(\Gamma_T^\varepsilon)} \leq \mu, \quad (5b)$$

and

$$\|(c^\varepsilon - M_1 e^{M_2 t})^+\|_{L^\infty(0, T; L^2(\Omega_\varepsilon^*))} + \|\nabla (c^\varepsilon - M_1 e^{M_2 t})^+\|_{L^2(\Omega_\varepsilon^*, T)} \leq \mu \varepsilon, \quad (6)$$

where the constant μ is independent of ε , $M_1 \geq \|c_0\|_{L^\infty(\Omega)}$, and $M_1 M_2 \geq |F(0)| + \|F'\|_{L^\infty} M_1 + \tilde{\mu} \|\beta\|_{L^\infty(\Omega \times Y_1)} \|r_b^\varepsilon\|_{L^\infty(\Gamma_T^\varepsilon)}$, with $\tilde{\mu}$ being the constant in the trace inequality (4).

Proof (Sketch) As in [30] the existence of a solution of the microscopic problem (1) and (2) for each fixed $\varepsilon > 0$ is obtained by applying fixed point arguments and Galerkin method.

Also, using the same arguments as in [30] we obtain that $c^\varepsilon(t, x) \geq 0$ for $(t, x) \in \Omega_{\varepsilon, T}^*$ and $r_l^\varepsilon(t, x) \geq 0$ for $(t, x) \in \Gamma_T^\varepsilon$, with $l = f, b$. To derive a priori estimates, we consider the structure of the microscopic equations. For non-negative solutions, by adding the equations for r_f^ε and r_b^ε , we obtain

$$\partial_t(r_f^\varepsilon + r_b^\varepsilon) = p(r_b^\varepsilon) - d_b r_b^\varepsilon - d_f r_f^\varepsilon.$$

Then the Lipschitz continuity of p and the non-negativity of r_f^ε and r_b^ε imply the boundedness of r_f^ε and r_b^ε on Γ_T^ε .

Considering c^ε as a test function in (3) and using the trace inequality (4), we obtain the estimates for c^ε . Testing (2) by $\partial_t r_f^\varepsilon$ and $\partial_t r_b^\varepsilon$, respectively, yields the estimates for the time derivatives of r_f^ε and r_b^ε . In the derivation of the a priori estimate for $\partial_t c^\varepsilon$ we use the equation for $\partial_t r_f^\varepsilon$ to estimate the non-linear term on the boundary Γ^ε , i.e.,

$$\begin{aligned} - \int_{\Gamma^\varepsilon} \alpha^\varepsilon(x) r_f^\varepsilon c^\varepsilon \partial_t c^\varepsilon d\sigma_x &= - \frac{1}{2} \frac{d}{dt} \int_{\Gamma^\varepsilon} \alpha^\varepsilon(x) r_f^\varepsilon |c^\varepsilon|^2 d\sigma_x \\ &\quad + \frac{1}{2} \int_{\Gamma^\varepsilon} \alpha^\varepsilon(x) (p(r_b^\varepsilon) - \alpha^\varepsilon(x) r_f^\varepsilon c^\varepsilon + \beta^\varepsilon(x) r_b^\varepsilon - d_f r_f^\varepsilon) |c^\varepsilon|^2 d\sigma_x \\ &\leq \frac{1}{2} \int_{\Gamma^\varepsilon} (\alpha^\varepsilon(x) p(r_b^\varepsilon) + \beta^\varepsilon(x) r_b^\varepsilon) |c^\varepsilon|^2 d\sigma_x - \frac{1}{2} \frac{d}{dt} \int_{\Gamma^\varepsilon} \alpha^\varepsilon(x) r_f^\varepsilon |c^\varepsilon|^2 d\sigma_x. \end{aligned}$$

Considering $(c^\varepsilon - M_1 e^{M_2 t})^+$ as a test function in (3), where M_1 and M_2 are as in the formulation of the lemma, we obtain

$$\begin{aligned} &\int_{\Omega_\varepsilon^*} |(c^\varepsilon(\tau) - M_1 e^{M_2 \tau})^+|^2 dx + \int_0^\tau \int_{\Omega_\varepsilon^*} M_1 M_2 e^{M_2 t} (c^\varepsilon - M_1 e^{M_2 t})^+ dx dt \\ &+ \int_0^\tau \left[\int_{\Omega_\varepsilon^*} |\nabla(c^\varepsilon - M_1 e^{M_2 t})^+|^2 dx + \varepsilon \int_{\Gamma^\varepsilon} \alpha^\varepsilon(x) r_f^\varepsilon c^\varepsilon (c^\varepsilon - M_1 e^{M_2 t})^+ d\sigma_x \right] dt \\ &\leq C_1 \int_0^\tau \left[\int_{\Omega_\varepsilon^*} F(c^\varepsilon) (c^\varepsilon - M_1 e^{M_2 t})^+ dx + \varepsilon \int_{\Gamma^\varepsilon} \beta^\varepsilon(x) r_b^\varepsilon (c^\varepsilon - M_1 e^{M_2 t})^+ d\sigma_x \right] dt \end{aligned}$$

for $\tau \in (0, T]$. Using the non-negativity and boundedness of β^ε and r_f^ε , along with the trace inequality (4), the last integral can be estimated as

$$\begin{aligned} &\varepsilon \int_0^\tau \int_{\Gamma^\varepsilon} \beta^\varepsilon r_b^\varepsilon (c^\varepsilon - M_1 e^{M_2 t})^+ d\sigma_x dt \\ &\leq \mu_1 \int_0^\tau \int_{\Omega_\varepsilon^*} (c^\varepsilon - M_1 e^{M_2 t})^+ dx dt + \varepsilon \mu_1 \int_0^\tau \int_{\Omega_\varepsilon^*} |\nabla(c^\varepsilon - M_1 e^{M_2 t})^+|^2 dx dt \\ &\leq \mu_1 \int_0^\tau \int_{\Omega_\varepsilon^*} (c^\varepsilon - M_1 e^{M_2 t})^+ dx dt + \mu_2 \delta \int_0^\tau \int_{\Omega_\varepsilon^*} |\nabla(c^\varepsilon - M_1 e^{M_2 t})^+|^2 dx dt + \mu_\delta \varepsilon^2 \end{aligned}$$

for any $\delta > 0$, where the constants μ_1 , μ_2 and μ_δ depend on $\|\beta\|_{L^\infty(\Omega \times Y_1)}$, $\|r_b^\varepsilon\|_{L^\infty(\Gamma_T^\varepsilon)}$ and on the transformation matrices R and K , but are independent of ε . More specifically, $\mu_1 = \tilde{\mu} \|\beta\|_{L^\infty(\Omega \times Y_1)} \|r_b^\varepsilon\|_{L^\infty(\Gamma_T^\varepsilon)}$. Using the non-negativity of c^ε and r_f^ε , the Lipschitz continuity of F , and the assumptions on M_1 and M_2 , and applying the Gronwall inequality yield estimate (6).

To show the uniqueness of a solution of the microscopic problem (1) and (2), we consider the equations for the difference of two solutions $(c_1^\varepsilon, r_{f,1}^\varepsilon, r_{b,1}^\varepsilon)$ and $(c_2^\varepsilon, r_{f,2}^\varepsilon, r_{b,2}^\varepsilon)$. The non-negativity of α^ε , $r_{f,j}^\varepsilon$, and c_j^ε , along with the boundedness of $r_{f,j}^\varepsilon$, ensures

$$\|r_{f,1}^\varepsilon(\tau) - r_{f,2}^\varepsilon(\tau)\|_{L^2(\Gamma^\varepsilon)}^2 \leq \mu \int_0^\tau \left[\sum_{l=f,b} \|r_{l,1}^\varepsilon - r_{l,2}^\varepsilon\|_{L^2(\Gamma^\varepsilon)}^2 + \|c_1^\varepsilon - c_2^\varepsilon\|_{L^2(\Gamma^\varepsilon)}^2 \right] dt.$$

Testing the sum of the equations for $r_{f,1}^\varepsilon - r_{f,2}^\varepsilon$ and $r_{b,1}^\varepsilon - r_{b,2}^\varepsilon$ by $r_{f,1}^\varepsilon + r_{b,1}^\varepsilon - r_{f,2}^\varepsilon - r_{b,2}^\varepsilon$ and using the estimate from above yield

$$\begin{aligned} \|r_{b,1}^\varepsilon(\tau) - r_{b,2}^\varepsilon(\tau)\|_{L^2(\Gamma^\varepsilon)}^2 &\leq \|r_{b,1}^\varepsilon(\tau) + r_{f,1}^\varepsilon(\tau) - r_{b,2}^\varepsilon(\tau) - r_{f,2}^\varepsilon(\tau)\|_{L^2(\Gamma^\varepsilon)}^2 \\ &\quad + \|r_{f,1}^\varepsilon(\tau) - r_{f,2}^\varepsilon(\tau)\|_{L^2(\Gamma^\varepsilon)}^2 \\ &\leq \mu_1 \int_0^\tau \sum_{l=f,b} \|r_{l,1}^\varepsilon - r_{l,2}^\varepsilon\|_{L^2(\Gamma^\varepsilon)}^2 dt + \mu_2 \int_0^\tau \|c_1^\varepsilon - c_2^\varepsilon\|_{L^2(\Gamma^\varepsilon)}^2 dt. \end{aligned}$$

Combining the last two inequalities and applying the Gronwall inequality imply the estimates for $\|r_{l,1}^\varepsilon(\tau) - r_{l,2}^\varepsilon(\tau)\|_{L^2(\Gamma^\varepsilon)}^2$, with $l = f, b$, in terms of $\|c_1^\varepsilon - c_2^\varepsilon\|_{L^2((0,\tau)\times\Gamma^\varepsilon)}^2$ for $\tau \in (0, T]$.

Considering $(c^\varepsilon - S)^+$, with some $S > 0$, as a test function in (3) and using the boundedness of r_f^ε and r_b^ε we obtain

$$\|(c^\varepsilon - S)^+\|_{L^\infty(0,T;L^2(\Omega_\varepsilon^*))} + \|\nabla(c^\varepsilon - S)^+\|_{L^2(\Omega_\varepsilon^*)} \leq \mu_1 S \left(\int_0^T |\Omega_\varepsilon^{*,S}(t)| dt \right)^{\frac{1}{2}},$$

where $S \geq \max\{\|c_0\|_{L^\infty(\Omega)}, \|\beta\|_{L^\infty(\Omega \times Y_1)} \|r_b^\varepsilon\|_{L^\infty(\Gamma_f^\varepsilon)}, |F(0)|\}$, μ_1 is some positive constant, and $\Omega_\varepsilon^{*,S}(t) = \{x \in \Omega_\varepsilon^* : c^\varepsilon(t, x) > S\}$. Then, Theorem II.6.1 in [20] yields the boundedness of c^ε in $(0, T) \times \Omega_\varepsilon^*$ for every fixed ε .

Considering now (3) for two solutions $(c_1^\varepsilon, r_{f,1}^\varepsilon, r_{b,1}^\varepsilon)$ and $(c_2^\varepsilon, r_{f,2}^\varepsilon, r_{b,2}^\varepsilon)$, we obtain the estimates for $\|c_1^\varepsilon - c_2^\varepsilon\|_{L^2(\Omega_\varepsilon^*, \tau)}^2$ and $\varepsilon \|c_1^\varepsilon - c_2^\varepsilon\|_{L^2((0,\tau)\times\Gamma^\varepsilon)}^2$ in terms of $\varepsilon \|r_{l,1}^\varepsilon - r_{l,2}^\varepsilon\|_{L^2((0,\tau)\times\Gamma^\varepsilon)}^2$, with $l = f, b$ and $\tau \in (0, T]$. Using the estimates for $\|r_{l,1}^\varepsilon(\tau) - r_{l,2}^\varepsilon(\tau)\|_{L^2(\Gamma^\varepsilon)}^2$ in terms of $\|c_1^\varepsilon - c_2^\varepsilon\|_{L^2((0,\tau)\times\Gamma^\varepsilon)}^2$, shown above, and applying the Gronwall inequality, we conclude that $r_{l,1}^\varepsilon = r_{l,2}^\varepsilon$ on $(0, T) \times \Gamma^\varepsilon$, with $l = f, b$, and $c_1^\varepsilon = c_2^\varepsilon$ in $(0, T) \times \Omega_\varepsilon^*$. \square

The assumptions on the non-periodic microstructure of Ω_ε^* and the regularity of the transformation matrices R and K ensure the following extension result.

Lemma 2 For $x_j^\varepsilon \in \Omega$, and $u \in W^{1,p}(Y_{x_j^\varepsilon}^*, K)$, with $p \in (1, +\infty)$ and $j \in \Xi_\varepsilon$, there exists an extension $\tilde{u} \in W^{1,p}(Y_{x_j^\varepsilon})$ from $Y_{x_j^\varepsilon}^*, K$ into $Y_{x_j^\varepsilon}$ such that

$$\|\tilde{u}\|_{L^p(Y_{x_j^\varepsilon})} \leq \mu \|u\|_{L^p(Y_{x_j^\varepsilon}^*, K)}, \quad \|\nabla \tilde{u}\|_{L^p(Y_{x_j^\varepsilon})} \leq \mu \|\nabla u\|_{L^p(Y_{x_j^\varepsilon}^*, K)}, \tag{7}$$

where μ depends on Y_1, Y_0, R and K and is independent of ε and $j \in \Xi_\varepsilon$.

For $u \in W^{1,p}(\Omega_\varepsilon^*)$ we have an extension $\tilde{u} \in W^{1,p}(\Omega)$ from Ω_ε^* into Ω such that

$$\|\tilde{u}\|_{L^p(\Omega)} \leq \mu \|u\|_{L^p(\Omega_\varepsilon^*)}, \quad \|\nabla \tilde{u}\|_{L^p(\Omega)} \leq \mu \|\nabla u\|_{L^p(\Omega_\varepsilon^*)}, \tag{8}$$

where μ depends on Y_1, Y_0, R and K and is independent of ε .

Proof (Sketch) The proof follows the same lines as in the periodic case, see e.g. [15, 19]. The only difference here is that the extension depends on the Lipschitz continuity of K and R and the uniform boundedness from above and below of $|\det K(x)|$ and $|\det R(x)|$ for all $x \in \overline{\Omega}$.

To show (8), we first consider the extension from $R_{x_j^\varepsilon}((Y_1 \setminus K_{x_j^\varepsilon} Y_0) + j)$ into $R_{x_j^\varepsilon}(Y_1 + j)$ and obtain the estimates in (7). Notice that due to the definition of $K(x)$, the fibre radius varies between different fibres in the plywood-like structure of the heart tissue, but is constant along each individual fibre. Thus, apart from the end parts of the fibres near $\partial\Omega$, we have to extend u only in the directions orthogonal to the fibres. In the definition of Ω_ε^* we consider those j that $\varepsilon R_{x_j^\varepsilon}(\overline{Y}_1 + j) \subset \Omega$. Hence the extension for the end parts of the fibres near $\partial\Omega$, i.e., for such $R_{x_j^\varepsilon}((Y_1 \setminus K_{x_j^\varepsilon} Y_0) + j)$ that $\bigcup_{m \in \{0,1\}^3} \varepsilon R_{x_j^\varepsilon}(\overline{Y}_1 + j \pm m) \cap \partial\Omega \neq \emptyset$, is also well-defined. Since $\det R(x) = 1$, and $0 < \rho_0^2 \leq |\det K(x)| \leq \rho_1^2 < \infty$ for all $x \in \overline{\Omega}$, we obtain that the constant μ in (7) is independent of x_j^ε , ε , and $j \in \Xi_\varepsilon$. Then scaling $R_{x_j^\varepsilon}((Y_1 \setminus K_{x_j^\varepsilon} Y_0) + j)$ and $R_{x_j^\varepsilon}(Y_1 + j)$ by ε and summing up over $j \in \Xi_\varepsilon$ in (7) imply (8).

In the case when the boundary $\partial\Omega$ crosses the fibres in a non-orthogonal way, we would obtain only a local extension to a subdomain $\Omega^\delta = \{x \in \Omega : \text{dist}(x, \partial\Omega) > \delta\}$ for any fixed $\delta > 0$. □

4 Derivation of Macroscopic Equations

To derive macroscopic equations for the microscopic problem posed in a domain with the non-periodic plywood-like microstructure, we approximate it by a problem defined in the domain with the corresponding locally periodic microstructure and apply the methods of locally periodic two-scale convergence (l-t-s) and l-p unfolding operator (see Appendix for the definitions and convergence results for l-t-s convergence and l-p unfolding operator). Notice that the regularity assumptions on the orientation angle γ are essential for the construction of an appropriate locally periodic microstructure for the non-periodic plywood-like structure.

To define the locally periodic microstructure related to the original non-periodic one, we consider, similarly to [8, 29], the partition covering of Ω by a family of open non-intersecting cubes $\{\Omega_n^\varepsilon\}_{1 \leq n \leq N_\varepsilon}$ of side ε^r , with $0 < r < 1$, such that

$$\Omega \subset \bigcup_{n=1}^{N_\varepsilon} \overline{\Omega}_n^\varepsilon \quad \text{and} \quad \Omega_n^\varepsilon \cap \Omega \neq \emptyset.$$

For each $x \in \mathbb{R}^3$, we consider a transformation matrix $D(x) \in \mathbb{R}^{3 \times 3}$ and assume that $D, D^{-1} \in \text{Lip}(\mathbb{R}^3; \mathbb{R}^{3 \times 3})$ and $0 < d_0 \leq |\det D(x)| \leq d_1 < \infty$ for all $x \in \overline{\Omega}$. The matrix D will be defined by the rotation matrix R and its derivatives and the specific form of D will be given later.

Then, the locally periodic microstructure is defined by considering a covering of Ω_n^ε by parallelepipeds $\varepsilon D_{x_n^\varepsilon} Y$ such that

$$\Omega_n^\varepsilon \subset \tilde{x}_n^\varepsilon + \bigcup_{\xi \in \Xi_n^\varepsilon} \varepsilon D_{x_n^\varepsilon}(\overline{Y} + \xi), \quad \text{where} \quad \Xi_n^\varepsilon = \{\xi \in \mathbb{Z}^3 : \varepsilon D_{x_n^\varepsilon}(Y + \xi) \cap \Omega_n^\varepsilon \neq \emptyset\},$$

and points $x_n^\varepsilon, \tilde{x}_n^\varepsilon \in \Omega_n^\varepsilon$, for $n = 1, \dots, N_\varepsilon$, are arbitrary chosen, but fixed. Here $Y = (0, 1)^3$, $D_x := D(x)$, and $D_{x_n^\varepsilon} = D(x_n^\varepsilon)$ for $1 \leq n \leq N_\varepsilon$. Then, the perforated domain with locally periodic microstructure is given by

$$\tilde{\Omega}_\varepsilon^* = \text{Int} \left(\bigcup_{n=1}^{N_\varepsilon} \Omega_n^{*,\varepsilon} \right) \cap \Omega, \quad \text{with } \Omega_n^{*,\varepsilon} = \left(\tilde{x}_n^\varepsilon + \bigcup_{\xi \in \Xi_n^\varepsilon} \varepsilon D_{x_n^\varepsilon} (\bar{Y}_{K_{x_n^\varepsilon}}^* + \xi) \right) \cap \bar{\Omega}_n^\varepsilon,$$

where $Y_{K_{x_n^\varepsilon}}^* = Y \setminus \bigcup_{j \in \{0,1\}^3} (\tilde{K}_{x_n^\varepsilon} Y_0 + j)$, with $\tilde{K}_{x_n^\varepsilon} = \tilde{K}(x_n^\varepsilon)$ for $n = 1, \dots, N_\varepsilon$, and the transformation matrix \tilde{K} will be specified later. We shall also denote

$$\hat{\Omega}_n^\varepsilon = \tilde{x}_n^\varepsilon + \text{Int} \left(\bigcup_{\xi \in \hat{\Xi}_n^\varepsilon} \varepsilon D_{x_n^\varepsilon} (\bar{Y} + \xi) \right) \quad \text{and} \quad \Lambda_\varepsilon^* = \tilde{\Omega}_\varepsilon^* \setminus \bigcup_{n=1}^{N_\varepsilon} \hat{\Omega}_n^\varepsilon,$$

where $\hat{\Xi}_n^\varepsilon = \{\xi \in \Xi_n^\varepsilon : \varepsilon D_{x_n^\varepsilon}(Y + \xi) \subset (\Omega_n^\varepsilon \cap \Omega)\}$. The boundaries of the locally periodic microstructure are defined as

$$\tilde{\Gamma}^\varepsilon = \bigcup_{n=1}^{N_\varepsilon} \Gamma_n^\varepsilon \cap \Omega, \quad \text{where } \Gamma_n^\varepsilon = \left(\tilde{x}_n^\varepsilon + \bigcup_{\xi \in \Xi_n^\varepsilon} \varepsilon D_{x_n^\varepsilon} (\tilde{\Gamma}_{x_n^\varepsilon, K} + \xi) \right) \cap \Omega_n^\varepsilon,$$

and

$$\hat{\Gamma}^\varepsilon = \bigcup_{n=1}^{N_\varepsilon} \left(\tilde{x}_n^\varepsilon + \bigcup_{\xi \in \hat{\Xi}_n^\varepsilon} \varepsilon D_{x_n^\varepsilon} (\tilde{\Gamma}_{x_n^\varepsilon, K} + \xi) \right),$$

where $\tilde{\Gamma}_{x_n^\varepsilon, K} = \tilde{K}_{x_n^\varepsilon} \Gamma$ and $\Gamma = \overline{\partial Y_0 \setminus \Gamma_{0,1}}$. For the problem analyzed here, we shall consider $\tilde{x}_n^\varepsilon = x_n^\varepsilon$.

The following calculations illustrate the motivation for the locally periodic approximation and determine formulas for the transformation matrices D and \tilde{K} . For $n = 1, \dots, N_\varepsilon$, we choose such $\kappa_n \in \mathbb{Z}^3$ that for $x_n^\varepsilon = R_{x_n^\varepsilon} \varepsilon \kappa_n$ we have $x_n^\varepsilon \in \Omega_n^\varepsilon$. In the definition of covering of Ω_n^ε by shifted parallelepipeds, we consider a numbering of $\xi \in \Xi_n^\varepsilon$ and write

$$\Omega_n^\varepsilon \subset x_n^\varepsilon + \bigcup_{j=1}^{I_n^\varepsilon} \varepsilon D_{x_n^\varepsilon} (\bar{Y} + \xi_j) \quad \text{for } \xi_j \in \Xi_n^\varepsilon.$$

Then for $1 \leq j \leq I_n^\varepsilon$ we consider $k_j^n = \kappa_n + \xi_j$ and $x_{k_j^n}^\varepsilon = R_{k_j^n} \varepsilon k_j^n$. Here $R_{k_j^n} := R_{x_{k_j^n}^\varepsilon}$ and $R_{\kappa_n} := R_{x_n^\varepsilon}$.

Using the regularity assumptions on the function γ and considering the Taylor expansion of R^{-1} around x_n^ε , i.e. around $\varepsilon \kappa_{n,3}$, we obtain

$$\begin{aligned} R_{k_j^n}^{-1}(x - x_{k_j^n}^\varepsilon) &= R_{\kappa_n}^{-1}x - \varepsilon k_j^n \\ &= R_{\kappa_n}^{-1}x + (R_{\kappa_n}^{-1})' x_n^\varepsilon \xi_{j,3} \varepsilon + (R_{\kappa_n}^{-1})'(x - x_n^\varepsilon) \xi_{j,3} \varepsilon + b(|\xi_{j,3} \varepsilon|^2)x - \varepsilon(\kappa_n + \xi_j) \\ &= R_{\kappa_n}^{-1}(x - x_n^\varepsilon) - \tilde{W}_{x_n^\varepsilon} \xi_j \varepsilon + (R_{\kappa_n}^{-1})'(x - x_n^\varepsilon) \xi_{j,3} \varepsilon + b(|\xi_{j,3} \varepsilon|^2)x, \end{aligned} \tag{9}$$

where $\tilde{W}_{x_n^\varepsilon} = \tilde{W}(x_n^\varepsilon)$ with $\tilde{W}(x) = (I - \nabla R^{-1}(\gamma(x_3))x)$. The notation of the gradient is understood as $\nabla R^{-1}(\gamma(x))x = \nabla_z (R^{-1}(\gamma(z))x)|_{z=x}$. Thus, for $x, x_n^\varepsilon \in \Omega_n^\varepsilon$, since $|x - x_n^\varepsilon| \leq C\varepsilon^r$, the distance between $R_{x_n^\varepsilon}^{-1}(x - x_n^\varepsilon) - \tilde{W}_{x_n^\varepsilon} \xi_j \varepsilon$ and $R_{k_j^n}^{-1}(x - x_{k_j^n}^\varepsilon)$ is of the order $\sup_{1 \leq j \leq I_n^\varepsilon} |\xi_j \varepsilon|^2 \sim \varepsilon^{2r}$.

This calculation together with the estimates below will ensure that the non-periodic plywood-like structure can be approximated by the corresponding locally periodic microstructure, comprising $\tilde{Y}_{x_n^\varepsilon}$ -periodic structure in each Ω_n^ε for $n = 1, \dots, N_\varepsilon$, and $|\Omega_n^\varepsilon| \sim \varepsilon^{3r}$ for an appropriate $r \in (0, 1)$. Here $\tilde{Y}_x = D(x)Y$, with $D(x) = R_x W(x)$, and

$$W(x) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & w(x) \\ 0 & 0 & 1 \end{pmatrix}, \tag{10}$$

where $w(x) = \gamma'(x_3)(\cos(\gamma(x_3))x_1 + \sin(\gamma(x_3))x_2)$ and $R_x = R(\gamma(x_3))$. The transformation matrix \tilde{K} is defined as $\tilde{K}(x) = W^{-1}(x)K(x)$ and the boundary of the muscle fibres in the locally periodic approximation is given by $\tilde{\Gamma}_x = D(x)\tilde{K}(x)\Gamma = R_x K(x)\Gamma$.

The definitions of R , W and γ ensure that the transformation matrices D and \tilde{K} are Lipschitz continuous, as well as $0 < d_0 \leq |\det D(x)| \leq d_1 < \infty$ and $0 < \rho_0^2 \leq |\det \tilde{K}(x)| \leq \rho_1^2 < \infty$ for all $x \in \bar{\Omega}$. Since ϑ is independent of the first variable, we consider in the definition of $W(x)$ the shift only in the second variable. Notice that if the original microstructure would be locally periodic, i.e. $R(\gamma(x_3)) = R(\gamma(x_{n,3}^\varepsilon))$ for $x \in \Omega_n^\varepsilon$ and some $x_n^\varepsilon \in \Omega_n^\varepsilon$, then the matrix W would be constant in each Ω_n^ε and we would obtain $D(x) = R(\gamma(x_3))$ for $x \in \Omega$.

In the approximation of the problem posed in the domain with the non-periodic plywood-like structure, we shall use the following lemma, proven in [6], that facilitate the estimate for the difference between the values of the characteristic function at two different points.

Lemma 3 ([6]) *Let χ_b be the characteristic function of the fibre of radius b , i.e., $\chi_b(x) = 1$ for $|\hat{x}| \leq b$ and $\chi_b(x) = 0$ for $|\hat{x}| > b$, with $\hat{x} = (x_2, x_3)$. Then*

$$\|\chi_b(x + \delta) - \chi_b(x)\|_{L^2(\Omega)}^2 \leq CbL|\delta| \quad \text{for all } b > 0 \text{ and } |\delta| \leq b,$$

where L is the length of the fibre and the constant C is independent of b , δ , and L .

Deriving estimates for the difference of solutions of the original microscopic problem and the corresponding locally periodic approximation and applying techniques of locally periodic homogenization, we obtain the following macroscopic equations for the microscopic problem (1) and (2).

Theorem 1 *A sequence of solutions of the microscopic problem (1) and (2) converges to a solution*

$$c \in L^2(0, T; H^1(\Omega)) \cap H^1(0, T; L^2(\Omega)) \quad \text{and} \quad r_l \in H^1(0, T; L^2(\Omega; L^2(\tilde{\Gamma}_x))),$$

where $l = f, b$, of the macroscopic equations

$$\begin{aligned} \theta(x)\partial_t c - \operatorname{div}(\mathcal{A}(x)\nabla c) &= \theta(x)F(c) + \frac{1}{|\tilde{Y}_x|} \int_{\tilde{\Gamma}_x} [\tilde{\beta}(x, y)r_b - \tilde{\alpha}(x, y)r_f c] d\sigma_y, \\ \mathcal{A}(x)\nabla c \cdot \mathbf{n} &= 0 \quad \text{on } \partial\Omega \times (0, T), \\ \partial_t r_f &= p(r_b) - \tilde{\alpha}(x, y)r_f c + \tilde{\beta}(x, y)r_b - d_f r_f, \\ \partial_t r_b &= \tilde{\alpha}(x, y)r_f c - \tilde{\beta}(x, y)r_b - d_b r_b, \end{aligned} \tag{11}$$

for $(t, x) \in (0, T) \times \Omega$ and $y \in \tilde{\Gamma}_x$, where the macroscopic diffusion coefficient \mathcal{A} is defined as

$$\mathcal{A}_{ij}(x) = \frac{1}{|\tilde{Y}_x|} \int_{\tilde{Y}_{x,K}^*} (A_{ij} + A_{ik}\partial_{y_k} w^j(x, y)) dy,$$

with w^j , for $j = 1, 2, 3$, which are solutions of the unit cell problems

$$\begin{aligned} \operatorname{div}(A(\nabla_y w^j + \mathbf{e}_j)) &= 0 \text{ in } \tilde{Y}_{y,K}^*, \\ A(\nabla_y w^j + \mathbf{e}_j) \cdot \mathbf{n} &= 0 \text{ on } \tilde{\Gamma}_x, \quad w^j \tilde{Y}_x - \text{periodic}, \int_{\tilde{Y}_x^*} w^j dy = 0. \end{aligned} \tag{12}$$

Here

$$\begin{aligned} \tilde{Y}_{x,K}^* &= D_x \left(Y \setminus \bigcup_{m \in \{0,1\}^3} (\tilde{K}_x Y_0 + m) \right), \quad \tilde{Y}_x = D_x Y, \\ \tilde{\Gamma}_x &= \bigcup_{m \in \{0,1\}^3} D_x (\tilde{K}_x \Gamma + m) \cap \tilde{Y}_x, \quad \theta(x) = \frac{|\tilde{Y}_{x,K}^*|}{|\tilde{Y}_x|}, \end{aligned}$$

where $Y = (0, 1)^3$, $R_x = R(\gamma(x_3))$, $D_x = R_x W_x$, $\tilde{K}_x = W_x^{-1} K_x$, with $W_x = W(x)$ defined by (10), and

$$\begin{aligned} \tilde{\alpha}(x, y) &= \sum_{m \in \mathbb{Z}^3} \alpha(x, R_x^{-1}(y - D_x m)), \\ \tilde{\beta}(x, y) &= \sum_{m \in \mathbb{Z}^3} \beta(x, R_x^{-1}(y - D_x m)). \end{aligned} \tag{13}$$

Proof Using calculations from above, we consider a domain with a locally periodic microstructure characterised by the periodicity cell $\tilde{Y}_{x_n^\varepsilon} = D_{x_n^\varepsilon} Y$ in each Ω_n^ε , with $n = 1, \dots, N_\varepsilon$ and the shift $x_n^\varepsilon \in \Omega_n^\varepsilon$ in the covering of Ω_n^ε by $D_{x_n^\varepsilon}(Y + \xi)$, with $\xi \in \Xi_n^\varepsilon$.

Then, the characteristic function of the extracellular space $\tilde{\Omega}_\varepsilon^*$ in a tissue with locally periodic microstructure is defined by $\chi_{\tilde{\Omega}_\varepsilon^*} = (1 - \chi_{\tilde{\Omega}_\varepsilon^f})\chi_\Omega$, where $\chi_{\tilde{\Omega}_\varepsilon^f}$ denotes the characteristic function of fibres

$$\chi_{\tilde{\Omega}_\varepsilon^f} = \sum_{n=1}^{N_\varepsilon} \chi_{\tilde{\Omega}_{n,f}^\varepsilon} \quad \text{and} \quad \chi_{\tilde{\Omega}_{n,f}^\varepsilon} = \sum_{\xi \in \Xi_n^\varepsilon} \vartheta(x_n^\varepsilon, R_{x_n^\varepsilon}^{-1}(x - x_n^\varepsilon - \varepsilon D_{x_n^\varepsilon} \xi) / \varepsilon) \chi_{\Omega_n^\varepsilon}.$$

The boundaries of the locally periodic microstructure are denoted by

$$\tilde{\Gamma}^\varepsilon = \bigcup_{n=1}^{N_\varepsilon} \bigcup_{\xi \in \Xi_n^\varepsilon} (x_n^\varepsilon + \varepsilon R_{x_n^\varepsilon} K_{x_n^\varepsilon} \Gamma + \varepsilon D_{x_n^\varepsilon} \xi) \cap \Omega.$$

Notice that non-periodic changes in the shape of the perforations (radius of muscle fibres) are approximated by the same transformation matrix $K(x)$. This is consistent with the results obtained in [10, 11, 23, 33]. However spatial changes in the periodicity are approximated by $D_x = R_x W_x$.

The reaction rates (binding and dissociation rates) are defined in terms of locally periodic microstructure in the following way

$$\begin{aligned} \tilde{\alpha}^\varepsilon(x) &= \sum_{n=1}^{N_\varepsilon} \sum_{\xi \in \Xi_n^\varepsilon} \alpha(x, (R_{x_n^\varepsilon}^{-1}(x - x_n^\varepsilon) - W_{x_n^\varepsilon} \varepsilon \xi) / \varepsilon) \chi_{\Omega_n^\varepsilon}, \\ \tilde{\beta}^\varepsilon(x) &= \sum_{n=1}^{N_\varepsilon} \sum_{\xi \in \Xi_n^\varepsilon} \beta(x, (R_{x_n^\varepsilon}^{-1}(x - x_n^\varepsilon) - W_{x_n^\varepsilon} \varepsilon \xi) / \varepsilon) \chi_{\Omega_n^\varepsilon}. \end{aligned}$$

To show that we can approximate the problem (1) and (2) by a microscopic problem defined in the domain with the locally periodic microstructure, we have to prove that the

difference between the characteristic function of the original domain $\chi_{\Omega_\varepsilon^*}$ and of the locally periodic perforated domain $\chi_{\tilde{\Omega}_\varepsilon^*}$ converges to zero strongly in $L^2(\Omega)$ as $\varepsilon \rightarrow 0$. Also, we have to show that the difference between boundary integrals and their locally periodic approximations converges to zero as $\varepsilon \rightarrow 0$. This will ensure that as $\varepsilon \rightarrow 0$ the sequence of solutions of the original microscopic problem (1)–(2) will converge to a solution of the macroscopic equations obtained by homogenization of the corresponding problem defined in the domain with locally periodic microstructure.

For the difference between $\chi_{\Omega_\varepsilon^*}$ and $\chi_{\tilde{\Omega}_\varepsilon^*}$, we have

$$\begin{aligned} & \int_{\Omega} |\chi_{\Omega_\varepsilon^*} - \chi_{\tilde{\Omega}_\varepsilon^*}|^2 dx \leq \mathcal{I}_1 + \mathcal{I}_2 \\ &= \sum_{n=1}^{N_\varepsilon} \int_{\Omega_n^\varepsilon} \sum_{j \in J_n^\varepsilon} \left| \vartheta(x_{k_j}^\varepsilon, R_{x_{k_j}^\varepsilon}^{-1}(x - x_{k_j}^\varepsilon)/\varepsilon) - \vartheta(x_n^\varepsilon, R_{x_n^\varepsilon}^{-1}(x - x_{k_j}^\varepsilon)/\varepsilon) \right|^2 dx \\ & \quad + \sum_{n=1}^{N_\varepsilon} \int_{\Omega_n^\varepsilon} \sum_{j \in J_n^\varepsilon} \left| \vartheta(x_n^\varepsilon, R_{x_n^\varepsilon}^{-1}(x - x_{k_j}^\varepsilon)/\varepsilon) - \vartheta(x_n^\varepsilon, (R_{x_n^\varepsilon}^{-1}(x - x_n^\varepsilon) - \varepsilon W_{x_n^\varepsilon} j)/\varepsilon) \right|^2 dx, \end{aligned}$$

where $x_n^\varepsilon = R_{x_n^\varepsilon} \varepsilon \kappa_n$, $x_{k_j}^\varepsilon = R_{x_{k_j}^\varepsilon} \varepsilon k_j$, with $k_j = \kappa_n + j$, and

$$J_n^\varepsilon = \left\{ j \in \mathbb{Z}^3 : \left[(x_{k_j}^\varepsilon + \varepsilon R_{x_{k_j}^\varepsilon} Y_1) \cup (x_n^\varepsilon + \varepsilon R_{x_n^\varepsilon} Y_1 + \varepsilon D_{x_n^\varepsilon} j) \right] \cap \Omega_n^\varepsilon \neq \emptyset \right\}.$$

We notice that $\varepsilon^3 |J_n^\varepsilon| \leq C \varepsilon^{3r}$ and $|N_\varepsilon| \leq C \varepsilon^{-3r}$. For the first integral, we have

$$\mathcal{I}_1 \leq \sum_{n=1}^{N_\varepsilon} \varepsilon^3 |J_n^\varepsilon| \|\nabla \rho\|_{L^\infty(\Omega)} \sup_{j \in J_n^\varepsilon} |x_n^\varepsilon - x_{k_j}^\varepsilon| \leq C \varepsilon^r.$$

To estimate the second integral, we use Lemma 3. Since in each Ω_n^ε the length of fibres is of order ε^r , applying estimate in Lemma 3, equality (9), and the estimates $N_\varepsilon \leq C \varepsilon^{-3r}$ and $|J_n^\varepsilon| \leq C \varepsilon^{3(r-1)}$, we conclude that

$$\mathcal{I}_2 \leq C \varepsilon^{3r-2}.$$

Thus, for $r \in (2/3, 1)$, we have $\mathcal{I}_1 \rightarrow 0$ and $\mathcal{I}_2 \rightarrow 0$ as $\varepsilon \rightarrow 0$.

To estimate the difference between boundary integral we have to extend c^ε , r_f^ε , and r_b^ε from Ω_ε^* to Ω . For c^ε , we can consider the extension as in Lemma 2. Then, using the extended \tilde{c}^ε and the fact that the reaction rates and the initial data are defined on whole Ω we can extend r_f^ε and r_b^ε to Ω as solutions of the ordinary differential equations with \tilde{c}^ε instead of c^ε

$$\begin{aligned} \partial_t \tilde{r}_f^\varepsilon &= p(\tilde{r}_b^\varepsilon) - \alpha^\varepsilon(x) \tilde{c}^\varepsilon \tilde{r}_f^\varepsilon + \beta^\varepsilon(x) \tilde{r}_b^\varepsilon - d_f \tilde{r}_f^\varepsilon && \text{in } (0, T) \times \Omega, \\ \partial_t \tilde{r}_b^\varepsilon &= \alpha^\varepsilon(x) \tilde{c}^\varepsilon \tilde{r}_f^\varepsilon - \beta^\varepsilon(x) \tilde{r}_b^\varepsilon - d_b \tilde{r}_b^\varepsilon && \text{on } (0, T) \times \Omega, \\ \tilde{r}_f^\varepsilon(0, x) &= r_{f0}^\varepsilon(x), \quad \tilde{r}_b^\varepsilon(0, x) = r_{b0}^\varepsilon(x) && \text{in } \Omega. \end{aligned} \tag{14}$$

The non-negativity of c^ε and the construction of the extension ensure that \tilde{c}^ε is non-negative. Then, in the same way as for r_f^ε and r_b^ε , using the properties of p and the non-negativity of the coefficients and initial data, we obtain the non-negativity of \tilde{r}_f^ε and \tilde{r}_b^ε . Thus, adding the equations for \tilde{r}_f^ε and \tilde{r}_b^ε , we obtain the boundedness of \tilde{r}_f^ε and \tilde{r}_b^ε in Ω_T , i.e.

$$\|\tilde{r}_f^\varepsilon\|_{L^\infty(\Omega_T)} + \|\tilde{r}_b^\varepsilon\|_{L^\infty(\Omega_T)} \leq C.$$

Differentiating equations in (14) with respect to x and using the estimate $\|\nabla \tilde{r}_b^\varepsilon\|_{L^2(\Omega)} \leq \|\nabla \tilde{r}_b^\varepsilon + \nabla \tilde{r}_f^\varepsilon\|_{L^2(\Omega)} + \|\nabla \tilde{r}_f^\varepsilon\|_{L^2(\Omega)}$, we obtain

$$\begin{aligned} &\|\nabla \tilde{r}_f^\varepsilon\|_{L^\infty(0,T;L^2(\Omega))} + \|\nabla \tilde{r}_b^\varepsilon\|_{L^\infty(0,T;L^2(\Omega))} \leq \mu_1 \|\nabla \tilde{c}^\varepsilon\|_{L^2(\Omega_T)} \\ &+ \varepsilon^{-1} \mu_2 \|\tilde{c}^\varepsilon\|_{L^2(\Omega_T)} + \varepsilon^{-1} \mu_3 \left[\|\tilde{r}_f^\varepsilon\|_{L^\infty(\Omega_T)} + \|\tilde{r}_b^\varepsilon\|_{L^\infty(\Omega_T)} \right] \leq \mu_4 \left(1 + \frac{1}{\varepsilon} \right), \end{aligned} \tag{15}$$

where the constants μ_j , with $j = 2, 3, 4$, depend on the derivatives of α, β , and μ_j , with $j = 1, 2, 3, 4$, are independent of ε . Hence, for each $\varepsilon > 0$, the extensions $\tilde{r}_f^\varepsilon, \tilde{r}_b^\varepsilon$, and \tilde{c}^ε are well-defined on the boundaries $\tilde{\Gamma}^\varepsilon$ of the locally periodic microstructure. In what follows we shall use the same notation for a function and for its extension.

Notice that ε^{-1} in the estimates for ∇r_f^ε and ∇r_b^ε will be compensated by ε in the estimate for the difference between neighbouring points in non-periodic and locally periodic domains, respectively, i.e.

$$|\varepsilon R_{x_k^\varepsilon} K_{x_k^\varepsilon} y - \varepsilon R_{x_n^\varepsilon} K_{x_n^\varepsilon} y| \leq C \varepsilon^{1+r} (1 + \|\gamma'\|_{L^\infty(\mathbb{R})}) (1 + \|\nabla K\|_{L^\infty(\Omega)}).$$

Then, for the boundary integrals, we have

$$\begin{aligned} &\varepsilon \left| \int_{\Gamma^\varepsilon} \alpha^\varepsilon r_f^\varepsilon c^\varepsilon \psi d\sigma_x^\varepsilon - \int_{\tilde{\Gamma}^\varepsilon} \tilde{\alpha}^\varepsilon r_f^\varepsilon c^\varepsilon \psi d\sigma_x^\varepsilon \right| \leq \mathcal{I}_3 + \mathcal{I}_4 \\ &= \varepsilon \sum_{n=1}^{N_\varepsilon} \sum_{j \in J_n^\varepsilon} \left| \int_{\varepsilon K_{x_k^\varepsilon}^\varepsilon \Gamma + x_k^\varepsilon} \alpha^\varepsilon r_f^\varepsilon c^\varepsilon \psi d\sigma_x^\varepsilon - \int_{\varepsilon K_{x_n^\varepsilon}^\varepsilon \Gamma + x_k^\varepsilon} \alpha^\varepsilon r_f^\varepsilon c^\varepsilon \psi d\sigma_x^\varepsilon \right| \\ &+ \varepsilon \sum_{n=1}^{N_\varepsilon} \sum_{j \in J_n^\varepsilon} \left| \int_{\varepsilon K_{x_n^\varepsilon}^\varepsilon \Gamma + x_k^\varepsilon} \alpha^\varepsilon r_f^\varepsilon c^\varepsilon \psi d\sigma_x^\varepsilon - \int_{\varepsilon K_{x_n^\varepsilon}^\varepsilon \Gamma + x_n^\varepsilon + \varepsilon D_{x_n^\varepsilon} j} \tilde{\alpha}^\varepsilon r_f^\varepsilon c^\varepsilon \psi d\sigma_x^\varepsilon \right| \end{aligned}$$

for $\psi \in C^1(\bar{\Omega}_T)$, where $K_x^R = R_x K_x$. Considering the regularity of K and R and the uniform boundedness from below and above of $|\det K|$, and using the trace estimate for the $L^2(\Gamma)$ -norm of a $H^\zeta(Y)$ -function, with $\zeta \in (1/2, 1)$, the first integral we can estimate as

$$\begin{aligned} \mathcal{I}_3 &\leq C_1 \varepsilon^d \sum_{n=1}^{N_\varepsilon} \sum_{j \in J_n^\varepsilon} \int_\Gamma \left| \alpha^\varepsilon r_f^\varepsilon(t, y_{k_j^\varepsilon}) c^\varepsilon(t, y_{k_j^\varepsilon}) - \alpha^\varepsilon r_f^\varepsilon(t, y_{k_n^\varepsilon}) c^\varepsilon(t, y_{k_n^\varepsilon}) \right| d\sigma_y + C_2 \varepsilon^r \\ &\leq C_3 \left[\varepsilon^{\frac{d+1}{2}} \sum_{n=1}^{N_\varepsilon} \sum_{j \in J_n^\varepsilon} \|c^\varepsilon\|_{L^2(\Gamma_{n,j}^\varepsilon)} \left[\int_\Gamma \left| \alpha^\varepsilon(y_{k_j^\varepsilon}) - \alpha^\varepsilon(y_{k_n^\varepsilon}) \right|^2 d\sigma_y \right]^{\frac{1}{2}} \right] \\ &+ C_4 \varepsilon^d \sum_{n=1}^{N_\varepsilon} \sum_{j \in J_n^\varepsilon} \left[\int_Y |c^\varepsilon(t, y_{k_j^\varepsilon}) - c^\varepsilon(t, y_{k_n^\varepsilon})|^2 + |r_f^\varepsilon(t, y_{k_j^\varepsilon}) - r_f^\varepsilon(t, y_{k_n^\varepsilon})|^2 dy \right. \\ &+ \int_Y \int_Y \frac{|[c^\varepsilon(t, y_{k_j^\varepsilon}^1) - c^\varepsilon(t, y_{k_j^\varepsilon}^2)] - [c^\varepsilon(t, y_{k_n^\varepsilon}^1) - c^\varepsilon(t, y_{k_n^\varepsilon}^2)]|^2}{|y^1 - y^2|^{2\zeta+d}} dy^1 dy^2 \\ &+ \left. \int_Y \int_Y \frac{|[r_f^\varepsilon(y_{k_j^\varepsilon}^1) - r_f^\varepsilon(y_{k_j^\varepsilon}^2)] - [r_f^\varepsilon(y_{k_n^\varepsilon}^1) - r_f^\varepsilon(y_{k_n^\varepsilon}^2)]|^2}{|y^1 - y^2|^{2\zeta+d}} dy^1 dy^2 \right]^{\frac{1}{2}} \\ &\times \left[\int_\Gamma (|c^\varepsilon(y_{k_j^\varepsilon})|^2 + |r_f^\varepsilon(y_{k_n^\varepsilon})|^2) d\sigma_y \right]^{\frac{1}{2}} + C_5 \varepsilon^r, \end{aligned}$$

where $d = \dim(\Omega) = 3$ and $\Gamma_{n,j}^\varepsilon = x_{k_j}^\varepsilon + \varepsilon K_{x_{k_j}^\varepsilon}^R \Gamma = x_{k_j}^\varepsilon + \varepsilon R_{x_{k_j}^\varepsilon} K(x_{k_j}^\varepsilon)\Gamma$, with $j \in J_n^\varepsilon$ and $n = 1, \dots, N_\varepsilon$. Here, we used the short notations

$$\begin{aligned} y_{k_j}^\varepsilon &= x_{k_j}^\varepsilon + \varepsilon K_{x_{k_j}^\varepsilon}^R y, & y_{\kappa_n^\varepsilon} &= x_{k_j}^\varepsilon + \varepsilon K_{x_n^\varepsilon}^R y, \\ y_{k_j^\varepsilon}^l &= x_{k_j}^\varepsilon + \varepsilon K_{x_{k_j}^\varepsilon}^R y^l, & y_{\kappa_n^\varepsilon}^l &= x_{k_j}^\varepsilon + \varepsilon K_{x_n^\varepsilon}^R y^l \end{aligned} \quad \text{for } l = 1, 2.$$

Using the regularity of γ , K , and α , and applying a priori estimates for c^ε and r_f^ε , together with (15), we obtain for $0 < \varsigma_1 < 1/2$, with $\varsigma + \varsigma_1 = 1$,

$$\begin{aligned} \int_0^T \mathcal{I}_3 dt &\leq \mu_1 \sum_{n=1}^{N_\varepsilon} \sum_{j \in J_n^\varepsilon} \varepsilon \int_0^T \left[\|\nabla r_f^\varepsilon\|_{L^2(Y_{k_j}^\varepsilon)} + \|\nabla c^\varepsilon\|_{L^2(Y_{k_j}^\varepsilon)} + \|\nabla \alpha^\varepsilon\|_{C(Y_{k_j}^\varepsilon)} \right] dt \\ &\quad \times \|\gamma'\|_{L^\infty(\mathbb{R})} \|\nabla K\|_{L^\infty(\Omega)} \left[\sup_{j \in J_n^\varepsilon} |x_{k_j}^\varepsilon - x_n^\varepsilon| + \sup_{j \in J_n^\varepsilon} |x_{k_j}^\varepsilon - x_n^\varepsilon|^{\varsigma_1} \right] + \mu_1 \varepsilon^r \\ &\leq \mu \varepsilon^{\varsigma_1 r}, \end{aligned}$$

where $Y_{k_j}^\varepsilon = x_{k_j}^\varepsilon + \varepsilon R_{x_{k_j}^\varepsilon} Y_1$. Conducting similar calculations as for \mathcal{I}_3 and using the fact that $|x_{k_j}^\varepsilon - x_n^\varepsilon - D_{x_n^\varepsilon} \varepsilon j| \leq C_1 |\varepsilon j|^2 \leq C_2 \varepsilon^{2r}$ yield

$$\begin{aligned} \int_0^T \mathcal{I}_4 dt &\leq \mu_1 \sum_{n=1}^{N_\varepsilon} \sum_{j \in J_n^\varepsilon} \int_0^T \left[\|\nabla r_f^\varepsilon\|_{L^2(\tilde{Y}_{n,j}^\varepsilon)} + \|\nabla c^\varepsilon\|_{L^2(\tilde{Y}_{n,j}^\varepsilon)} + \|\nabla \tilde{\alpha}^\varepsilon\|_{C(\tilde{Y}_{n,j}^\varepsilon)} \right] dt \\ &\quad \times \left(\varepsilon^{1+r} + \varepsilon^\varsigma \left| R_{x_{k_j}^\varepsilon}^{-1}(x - x_{k_j}^\varepsilon) - R_{x_n^\varepsilon}^{-1}(x - x_n^\varepsilon) - W_{x_n^\varepsilon} \varepsilon j \right|^{\varsigma_1} \right) \\ &\leq \mu \left[\varepsilon^{(2r-1)\varsigma_1} + \varepsilon^r \right], \end{aligned}$$

where $\varsigma + \varsigma_1 = 1$ and $\tilde{Y}_{n,j}^\varepsilon = x_n^\varepsilon + \varepsilon R_{x_n^\varepsilon} Y_1 + \varepsilon D_{x_n^\varepsilon} j$. In a similar way, we obtain the estimates for the difference of the other two boundary integrals. Thus, we conclude that for $r \in (1/2, 1)$, the difference between the boundary integrals for non-periodic and locally periodic microstructures converges to zero as $\varepsilon \rightarrow 0$.

Notice that since the extensions of c^ε and r_l^ε , with $l = f, b$, are in $L^2(0, T; H^1(\Omega))$ and $\partial\Omega$ is Lipschitz continuous, we can extend them to $L^2(0, T; H^1(\Omega_1))$ with $\Omega \subset \Omega_1$.

Now, we rewrite the weak formulation of the microscopic equation for c^ε as

$$\begin{aligned} &\langle \partial_t c^\varepsilon - F(c^\varepsilon), \phi \chi_{\Omega_\varepsilon^*} \rangle_{\Omega_T} + \langle A \nabla c^\varepsilon, \nabla \phi \chi_{\Omega_\varepsilon^*} \rangle_{\Omega_T} - \varepsilon \langle \beta^\varepsilon r_b^\varepsilon - \alpha^\varepsilon c^\varepsilon r_f^\varepsilon, \phi \rangle_{\Gamma_T^\varepsilon} \\ &= \left[\langle \partial_t c^\varepsilon - F(c^\varepsilon), \phi \chi_{\tilde{\Omega}_\varepsilon^*} \rangle_{\Omega_T} + \langle A \nabla c^\varepsilon, \nabla \phi \chi_{\tilde{\Omega}_\varepsilon^*} \rangle_{\Omega_T} - \varepsilon \langle \tilde{\beta}^\varepsilon r_b^\varepsilon - \tilde{\alpha}^\varepsilon c^\varepsilon r_f^\varepsilon, \phi \rangle_{\tilde{\Gamma}_T^\varepsilon} \right] \\ &\quad + \left[\langle \partial_t c^\varepsilon - F(c^\varepsilon), \phi (\chi_{\Omega_\varepsilon^*} - \chi_{\tilde{\Omega}_\varepsilon^*}) \rangle_{\Omega_T} + \langle A \nabla c^\varepsilon, \nabla \phi (\chi_{\Omega_\varepsilon^*} - \chi_{\tilde{\Omega}_\varepsilon^*}) \rangle_{\Omega_T} \right] \\ &\quad - \varepsilon \left[\langle \beta^\varepsilon r_b^\varepsilon - \alpha^\varepsilon c^\varepsilon r_f^\varepsilon, \phi \rangle_{\Gamma_T^\varepsilon} - \langle \tilde{\beta}^\varepsilon r_b^\varepsilon - \tilde{\alpha}^\varepsilon c^\varepsilon r_f^\varepsilon, \phi \rangle_{\tilde{\Gamma}_T^\varepsilon} \right] = I_1 + I_2 + I_3 \end{aligned}$$

for $\phi \in C^1(\overline{\Omega_T})$. Notice that we use the same notation for c^ε , r_f^ε , and r_b^ε and their extensions.

The estimates for \mathcal{I}_1 , \mathcal{I}_2 , \mathcal{I}_3 , and \mathcal{I}_4 , shown above, and similar calculations for the difference of the boundary integrals of $\beta^\varepsilon r_b^\varepsilon$ and $\tilde{\beta}^\varepsilon r_b^\varepsilon$, respectively, imply that $I_2 \rightarrow 0$ and

$I_3 \rightarrow 0$ as $\varepsilon \rightarrow 0$. Thus we obtain that

$$\begin{aligned} 0 &= \lim_{\varepsilon \rightarrow 0} \left[\langle \partial_t c^\varepsilon - F(c^\varepsilon), \phi \chi_{\Omega_\varepsilon^*} \rangle_{\Omega_T} + \langle A \nabla c^\varepsilon, \nabla \phi \chi_{\Omega_\varepsilon^*} \rangle_{\Omega_T} - \varepsilon \langle \beta^\varepsilon r_b^\varepsilon - \alpha^\varepsilon c^\varepsilon r_f^\varepsilon, \phi \rangle_{\Gamma_T^\varepsilon} \right] \\ &= \lim_{\varepsilon \rightarrow 0} \left[\langle \partial_t c^\varepsilon - F(c^\varepsilon), \phi \chi_{\tilde{\Omega}_\varepsilon^*} \rangle_{\Omega_T} + \langle A \nabla c^\varepsilon, \nabla \phi \chi_{\tilde{\Omega}_\varepsilon^*} \rangle_{\Omega_T} - \varepsilon \langle \tilde{\beta}^\varepsilon r_b^\varepsilon - \tilde{\alpha}^\varepsilon c^\varepsilon r_f^\varepsilon, \phi \rangle_{\tilde{\Gamma}_T^\varepsilon} \right]. \end{aligned}$$

Similarly, we obtain that

$$\begin{aligned} 0 &= \lim_{\varepsilon \rightarrow 0} \varepsilon \langle \partial_t r_f^\varepsilon - p(r_b^\varepsilon) + \alpha^\varepsilon c^\varepsilon r_f^\varepsilon - \beta^\varepsilon r_b^\varepsilon + d_f r_f^\varepsilon, \phi \rangle_{\Gamma_T^\varepsilon} \\ &= \lim_{\varepsilon \rightarrow 0} \varepsilon \langle \partial_t r_f^\varepsilon - p(r_b^\varepsilon) + \tilde{\alpha}^\varepsilon c^\varepsilon r_f^\varepsilon - \tilde{\beta}^\varepsilon r_b^\varepsilon + d_f r_f^\varepsilon, \phi \rangle_{\tilde{\Gamma}_T^\varepsilon}, \\ 0 &= \lim_{\varepsilon \rightarrow 0} \varepsilon \langle \partial_t r_b^\varepsilon - \alpha^\varepsilon c^\varepsilon r_f^\varepsilon + \beta^\varepsilon r_b^\varepsilon + d_b r_b^\varepsilon, \phi \rangle_{\Gamma_T^\varepsilon} \\ &= \lim_{\varepsilon \rightarrow 0} \varepsilon \langle \partial_t r_b^\varepsilon - \tilde{\alpha}^\varepsilon c^\varepsilon r_f^\varepsilon + \tilde{\beta}^\varepsilon r_b^\varepsilon + d_b r_b^\varepsilon, \phi \rangle_{\tilde{\Gamma}_T^\varepsilon} \quad \text{for } \phi \in C^1(\bar{\Omega}_T). \end{aligned}$$

The definition of $\tilde{\Omega}_\varepsilon^*$, $\tilde{\Gamma}^\varepsilon$, $\tilde{\alpha}^\varepsilon$, and $\tilde{\beta}^\varepsilon$ implies that the original non-periodic problem is approximated by equations posed in a domain with locally periodic microstructure. Hence we can apply the methods of locally periodic two-scale convergence (l-t-s) and l-p unfolding operator to derive the limit equations.

Using the extension of c^ε , we have that the sequences $\{c^\varepsilon\}$, $\{\nabla c^\varepsilon\}$ and $\{\partial_t c^\varepsilon\}$ are defined on Ω_T and we can determine $\mathcal{T}_\mathcal{L}^\varepsilon(c^\varepsilon)$, $\mathcal{T}_\mathcal{L}^\varepsilon(\nabla c^\varepsilon)$, $\partial_t \mathcal{T}_\mathcal{L}^\varepsilon(c^\varepsilon)$, and $\mathcal{T}_\mathcal{L}^{b,\varepsilon}(c^\varepsilon)$. The properties of $\mathcal{T}_\mathcal{L}^\varepsilon$ and $\mathcal{T}_\mathcal{L}^{b,\varepsilon}$ together with estimates (5a)–(5b) ensure

$$\begin{aligned} \|\mathcal{T}_\mathcal{L}^\varepsilon(c^\varepsilon)\|_{L^2(\Omega_T \times Y)} + \|\mathcal{T}_\mathcal{L}^\varepsilon(\nabla c^\varepsilon)\|_{L^2(\Omega_T \times Y)} + \|\partial_t \mathcal{T}_\mathcal{L}^\varepsilon(c^\varepsilon)\|_{L^2(\Omega_T \times Y)} &\leq C, \\ \|\mathcal{T}_\mathcal{L}^{b,\varepsilon}(c^\varepsilon)\|_{L^2(\Omega_T \times \Gamma)} + \sum_{j=f,b} \|\mathcal{T}_\mathcal{L}^{b,\varepsilon}(r_j^\varepsilon)\|_{H^1(0,T;L^2(\Omega \times \Gamma))} &\leq C. \end{aligned}$$

Then, the convergence results for the l-p unfolding operator and l-t-s convergence, see [29, 30] or Appendix, imply that there exist subsequences (denoted again by c^ε , r_f^ε and r_b^ε) and the functions $c \in L^2(0, T; H^1(\Omega))$, $\partial_t c \in L^2(\Omega_T)$, $c_1 \in L^2(\Omega_T; H^1_{\text{per}}(\tilde{Y}_x))$, $r_f, r_b \in H^1(0, T; L^2(\Omega; L^2(\tilde{\Gamma}_x)))$ such that

$$\begin{aligned} \mathcal{T}_\mathcal{L}^\varepsilon(c^\varepsilon) &\rightarrow c && \text{strongly in } L^2(\Omega_T; H^1(Y)), \\ \partial_t \mathcal{T}_\mathcal{L}^\varepsilon(c^\varepsilon) &\rightarrow \partial_t c && \text{weakly in } L^2(\Omega_T \times Y), \\ \mathcal{T}_\mathcal{L}^\varepsilon(\nabla c^\varepsilon) &\rightarrow \nabla c + D_x^{-T} \nabla_{\tilde{y}} c_1(\cdot, D_x \cdot) && \text{weakly in } L^2(\Omega_T \times Y), \\ \mathcal{T}_\mathcal{L}^{b,\varepsilon}(c^\varepsilon) &\rightarrow c && \text{strongly in } L^2(\Omega_T; L^2(\Gamma)), \\ r_j^\varepsilon &\rightarrow r_j, \quad \partial_t r_j^\varepsilon \rightarrow \partial_t r_j && \text{l-t-s, } r_j, \partial_t r_j \in L^2(\Omega_T; L^2(\tilde{\Gamma}_x)), \\ \mathcal{T}_\mathcal{L}^{b,\varepsilon}(r_j^\varepsilon) &\rightarrow r_j(\cdot, \cdot, D_x \tilde{K}_x \cdot) && \text{weakly in } L^2(\Omega_T \times \Gamma), \\ \partial_t \mathcal{T}_\mathcal{L}^{b,\varepsilon}(r_j^\varepsilon) &\rightarrow \partial_t r_j(\cdot, \cdot, D_x \tilde{K}_x \cdot) && \text{weakly in } L^2(\Omega_T \times \Gamma), \quad j = f, b. \end{aligned} \tag{16}$$

The coefficients $\tilde{\alpha}^\varepsilon$ and $\tilde{\beta}^\varepsilon$ can be defined as locally periodic approximations of $\tilde{\alpha}$ and $\tilde{\beta}$, given by (13), i.e.,

$$\tilde{\alpha}^\varepsilon = \mathcal{L}^\varepsilon(\tilde{\alpha}), \quad \tilde{\beta}^\varepsilon = \mathcal{L}^\varepsilon(\tilde{\beta}) \quad \text{with } \tilde{x}_n^\varepsilon = x_n^\varepsilon.$$

See Appendix or [29] for the definition of the locally periodic approximation \mathcal{L}^ε . The regularity assumptions on α, β, K , and γ ensure that $\tilde{\alpha}, \tilde{\beta} \in C(\bar{\Omega}; C_{\text{per}}(\tilde{Y}_x))$.

Considering $\psi^\varepsilon(t, x) = \psi_1(t, x) + \varepsilon \mathcal{L}_\rho^\varepsilon(\psi_2)(t, x)$, with $\psi_1 \in C^1(\bar{\Omega}_T)$ and $\psi_2 \in C^1_0(\Omega_T; C^1_{\text{per}}(\tilde{Y}_x))$, as a test function in (3) (see Appendix or [29] for the definition of $\mathcal{L}_\rho^\varepsilon$)

and applying 1-p unfolding operator and 1-p boundary unfolding operator imply

$$\begin{aligned} & \langle \mathcal{T}_L^\varepsilon(\chi_{\Omega_\varepsilon}^\varepsilon) \partial_t \mathcal{T}_L^\varepsilon(c^\varepsilon), \mathcal{T}_L^\varepsilon(\psi^\varepsilon) \rangle_{\Omega_T \times Y} + \langle A \mathcal{T}_L^\varepsilon(\chi_{\Omega_\varepsilon}^\varepsilon) \mathcal{T}_L^\varepsilon(\nabla c^\varepsilon), \mathcal{T}_L^\varepsilon(\nabla \psi^\varepsilon) \rangle_{\Omega_T \times Y} \\ &= \langle \mathcal{T}_L^\varepsilon(\chi_{\Omega_\varepsilon}^\varepsilon) F(\mathcal{T}_L^\varepsilon(c^\varepsilon)), \mathcal{T}_L^\varepsilon(\psi^\varepsilon) \rangle_{\Omega_T \times Y} \\ &+ \left\langle \sum_{n=1}^{N_\varepsilon} \frac{\sqrt{g_{x_n^\varepsilon}}}{\sqrt{g} |\tilde{Y}_{x_n^\varepsilon}|} \left[\mathcal{T}_L^{b,\varepsilon}(\tilde{\beta}^\varepsilon r_b^\varepsilon) - \mathcal{T}_L^{b,\varepsilon}(\tilde{\alpha}^\varepsilon) \mathcal{T}_L^{b,\varepsilon}(c^\varepsilon) \mathcal{T}_L^{b,\varepsilon}(r_f^\varepsilon) \right] \chi_{\Omega_n^\varepsilon}, \mathcal{T}_L^{b,\varepsilon}(\psi^\varepsilon) \right\rangle_{\Omega_T \times \Gamma} \\ &- \langle A \nabla c^\varepsilon, \nabla \psi^\varepsilon \rangle_{\Lambda_\varepsilon^*, T} + \langle F(c^\varepsilon) - \partial_t c^\varepsilon, \psi^\varepsilon \rangle_{\Lambda_\varepsilon^*, T} + \varepsilon \langle \tilde{\beta}^\varepsilon r_b^\varepsilon - \tilde{\alpha}^\varepsilon c^\varepsilon r_f^\varepsilon, \psi^\varepsilon \rangle_{\tilde{\Gamma}_\varepsilon \setminus \hat{\Gamma}_\varepsilon, T} + \delta(\varepsilon), \end{aligned}$$

where $\delta(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$, $\chi_{\Omega_\varepsilon}^\varepsilon = \mathcal{L}_0^\varepsilon(\chi_{\tilde{Y}_{x,K}^*})$ and $\chi_{\tilde{Y}_{x,K}^*}$ is the characteristic function of $\tilde{Y}_{x,K}^*$ extended \tilde{Y}_x -periodically to \mathbb{R}^3 .

The regularity assumptions on γ and K ensure that $\chi_{\tilde{Y}_{x,K}^*} \in L^\infty(\bigcup_{x \in \Omega} \{x\} \times \tilde{Y}_x)$ and $\chi_{\tilde{Y}_{x,K}^*} \in C(\bar{\Omega}; L^p_{\text{per}}(\tilde{Y}_x))$ for $p \in (1, +\infty)$.

Applying the results from [30] we obtain $\mathcal{T}_L^\varepsilon(\chi_{\Omega_\varepsilon}^\varepsilon)(x, \tilde{y}) \rightarrow \chi_{\tilde{Y}_{x,K}^*}(x, D_x \tilde{y})$ in $L^p(\Omega \times Y)$ and $\mathcal{T}_L^{b,\varepsilon}(\tilde{\beta}^\varepsilon)(x, \tilde{y}) \rightarrow \tilde{\beta}(x, D_x \tilde{K}_x \tilde{y})$, $\mathcal{T}_L^{b,\varepsilon}(\tilde{\alpha}^\varepsilon)(x, \tilde{y}) \rightarrow \tilde{\alpha}(x, D_x \tilde{K}_x \tilde{y})$ in $L^p(\Omega \times \Gamma)$ for $p \in (1, +\infty)$, as $\varepsilon \rightarrow 0$.

Using the a priori estimates for c^ε , r_f^ε and r_b^ε , the strong convergence of $\mathcal{T}_L^\varepsilon(c^\varepsilon)$ in $L^2(\Omega_T; H^1(Y))$, the strong convergence and boundedness of $\mathcal{T}_L^{b,\varepsilon}(\tilde{\alpha}^\varepsilon)$, the weak convergence and boundedness of $\mathcal{T}_L^{b,\varepsilon}(r_f^\varepsilon)$, together with the regularity of D , γ , and K , and the strong convergence of $\mathcal{T}_L^{b,\varepsilon}(\psi^\varepsilon)$, we obtain

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \left\langle \sum_{n=1}^{N_\varepsilon} \frac{\sqrt{g_{x_n^\varepsilon}}}{\sqrt{g} |\tilde{Y}_{x_n^\varepsilon}|} \mathcal{T}_L^{b,\varepsilon}(\tilde{\alpha}^\varepsilon) \mathcal{T}_L^{b,\varepsilon}(c^\varepsilon) \mathcal{T}_L^{b,\varepsilon}(r_f^\varepsilon) \chi_{\Omega_n^\varepsilon}, \mathcal{T}_L^{b,\varepsilon}(\psi^\varepsilon) \right\rangle_{\Omega_T \times \Gamma} \\ &= \left\langle \frac{\sqrt{g_x}}{\sqrt{g} |\tilde{Y}_x|} \tilde{\alpha}(x, D_x \tilde{K}_x \tilde{y}) c(t, x) r_f(t, x, D_x \tilde{K}_x \tilde{y}), \psi_1(t, x) \right\rangle_{\Omega_T \times \Gamma}. \end{aligned}$$

Similar arguments along with the Lipschitz continuity of F and the strong convergence of $\mathcal{T}_L^\varepsilon(\chi_{\Omega_\varepsilon}^\varepsilon)$ ensure

$$\langle \mathcal{T}_L^\varepsilon(\chi_{\Omega_\varepsilon}^\varepsilon) F(\mathcal{T}_L^\varepsilon(c^\varepsilon)), \mathcal{T}_L^\varepsilon(\psi^\varepsilon) \rangle_{\Omega_T \times Y} \rightarrow \langle \chi_{\tilde{Y}_{x,K}^*}(x, D_x \tilde{y}) F(c), \psi_1 \rangle_{\Omega_T \times Y}$$

as $\varepsilon \rightarrow 0$. Using the convergence results (16), the strong convergence of $\mathcal{T}_L^\varepsilon(\psi^\varepsilon)$ and $\mathcal{T}_L^\varepsilon(\nabla \psi^\varepsilon)$ and the fact that $|\Lambda_\varepsilon^*| \rightarrow 0$ as $\varepsilon \rightarrow 0$, taking the limit as $\varepsilon \rightarrow 0$, and considering the change of variables $y = D_x \tilde{y}$ for $\tilde{y} \in Y$ and $y = D_x \tilde{K}_x \tilde{y}$ for $\tilde{y} \in \Gamma$, we obtain

$$\begin{aligned} & \langle |\tilde{Y}_x|^{-1} c, \psi_1 \rangle_{\tilde{Y}_{x,K}^* \times \Omega_T} + \langle |\tilde{Y}_x|^{-1} A(\nabla c + \nabla_y c_1), \nabla \psi_1 + \nabla_y \psi_2 \rangle_{\tilde{Y}_{x,K}^* \times \Omega_T} \\ &+ \langle |\tilde{Y}_x|^{-1} [\tilde{\alpha}(x, y) r_f c - \tilde{\beta}(x, y) r_b], \psi_1 \rangle_{\tilde{\Gamma}_x \times \Omega_T} = \langle |\tilde{Y}_x|^{-1} F(c), \psi_1 \rangle_{\tilde{Y}_{x,K}^* \times \Omega_T}. \end{aligned}$$

Considering $\psi_1(t, x) = 0$ for $(t, x) \in \Omega_T$, we obtain

$$c_1(t, x, y) = \sum_{j=1}^3 \partial_{x_j} c(t, x) w^j(x, y),$$

where w^j are solutions of (12). Choosing $\psi_2(t, x, y) = 0$ for $(t, x) \in \Omega_T$ and $y \in \tilde{Y}_x$ yields the macroscopic equation for c .

Using the strong convergence of $\mathcal{T}_{\mathcal{L}}^{b,\varepsilon}(c^\varepsilon)$ in $L^2(\Omega_T; L^2(\Gamma))$, estimates (5a)–(5b) and (6), and the Lipschitz continuity of p we obtain that $\{\mathcal{T}_{\mathcal{L}}^{b,\varepsilon}(r_j^\varepsilon)\}$ is a Cauchy sequence in $L^2(\Omega_T; L^2(\Gamma))$ for $j = f, b$, and hence upto a subsequence, $\mathcal{T}_{\mathcal{L}}^{b,\varepsilon}(r_j^\varepsilon) \rightarrow r_j(\cdot, \cdot, D_x \tilde{K}_x \cdot)$ strongly in $L^2(\Omega_T; L^2(\Gamma))$. Then applying the l-p boundary unfolding operator to the equations on $\tilde{\Gamma}^\varepsilon$ and taking the limit as $\varepsilon \rightarrow 0$ we obtain the equations for r_f and r_b . The proof of the uniqueness of a solution of the macroscopic problem is similar to the corresponding proof for the microscopic problem, and hence the convergence of the whole sequences of solutions of the microscopic problem follows. \square

Remark 1 Notice that for the proof of the homogenization results it is sufficient to have a local extension of c^ε from Ω_ε^* to Ω^δ , with $\Omega^\delta = \{x \in \Omega : \text{dist}(x, \partial\Omega) > \delta\}$ for any fixed $\delta > 0$, and hence, the local strong convergence of $\mathcal{T}_{\mathcal{L}}^\varepsilon(c^\varepsilon)$, i.e., the strong convergence in $L^2(0, T; L^2_{\text{loc}}(\Omega; H^1(Y)))$.

Remark 2 For numerical computations of the cell problems (12) and the ordinary differential equations (11), defining the dynamics of receptor densities, approaches from the two-scale finite element method [24] or the heterogeneous multiscale method [1, 2, 17, 18] can be applied.

Acknowledgments This research was supported by the EPSRC First Grant EP/K036521/1.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Appendix: Definition and Convergence Results for the l-t-s Convergence and l-p Unfolding Operator

We shall consider the space $C(\bar{\Omega}; C_{\text{per}}(\tilde{Y}_x))$ given in a standard way, i.e. for any $\tilde{\psi} \in C(\bar{\Omega}; C_{\text{per}}(Y))$ the relation $\psi(x, y) = \tilde{\psi}(x, D_x^{-1}y)$ with $x \in \Omega$ and $y \in \tilde{Y}_x$ yields $\psi \in C(\bar{\Omega}; C_{\text{per}}(\tilde{Y}_x))$. In the same way the spaces $L^p(\Omega; C_{\text{per}}(\tilde{Y}_x))$, $L^p(\Omega; L^q_{\text{per}}(\tilde{Y}_x))$ and $C(\bar{\Omega}; L^q_{\text{per}}(\tilde{Y}_x))$, for $1 \leq p \leq \infty$, $1 \leq q < \infty$, are given.

Due to the assumptions on D , i.e. $D \in \text{Lip}(\bar{\Omega})$ and $0 < d_0 \leq |\det D(x)| \leq d_1 < \infty$ for all $x \in \bar{\Omega}$, we obtain that the function $u : \Omega \rightarrow C(\tilde{Y}_x)$ is well-defined. The separability of $C_{\text{per}}(\tilde{Y}_x)$ for each $x \in \Omega$ and the Weierstrass approximation for continuous functions $u : \Omega \rightarrow C_{\text{per}}(\tilde{Y}_x)$ ensure the separability of $C(\bar{\Omega}; C_{\text{per}}(\tilde{Y}_x))$. Also, we have the following relation for the norm $\|\psi\|_{C(\bar{\Omega}; C_{\text{per}}(\tilde{Y}_x))} := \sup_{x \in \bar{\Omega}} \sup_{y \in \tilde{Y}_x} |\psi(x, y)|$:

$$\|\psi\|_{C(\bar{\Omega}; C_{\text{per}}(\tilde{Y}_x))} = \sup_{x \in \bar{\Omega}} \sup_{y \in \tilde{Y}_x} |\tilde{\psi}(x, D_x^{-1}y)| = \sup_{x \in \bar{\Omega}} \sup_{\tilde{y} \in Y} |\tilde{\psi}(x, \tilde{y})|.$$

The assumptions on D and K ensure that

$$\begin{aligned} L^2(\Omega; H^1(\tilde{Y}_x)) &= \{u \in L^2(\Omega; L^2(\tilde{Y}_x)), \nabla_y u \in L^2(\Omega; L^2(\tilde{Y}_x)^3)\} \quad \text{and} \\ L^2(\Omega; L^2(\tilde{\Gamma}_x)) &= \{u : \cup_{x \in \Omega} (\{x\} \times \tilde{\Gamma}_x) \rightarrow \mathbb{R} \text{ measurable, with} \\ &\quad u(x) \in L_2(\tilde{\Gamma}_x) \text{ for a.e. } x \in \Omega \text{ and } \int_{\Omega} \|u\|_{L^2(\tilde{\Gamma}_x)}^2 dx < \infty\} \end{aligned}$$

are well-defined, separable Hilbert spaces, see e.g. [16, 26, 32].

Consider $\psi \in C(\bar{\Omega}; C_{\text{per}}(\tilde{Y}_x))$ and corresponding $\tilde{\psi} \in C(\bar{\Omega}; C_{\text{per}}(Y))$. As a locally periodic (l-p) approximation of ψ , we name $\mathcal{L}^\varepsilon : C(\bar{\Omega}; C_{\text{per}}(\tilde{Y}_x)) \rightarrow L^\infty(\Omega)$ given by, see [29],

$$(\mathcal{L}^\varepsilon \psi)(x) = \sum_{n=1}^{N_\varepsilon} \tilde{\psi} \left(x, \frac{D_{x_n^\varepsilon}^{-1}(x - \tilde{x}_n^\varepsilon)}{\varepsilon} \right) \chi_{\Omega_n^\varepsilon}(x) \quad \text{for } x \in \Omega. \tag{17}$$

We consider also the map $\mathcal{L}_0^\varepsilon : C(\bar{\Omega}; C_{\text{per}}(\tilde{Y}_x)) \rightarrow L^\infty(\Omega)$ defined for $x \in \Omega$ as

$$(\mathcal{L}_0^\varepsilon \psi)(x) = \sum_{n=1}^{N_\varepsilon} \psi \left(x_n^\varepsilon, \frac{x - \tilde{x}_n^\varepsilon}{\varepsilon} \right) \chi_{\Omega_n^\varepsilon}(x) = \sum_{n=1}^{N_\varepsilon} \tilde{\psi} \left(x_n^\varepsilon, \frac{D_{x_n^\varepsilon}^{-1}(x - \tilde{x}_n^\varepsilon)}{\varepsilon} \right) \chi_{\Omega_n^\varepsilon}(x).$$

If we choose $\tilde{x}_n^\varepsilon = D_{x_n^\varepsilon} \varepsilon k$ for some $k \in \mathbb{Z}^3$, then the periodicity of $\tilde{\psi}$ implies

$$(\mathcal{L}^\varepsilon \psi)(x) = \sum_{n=1}^{N_\varepsilon} \tilde{\psi} \left(x, \frac{D_{x_n^\varepsilon}^{-1} x}{\varepsilon} \right) \chi_{\Omega_n^\varepsilon}(x), \quad (\mathcal{L}_0^\varepsilon \psi)(x) = \sum_{n=1}^{N_\varepsilon} \tilde{\psi} \left(x_n^\varepsilon, \frac{D_{x_n^\varepsilon}^{-1} x}{\varepsilon} \right) \chi_{\Omega_n^\varepsilon}(x)$$

for $x \in \Omega$, see e.g. [29] for more details. In the similar way, we define $\mathcal{L}^\varepsilon \psi$ and $\mathcal{L}_0^\varepsilon \psi$ for ψ in $C(\bar{\Omega}; L^q_{\text{per}}(\tilde{Y}_x))$ or $L^p(\Omega; C_{\text{per}}(\tilde{Y}_x))$.

We define also a regular approximation of $\mathcal{L}^\varepsilon \psi$

$$(\mathcal{L}_\rho^\varepsilon \psi)(x) = \sum_{n=1}^{N_\varepsilon} \tilde{\psi} \left(x, \frac{D_{x_n^\varepsilon}^{-1} x}{\varepsilon} \right) \phi_{\Omega_n^\varepsilon}(x) \quad \text{for } x \in \Omega,$$

where $\phi_{\Omega_n^\varepsilon}$ are approximations of $\chi_{\Omega_n^\varepsilon}$ such that $\phi_{\Omega_n^\varepsilon} \in C_0^\infty(\Omega_n^\varepsilon)$ and

$$\sum_{n=1}^{N_\varepsilon} |\phi_{\Omega_n^\varepsilon} - \chi_{\Omega_n^\varepsilon}| \rightarrow 0 \text{ in } L^2(\Omega), \quad \|\nabla^m \phi_{\Omega_n^\varepsilon}\|_{L^\infty(\mathbb{R}^d)} \leq C \varepsilon^{-\rho m} \text{ for } 0 < r < \rho < 1.$$

We recall here the definition of locally periodic two-scale (l-t-s) convergence and l-p unfolding operator, see [29, 30] for details.

Definition 2 ([29]) Let $u^\varepsilon \in L^p(\Omega)$ for all $\varepsilon > 0$ and $1 < p < \infty$. We say the sequence $\{u^\varepsilon\}$ converges l-t-s to $u \in L^p(\Omega; L^p(\tilde{Y}_x))$ as $\varepsilon \rightarrow 0$ if $\|u^\varepsilon\|_{L^p(\Omega)} \leq C$ and for any $\psi \in L^q(\Omega; C_{\text{per}}(\tilde{Y}_x))$

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega} u^\varepsilon(x) \mathcal{L}^\varepsilon \psi(x) dx = \int_{\Omega} \int_{\tilde{Y}_x} u(x, y) \psi(x, y) dy dx,$$

where \mathcal{L}^ε is the l-p approximation of ψ and $1/p + 1/q = 1$.

Definition 3 ([30]) A sequence $\{u^\varepsilon\} \subset L^p(\tilde{\Gamma}^\varepsilon)$, with $1 < p < \infty$, is said to converge locally periodic two-scale (l-t-s) to $u \in L^p(\Omega; L^p(\tilde{\Gamma}_x))$ if

$$\varepsilon \|u^\varepsilon\|_{L^p(\tilde{\Gamma}^\varepsilon)}^p \leq C$$

and for any $\psi \in C(\bar{\Omega}; C_{\text{per}}(\tilde{Y}_x))$

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \int_{\tilde{\Gamma}^\varepsilon} u^\varepsilon(x) \mathcal{L}^\varepsilon \psi(x) d\sigma_x = \int_{\Omega} \frac{1}{|\tilde{Y}_x|} \int_{\tilde{\Gamma}_x} u(x, y) \psi(x, y) d\sigma_y dx,$$

where \mathcal{L}^ε is the l-p approximation of ψ defined in (17).

Lemma 4 ([30]) For $\psi \in C(\overline{\Omega}; C_{\text{per}}(\tilde{Y}_x))$ and $1 \leq p < \infty$, we have that

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \int_{\tilde{\Gamma}^\varepsilon} |\mathcal{L}^\varepsilon \psi(x)|^p d\sigma_x = \int_{\Omega} \frac{1}{|\tilde{Y}_x|} \int_{\tilde{\Gamma}_x} |\psi(x, y)|^p d\sigma_y dx.$$

Definition 4 ([30]) For any Lebesgue-measurable on Ω function ψ the locally periodic unfolding operator (1-p unfolding operator) $\mathcal{T}_{\mathcal{L}}^\varepsilon$ is defined as

$$\mathcal{T}_{\mathcal{L}}^\varepsilon(\psi)(x, y) = \sum_{n=1}^{N_\varepsilon} \psi \left(\varepsilon D_{x_n^\varepsilon} \left[D_{x_n^\varepsilon}^{-1} x / \varepsilon \right]_Y + \varepsilon D_{x_n^\varepsilon} y \right) \chi_{\hat{\Omega}_n^\varepsilon}(x)$$

for $x \in \Omega$ and $y \in Y$.

The definition implies that $\mathcal{T}_{\mathcal{L}}^\varepsilon(\phi)$ is Lebesgue-measurable on $\Omega \times Y$ and is zero for $x \in \Lambda^\varepsilon$, where $\Lambda^\varepsilon = \bigcup_{n=1}^{N_\varepsilon} (\Omega_n^\varepsilon \setminus \hat{\Omega}_n^\varepsilon) \cap \Omega$.

Definition 5 ([30]) For any Lebesgue-measurable on $\tilde{\Gamma}^\varepsilon$ function ψ the 1-p boundary unfolding operator $\mathcal{T}_{\mathcal{L}}^{b,\varepsilon}$ is defined as

$$\mathcal{T}_{\mathcal{L}}^{b,\varepsilon}(\psi)(x, y) = \sum_{n=1}^{N_\varepsilon} \psi \left(\varepsilon D_{x_n^\varepsilon} \left[D_{x_n^\varepsilon}^{-1} x / \varepsilon \right]_Y + \varepsilon D_{x_n^\varepsilon} \tilde{K}_{x_n^\varepsilon} y \right) \chi_{\hat{\Omega}_n^\varepsilon}(x)$$

for $x \in \Omega$ and $y \in \Gamma$.

These definitions provide a generalization of the periodic unfolding operator and periodic boundary unfolding operator introduced in [12–14] to locally periodic microstructures.

Theorem 2 ([30]) For a sequence $\{w^\varepsilon\} \subset L^p(\Omega)$, with $p \in (1, \infty)$, satisfying

$$\|w^\varepsilon\|_{L^p(\Omega)} + \varepsilon \|\nabla w^\varepsilon\|_{L^p(\Omega)} \leq C$$

there exist a subsequence (denoted again by $\{w^\varepsilon\}$) and $w \in L^p(\Omega; W_{\text{per}}^{1,p}(\tilde{Y}_x))$ such that

$$\begin{aligned} \mathcal{T}_{\mathcal{L}}^\varepsilon(w^\varepsilon) &\rightharpoonup w(\cdot, D_x \cdot) && \text{weakly in } L^p(\Omega; W^{1,p}(Y)), \\ \varepsilon \mathcal{T}_{\mathcal{L}}^\varepsilon(\nabla w^\varepsilon) &\rightharpoonup D_x^{-T} \nabla_y w(\cdot, D_x \cdot) && \text{weakly in } L^p(\Omega \times Y). \end{aligned}$$

Theorem 3 ([30]) For a sequence $\{w^\varepsilon\} \subset W^{1,p}(\Omega)$, with $p \in (1, \infty)$, that converges weakly to w in $W^{1,p}(\Omega)$, there exist a subsequence (denoted again by $\{w^\varepsilon\}$) and a function $w_1 \in L^p(\Omega; W_{\text{per}}^{1,p}(\tilde{Y}_x))$ such that

$$\begin{aligned} \mathcal{T}_{\mathcal{L}}^\varepsilon(w^\varepsilon) &\rightharpoonup w && \text{weakly in } L^p(\Omega; W^{1,p}(Y)), \\ \mathcal{T}_{\mathcal{L}}^\varepsilon(\nabla w^\varepsilon)(\cdot, \cdot) &\rightharpoonup \nabla_x w(\cdot) + D_x^{-T} \nabla_y w_1(\cdot, D_x \cdot) && \text{weakly in } L^p(\Omega \times Y). \end{aligned}$$

Theorem 4 ([30]) For a sequence $\{w^\varepsilon\} \subset L^p(\tilde{\Gamma}^\varepsilon)$, with $p \in (1, \infty)$, satisfying

$$\varepsilon \|w^\varepsilon\|_{L^p(\tilde{\Gamma}^\varepsilon)}^p \leq C$$

there exist a subsequence (denoted again by $\{w^\varepsilon\}$) and $w \in L^p(\Omega; L^p(\tilde{\Gamma}_x))$ such that

$$w^\varepsilon \rightarrow w \quad \text{locally periodic two-scale (l-t-s)}.$$

Theorem 5 ([30]) *Let $\{w^\varepsilon\} \subset L^p(\tilde{\Gamma}^\varepsilon)$ with $\varepsilon \|w^\varepsilon\|_{L^p(\tilde{\Gamma}^\varepsilon)}^p \leq C$, where $p \in (1, \infty)$. The following assertions are equivalent*

- (i) $w^\varepsilon \rightarrow w$ *l-t-s*, $w \in L^p(\Omega; L^p(\tilde{\Gamma}_x))$.
- (ii) $\mathcal{T}_{\mathcal{L}}^{b,\varepsilon}(w^\varepsilon) \rightharpoonup w(\cdot, D_x \tilde{K}_x \cdot)$ *weakly in $L^p(\Omega \times \Gamma)$.*

Theorems 4 and 5 imply that for $\{w^\varepsilon\} \subset L^p(\tilde{\Gamma}^\varepsilon)$ with $\varepsilon \|w^\varepsilon\|_{L^p(\tilde{\Gamma}^\varepsilon)}^p \leq C$ we have the weak convergence of $\{\mathcal{T}_{\mathcal{L}}^{b,\varepsilon}(w^\varepsilon)\}$ in $L^p(\Omega \times \Gamma)$, where $p \in (1, \infty)$.

References

1. Abdulle, A.: The finite element heterogeneous multiscale method: a computational strategy for multi-scale PDEs. *GAKUTO Int. Ser. Math. Sci. Appl.* **31**, 133–181 (2009)
2. Abdulle, A., Weinan, E., Engquist, B., Vanden-Eijnden, E.: The heterogeneous multiscale method. *Acta Numer.* **21**, 1–87 (2012)
3. Alexandre, R.: Homogenization and $\theta - 2$ convergence. *Proc. Roy. Soc. Edinb.* **127A**, 441–455 (1997)
4. Belhadj, M., Cancès, E., Gerbeau, J.-F., Mikelić, A.: Homogenization approach to filtration through a fibrous medium. *INRIA* **5277** (2004)
5. Belyaev, A.G., Pyatnitskii, A.L., Chechkin, G.A.: Asymptotic behavior of a solution to a boundary value problem in a perforated domain with oscillating boundary. *Sib. Math. J.* **39**, 621–644 (1998)
6. Briane, M.: Homogénéisation de matériaux fibrés et multi-couches. PhD thesis, Université Paris VI (1990)
7. Briane, M.: Three models of non periodic fibrous materials obtained by homogenization. *RAIRO Modél Math. Anal. Numér.* **27**, 759–775 (1993)
8. Briane, M.: Homogenization of a non-periodic material. *J. Math. Pures Appl.* **73**, 47–66 (1994)
9. Chavarría-Krauser, A., Ptashnyk, M.: Homogenization approach to water transport in plant tissues with periodic microstructures. *Math. Model. Nat. Phenom.* **8**, 80–111 (2013)
10. Chechkin, G.A., Piatnitski, A.L.: Homogenization of boundary-value problem in a locally periodic perforated domain. *Appl. Anal.* **71**, 215–235 (1999)
11. Chenais, D., Mascarenhas, M.L., Trabuco, L.: On the optimization of non periodic homogenized microstructures. *RAIRO Modél Math. Anal. Numér.* **31**, 559–597 (1997)
12. Cioranescu, D., Damlamian, A., Griso, G.: The periodic unfolding method in homogenization. *SIAM J. Math. Anal.* **40**, 1585–1620 (2008)
13. Cioranescu, D., Donato, P., Zaki, R.: The periodic unfolding method in perforated domains. *Port. Math.* **63**, 467–496 (2006)
14. Cioranescu, D., Damlamian, A., Donato, P., Griso, G., Zaki, R.: The periodic unfolding method in domains with holes. *SIAM J. Math. Anal.* **44**, 718–760 (2012)
15. Cioranescu, D., Saint Jean Paulin, J.: *Homogenization of Reticulated Structures*. Springer-Verlag, New York (1999)
16. Dixmier, J.: *Von Neumann Algebras*. North-Holland, Amsterdam (1981)
17. E, W., Engquist, B.: The heterogeneous multiscale methods. *Commun. Math. Sci.* **1**, 87–132 (2003)
18. Efendiev, Y., Hou, T.Y.: *Multiscale Finite Element Methods: Theory and Applications*. Springer, New York (2009)
19. Hornung, J., Jäger, W.: Diffusion, convection, adsorption, and reaction of chemicals in porous media. *J. Differ. Equ.* **92**, 199–225 (1991)
20. Ladyženskaja, O.A., Solonnikov, V.A., Ural'ceva, N.N.: *Linear and Quasi-Linear Equations of Parabolic Type*. American Mathematical Society, Providence, RI (1968)
21. Marciniak-Czochra, A., Ptashnyk, M.: Derivation of a macroscopic receptor-based model using homogenization techniques. *SIAM J. Math. Anal.* **40**, 215–237 (2008)
22. Mascarenhas, M.L., Poliševski, D.: The warping, the torsion and the Neumann problems in a quasi-periodically perforated domain. *RAIRO Modél Math. Anal. Numér.* **28**, 37–57 (1994)
23. Mascarenhas, M.L.: Homogenization problems in locally periodic perforated domains. In: Ciarlet, P.G., Trabuco, L., Viano, J. (eds.) *Asymptotic Methods for Elastic Structures: Proceedings of the International Conference, Lisbon, Portugal, 1993*, pp. 141–149. Walter de Gruyter, Berlin (1995)
24. Matache, A.-M., Schwab, C.: Two-scale FEM for homogenisation problems. *ESAIM: M2AN* **36**, 537–572 (2002)

25. McCulloch, A.D.: Cardiac biomechanics. In: Bronzino, J. (ed.) *The Biomedical Engineering Handbook*. 2nd edn. CRC Press LLC, Florida (2000). Chapter 28
26. Meier, S.A.: Two-scale models of reactive transport in porous media involving microstructural changes. Ph.D. Thesis. University of Bremen, Bremen, Germany (2008)
27. Nguetseng, G.: A general convergence result for a functional related to the theory of homogenization. *SIAM J. Math. Anal.* **20**, 608–623 (1989)
28. Peskin, C.S.: Fiber architecture of the left ventricular wall: An asymptotic analysis. *Commun. Pure. Appl. Math.* **42**, 79–113 (1989)
29. Ptashnyk, M.: Two-scale convergence for locally-periodic microstructures and homogenization of plywood structures. *SIAM Multiscale Model Simul.* **11**, 92–117 (2013)
30. Ptashnyk, M.: Locally periodic unfolding method and two-scale convergence on surfaces of locally periodic microstructures. *SIAM Multiscale Model Simul.* **13**, 1061–1105 (2015)
31. Shkoller, S.: An approximate homogenization scheme for nonperiodic materials. *Comput. Math. Appl.* **33**, 15–34 (1997)
32. Showalter, R.E., Walkington, N.J.: Micro-structure models of diffusion in fissured media. *J. Math. Anal. Appl.* **155**, 1–20 (1991)
33. van Noorden, T.L., Muntean, A.: Homogenisation of a locally periodic medium with areas of low and high diffusivity. *Eur. J. Appl. Math.* **22**, 493–516 (2011)
34. Wheeler-Jones, C.P.D.: Cell signalling in the cardiovascular system: an overview. *Heart* **91**, 1366–1374 (2005)