



**University of Dundee**

## **Insights into the ion-coupling mechanism in the MATE transporter NorM-VC**

Krah, Alexander; Zachariae, Ulrich

*Published in:*  
Physical Biology

*DOI:*  
[10.1088/1478-3975/aa5ee7](https://doi.org/10.1088/1478-3975/aa5ee7)

*Publication date:*  
2017

*Document Version*  
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

### *Citation for published version (APA):*

Krah, A., & Zachariae, U. (2017). Insights into the ion-coupling mechanism in the MATE transporter NorM-VC. *Physical Biology*, 14, [045009]. <https://doi.org/10.1088/1478-3975/aa5ee7>

### **General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Supporting Information

### Insights into the ion-coupling mechanism in the MATE transporter NorM-VC

Alexander Krah<sup>1,2\*</sup> & Ulrich Zachariae<sup>1,2</sup>

<sup>1</sup>Computational Biology, School of Life Sciences, University of Dundee, Dundee, DD1 5EH,  
UK

<sup>2</sup>Physics, School of Science and Engineering, University of Dundee, Nethergate, Dundee, DD1  
4NH, UK




Figure S1: a) Distribution of ion binding distances in the state E255<sub>deprot</sub>/D371<sub>prot</sub> using 10 kJ/(mol\*nm<sup>2</sup>) backbone restraints. b) Distance distributions within the binding site in an unrestrained simulation. In both simulations no electric field was applied.




Figure S2: Potential of Mean Force (PMF) of Na<sup>+</sup> ion translocation from the binding site to the bulk solution in the E255<sub>prot</sub>/D371<sub>deprot</sub> state. The free energy difference is 11.2 $\pm$  1.0 kcal/mol. This free energy difference is not in good agreement with experimental data (5.6 kcal/mol).