

Flight Lieutenant Peach's observations on Burning Feet Syndrome in Far Eastern Prisoners of War 1942-1945.

Journal:	<i>QJM: An International Journal of Medicine</i>
Manuscript ID	QJM-2016-547.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	24-Aug-2016
Complete List of Authors:	Roccroft, Nathaniel; Royal Navy, Medicine Mathew, Emily; Imperial College London, Department of Bioengineering and Research Fellow, Department of Surgery and Cancer Parkes, Meg; University of Liverpool, School of Tropical Medicine Frankland, William; N/A, N/A Gill, Geoff; University of Liverpool, School of Medicine Bouhassira, Didier ; Universite Versailles Saint-Quentin-en-Yvelines UFR medicale Paris Ile de France Ouest, Centre de Traitement et d'Evaluation de la Douleur Rice, Andrew; Imperial College London, Faculty of Medicine, Department of Surgery & Cancer
Keywords:	Neurology, Miscellaneous/other


 SCHOLARONE™
Manuscripts

Manuscript title:

Flight Lieutenant Peach's observations on Burning Feet Syndrome in Far Eastern Prisoners of War 1942-1945.

Authors:

Nathaniel T Roocroft¹, Emily Mayhew², Meg Parkes³, William A Frankland⁴, Geoffrey V Gill⁵, Didier Bouhassira⁶, Andrew SC Rice⁷

1. Nathaniel T Roocroft MB BS, BSc. Royal Navy General Duties Medical Officer, RFA ARGUS, United Kingdom.

2. Emily Mayhew PhD, Historian in Residence, Department of Bioengineering and Research Fellow, Department of Surgery and Cancer, Imperial College, London, United Kingdom.

3. Margaret (Meg) Helen Parkes, MPhil, University of Liverpool, Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, L3 5QA, UK

4. William A Frankland MA, DM, FRCP. Consultant Allergist, 46 Devonshire Close, London W1G 7BG.

5. Geoffrey V Gill MA, MSc, MD, PhD, FRCP, DTM&H. Emeritus Professor of International Medicine, Liverpool School of Tropical Medicine and University of Liverpool, United Kingdom & Consultant Physician, Aintree University Hospital, Liverpool, United Kingdom.

6. Didier Bouhassira MD PhD. Professor of Neurology, INSERM U-987, Centre d'Evaluation et de Traitement de la Douleur, Hôpital Ambroise Paré, APHP, Boulogne-Billancourt, France.

7. Andrew SC Rice, MD, FRCP, FRCA, FFPMRCA. Professor of Pain Research, Department of Surgery and Cancer, Imperial College, London, United Kingdom & Hon. Consultant in

1
2
3 Pain Medicine, Chelsea and Westminster Hospital NHS Foundation Trust, London, United
4
5 Kingdom.
6
7

8 **Corresponding Author address:**
9

10 Nathaniel Roocroft

11
12 Medical Officer

13
14 RFA ARGUS

15
16 BFPO 433
17
18
19

20 **Corresponding author email address:**
21

22 nroocroft@gmail.com
23
24

25 **Keywords:**
26

27 Neuropathic pain, Burning Feet Syndrome, Far Eastern Prisoner of War, Nutritional

28
29 Neuropathy, Peripheral Neuropathy.
30
31
32

33 **Word Count:**
34

35 3734 words.
36
37

38 **Cover Letter Abbreviated Summary**
39

40 This historical review analyses 'Burning Feet Syndrome', a condition suffered by Far Eastern

41
42 Prisoners of War in the Second World War. Case records from RAF doctor Nowell Peach,

43
44 written at the time, are retrospectively assessed against modern diagnostic criteria to

45
46 determine if the syndrome can be retrospectively classed as neuropathic pain.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

In the Asia-Pacific theatre of World War II, an estimated 132,000 Allied military personnel were captured by Japanese forces and incarcerated as Far East Prisoners of War (FEPOW)¹. The mortality rate of European and Australian FEPOW was approximately 27%, seven times that of Allied POW held by the Germans². Morbidity and mortality were high amongst FEPOW as a result of arduous physical labour in the adverse environment of tropical jungle, compounded by a policy of brutality, starvation, neglect and consequent disease. This maltreatment occurred throughout the region and was inflicted on civilian and military detainees alike, with the building of the notorious 'Death Railway' between Thailand and Burma (Siam) one of the best known events. Whilst the majority of prisoners were local Asian civilians, the plight of British, Australian and Dutch military captives has been recorded in most detail, most of whom were captured during the 1941-2 South East Asia campaigns which culminated in the surrender of Singapore.

The focus of this historical review is "Burning Feet Syndrome" (BFS), which afflicted between 10-33% of prisoners³. Although retrospective accounts of BFS were published by surviving FEPOW medical officers in the immediate post-war period^{4, 5, 6, 7}, these were largely based on the authors' post captivity recollections as no contemporaneous medical records of individual cases were thought to have survived. This situation changed in 2007 when the late Nowell Peach was interviewed as part of the Liverpool School of Tropical Medicine FEPOW oral history study^{1, 8}. He revealed the existence of a series of case notes recording 54 cases of BFS, including detailed symptom descriptions and neurological examinations, which he had recorded whilst in captivity in Java. This review is primarily an analysis of the data in Peach's case records, in particular it reassesses its neuropathic pain aspects using 21st century diagnostic criteria for painful peripheral neuropathy.

Hardship and malnutrition

Following capture, FEPOW were distributed to camps across Asia, where unlike POW in other theatres who were protected by the Geneva conventions, ‘humiliation, gross under-feeding, and savage regimentation were the daily routine’⁵.

The Australian medical officer Sir Edward ‘Weary’ Dunlop wrote of his experiences, indicating that ‘only those who were present can comprehend the seeming hopelessness of it as their bodies wasted and their friends died’. The combination of extreme malnutrition, poor sanitation, a hostile jungle environment, and the high physical workload provided conditions in which diseases were rife. Dunlop characterised the battle against sickness as a ‘new war’, ‘for physical and moral survival’⁹.

The combined effects of deliberate neglect by their captors, and imprisonment in remote areas at the end of supply lines led to severe multifactorial malnutrition. The diet was almost entirely rice-based; grossly deficient in proteins, fats, and vitamins, especially of the B-complex^{5, 7, 10}. Medical officers repeatedly raised the issue with the Japanese, and despite their admission that rations were insufficient, little heed was paid to requests⁷. Vitamins were seen by their captors as inessential ‘luxuries’⁵.

Clinical avitaminosis was anticipated but took around 4 months for signs to appear across the camps⁷. Dunlop reported that the dermatological condition pellagra was the most common disorder, and beriberi occurred in all forms, yet scurvy and vitamin A deficiency were uncommon⁵. In conversation in September 2015, Frankland however recalled only two cases of pellagra in his three and half years of captivity, although he was held in a camp where vegetables were grown for much of this time. From around July 1942 Allied medical officers found FEPOW presenting with a new and unique set of BFS symptoms, which appeared to differ from recognised and more common nutritional disorders.

Burning Feet Syndrome

BFS had been described in situations of malnutrition since 1826¹¹. Known to FEPOW by a variety of colloquial names including ‘happy feet’⁶ and ‘electric feet’¹⁰, the term BFS generally only started to appear in post-war medical literature. Hitherto, BFS had received sporadic attention, generally viewed as an ‘obscure tropical affliction consigned to a few lines of small print in tropical disease texts’³.

Referring to the burden of illness in FEPOW, Simpson wrote that ‘diseases were so gross and numerous in a single individual, that the picture of a single malady was hopelessly confused’⁷. However, a clear picture of BFS is presented in the various post-war accounts, which are remarkably consistent when it is considered that camps were located across Asia and written by doctors with little to no communication with each other, or the outside world. Dunlop gave a succinct symptomatic description of BFS as an ‘intense burning of the feet and an exquisite sensitiveness which scarcely enables them to walk, and they cannot sleep or get any rest’⁵.

Usually occurring 3-5 months after capture, and often following a debilitating bout of infection such as dysentery or malaria⁴, the condition usually started with a dull ache in the feet, specifically over the metatarsal region, and progressed to a ‘severe burning sensation’ and hyperaesthesia⁶. This was followed by a more severe, but intermittent, ‘sharp’ or ‘shooting’ type pain, ‘like toothache’¹² which radiated into the heels and sometimes up the shins⁴. The pains were always distal and symmetrical in distribution and a small minority of cases (6.4%) reported tingling or numbness in their hands^{4,7}.

Glusman commented that the sensory gain phenomena of plantar hyperesthesia and hyperalgesia were universal ‘in all but the earliest and mildest cases’³ and Cruickshank recorded hypersensitivity to pinprick and light touch in about 22.2%⁴. There are multiple

1
2
3 accounts describing how patients would often be found ‘shifting their position from one foot
4
5 to the other’ when standing, ‘usually wincing with pain as each foot touches the ground’⁶.

6
7
8 Simpson described how the hyperaesthesia was ‘superficial in some cases, but usually deep
9
10 muscle tenderness’. With progression, the hyperaesthesia ‘tended to be replaced by
11
12 hypoaesthesia’, often with a patchy distribution of the two⁷.

13
14
15
16 Apart from occasional excessive sweating^{4,5}, accounts generally described no changes in the
17
18 appearance or circulation to the feet when compared to controls. Preservation of tendon
19
20 reflexes was the rule; however there were differences between reports, with some finding an
21
22 increase in deep reflexes^{4,7,12}.

23
24
25 As is the case for many painful neuropathies, pain was particularly troublesome at night and
26
27 often associated with insomnia¹⁰. Patients were described as sitting in a squatting position,
28
29 gripping their toes and massaging their feet ‘for hours on end’^{4,6}. During the day, some men
30
31 reported being comparatively free from pain⁷. Relief was sought by some through immersing
32
33 the feet in buckets of water, whilst others reported that walking helped. These men would
34
35 often walk a ‘long nocturnal vigil’ to minimise the pain, with a characteristic gait
36
37 ‘reminiscent of a man with bad corns’⁷.

38 39 40 41 *Cause of BFS*

42
43
44 No longer an ‘obscure tropical affliction’, BFS was becoming a significant health problem in
45
46 POW camps across the region. Simpson, who found up to 300 of between 1500 to 2000 men
47
48 affected, wrote how ‘in the average case of this syndrome, the malady dominated the whole
49
50 man’s life’⁷. Large numbers of BFS patients were found also found by Cruickshank in
51
52 Singapore¹² and Page in Hong Kong and the Philippines⁶. Others reported a much lower
53
54 prevalence: in 551 cases of dietary associated scrotal dermatitis seen in Changi camp in
55
56
57
58
59
60

1
2
3 Singapore, Frankland, noted only six cases with painful feet¹³. In 1943 he relocated to a camp
4
5 at Blakang Mati (an island off Singapore, now called Sentosa), and in personal
6
7 communication (September 2015) he recalls few cases of burning feet or scrotal dermatitis,
8
9 perhaps attributable to FEPOW being allowed to grow their own vegetables.
10

11
12 Whilst most medical officers had not experienced BFS previously, its presentation as a
13
14 painful dysaesthetic polyneuropathy was described 'remarkably consistently by different
15
16 authors', often in distinct terms from the peripheral paraesthesiae of thiamine deficient beri-
17
18 beri^{10,12,14}. Ascribing a single aetiology for BFS in FEPOW suffering from severe starvation,
19
20 multiple avitaminoses, and co-existing tropical infections was difficult, and no overall
21
22 consensus exists. Nevertheless, most doctors reached the conclusion that the poor diet,
23
24 resulting in a specific B-vitamin deficiency, was to blame, largely because BFS often
25
26 presented alongside signs of other B-vitamin deficiency diseases which were well recognised
27
28 by the doctors. Glusman recorded 58% of BFS patients had an associated vitamin deficiency,
29
30 reporting an 'invariable' association with malnutrition³.
31
32
33

34
35 Simpson noticed other factors which supported this view. The condition was less common in
36
37 officers who possessed money and could supplement their diet with eggs and Kacang Hijau
38
39 beans⁷. Page and Glusman both noted that in groups of FEPOW who received barley in
40
41 addition to their rice ration, there was a 'paucity of new patients, and a low rate of relapse'^{3,6}.
42
43
44

45
46 Compared to the other doctors who wrote about BFS, Cruickshank recognised the condition
47
48 because he was aware of a study by Landor and Pallister in 1935, in which inmates in
49
50 Malayan jails suffering from BFS reported resolution of symptoms once Marmite (a yeast
51
52 extract) and autoclaved yeast, rich in the heat stable factors of the vitamin B-complex, were
53
54 added to their diet^{4,15}.
55
56
57
58
59
60

1
2
3 Once the link to dietary deficiency had been made, the next obstacle was to determine the
4 exact cause. But with scarce resources, and other nutritional deficiencies requiring treatment,
5 rigorous trials of individual vitamins proved challenging.
6
7
8

9 10 *Research*

11
12 Archibald Cochrane had conducted his first successful controlled trial in a POW camp at
13 Salonika, Greece, when he demonstrated that prisoners with wet beri-beri given B-vitamins in
14 the form of yeast reported symptomatic improvement¹⁶. Rudimentary trials of therapy with
15 individual B-vitamins for BFS were attempted in the Far East.
16
17
18
19

20
21 Parenterally or intrathecally administered Thiamine (B1) was investigated to ascertain its
22 effect in the treatment of BFS, although because of limited supplies the doses were probably
23 subtherapeutic^{4,6}. The limited thiamine available was prioritised for cases of ‘frank beriberi’
24 because of the prevailing belief of effectiveness. However, in six cases of BFS treated by
25 Cruickshank there was no recorded improvement⁴, and Glusman found it ‘not effective’³.
26
27
28
29

30
31 Nicotinic acid (B3) was tried and whilst effective at treating pellagra, which regularly
32 coexisted, appeared ineffective for BFS³. However, Cruickshank reported that 98/119
33 patients treated with intravenous Nicotinic acid for 5-10 consecutive days showed at least
34 some improvement⁴. Churchill and Frankland both reported similar findings in a small
35 number of cases^{10,13}.
36
37
38
39

40
41 A paper from India reported that pantothenic acid (B5) rapidly cured 53 patients of BFS¹⁷.
42 However, a 1957 clinical trial conducted in a rural community in Ceylon (Sri Lanka) found
43 no benefit from pantothenic acid in BFS¹⁸.
44
45
46
47

48 49 *Treatment and prognosis*

50
51
52
53
54
55
56
57
58
59
60

1
2
3 Treatment of BFS varied and depended largely on the meagre resources available in
4 individual POW camps. In general, when permitted by the Japanese, patients avoided duties
5 involving standing or walking. In severe cases bedrest was advised, however the temporary
6 symptomatic relief afford by walking, especially at night, made this difficult to enforce.
7
8 When the Japanese started to relocate many POW to labour camps after 1943 follow up of
9 patients became impossible.
10
11
12
13
14
15
16

17 Once the link to B-vitamins was made, sources such as rice-polishings, soya beans, and green
18 leaves were used to supplement diets when possible⁴. If appropriate nutrition could be
19 provided, prognosis of BFS was good. By July 1943, after an improvement in diet in
20 Cruickshank's camp at Changi (Singapore), the disease had almost disappeared⁴. In his series
21 of 27 cases however, Churchill reported that oral administration of Marmite and nicotinic
22 acid were not associated with symptomatic improvement, but in 'milder cases rice-polishings
23 seemed to be effective'¹⁰. Simpson reported that a daily administration of Kacang Hijau
24 beans (a legume used extensively in the Far East) improved symptoms in 4-6 weeks⁷, and that
25 after a week or two of general dietary supplementation with these beans no further new cases
26 of BFS developed.
27
28
29
30
31
32
33
34
35
36
37
38
39

40 After liberation in 1945 symptoms of nutritional neuropathies soon resolved in most FEPOW,
41 however BFS symptoms were reported to persist in a small group of patients after repatriation
42 despite resumption of an adequate diet⁴. A 1982 follow up study investigated the health of
43 898 surviving FEPOW¹⁴. Of these, 49 (5.5%) were classified as having a persistent
44 nutritional neuropathy attributable to their FEPOW period. A further 38 (4.2%), had possible
45 signs of asymptomatic nutritional neurological damage. 24 of the 49 patients had a peripheral
46 neuropathy, of whom 11 (1.2%) had symptoms consistent with persisting BFS. Of the 24
47 patients, the neuropathy was mainly sensory in eight, motor in six, and mixed in 10, with the
48 lower rather than upper limbs usually affected. Tricyclic antidepressants were tried as a
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 therapeutic intervention, and ‘sometimes relieved the pain, at least partly’¹. Table 1 shows
4
5 two examples of cases which illustrate these presentations¹⁴.
6
7

8 *Nowell Peach*

9

10
11 Several retrospective accounts of BFS were published in the immediate post-war period, and
12
13 whilst detailing with remarkable consistency the clinical presentation of BFS, the conditions
14
15 of captivity meant the authors were unable to support these accounts with contemporaneous
16
17 documentation or medical record⁹. Even when records were made, the lack of cooperation
18
19 from the Japanese meant it was unlikely they would survive captivity. Glusman recorded
20
21 detailed neurological examinations but his original case records were destroyed³.
22
23

24
25 However, there was one exception. Flight Lieutenant Nowell Peach was a medical officer
26
27 serving with the Royal Air Force Volunteer Reserve who had been taken prisoner in Java at
28
29 the end of February 1942⁸. He recorded detailed clinical notes on a form, including
30
31 neurological examination, of 54 FEPOW presenting with BFS at Tandjong Priok camp
32
33 (Jakarta, Indonesia) between November 1942 and March 1943.
34
35

36
37 Figures 1, 2, and 3 show three examples of these unique records, which he managed to
38
39 secrete from the Japanese. Peach survived the war and became a General Practitioner
40
41 Surgeon. He revealed the existence of the original notes when interviewed by the FEPOW
42
43 oral history project half a century later. The history and some medical aspects of these
44
45 records have been reviewed before¹²; here we analyse the pain and related symptoms and
46
47 signs which Peach reported.
48
49

50
51 Peach’s records comprise cases seen between late 1942 and early 1943 which coincided with
52
53 the period when ‘clearest and least complicated picture’ of BFS was seen⁵. BFS was first
54
55 recognised in July 1942, the incidence increased over the summer and had become
56
57
58
59
60

1
2
3 widespread by October 1942. However, from 1943 there was a generalised movement of
4
5 FEPOW to provide slave labour for the large scale construction projects such as the railways.
6
7 Simpson explained that during ‘this hectic, hopeless period, it was impossible to follow up in
8
9 detail the fate of these cases’. After the completion of these projects in 1944/45, the
10
11 survivors were ‘fed better and worked for shorter periods’, and knowledge of the prophylactic
12
13 value of diet prevented further cases of BFS⁷.
14
15

16
17 In order to elucidate BFS, we have analysed 53 of Peach’s case files (one being omitted due
18
19 to illegibility). The patients had a mean age of 29 years (range 20-41) and had experienced
20
21 symptoms for a mean of 13 weeks (range 2-56). Forty-seven (89%) patients reported pain
22
23 intensity categorised as ‘moderate’ or ‘severe’. Peach later recorded that 26 (49%) of cases
24
25 were “improving” with only five (9%) progressing.
26
27

28
29 Appetite was recorded as ‘normal’ in 34 (64%) patients, with the same number being given a
30
31 full ration; 25 (47%) were receiving extra food, including milk and fruit. The documentation
32
33 of appetite, ration and additional nutrition, alongside the clinical picture, demonstrate that
34
35 Peach recognised the link between nutrition and BFS. He notes in three case files that the
36
37 condition had improved after dietary supplementation with eggs and soya beans.
38
39

40
41 Thirty-one (58%) cases were recorded as having weight loss, and other features of nutritional
42
43 deficiency were often recorded in many patients, the most frequent being those in the lips,
44
45 scrotum, tongue and eyes, matching the reports from other camps. A history of tropical
46
47 infection was frequently found in BFS cases with 13 (25%) recording malaria, 24 (45%)
48
49 dengue fever and 23 (43%) dysentery. Peach noted that in six (11%) cases the presentation of
50
51 BFS coincided with an infection.
52
53

54
55 *Symptoms and physical examination*
56
57
58
59
60

1
2
3 Peach wrote a brief symptom descriptor summary for each case. These are 'free text', and
4
5 thus do not constitute the current systematic approach of recording neuropathic pain
6
7 descriptors^{19,20,21}. Nevertheless, these descriptors do afford insight into the symptoms
8
9 reported by FEPOW patients with BFS and are similar to those described in other reports
10
11 (Table 2).
12
13

14
15 A unique aspect of Peach's files was a brief neurological examination, of which the
16
17 frequency of neurological signs is recorded in Table 3. He assessed tendon reflexes using a
18
19 tendon hammer made for him by Army engineers (Figure 4). The solitary example of
20
21 hyperaesthesia recorded was found on the soles and the tips of the fingers and toe, and
22
23 presented alongside reduced sensation in the legs.
24
25

26 27 *Neuropathic Pain*

28
29 Although these FEPOW may have also suffered feet pain from another origin (e.g.
30
31 osteoarthritis), this is unlikely to have been the case in most patients, and based on the
32
33 description and location it might be thought intuitively that the pain reported in BFS was
34
35 strongly suggestive of neuropathic pain. We scrutinised Peach's records to ascertain to what
36
37 extent this would be the case when judged by current diagnostic criteria (Figure 5)²².
38
39

40
41 Neuropathic pain is defined by the International Association for the Study of Pain (IASP) as
42
43 'pain caused by a lesion or disease of the somatosensory system'²³. In practice, applying this
44
45 definition essentially revolves around two questions: firstly, whether there is a lesion or
46
47 disease of the somatosensory system? Secondly, is the pain neuropathic? The second question
48
49 is often addressed by determining whether the pain is located in a neuroanatomically
50
51 plausible location and by the presence of certain symptom descriptors known to be associated
52
53 with neuropathic pain. To augment the clinical utility of the grading system it follows the
54
55
56
57
58
59
60

1
2
3 pattern of a conventional clinical consultation. Herein we have attempted, for the first time, to
4
5 apply this grading system in a historical context.
6
7

8 For “possible” neuropathic pain there is a requirement for: 1. A history of relevant
9
10 neurological lesion or disease and 2. The pain distribution is neuroanatomically plausible.
11
12 The history of severe malnutrition clearly raises the possibility of a relevant neurological
13
14 disease in the form of a peripheral polyneuropathy, particularly given the distal symmetrical
15
16 pattern of symptoms implicit in Peach’s files. The grading system also permits the use of
17
18 symptom descriptors which, whilst not pathognomonic for neuropathic pain, are strongly
19
20 associated with it. A range of questionnaires have been designed and validated for this
21
22 purpose^{20,21}. The Neuropathic Pain Symptom Inventory (NPSI) is a symptom inventory
23
24 which evaluates the presence and severity of 10 different neuropathic pain descriptors¹⁹.
25
26 Whilst it is not possible to fully utilise any of these instruments retrospectively because not
27
28 all the required questions were asked or quantified by Peach, it is possible to ask to what
29
30 extent the symptom descriptors recorded by Peach also feature in the NPSI or the DN4 pain
31
32 questionnaire. 52/53 (98%) of Peach’s files document either “pain” or “ache” and 49 (92%)
33
34 had at least one of the DN4 descriptors of ‘burning’ or ‘electric shock’ recorded. Peach
35
36 recorded most frequently the NPSI descriptors of ‘shooting’ in 38 (72%) of patients, ‘aching’
37
38 in 37 (70%) and ‘burning’ in 30 (57%). According to the validation paper for the NPSI¹⁹, the
39
40 symptom quality of ‘burning’ is reported by 70% of patients with neuropathic pain and
41
42 symptoms similar to ‘shooting’ (‘stabbing’ or ‘electric shocks’) in ~60%; with ‘aching’ not
43
44 included due to it being frequently described in other types of pain. ‘Burning’ or ‘shooting’
45
46 (or similar) also feature in all 5 of the major diagnostic symptom questionnaires for
47
48 neuropathic pain²⁰. Four reports of BFS in FEPOWs all described patients suffering ‘burning’
49
50 and ‘shooting pain’; for example Cruickshank documented 58.6% of the 500 cases he
51
52 assessed reported ‘sharp’, ‘shooting’ pain^{3,4,6,7}. The pain symptom descriptors which feature
53
54
55
56
57
58
59
60

1
2
3 in Peach's and other accounts are also prominent in modern day profiles of patients with
4
5 painful neuropathies including HIV-associated sensory neuropathy²⁴ and diabetic neuropathy,
6
7 which has been reported as "distal, symmetrical, often associated with nocturnal
8
9 exacerbations, and commonly described as prickling, deep aching, sharp like an electric
10
11 shock, and burning"²⁵; corresponding with the descriptions given of BFS.

12
13
14 For a grading of "probable" neuropathic pain there is an additional requirement relating to
15
16 sensory examination: "sensory signs in the same neuroanatomically plausible distribution as
17
18 pain"²². In Peach's report 37 (70%) patients had "pain" recorded as a symptom descriptor of
19
20 whom only 6 (16%) had sensory abnormalities revealed by his physical examination. Sixteen
21
22 (30%) patients had both pain and paraesthesiae recorded of whom 10 (62.5%) of those had
23
24 sensory abnormalities recorded. It might be reasonably expected that a higher proportion of
25
26 these patients might have had detectable abnormalities in sensation. However, Peach was not
27
28 a trained neurologist and was conducting his examinations in adverse conditions with
29
30 improvised instruments. For a grading of "definite" neuropathic pain a diagnostic test
31
32 confirming a lesion or disease of the somatosensory system is required- clearly beyond what
33
34 was feasible in captivity.
35
36
37
38
39

40 The recent NeuPSIG criteria were developed to exclude cases without evidence of a
41
42 neurological lesion from clinical examination or laboratory test, and this situation is found in
43
44 many clinical conditions (i.e. small fiber neuropathies, radiculopathies), where it could be
45
46 difficult to reach the level of "probable" neuropathic pain without using sophisticated clinical
47
48 or laboratory tests. Therefore, on the basis of the foregoing discussion the pain suffered by
49
50 the majority of Peach's patients with BFS was possibly neuropathic in origin, but it is not
51
52 appropriate to ascribe a level of certainty of "probable" in most cases.
53
54

55
56
57 *Conclusion*
58
59
60

1
2
3 Flight Lieutenant Nowell Peach's long hidden clinical records of individual BFS patients
4
5 have permitted scrutiny of whether their pain was likely to be neuropathic in origin when
6
7 analysed using 21st century standards and definitions. When analysed alongside other papers
8
9 from the era, a number of conclusions can be drawn. The aetiology of BFS is almost certainly
10
11 nutritional, but the exact dietary factor is less certain. FEPOW were grossly malnourished
12
13 with multiple vitamin deficiencies. BFS is described as a unique presentation, seen both
14
15 alongside, and independently from, other deficiencies of B-vitamins. The limited trials which
16
17 were conducted in the POW camps did not draw any positive results about a specific vitamin
18
19 deficiency. However, when treated with a general improvement in diet, or food sources rich
20
21 in multiple B-vitamins, the condition improved. It is therefore most likely that BFS, a
22
23 condition which can now be described as a "possible" neuropathic pain syndrome,
24
25 represented a deficiency of multiple B-vitamins. After seventy years, the data painstakingly
26
27 gathered under the worst of circumstances, by a determined medic, have proved to be of
28
29 interest and value in the modern diagnostics of neuropathic pain.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legend

Figure 1: An example of a case record written by Nowell Peach as a FEPOW in 1942.

The complaint reads, 'Constant ache in distal part of foot, with intermittent pain of a shooting nature. Any change of occupation gives temporary relief from the shooting pain, which came on again after about an hour in any one position. No superficial burning. Shooting pain is "like a dentist touching a nerve".'

Figure 2: An example of a case record written by Nowell Peach as a FEPOW in 1942.

The complaint reads, 'Burning under metatarsal heads and heels, and shooting pains in feet. Pins and needles in fingers.'

Figure 3: An example of a case record written by Nowell Peach as a FEPOW in 1942.

The complaint reads, 'Burning of soles and aching in toes. Continual. Worse at night. Not relieved by walking. Feels as though he has tight boots on'.

Figure 4: Surgical instruments used by FEPOW in captivity, including Nowell Peach's homemade tendon hammer inscribed with 'Java 1942'.

Figure 5: Diagram showing current criteria required for diagnosing neuropathic pain.

References

1. Parkes M, GV Gill. *Captive Memories: Far East POWs & Liverpool School of Tropical Medicine*. 1st ed. Palatine; 2015.
2. Tanaka Y. *Hidden Horrors: Japanese War Crimes In World War II*. 1st ed. Westview Press; 1997.
3. Glusman M. The Syndrome of "Burning Feet" (Nutritional Melalgia) as a Manifestation of Nutritional Deficiency. *American Journal of Medicine*. 1947; 3: 211-233.
4. Cruickshank E. Painful feet in prisoners of war in the Far East. *The Lancet*. 1946; 248: 369-372.
5. Dunlop E. Medical Experiences in Japanese Captivity. *British Medical Journal*. 1946; 2: 481-486.
6. Page J. Painful-feet syndrome among prisoners of war in the Far East. *British Medical Journal*. 1946; 2: 260-262.
7. Simpson J. A syndrome of painful feet and retrobulbar neuritis occurring amongst British prisoners of war in the Far East. MD Thesis, UK: University of Durham; 1946.
8. Parkes M. Alfred Nowell Hamilton Peach. *British Medical Journal*. 2012; 344: e17649.
9. Dunlop E. *The War Diaries of Weary Dunlop*. 1st Ed. Australia: Thomas Nelson; 1986.
10. Churchill M. Dietary deficiency diseases among prisoners of war. *Journal of the Royal Army Medical Corps*. 1945; 85: 294-298.
11. Grierson J. On the burning in the feet of natives. *Transactions of the Medical and Physical Society of Calcutta*. 1826; 2: 275-280.
12. Welch E, Peach N, Parkes M, Gill GV. Burning feet syndrome: An old tropical syndrome revisited. *Ann Trop Med Public Health*. 2013; 6: 65-70.
13. Frankland A. Deficiency scrotal dermatitis in P.O.Ws in the Far East. *British Medical Journal*. 1948; 1: 1023-6.
14. Gill GV, Bell DR. Persisting nutritional neuropathy amongst former war prisoners. *Journal of Neurology, Neurosurgery, and Psychiatry*. 1982; 45(10): 861-865.
15. Landor JV, Pallister RA. Avitaminosis B2. *Transactions of the Royal Society of Medicine and Hygiene*. 1935; 29: 121-134.
16. Gerris J. The legacy of Archibald Cochrane: from authority based towards evidence based

- 1
2
3 medicine. Facts, Views and Vision. Issues in Obstetrics, Gynaecology and Reproductive Health.
4
5 2011; 3: 233-237.
6
7 17. Gopalan C. The burning feet syndrome. Indian Medical Gazette. 1946;81:22-26.
8
9 18. Bibile SW, Lionel NDW, Dunuwille R, Perera G. Pantothenol and the burning feet syndrome.
10
11 British Journal of Nutrition. 1957;11: 434-439.
12
13 19. Bouhassira D, Attal N, Fermanian J, Alchaar H, Gautron M, Masquelier E et al. Development and
14
15 validation of the Neuropathic Pain Symptom Inventory. Pain. 2004;108: 248-57.
16
17 20. Bennett MI, Attal N, Backonja MM, Baron R, Bouhassira D, Freynhagen R et al.. Using screening
18
19 tools to identify neuropathic pain. Pain. 2007; 127:199-203.
20
21 21. Haanpää M, Attal N, Backonja M, Baron R, Bennett M, Bouhassira D et al. NeuPSIG guidelines
22
23 on neuropathic pain assessment. Pain. 2011; 152(1): 14–27.
24
25 22. Finnerup NB, Haroutounian S, Kamerman P, Baron R, Bennett DL, Bouhassira D, et al.
26
27 Neuropathic pain: An updated grading system for research and clinical practice. Pain. Epub 2016 Apr
28
29 28.
30
31 23. Jensen TS, Baron R, Haanpää M, Kalso E, Loeser JD, Rice AS et al.. A new definition of
32
33 neuropathic pain. Pain. 2011; 152(10): 2204-2205.
34
35 24. Phillips TJ, Brown M, Ramirez JD, Perkins J, Woldeamanuel YW, Williams AC et al..Sensory,
36
37 psychological, and metabolic dysfunction in HIV-associated peripheral neuropathy: A cross-sectional
38
39 deep profiling study. PAIN. 2014; 155(9): 1846-1860.
40
41 25. Tesfaye S, Boulton AJ, Dyck PJ, Freeman R, Horowitz M, Kempner P et al. Diabetic neuropathies:
42
43 update on definitions, diagnostic criteria, estimation of severity, and treatments. Diabetes Care. 2010;
44
45 33: 2285–93.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Two cases of persistent symptomatic nutritional neurological neuropathies in FEPOW followed up in 1982 and published in the Journal of Neurology, Neurosurgery, and Psychiatry by Gill and Bell.

Case One	This 62-year-old-man was assessed 35 years after release, in 1980. He was interned in Java and the Celebes where he suffered malaria, dysentery, wet beriberi and paraesthesiae of the legs with loss of sensation of ground contact. His vision was also blurred. He regained good health on release, but paraesthesiae of his feet continued. When examined at Liverpool he had reduced pain and temperature sensation bilaterally, to mid-thigh level. Nerve conduction studies confirmed a 'patchy peripheral nerve lesion consistent with nutritional neuropathy'.
Case Two	A 57-year-old ex-FEPOW was seen in 1975. He had been interned in Rangoon during the war, had suffered chronic dysentery, and lost 32 kg in weight. He developed paraesthesiae peripherally, burning of the hands and legs, unsteadiness in the dark, deafness and poor vision. After the war his symptoms gradually improved, but paraesthesiae, unsteadiness and burning feet persisted. He was regularly kept awake at night by the pains in his feet, and he would walk around the house to gain relief. Examination revealed marked signs of a sensory neuropathy in the legs, as well as evidence of posterior column loss.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2: Table showing frequency of symptom descriptors in case records recorded by Nowell Peach in FEPOW. Results are recorded in absolute number, and in brackets as a percentage.

Location of pain	Feet	47 (89%)
	Metatarsals	20 (38%)
	Legs	19 (26%)
	Toes	13 (25%)
	Hands	7 (13%)
	Shooting	38 (72%)
Character of pain	Aching	37 (70%)
	Burning	30 (57%)
	During night	8 (15%)
Pain worsens	With stillness or prolonged standing	3 (6%)
	With walking	10 (19%)
Pain improves	Being preoccupied with work	3 (6%)
	Immersing feet in cold water	2 (4%)

Table 3: Table showing frequency of neurological signs in 53 case records recorded by Nowell Peach in FEPOW. Results are recorded in absolute number, and in brackets as a percentage.

Sensation	Anaesthesia to pin prick	8 (15%)
	Anaesthesia to light touch	5 (9)
	Anaesthesia to pressure	3 (6%)
	Hyperaesthesia	1 (2%)
Motor	Weakness	6 (12%)
	Hyperreflexia knee reflex	4 (8%)
	Hyporeflexia knee reflex	9 (17%)
	Hyperreflexia ankle reflex	1 (2%)
	Hyporeflexia ankle reflex	14 (26%)

Complaint: <i>Burning under metatarsal heads & heels, and shooting pains in feet. Pins & needles in fingers describes the pain in feet as being like that after freezing. Standing on cold floor makes worse. Eased by pouring on cold water.</i>		
Duration: <i>14 months.</i>	Intensity: <i>Medium.</i>	Progress: <i>Improving.</i>
Other Deficiency Signs:		
Lips - <i>at time of onset.</i> Tongue - Nails - Mouth - Throat -	Scrotum - Nails - Nasolabial - Pigment <i>Forehead of face.</i> Anæmia - Oedema - <i>And slightly.</i>	Loss of Weight - Eyes <i>Vision good. Hms 5/10. No optic atrophy.</i> Aesthesia + Paresis + Giddiness -
M. Malaria + 3. (<i>? type</i>) Dysentery <i>Bacillary.</i>	V.D. - Dengue +	Others - Smoking <i>Nif.</i>
Appetite: <i>Good.</i> Diet: <i>Rice Full Ration.</i>	Indigestion: <i>Nif.</i>	Stools: <i>N.</i> Extras <i>5/12. Milk. Meat. Yeast. Fruit.</i>
Feet (Condition)		
Deformity: <i>Nif.</i> Vasculature: <i>N.</i> Pulse, Ant. Tib. <i>Palp.</i>	Skin: <i>N.</i> Sweating: <i>N.</i>	
Sensory System		
ANAESTHESIA (Pin Prick: <i>hypoaesthesia inner side of sole. Otherwise N.</i>) (Stroke : <i>N.</i>) (Pressure : <i>N.</i>)	PARAESTHESIA (Superficial: <i>Nif.</i>) (Deep: <i>Nif.</i>)	
Motor System:		
KJ: - AJ: - Wasting: <i>Nif now. Was + at onset.</i> Paresis: <i>gait normal now, was off legs.</i>	CNS Plantar: <i>N.</i> Romberg: <i>N.</i> Pupil: <i>N.</i> Nystagmus: <i>N.</i>	
Cardiovascular Pulse: (Rate: <i>72.</i>) (Rhythm: <i>Reg.</i>) (Tension: <i>N.</i>) Femoral Artery: <i>No tones</i>		

Figure 1: An example of a case record written by Nowell Peach as a FEPOW in 1942. The complaint reads, 'Constant ache in distal part of foot, with intermittent pain of a shooting nature. Any change of occupation gives temporary relief from the shooting pain, which came on again after about an hour in any one position. No superficial burning. Shooting pain is "like a dentist touching a nerve"'.
No superficial burning. Shooting pain is "like a dentist touching a nerve".

223x184mm (72 x 72 DPI)

<u>Complaint.</u> Constant ache in distal part of foot, with intermittent pain of a shooting nature. Any change of occupation gives temporary relief from the shooting pain, which came on again after about an hour in any one position. No superficial burning. Shooting pain is "like dentist touching a nerve".		
<u>Duration:</u> 6 weeks.	<u>Intensity:</u> Severe at times.	<u>Progress:</u> Stationary.
<u>Other Deficiency Signs:</u>		
<u>Lips.</u>	} N.	<u>Scrotum.</u> Eczema 6 months.
<u>Tongue.</u>		<u>Nails.</u> N.
<u>Gums.</u>		<u>Nasolabial.</u> N.
<u>Mouth.</u>		<u>Pigment.</u> Nil
<u>Throat.</u>		<u>Anaemia.</u> Nil
		<u>Oedema.</u> Nil
<u>P.H.</u> Malaria -		<u>V.D.</u> -
		<u>Dengue</u> "Wors" 12/day until 4/20, probably because of dentistry
		<u>Others.</u> Dyspepsia -
<u>Dysentery</u> -		
<u>Appetite:</u> Poor.	<u>Indigestion:</u> +	<u>Stools:</u> N.
<u>Diet:</u> Rice 1/2 Ration.	<u>Extras</u> Extra Stew ration, Eggs 2/week until 2 months ago	
<u>Feet (Condition)</u>		
<u>Deformity</u> Nil		<u>Skin</u> N.
<u>Vascularity</u> Nil		<u>Sweating</u> N.
<u>PULSE (ANTI TIB.)</u> Nil		
<u>Sensory System</u>		
<u>ANAESTHESIA</u> (Pin Prick. Nil) } Anaesthesia		<u>PARAESTHESIA</u> Superficial. Drows to metatarsal heads.
(Stroke. Nil)		Deep. N.
(Pressure. N.)		
<u>Motor System.</u> KJ N		<u>GNS</u> Plantar N.
AJ N		Rhomberg N.
Wasting N		Pupil N.
Paresis N		Nystagmus N.
<u>Cardiovascular.</u> (Rate. 112.		<u>Size.</u> N.
(Rhythm. R.		<u>Heart</u> Sounds. N.
(Tension. N.		<u>Exercise Tolerance.</u> N.

Figure 2: An example of a case record written by Nowell Peach as a FEPOW in 1942. The complaint reads, 'Burning under metatarsal heads and heels, and shooting pains in feet. Pins and needles in fingers. Describes pain in feet as being like that after freezing. Standing on cold floor makes worse. Eased by pouring on cold water'.!! +

226x216mm (72 x 72 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

Complaint: *Burning of soles & aching in toes. Continual. Worse at night. Not relieved by walking. Feels as though he has tight boots on.*

Duration: *2 months.* Intensity: *Severe.* Progress: *Progressing.*

Other Deficiency Signs:

Lips: <i>Perfect.</i>	Oedema: -	Scrotum: -	Eyes: -
Tongue: <i>Marginal glossitis</i>	Nails: -	Nasolabial: -	Asthemia: +.
Mouth: <i>+</i>	Pigment: -	Anaemia: -	Paresis: -
Gums: <i>N. -</i>			Giddiness: -
Throat: <i>N. -</i>			Loss of Weight: <i>> 1 stone.</i>

P.H. *Malaria + 8 times.* V.D. - *Others.*
Dysentery. - *Dengue. 10/12 ago.* *Smoking. 1/2 mod.*

Appetite: *Good.* Indigestion: - *Stools: N.*

Diet. *Rice. Full ration.* Extras. *eg 3 times a week.*

Feet (Condition)

Deformity: *? 1st flattening.* Skin: *N.*

Vascularity: *N.* Sweating: *N.*

Pulse. Ant. Tib. *N.*

Sensory System:

ANAESTHESIA { *Pin Prick: {not felt in foot}* } Superficial *N.*
 { *Stroke: {base of legs + wrist}* }
 { *Pressure: N.* } PARAESTHESIA Deep *Calves.*

Motor System

KJ: <i>N.</i>	CNS
AJ: <i>N.</i>	Plantar: <i>N.</i>
Wasting: <i>? calves.</i>	Rhomberg: <i>N.</i>
Paresis: -	Pupil: <i>N.</i>
	Nystagmus: <i>N.</i>

Cardiovascular, Pulse: (Rate: *84*) Heart: (Size: *N.*)
 (Rhythm: *R.*) (Sounds: *N.*)
 (Tension: *Low.*) Exercise Tolerance: *N.*

Femoral Artery: *N.*

34 Figure 3: An example of a case record written by Nowell Peach as a FEPOW in 1942. The complaint reads,
 35 'Burning of soles and aching in toes. Continual. Worse at night. Not relieved by walking. Feels as though he
 36 has tight boots on'.!! †

37 220x176mm (72 x 72 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Figure 4: Surgical instruments used by FEPOW in captivity, including Nowell Peach's homemade tendon hammer inscribed with 'Java 1942'.

152x95mm (200 x 200 DPI)

ew Only




Figure 5: Diagram showing current criteria required for diagnosing neuropathic pain.!! †

186x158mm (72 x 72 DPI)