

Are rheumatoid factor, anti-citrullinated, and anti-carbamylated protein antibodies linked by post-translational modification of IgG?

Are rheumatoid factor, anti-citrullinated, and anti-carbamylated protein antibodies linked by post-translational modification of IgG?

David Hutchinson MD^{1 2}, Daniel Murphy BmBs¹, Alex Clarke BSc² and Paul Eggleton MPhil, PhD²

¹ Department of Rheumatology, Royal Cornwall Hospital Trust, Truro, Cornwall TR1 3LQ

² University of Exeter Medical School, Exeter, Devon EX1 2LU

Competing interests

The authors declare they have no competing interests.

Acknowledgements

Support for this project was provided by the Cornwall Arthritis Trust.

Each individual named as author meets your criteria for authorship. It has not been, as is not being, submitted elsewhere. There are no financial conflicts of interest.

Correspondence to:

Dr. Daniel Murphy,

Department of Rheumatology, Royal Cornwall Hospital Trust, Truro, Cornwall TR1 3LQ.

E-mail: Daniel.Murphy8@nhs.net

Phone: 01872 250000

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as an 'Accepted Article', doi: 10.1002/art.39832

© 2016 American College of Rheumatology

Received: May 31, 2016; Accepted: Jul 28, 2016

This article is protected by copyright. All rights reserved.

Are rheumatoid factor, anti-citrullinated, and anti-carbamylated protein antibodies linked by post-translational modification of IgG?

To the editor. We read with interest the recent article by Koppejan et al [1] where a strong association between rheumatoid arthritis (RA) and triple positivity for rheumatoid factor (RF), anti-citrullinated protein antibodies (anti-CCP) and anti-carbamylated protein antibodies (anti-CarP) (odds ratio (OR) 194 95% CI (23-1609), $p < 0.0001$) was observed. The authors suggest that these findings offer new insights into the evolution of autoimmunity in preclinical RA.

We suggest that misfolding of immunoglobulin G (IgG) in the lung has the potential to initiate seropositive RA, sequentially generating the three RA autoantibodies observed in the above study. We propose a pathway relevant to smokers and individuals carrying the HLA class II shared epitope (SE), explaining the very high prevalence of smoking and the SE noted in RA patients [1]. Immunoglobulin G is a plausible “universal” antigen for all three known RA autoantibodies, as although there are a plethora of proteins that can act as antigens to both anti-CCP and anti-CarP antibodies, RF only has one antigen.

Rheumatoid factor stage: smoking-induced bronchial associated lymphoid tissue (BALT) and an increased generation of IgG

Smoking associates with the development of BALT, a process associated with seropositive RA [2]. This plasma cell abundance potentiates increased IgG production, evidenced by OR for the development of RF-positive RA in the lowest compared to the highest tertiles of IgG distribution in individuals prior to disease development being 2.22 (95% CI 1.16, 4.26) [3]. IgA RF is present in the sera of individuals who subsequently develop seropositive RA up to 15 years before disease development and is observed

Are rheumatoid factor, anti-citrullinated, and anti-carbamylated protein antibodies linked by post-translational modification of IgG?

prior to both ACPA and anti-CarP [4]. We propose that this initial autoantibody response may represent a signal of increased lung IgG production, with IgA RF antibodies responding to native IgG protein.

Citrullinated IgG heavy chain (clgGH) generation stage: misfolding induced by citrullination

Protein misfolding is enhanced by citrullination [5] and IgG misfolding has been suggested as a model of RA autoimmunity with the generation of IgGH, a 50 kilodalton (kDa) post-translationally modified IgG heavy chain fragment [6]. Growth arrest and DNA damage-inducible gene 34 has an important role in the unfolded protein response and significantly relates to the presence of ACPA in RA [7], suggesting ACPA generation associates with protein misfolding. Given that the lung is a site of citrullination [8], we suggest that the intracellular citrullination of IgGH occurs in lung plasma cells resulting in the development of the novel neoantigen clgGH.

Anti-citrullinated protein antibody stage: shared epitope dependant pathway of clgGH cellular escape and extra-cellular antigenicity

Under normal circumstances misfolded proteins are processed to peptides. Interestingly, it has been suggested the transport of IgGH out of cells occurs as a complex with SE alleles as this specific complex is recognised by autoantibodies occurring in RF-positive RA patient's sera, but not by autoantibodies in RF-positive sera from individuals without RA [6]. We suggest further that the process of IgGH-SE binding is greatly enhanced by citrullination, given that citrullination of calreticulin enhances SE

Are rheumatoid factor, anti-citrullinated, and anti-carbamylated protein antibodies linked by post-translational modification of IgG?

binding by 10000-fold compared to the native protein [9]. Once the clgGH is extracellular, specific ACPA are likely to develop, with cross reactivity giving rise to a positive anti-CCP result.

Anti-carbamylated protein antibody stage: smoking induced carbamylation of clgGH

Cigarette smoking associates with raised serum levels of thiocyanate and myeloperoxidase-catalyzed oxidation of thiocyanate results in increased carbamylation in smokers [10]. Recently, vimentin carbamylation was noted in chronic obstructive pulmonary disease lung samples [8]. Ergo, we suggest that IgGH is also carbamylated in the lung. Importantly the carbamylation of clgGH may greatly diminish the clearance of this antigen locally by the classical complement pathway as carbamylation of IgG abrogates activation of the classical complement pathway [11]. This process may give rise to large quantities of uncleared carbamylated clgGH entering the circulation and generation of a positive anti-CarP response.

RA initiation stage: Circulation of carbamylated clgGH

The synovial fluid of RA patients has been observed to contain specific IgG which has the potential to stimulate an intense antibody formation when injected into mice which is not observed with native IgG [12]. This response was characterized by high and sustained levels of IgG1 antibodies with RF activity. We propose that the generation in the lung of carbamylated clgGH leads to circulation of this antigen with the potential to enter the joint and suggest that carbamylated clgGH initiates a cascade of B-cell

Are rheumatoid factor, anti-citrullinated, and anti-carbamylated protein antibodies linked by post-translational modification of IgG?

activation, with the development of tertiary lymphoid tissue, further IgG misfolding and RA development, explaining the authors observation that when all three autoantibodies are present there is a strong association with RA.

References

1. Koppejan H, Trouw LA, Sokolove J, Lahey LJ, Huizinga TJ, Smolik IA, et al. Anti-carbamylated protein antibodies in rheumatoid Arthritis, first degree relatives and controls: comparison to anti-citrullinated protein antibodies. *Arthritis Rheum* 2016 doi: 10.1002/art.39664. [Epub ahead of print]
2. Rangel-Moreno J, Hartson L, Navarro C, Gaxiola M, Selman M, Randall TD. Inducible bronchus-associated lymphoid tissue (iBALT) in patients with pulmonary complications of rheumatoid arthritis. *J Clin Invest* 2006;116:3183-94.
3. Aho K, Heliövaara M, Knekt P, Reunanen A, Aromaa A, Leino A, et al: Serum immunoglobulins and the risk of rheumatoid arthritis. *Ann Rheum Dis* 1997;56:351-56.
4. Brink M, Hansson M, Mathsson-Alm L, Wijayatunga P, Verheul MK, Trouw LA, et al. Rheumatoid factor isotypes in relation to antibodies against citrullinated peptides and carbamylated proteins before the onset of rheumatoid arthritis. *Arthritis Res Ther* 2016;18:43.

Are rheumatoid factor, anti-citrullinated, and anti-carbamylated protein antibodies linked by post-translational modification of IgG?

5. Tarcsa E, Marekov LN, Mei G, Melino G, Lee SC, Steinert PM. Protein unfolding by peptidylarginine deiminase. Substrate specificity and structural relationships of the natural substrates trichohyalin and filaggrin. *J Biol Chem* 1996;271:30709–16.

6. Jin H, Arase N, Hirayasu K, Kohyama M, Suenaga T, Saito F et al. Autoantibodies to IgG/HLA class II complexes are associated with rheumatoid arthritis susceptibility. *Proc Natl Acad Sci USA* 2014;111:3787-92.

7. Clavarino G, Adriouach S, Quesada JL, Clay M, Chevreau M, Trocme C, et al. Unfolded protein response gene GADD34 is overexpressed in rheumatoid arthritis and related to the presence of circulating anti-citrullinated protein antibodies. *Autoimmunity* 2016;49:172-8.

8. Lugli EB, Correia RE, Fischer R, Lundberg K, Bracke KR, Montgomery AB, et al. Expression of citrulline and homocitrulline residues in the lungs of non-smokers and smokers: implications for autoimmunity in rheumatoid arthritis. *Arthritis Res Ther* 2015;17:9.

9. Ling S, Cline EN, Haug TS, Fox DA, Holoshitz J. Citrullinated calreticulin potentiates rheumatoid arthritis shared epitope signaling. *Arthritis Rheum* 2013;65:618-26.

10. Wang Z, Nicholls SJ, Rodriguez ER, Kumm O, Horkko S, Barnard J, et al. Protein carbamylation links inflammation, smoking, uremia and atherogenesis. *Nat Med* 2007;13:1176-84.

Are rheumatoid factor, anti-citrullinated, and anti-carbamylated protein antibodies linked by post-translational modification of IgG?

11. Koro C, Bielecka E, Dahl-Knudsen A, Enghild JJ, Scavenius C, Brun JG, et al.

Carbamylation of immunoglobulin abrogates activation of the classical complement pathway. *Eur J Immunol* 2014;44:3403-12.

12. Abedi-Valugerdi M, Ridderstad A, al-Balaghi S, Möller E. Human IgG rheumatoid

factors and RF-like immune complexes induce IgG1 rheumatoid factor production in mice. *Scand J Immunol* 1995;41:575-82.

Accepted Article