

## Accepted Manuscript

Title: Protective effects of exogenous and endogenous hydrogen sulfide in mast cell-mediated pruritus and cutaneous acute inflammation in mice

Author: L. Rodrigues E. Ekundi-Valentim J. Florenzano A.R.A Cerqueira T.P. Schmidt K.T. Santos A.G. Soares S.A. Teixeira M.T.C.P. Ribela G. De Nucci M. Wood M. Whiteman M.N. Muscará S.K.P. Costa



PII: S1043-6618(16)30529-1  
DOI: <http://dx.doi.org/doi:10.1016/j.phrs.2016.11.006>  
Reference: YPHRS 3389

To appear in: *Pharmacological Research*

Received date: 31-5-2016  
Revised date: 20-10-2016  
Accepted date: 9-11-2016

Please cite this article as: Rodrigues L, Ekundi-Valentim E, Florenzano J, Cerqueira ARA, Schmidt TP, Santos KT, Soares AG, Teixeira SA, Ribela MTCP, De Nucci G, Wood M, Whiteman M, Muscará MN, Costa S.K.P. Protective effects of exogenous and endogenous hydrogen sulfide in mast cell-mediated pruritus and cutaneous acute inflammation in mice. *Pharmacological Research* <http://dx.doi.org/10.1016/j.phrs.2016.11.006>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**RESEARCH PAPER**

Protective effects of exogenous and endogenous hydrogen sulfide in mast cell-mediated pruritus and cutaneous acute inflammation in mice

Rodrigues L<sup>1</sup>, Ekundi-Valentim E<sup>1</sup>, Florenzano J<sup>1</sup>, Cerqueira ARA<sup>1</sup>, Schmidt TP<sup>1</sup>, Santos KT<sup>1</sup>, Soares AG<sup>1</sup>, Teixeira SA<sup>1</sup>, Ribela MTCP<sup>2</sup>, De Nucci G<sup>1</sup>, Wood M<sup>3</sup>, Whiteman M<sup>3</sup>, Muscará MN<sup>1</sup>, Costa SKP<sup>1\*</sup>

<sup>1</sup>Department of Pharmacology, Institute of Biomedical Sciences, University of Sao Paulo, Sao Paulo, Brazil.

<sup>2</sup>Department of Biotechnology, Institute of Nuclear and Energetic Research (IPEN), Sao Paulo, Brazil.

<sup>3</sup> University of Exeter Medical School, Exeter, UK.

**Correspondence to:**

\*Soraia K P Costa, PhD

Dept. of Pharmacology – ICB, University of São Paulo


Av. Prof. Lineu Prestes, 1524 - Room 326,

Butantan, 05508-900, São Paulo, SP, Brazil.

Telephone / fax: +55 (0)11 3091-7320

E-mail:skcosta@usp.br

## Graphical abstract



## ABSTRACT

The recently described ‘gasomediator’ hydrogen sulfide (H<sub>2</sub>S) has been involved in pain mechanisms, but its effect on pruritus, a sensory modality that similarly to pain acts as a protective mechanism, is poorly known and controversial. The effects of the slow-releasing (GYY4137) and spontaneous H<sub>2</sub>S donors (Na<sub>2</sub>S and Lawesson’s reagent, LR) were evaluated in histamine and compound 48/80 (C48/80)-dependent dorsal skin pruritus and inflammation in male BALB/c mice. Animals were intradermally (i.d.) injected with C48/80 (3 μg/site) or histamine (1 μmol/site) alone or co-injected with Na<sub>2</sub>S, LR or GYY4137 (within the 0.3 - 100 nmol range). The involvement of endogenous H<sub>2</sub>S and K<sub>ATP</sub> channel-dependent mechanism were also evaluated. Pruritus was assessed by the number of scratching bouts, whilst skin inflammation was evaluated by the

extravascular accumulation of intravenously injected  $^{125}\text{I}$ -albumin (plasma extravasation) and myeloperoxidase (MPO) activity (neutrophil recruitment). Histamine or C48/80 significantly evoked itching behavior paralleled by plasma extravasation and increased MPO activity.  $\text{Na}_2\text{S}$  and LR significantly ameliorated histamine or C48/80-induced pruritus and inflammation, although these effects were less pronounced or absent with GYY4137. Inhibition of endogenous  $\text{H}_2\text{S}$  synthesis exacerbated C48/80-induced responses, whereas the blockade of  $\text{K}_{\text{ATP}}$  channels by glibenclamide did not. High-performance liquid chromatography coupled to tandem mass spectrometry (HPLC-MS/MS) revealed that  $\text{Na}_2\text{S}$  and LR, but not GYY4137, significantly attenuated C48/80-induced histamine release from rat peritoneal mast cell in vitro. We provide first evidences that  $\text{H}_2\text{S}$  exerted protective effect against acute pruritus mediated via histaminergic pathways in murine skin, thus making of  $\text{H}_2\text{S}$  donors a potential alternative/complementary therapy for treatment of acute pruritus.

**Keywords:** hydrogen sulfide; pruritus; skin;  $\text{Na}_2\text{S}$ ; Lawesson's reagent; GYY4137

## 1. Introduction

Pruritus (itch) is an autonomous pain-independent sensation that, similarly to pain, acts as a distressing physiological self-protective mechanism in both humans and animals. This response greatly affects life quality and can be triggered by inflammatory skin diseases, systemic diseases, neuropathic conditions and psychogenic disorders. According to the etiology, it may be acute or chronic (duration longer than 6 weeks), localized or generalized [1], [2]. A range of mediators, such as histamine [3], prostaglandins [4], serotonin [5], bradykinin [6], cytokines [7], endothelin-1 [8], leukotrienes [9], proteases [10,11], neuropeptides [12] and opioids [13] orchestrate this response by acting on their receptors located on the nerve terminals. Pruritus (scratching behavior) is also a common symptom that results from insect bites, and can be experimentally induced in animals by the intradermal (i.d.) injection of insect saliva or venom toxins [12,14,15].

The management of pruritus, either dependent or independent of histaminergic pathways, is always recommended when removal of the trigger factors does not control the itch or perhaps unknown. Topical (e.g. anaesthetics, antihistamines, steroids, calcineurin inhibitors and capsaicin cream) or systemic approach (e.g. antihistamines, antidepressants and immunosuppressors) are normally prescribed according to the etiology [2].

Interestingly, over the last ten years, hydrogen sulfide ( $H_2S$ ) a new mediator that belongs to the class of endogenous gases such as nitric oxide (NO) and carbon monoxide (CO), has emerged and brought about divergent findings regarding its role in acute / chronic inflammatory responses and nociception [16–19]. However, the use of slow-releasing  $H_2S$  donors (such as SG-1002, diallyl trisulfide and GYY4137) and hybrid  $H_2S$ -releasing non steroidal anti-inflammatory compounds (such as the naproxen derivative ATB-346)

strengthens the beneficial therapeutical effects of H<sub>2</sub>S in articular inflammation [20], colorectal cancer [21], periodontitis [22] and pain [23] with additional gastrointestinal safety [24]. More recently, low serum levels of H<sub>2</sub>S has been associated with psoriasis [25], a disease often associated with pruritus [26]. In contrast, Wang and co-workers [27] showed that the i.d. injection of high doses of NaHS or Na<sub>2</sub>S, but not GYY4137, evoked a dose-dependent scratching behavior in mice, which is possibly related to the H<sub>2</sub>S releasing rate. Considering that the results on the effects of H<sub>2</sub>S on itch behavior are rather limited and controversial, this study was carried out to evaluate whether slow and spontaneous-H<sub>2</sub>S releasing donors, used at low doses, are able to reduce acute pruritus and the related cutaneous inflammation mediated by histamine.

## **2. Material and methods**

### *2.1. Animals*

Male BALB/c mice (20 - 30 g) and Wistar rats (180 - 200 g) were obtained from the local animal care facilities and housed in groups (up to five animals per cage) under standard controlled conditions (22°C; 12/12 h light/dark cycle) with free access to commercial rodent chow and water. All the experimental protocols were approved by the local ethics committee (CEUA-ICB; protocol n°. 33, pgs. 85, book no. 02/2010), in accordance with the guidelines from the Brazilian Council for Control of Animal Experimentation (CONCEA) and the Directive 2010/63/EU, comprising with the Animal Welfare Act.

### *2.2. Induction of pruritus (itching) in the mouse dorsal skin*

Mice were transiently anaesthetized with inhaled isoflurane (3% v/v in O<sub>2</sub>) and the rostral part of the back ( $\cong$  2 cm) near to the neck was shaved. Histamine (1  $\mu$ mol/site), C48/80 (3  $\mu$ g/site) or its corresponding vehicle Tyrode were i.d. injected, in a volume of 50  $\mu$ l, alone or in combination with 0.3-10 nmol/site of Na<sub>2</sub>S, Lawesson's reagent (LR; both, spontaneous H<sub>2</sub>S donors) or the slow-release H<sub>2</sub>S donor GYY4137. Mice were individually placed into a perspex transparent box (12x20x17 cm; Insight, Brazil) in a quiet room adapted with video camera (Sony HDR-PJ230), where the mice were daily acclimatized for 40 min during the two days previous to the experiments. A maximum of four mice were simultaneously recorded during the same period and the number of scratching bouts were counted as detailed in [12]. The number of scratching bouts was expressed either as absolute countings or as percentage values determined in 40 min. In all the experiments, the investigator who quantified the scratching behavior was unaware of the experimental group identities.

### *2.3. Assessment of dorsal cutaneous plasma extravasation*

Mice were anesthetized with urethane (2.5 g/kg; i.p.), the rostral back shaved, and 100  $\mu$ l of <sup>125</sup>I-bovine serum albumin (<sup>125</sup>I-BSA, 0.037 MBq) was intravenously (i.v.) injected via the tail vein. Histamine (30 nmol/site), C48/80 (3  $\mu$ g/site) or Tyrode were i.d. injected alone or co-injected with Na<sub>2</sub>S, LR or GYY4137 (1–100 nmol/site) throughout six randomized skin sites as previously described [28]. The results were expressed as  $\mu$ l of plasma per g of tissue or percentage based on the control values (obtained with either histamine or C48/80 alone).

### *2.4. Pharmacological treatments*

To investigate the involvement of  $K_{ATP}$  channel in  $H_2S$  donors-mediated protective effects, a set of mice was pretreated (- 30 min), via intraperitoneal (i.p.), with the  $K_{ATP}$  channel blocker, glibenclamide (10 or 30 mg/kg, i.p. [29]) or its corresponding vehicle carboxymethylcellulose (CMC; 0.1 ml, i.p.). In order to establish the effective dose of glibenclamide, another group of mice was pretreated (- 30 min) with glibenclamide 10 or 30 mg/kg and then i.d. injected with the  $K_{ATP}$  channel opener, pinacidil (10 - 30 nmol/site; i.d.). In order to assess the role of endogenous  $H_2S$  in histamine-induced skin pruritus and skin inflammation, two independent groups of mice were pretreated (- 60 min; i.p.) with the CSE and CBS inhibitors  $\beta$ -cyanoalanine (BCA, 50 mg/kg) and aminooxyacetic acid (AOAA, 20 mg/kg), a CSE and CBS inhibitors, respectively.

## 2.5. Biochemical analysis

### 2.5.1. Measurement of myeloperoxidase (MPO) activity

Mice were anaesthetized with isoflurane and i.d. injected with the test agents, as described above (item 2.3), and four hour later they were killed via an overdose of urethane followed by cervical dislocation. The injected skin sites were removed, and the myeloperoxidase (MPO) activity was measured as previously described [30]. The results were expressed as units of MPO per mg of protein (or percentage).

### 2.5.2. Production of $H_2S$ by mouse dorsal skin

The endogenous  $H_2S$  production in the naïve mouse dorsal skin was carried out based on the formation of lead sulfide, according to [31]. Briefly, skin, brain and liver were excised and homogenized. After centrifugation (10,000 g, 10 min, 4°C), the



obtained supernatants (400 µg protein) was incubated with substrates (L-cysteine 10 mM and pyridoxal-5'-phosphate 2 mM) for 2h 30min at 37 °C. The dark dots densities on the lead acetate white paper strips (12x8 cm) placed over the 96-wells microplate were analyzed from the digitalized images using the software ImageJ (NIH, USA). Hydrogen sulfide concentrations were extrapolated from a calibration curve generated with NaHS (1.95 - 2000 µM).

#### *2.6. Rat mast cell isolation and quantification of in vitro histamine release by HPLC-MS/MS*

Rats (n=6) were exsanguinated under deep isoflurane anaesthesia (5% v/v in O<sub>2</sub>), mast cells were isolated from the peritoneal cavity and purified (95%) by Percoll gradients (as determined by Cytospin<sup>®</sup> preparations stained with May-Grünwald Giemsa, and trypan blue dye exclusion). Briefly, to 0.5 ml mast cell aliquots (4x10<sup>5</sup> cells/ml) were simultaneously added compound 48/80 (1 µg/ml) and the test agents (Na<sub>2</sub>S, LR or GYY4137 at 100–1000 µM) and incubated at 37°C during 15 min. The amount of histamine released was quantified by high performance liquid chromatography coupled to an electrospray tandem mass spectrometry (HPLC-MS/MS), as described previously [32].

#### *2.7. Drugs and reagents*

Lawesson's reagent (2,4-bis[4-methoxyphenyl]-1,3,2,4-dithiadiphosphatane 2,4-disulphide), histamine (2-[1H-imidazol-4-yl]ethanamine), pyridoxal 5'-phosphate, L-cysteine, glibencamide (5-Chloro-N-[4-(cyclohexylureidosulfonyl) phenethyl]-2-methoxybenzamide), phenylmethanesulfonyl fluoride (PMSF), o-dianisidine dihydrochloride (3,3'-dimethoxybenzidine dihydrochloride), aminoxyacetic acid

(AOAA), urethane (carbamic acid ethyl ester), HTAB (hexadecyl trimethylammonium bromide), trypan blue and compound 48/80 (N-methyl-p-methoxyphenethylamine) were purchased from Sigma Chemical Co. (St Louis, MO, USA). CMC (carboxymethylcellulose) was obtained from Cromoline Química Fina Ltda (Diadema, São Paulo, Brazil). Percoll and Na<sub>2</sub>S.9H<sub>2</sub>O were purchased from GE Healthcare Bio-Sciences (Uppsala, Sweden) and Dinâmica Química Contemporânea Ltda (Diadema, São Paulo, Brazil), respectively. Isoflurane (1-chloro-2,2,2-trifluoroethyl difluoromethyl ether), hydrogen peroxide and lead acetate were purchased from Cristália (Itapira, São Paulo, Brazil) and Labsynth® (Diadema, São Paulo, Brazil), respectively. BCA ( $\beta$ -cyanoalanine) was obtained from Cayman Chemical Company (USA) and GYY4137 was synthesized in house, as described in [33].

## 2.8. Statistical analysis

Data are expressed as mean  $\pm$  SEM for n animals. Differences among the groups were analyzed by one-way ANOVA followed by Bonferroni or Dunnett's test for multiple comparisons, using the software GraphPad Prism (version 4.0, San Diego, CA, USA). Values of P lower than 0.05 were taken as significant.


## 3. Results

### 3.1. Histamine or C48/80-induced pruritus is reduced by H<sub>2</sub>S donors

As shown in Figure 1, the i.d. injection of histamine (1  $\mu$ mol/site) resulted in significant increase of pruritus in the dorsal skin compared with the Tyrode injected group. The co-injection of histamine with Na<sub>2</sub>S (1 and 3 nmol/site,  $P < 0.05$ ; Fig. 1A), LR (3 and 10 nmol/site,  $P < 0.05$ ; Fig. 1B) and GYY4137 (1 nmol/site,  $P < 0.05$ ; Fig. 1C)

reduced the number of scratching bouts in a dose-dependent manner compared to histamine alone, except that at a higher doses both Na<sub>2</sub>S and GYY4137 failed to evoke this effect. The i.d. injection of higher dose of Na<sub>2</sub>S (10 nmol/site), LR (10 nmol/site) or GYY4137 (10 nmol/site) by itself did not evoke scratching behavior (Fig. 1A-C).

Similarly, the i.d. injection of C48/80 (3 µg/site), a mast cells degranulator, also evoked a marked number of scratching bouts in comparison with Tyrode ( $P < 0.05 - P < 0.001$ ; Fig. 2A-C). This response was significantly inhibited, but not in a dose-dependent fashion, by the co-injection with increasing doses of Na<sub>2</sub>S (1, 3 and 10 nmol/site,  $P < 0.05$ ; Fig 2A) or LR (0.3 - 10 nmol/site,  $P < 0.05 - P < 0.001$ ; Fig 2B). GYY4137 i.d. injected in all tested doses (0.3 - 10 nmol/site) failed to significantly inhibit C48/80-induced pruritus (Fig 2C). None of the H<sub>2</sub>S donors produced a significant scratching behavior compared to Tyrode injected group.



**Fig. 1.** Dose response relationship for H<sub>2</sub>S-releasing donors on histamine-induced scratching bouts. The scratching bouts evoked by i.d. injection of histamine alone or co-injected with Na<sub>2</sub>S (0.3 - 10 nmol/site., *n* = 5 - 10), LR (0.3 - 10 nmol/site., *n* = 5 - 8) and GYY4137 (0.3 - 10 nmol/site., *n* = 5 - 6) are illustrated on panels A, B and C, respectively.

Independent groups of mice were i.d. injected only with H<sub>2</sub>S donors at higher doses. Dashed line represents the pruritus induced by i.d. injection of vehicle, the Tyrode solution. Data are expressed as mean  $\pm$  SEM. \*\* $P < 0.01$  –\*\*\* $P < 0.001$  vs. Tyrode, # $P < 0.05$  vs. histamine (One-way ANOVA followed by the Dunnett's test).



**Fig. 2.** Dose-response curves for H<sub>2</sub>S-releasing donors on C48/80-induced scratching bouts. Panels (A), (B) and (C) show the percentage of scratching bouts evoked by C48/80 alone (3 µg/site; i.d) and co-injected with Na<sub>2</sub>S (0.3 - 10 nmol/site, *n* = 5–8), LR (0.3 - 10 nmol/site, *n* = 5–7) and GYY4137 (0.3 - 10 nmol/site, *n* = 7), respectively. A set of mice received only i.d. injection of the H<sub>2</sub>S donors at higher dose. Dashed line represents the pruritus evoked by i.d. injection of Tyrode. Data are expressed as mean ± SEM. \**P* < 0.05 – \*\*\**P* < 0.001 vs. Tyrode, #*P* < 0.05 – ###*P* < 0.001 vs. C48/80 alone (One-way ANOVA followed by the Dunnett’s test).


### 3.2. Effects of H<sub>2</sub>S-releasing donors on dorsal skin plasma extravasation and neutrophil influx induced by amines (increased MPO activity)

The i.d. injection of histamine (30 nmol/site) induced a significant (*P* < 0.001) plasma protein extravasation in the mouse dorsal skin in comparison with Tyrode (Fig. 3). This response was dose-dependently reduced by the co-injection of Na<sub>2</sub>S (3 - 100 nmol/site, *P* < 0.05; Fig. 3A) and LR (1 - 10 nmol/site, *P* < 0.05; Fig. 3B), but unaffected by GYY4137 (Fig. 3C). Neither Tyrode nor H<sub>2</sub>S donors injected i.d. produced a significant increase in plasma extravasation.

The i.d. injection of C48/80 (3 µg/site) also resulted in significant (*P* < 0.001) amount of plasma extravasation in comparison with Tyrode (Fig. 4). Co-injected with H<sub>2</sub>S donors Na<sub>2</sub>S and LR, the plasma extravasation induced by C48/80 was significantly reduced (*P* < 0.05) at doses of 30 and 100 nmol/site (Fig. 4A and B). The simultaneous injection of GYY4137 with histamine failed to significantly affect the plasma extravasation induced by C48/80 (Fig. 4C). None of H<sub>2</sub>S donors at a higher dose (100


nmol/site) increased significantly the microvascular permeability when i.d. injected alone in the mouse dorsal skin.

After 4 hours i.d injection of C48/80 in the mouse dorsal skin (3  $\mu$ g/site), a marked ( $P < 0.001$ ) increase in MPO activity was noticed compared with Tyrode (Fig. 5). The i.d. co-injection of Na<sub>2</sub>S (3, 10 and 30 nmol/site; Fig. 5A), LR (10, 30 and 100 nmol/site; Fig. 5B) or GYY4137 (100 nmol/site; Fig. 5C) led to a significant ( $P < 0.05$  -  $P < 0.001$ ) inhibitory effect on C48/80-induced increased MPO activity. At a higher dose, Na<sub>2</sub>S (100 nmol/site) failed to inhibit C48/80-induced increased MPO activity compared to this compound alone (Fig. 5A). H<sub>2</sub>S donors had no effects when i.d. injected alone.






**Fig. 3.** Dose-response curves of the H<sub>2</sub>S donors on plasma extravasation induced by i.d. injection of histamine. Panels (A), (B) and (C) show the co-injection effects of Na<sub>2</sub>S (1 - 100 nmol/site, *n* = 5-7), LR (1 - 100 nmol/site, *n* = 5) and GYY4137 (1 - 100 nmol/site, *n* = 7-10) on histamine (30 nmol/site)-induced plasma extravasation. At higher dose, H<sub>2</sub>S donors i.d. injected induced similar plasma extravasation to that produced by its vehicle Tyrode (dashed line). Data are expressed as mean ± SEM. \*\*\**P* < 0.001 vs. Tyrode, #*P* < 0.05 vs. Histamine alone (One-way ANOVA followed by the Dunnett's test).




**Fig. 4.** Dose-response relationship between H<sub>2</sub>S-releasing donors on C48/80-induced cutaneous plasma extravasation. Panels (A), (B) and (C) illustrate response evoked by i.d. injection of C48/80 (3 µg/site) alone and co-injected with Na<sub>2</sub>S (1 - 100 nmol/site., *n* = 6), LR (1 - 100 nmol/site., *n* = 8) and GYY4137 (1 - 100 nmol/site., *n* = 10), respectively. At higher dose, the i.d. of H<sub>2</sub>S donors induced similar plasma extravasation to that produced by its vehicle Tyrode (dashed line). Data are expressed as mean ± SEM. \*\*\**P* < 0.001 vs. Tyrode, #*P* < 0.05 vs. C48/80 alone (One-way ANOVA followed by the Dunnett's test).



**Fig. 5.** Effects of H<sub>2</sub>S donors on C48/80-induced increased MPO activity. Panels (A), (b) and (C) show the MPO activity evoked by i.d. injection of C48/80 (3 µg/site) alone and co-injected with Na<sub>2</sub>S (1-100 nmol/site.,  $n = 6$ ), LR (1-100 nmol/site.,  $n = 10$ ) and GYY4137 (1-100 nmol/site.,  $n = 8$ ), respectively, 4 hours later. At higher doses, H<sub>2</sub>S donors evoked similar response to that evoked by its vehicle Tyrode (dashed line). Data are expressed as mean  $\pm$  SEM. \*\*\* $P < 0.001$  vs. Tyrode, # $P < 0.05$  - ### $P < 0.001$  vs. C48/80 alone (One-way ANOVA followed by the Dunnett's test).

### 3.3. Role of endogenous H<sub>2</sub>S in C48/80-induced both pruritus and neutrophil influx


As expected, the i.d. injection of C48/80 (3 µg/site) induced a significant increase on the number of scratching bouts (Fig. 6A and B) and MPO activity (Fig. 6C and D). The pretreatment of mice with the CSE inhibitor BCA (50 mg/kg; i.p., - 60 min) significantly exacerbated ( $P < 0.05$ ) C48/80-induced pruritus and MPO activity compared to control group pretreated with saline (Fig. 6A and 6C), whilst the pretreatment of animals with the nonselective CBS inhibitor AOAA (20 mg/kg; i.p., - 60 min) did not. The i.d. injection of Tyrode in mice dorsal skin did not evoke a significant increase in MPO activity; however, the pretreatment of mice with BCA or AOAA resulted in a significant increase of MPO activity compared to Tyrode response in saline-pretreated mice (Fig. 6C and D).



**Fig. 6.** Endogenous blockade of H<sub>2</sub>S synthesis potentiated C48/80-induced pruritus and MPO activity in mouse dorsal skin. Panels (A) and (B) show the effects of pretreatment with either the CSE inhibitor,  $\beta$ -cyanoalanine (BCA, 50 mg/kg., -60 min) or the nonselective CBS inhibitor aminooxyacetic acid (AOAA, 20 mg/kg., -60 min) in the pruritus evoked by C48/80 (n=5-8). Panels (C) and (D) show the same effects of the same treatments in C48/80-induced MPO activity (n=5-8). Data are expressed as mean  $\pm$  S.E.M. for n animals. \*  $P < 0.05$  and \*\*\*  $P < 0.001$  vs. Tyrode. ##  $P < 0.01$  and ###  $P < 0.001$  vs. saline pretreated mice. (One-way ANOVA followed by Bonferroni's multiple comparison test).

#### 3.4 Determination of H<sub>2</sub>S in mouse dorsal skin

The naïve mouse skin, similarly to the brain, produce a significant and equivalent amount of H<sub>2</sub>S ( $\cong 0.3$  nmol / mg of protein/min), whereas a marked H<sub>2</sub>S generation was generated in the liver ( $\cong 1.2$  nmol / mg of protein/min) of naïve mice (Fig. 7).




**Fig. 7.** Constitutive H<sub>2</sub>S generation in the naïve skin, liver and brain of mice. Panel A shows the H<sub>2</sub>S generation in the skin, liver and brain of naïve animals, expressed as nmol of H<sub>2</sub>S per mg of protein/min, and panel B illustrates the representative corresponding skin, liver and brain dark dots densities on the lead acetate white paper strip. Data are expressed as mean  $\pm$  S.E.M. for n = 4 animals.

### 3.5. Histamine release from mast cells activation is reduced by H<sub>2</sub>S-releasing donors

C48/80 (1  $\mu$ g/ml) markedly stimulated the release of histamine ( $P < 0.001$ ) from rat peritoneal mast cells compared with histamine spontaneously released from mast cells

treated only with buffer (KRP; Fig. 8). The concomitant incubation of mast cells with C48/80 and Na<sub>2</sub>S (100 and 1000 μM;  $P < 0.05 - P < 0.01$ ) or LR (500 - 1000 μM;  $P < 0.05 - P < 0.001$ ) resulted in significant decrease of histamine release from these cells. In all tested concentrations, GYY4137 did not prevent histamine release from C48/80-induced mast cell degranulation (Fig. 8). The % of histamine released from mast cells incubated with H<sub>2</sub>S donors alone matched with KRP value (Fig. 8).




**Fig. 8.** Spontaneous, but not slow, H<sub>2</sub>S donors reduce C48/80-induced mast cell degranulation. Black bar illustrates the % of histamine released from mast cells treated with C48/80 (1 μg/ml) in krebs ringer phosphate solution (KRP). Cross-hatched bars show the % of histamine release from mast cells in response to C48/80 plus Na<sub>2</sub>S, LR or GYY4137 (100 - 1000 μM). Data are expressed as % of histamine release. Values are presented as mean ± SEM of three independent experiments. \*\*\* $P < 0.001$  vs. KRP (basal release), # $P < 0.05$ - ##  $P < 0.01$ -###  $P < 0.001$  vs. C48/80 alone (One-way ANOVA followed by Bonferroni's multiple comparison test).



### 3.6. *Lack of involvement of $K_{ATP}$ channels in the antipruritic effect of $H_2S$*

The pretreatment of mice with glibenclamide (10 or 30 mg/kg; i.p., - 30 min) significantly inhibited, at both doses, the  $K_{ATP}$  channel opener pinacidil (10 and 30 nmol/site)-induced exacerbation of plasma extravasation evoked by histamine (Fig. 9A), thus suggesting an effective blockade of this channel. The pretreatment of animals with 10 mg/kg of glibenclamide did not significantly affect histamine-induced plasma extravasation (Fig. 9B) or pruritus (Fig. 9C) compared to histamine responses in animals pretreated with vehicle CMC. Likewise, glibenclamide treatment did not alter the protective effect of  $Na_2S$  (3 or 30 nmol/site) against histamine-induced plasma extravasation or pruritus (Fig. 9B-C).



**Fig. 9.** Blockade of  $K_{ATP}$  channels failed to reverse  $Na_2S$ -induced protective effect against histamine-induced plasma extravasation and pruritus. Panel (A) shows the effect of glibenclamide (10 or 30 mg/kg., i.p., - 30 min) or its vehicle (CMC) on pinacidil (10 or 30 nmol/site)-induced potentiating of plasma extravasation in response to i.d. of histamine (30 nmol/site;  $n=5$ ), whereas panels (B) and (C) show the effects of histamine in glibenclamide pretreated group (10 mg/kg., i.p., - 30 min) in addition to  $Na_2S$  (30 nmol/site.,  $n = 5 - 6$ ) and pruritus (1  $\mu$ mol/site.,  $n = 5 - 10$ ). Data are presented as mean  $\pm$  S.E.M. for  $n$  animals. \*\*  $P < 0.01$  - \*\*\*  $P < 0.001$  vs. Tyrode + CMC, ##  $P < 0.01$  - ###  $P < 0.001$  vs. histamine + CMC, + $P < 0.05$  - +++ $P < 0.001$  vs histamine + pinacidil. (One-way ANOVA followed by Bonferroni's test).

## Discussion

The major finding in this study is to show that the spontaneous  $H_2S$  donors  $Na_2S$  and LR significantly ameliorated pruritus and the acute cutaneous inflammation induced by histamine and C48/80 in the mouse dorsal skin, being this effect less pronounced (or absent) in mice treated with the slow  $H_2S$ -releasing donor GYY4137, whilst the endogenous blockade of  $H_2S$  biosynthesis aggravates skin inflammation and pruritus.

Histamine-induced plasma extravasation (oedema) is mainly processed by activation of  $H_1$ -histaminergic receptors and, to a lesser extent,  $H_2$  receptors expressed in the skin vessels of both rodents [34,35] and humans [36,37]. Herein, the simultaneous i.d. injection of  $H_2S$  donors  $Na_2S$  or LR within the 0.3-10 nmoles/site dose range, significantly inhibited pruritus and plasma extravasation produced by histamine, even though it did not follow a clear dose-dependent response. Instead, a trend to biphasic

pattern in the scratching behavior (pruritus) and plasma protein extravasation was observed with these H<sub>2</sub>S donors, in particular with Na<sub>2</sub>S.

Even though we have used smaller doses of Na<sub>2</sub>S compared to previous study [27], it is possible that as this compound instantaneously delivers H<sub>2</sub>S, the higher tested dose (100 nmol/site) led to an immediate non-physiologically concentration of H<sub>2</sub>S in the local microcirculation, which may be distinct from that seen in the dorsal skin. In line with this, we provide the first evidence that the mouse dorsal healthy skin constitutively produces low amounts of H<sub>2</sub>S ( $\cong$  0.3 nmol of H<sub>2</sub>S per mg of protein per min), similarly to that produced by the brain but less than that produced by the liver. Thus, these results point to the same suggestion made by previous studies, in terms that the application of high doses of a fast H<sub>2</sub>S-releasing donors can rapidly lead to elevated concentrations of this mediator, which in turn may result in toxic effects, rather than the beneficial ones seen with low amounts of H<sub>2</sub>S [33,38–41].

Histamine exerts its inflammatory effects, in part, through NO generation *in situ* [35], since the blockade of NOS significantly inhibited histamine-induced plasma extravasation in human nasal airway [42]. Interestingly, Ali and co-workers [43] demonstrated that low concentration of H<sub>2</sub>S or spontaneous H<sub>2</sub>S donors reversed histamine, but not isoprenaline-induced vasodilatation, possibly via a direct chemical reaction between H<sub>2</sub>S and NO, which may lead to the formation of a nitrosothiol [44] or even nitroxyl species (NO-/HNO) [45] of poor vasorelaxant activities.

Similarly to histamine, the cationic secretagogue C48/80 also induces a marked pruritus behavior and skin inflammation characterized by plasma extravasation and neutrophil influx, except for the more pronounced response observed with the latter when applied at similar doses mainly due to the exocytosis of multiple preformed mediators from mast cells (e.g. histamine, 5-HT, etc.) on both blood vessels and neurons, via

interactions with their respective receptors. In fact, C48/80 effects can be significantly inhibited by antagonists of histamine H1 receptors, 5-HT or substance P (NK1) receptors [36,46,47].

C48/80-induced plasma extravasation and pruritus was partially, but significantly, inhibited by the simultaneous co-injection of the spontaneous H<sub>2</sub>S donors Na<sub>2</sub>S or LR, whereas neither histamine nor C48/80-induced plasma leakage was significantly affected by the slow H<sub>2</sub>S-releasing donor GYY4137. It is thus possible that a significant amount of H<sub>2</sub>S immediately available in the microvascular bed during the initial phases of the vascular response is necessary to interact with NO generated by histamine (or the histamine releaser C48/80), and consequently inhibit its potentiating action on microvascular permeability. Indeed, it is well established that plasma extravasation occurs immediately after the i.d. injection of chemical mediators (such as histamine), whereas the leukocyte recruitment into the cutaneous tissue takes longer periods of time (>3 h). In agreement, our results show that 4 hours after the i.d. injection of C48/80 a marked neutrophil influx occurs (measured as MPO activity), and this response was effectively reduced by all the H<sub>2</sub>S releasing donors. It is thus possible that the GYY4137 compound may have released enough H<sub>2</sub>S along this time, which in turn can counteract with the dynamic of leukocyte influx in the microcirculation in response to C48/80.

A direct stabilizing effect of H<sub>2</sub>S on mast cells degranulation could explain the reduced leukocyte influx in response to C48/80. In order to test this hypothesis we performed *in vitro* experiments using rat peritoneal mast cells stimulated with C48/80 and measured the release of histamine. In fact, we observed that Na<sub>2</sub>S and LR, but not GYY4137 prevented mast cell degranulation. In agreement with our data, Zanardo and co-workers [48] have shown that spontaneous H<sub>2</sub>S donors, such as Na<sub>2</sub>S, NaHS and LR, markedly inhibited carrageenan-induced leukocyte adhesion and infiltration in a rodent

air pouch model. Similar results were obtained by Ekundi-Valentim and co-workers [16,20] in the model of carrageenan-induced synovitis in rats, and by Fiorucci and co-workers [49] in the acetyl salicylic acid-induced gastric injury in rats. Reduced expression of the adhesion molecules ICAM-1, VCAM-1, LFA-1, P-selectin, E-selectin in both endothelium and leukocytes has been suggested as the underlying mechanism of H<sub>2</sub>S-mediated inhibition of leukocyte influx [40,48–50].

The activation of K<sub>ATP</sub> channels by H<sub>2</sub>S has been shown as the mechanism underlying the inhibitory effects of this mediator in a variety of experimental approaches, including aspirin-induced leukocyte adherence in mesenteric venules [48] and high glucose-induced cardiac cells injury [51]. However, this not seem to be the cause beyond the protective effects of the spontaneous H<sub>2</sub>S donors against histamine-induced pruritus and skin inflammation, as pretreatment of mice with glibenclamide, a selective K<sub>ATP</sub> channels blocker, did not antagonize the protective effects of H<sub>2</sub>S, even when used at doses that abolish the vasodilatation induced in the mouse dorsal skin by the K<sub>ATP</sub> channel opener pinacidil.

Compound 48/80 has been largely used as an IgE independent activator of mast cells, which mainly due to histamine release (and activation of peripheral H<sub>1</sub> and H<sub>4</sub> receptors), results in sensitization of afferent nerves [52,53]. Similarly to the spontaneous H<sub>2</sub>S donor Na<sub>2</sub>S, GYY4137 inhibited histamine-induced pruritus; however, this was not the case with C48/80-induced pruritus. Thus, either the intensity of response evoked by histamine released from C48/80 was higher than exogenous histamine when paralleled by the low availability of H<sub>2</sub>S released from GYY4137 *in situ* on the peripheral afferent neurons or it is possibly that C48/80 produces scratching behavior that somehow depends on functional changes evoked by direct activation of pruriceptors. While direct excitatory effects of C48/80 on dorsal root ganglion (DRG), enteric neurons and visceral afferents

can occur independently of mast cell activation [54], other works show that the mechanism involved in C48/80-induced nociception is mediated by a cascade activation that starts after mast cell activation, including the release of mediators that can activate these nociceptors and promote pain [55].

Curiously, and in contrast to our findings, a recent work by Wang and co-workers [27] has shown that the i.d. injection of mice with the spontaneous H<sub>2</sub>S donors NaHS and Na<sub>2</sub>S, at mM range (7 – 280 and 1.7 – 67, respectively), led to a marked itching behavior and touch-evoked itching (allokinesis), while inhibition of endogenous H<sub>2</sub>S synthesis by pretreatment of the mice with CSE and CBS inhibitors reduced C48/80-induced pruritus. Previously, Elies and co-workers [56] showed that NaHS at low concentrations (100-300 μM) markedly inhibits Ca<sup>2+</sup><sub>v3.2</sub> T-type calcium channel in HEK293 cells, while NaHS at high concentrations (3-10 mM) led to a marked activation of these channels, possibly due to the resultant high Na<sup>+</sup> concentrations. In line with these findings, we speculated that the presence of these high Na<sup>+</sup> concentrations may explain the discrepancies between our results and those shown in Wang's paper. We also show that inhibition of endogenous H<sub>2</sub>S biosynthesis by BCA, but not by AOAA, resulted in enhanced C48/80-induced scratching behavior in addition to increased MPO activity. Again, the C48/80-induced pruritus results are in contrast with those produced by Wang and co-workers [27], but the increment in MPO activity are agreement with previous work by Zanardo and co-workers [48], in which they showed that animals treated with BCA exhibited enhanced leukocyte adherence and infiltration in an air pouch model [47]. Thus, it is possible that the administration route used by these authors (i.d.) may account for the discrepancies considering that local desensitization effects may be produced by these drugs independently of H<sub>2</sub>S generation.

In addition to potentiating the scratching behavior, H<sub>2</sub>S synthesis blockade by BCA or AOAA has also enhanced neutrophil influx into the dorsal skin, under basal conditions. Furthermore, the non-selective inhibition of CBS/CSE by AOAA increased the basal neutrophil influx to an extent similar to that induced by C48/80, thus reflecting the protective role of endogenous H<sub>2</sub>S in the normal skin physiology.

## Conclusions

Altogether the results shown in this study provide the first evidence that endogenous H<sub>2</sub>S, as well as the administration of low doses of spontaneous H<sub>2</sub>S donors, such as LR and Na<sub>2</sub>S, exert protective roles against histamine/mast cell-mediated acute pruritus and inflammation of the dorsal mouse skin, independently of H<sub>2</sub>S action on K<sub>ATP</sub> channels.

## Conflicts of interest

We have no conflict of interest to declare.

## Acknowledgements

Sao Paulo Research Foundation (Fapesp grant numbers: 2013/04.151-3, 2014/15.576-8, 2014/24.518-1) and CNPq (grant number: 163278/2012-1). GDN, MNM and SKPC are recipients of fellowships from the National Council for Scientific and Technological Development (CNPq). We thank Irene M Gouvea, Flávia B de Lira and Mauro Sucupira for their technical support.

## References

- [1] S. Ständer, E. Weisshaar, T. Mettang, J.C. Szepietowski, E. Carstens, A. Ikoma, et al., Clinical classification of itch: a position paper of the International Forum for the



- Study of Itch., *Acta Dermato-Venereologica*. 87 (2007) 291–4.
- [2] T.A. Leslie, M.W. Greaves, G. Yosipovitch, Current topical and systemic therapies for itch., *Handbook of Experimental Pharmacology*. 226 (2015) 337–56.
- [3] P. Sikand, S.G. Shimada, B.G. Green, R.H. LaMotte, Sensory responses to injection and punctate application of capsaicin and histamine to the skin., *Pain*. 152 (2011) 2485–94.
- [4] M. Schmelz, R. Schmidt, C. Weidner, M. Hilliges, H.E. Torebjork, H.O. Handwerker, Chemical response pattern of different classes of C-nociceptors to pruritogens and algogens., *Journal of Neurophysiology*. 89 (2003) 2441–8.
- [5] T. Akiyama, M.I. Carstens, E. Carstens, Enhanced scratching evoked by PAR-2 agonist and 5-HT but not histamine in a mouse model of chronic dry skin itch., *Pain*. 151 (2010) 378–83.
- [6] M. Hosogi, M. Schmelz, Y. Miyachi, A. Ikoma, Bradykinin is a potent pruritogen in atopic dermatitis: a switch from pain to itch., *Pain*. 126 (2007) 16–23.
- [7] K. Kasutani, E. Fujii, S. Ohyama, H. Adachi, M. Hasegawa, H. Kitamura, et al., Anti-IL-31 receptor antibody is shown to be a potential therapeutic option for treating itch and dermatitis in mice., *British Journal of Pharmacology*. 171 (2014) 5049–58.
- [8] D.S. McQueen, M.A. Noble, S.M. Bond, Endothelin-1 activates ETA receptors to cause reflex scratching in BALB/c mice., *British Journal of Pharmacology*. 151 (2007) 278–84.
- [9] E.S. Fernandes, C.T. Vong, S. Quek, J. Cheong, S. Awal, C. Gentry, et al., Superoxide generation and leukocyte accumulation: key elements in the mediation of leukotriene B<sub>4</sub>-induced itch by transient receptor potential ankyrin 1 and transient receptor potential vanilloid 1., *FASEB Journal : Official Publication of the Federation of American Societies for Experimental Biology*. 27 (2013) 1664–73.
- [10] R. Costa, D.M. Marotta, M.N. Manjavachi, E.S. Fernandes, J.F. Lima-Garcia, A.F. Paszcuk, et al., Evidence for the role of neurogenic inflammation components in trypsin-elicited scratching behaviour in mice., *British Journal of Pharmacology*. 154 (2008) 1094–103.
- [11] K. Tsujii, T. Andoh, J.-B.B. Lee, Y. Kuraishi, Activation of proteinase-activated receptors induces itch-associated response through histamine-dependent and -independent pathways in mice., *Journal of Pharmacological Sciences*. 108 (2008) 385–8.
- [12] S.K. Costa, A. Starr, S. Hyslop, D. Gilmore, S.D. Brain, How important are NK1 receptors for influencing microvascular inflammation and itch in the skin? Studies using *Phoneutria nigriventer* venom., *Vascular Pharmacology*. 45 (2006) 209–14.
- [13] R.G. Jannuzzi, Nalbuphine for Treatment of Opioid-induced Pruritus: A Systematic Review of Literature., *The Clinical Journal of Pain*. 32 (2016) 87–93.
- [14] A. Ikoma, M. Steinhoff, S. Ständer, G. Yosipovitch, M. Schmelz, The neurobiology of itch., *Nature Reviews. Neuroscience*. 7 (2006) 535–47.
- [15] B. Morfín Maciel, [Itching caused by insects]., *Revista Alergia Mexico (Tecamachalco, Puebla, Mexico : 1993)*. 50 (2003) 154–60.
- [16] E. Ekundi-Valentim, K.T. Santos, E.A. Camargo, A. Denadai-Souza, S.A. Teixeira, C.I. Zanoni, et al., Differing effects of exogenous and endogenous hydrogen sulphide in carrageenan-induced knee joint synovitis in the rat., *British Journal of Pharmacology*. 159 (2010) 1463–74.
- [17] L.R. Benetti, D. Campos, S.A. Gurgueira, A.E. Vercesi, C.E. Guedes, K.L. Santos, et al., Hydrogen sulfide inhibits oxidative stress in lungs from allergic mice in vivo., *European Journal of Pharmacology*. 698 (2013) 463–9.
- [18] E. Distrutti, S. Cipriani, B. Renga, A. Mencarelli, M. Migliorati, S. Cianetti, et

- al., Hydrogen sulphide induces micro opioid receptor-dependent analgesia in a rodent model of visceral pain., *Molecular Pain*. 6 (2010) 36.
- [19] A.F. Donatti, R.M. Araujo, R.N. Soriano, L.U. Azevedo, C.A. Leite-Panissi, L.G. Branco, Role of hydrogen sulfide in the formalin-induced orofacial pain in rats., *European Journal of Pharmacology*. 738 (2014) 49–56.
- [20] E. Ekundi-Valentim, F.P. Mesquita, K.T. Santos, M.A. de Paula, J. Florenzano, C.I. Zanoni, et al., A comparative study on the anti-inflammatory effects of single oral doses of naproxen and its hydrogen sulfide (H<sub>2</sub>S)-releasing derivative ATB-346 in rats with carrageenan-induced synovitis., *Medical Gas Research*. 3 (2013) 24.
- [21] W. Elsheikh, R.W. Blackler, K.L. Flannigan, J.L. Wallace, Enhanced chemopreventive effects of a hydrogen sulfide-releasing anti-inflammatory drug (ATB-346) in experimental colorectal cancer., *Nitric Oxide : Biology and Chemistry / Official Journal of the Nitric Oxide Society*. 41 (2014) 131–7.
- [22] B.S. Herrera, L.S. Coimbra, A.R. da Silva, S.A. Teixeira, S.K. Costa, J.L. Wallace, et al., The H<sub>2</sub>S-releasing naproxen derivative, ATB-346, inhibits alveolar bone loss and inflammation in rats with ligature-induced periodontitis., *Medical Gas Research*. 5 (2015) 4.
- [23] M. Chattopadhyay, R. Kodela, P.L. Duvalsaint, K. Kashfi, Gastrointestinal safety, chemotherapeutic potential, and classic pharmacological profile of NOSH-naproxen (AVT-219) a dual NO- and H<sub>2</sub>S-releasing hybrid., *Pharmacology Research & Perspectives*. 4 (2016) e00224.
- [24] O. Sulaieva, J.L. Wallace, Gaseous mediator-based anti-inflammatory drugs., *Current Opinion in Pharmacology*. 25 (2016) 1–6.
- [25] A.K. Alshorafa, Q. Guo, F. Zeng, M. Chen, G. Tan, Z. Tang, et al., Psoriasis is associated with low serum levels of hydrogen sulfide, a potential anti-inflammatory molecule., *The Tohoku Journal of Experimental Medicine*. 228 (2012) 325–32.
- [26] E. Carstens, T. Akiyama, *Itch: Mechanisms and Treatment*, CRC Press/Taylor & Francis. (2014).
- [27] X.-L.L. Wang, B. Tian, Y. Huang, X.-Y.Y. Peng, L.-H.H. Chen, J.-C.C. Li, et al., Hydrogen sulfide-induced itch requires activation of Cav3.2 T-type calcium channel in mice., *Scientific Reports*. 5 (2015) 16768.
- [28] S.K. Costa, G. De Nucci, E. Antunes, S.D. Brain, Involvement of vanilloid receptors and purinoceptors in the Phoneutria nigriventer spider venom-induced plasma extravasation in rat skin., *European Journal of Pharmacology*. 391 (2000) 305–15.
- [29] S.K. Costa, H. Moreno, S.D. Brain, G. De Nucci, E. Antunes, The effect of Phoneutria nigriventer (armed spider) venom on arterial blood pressure of anaesthetised rats., *Eur. J. Pharmacol.* 298 (1996) 113–20.
- [30] L.M.M. Yshii, G.H. Souza, E.A. Camargo, M.N. Eberlin, M.T. Ribela, M.N. Muscará, et al., Characterization of the mechanisms underlying the inflammatory response to *Polistes lanio lanio* (paper wasp) venom in mouse dorsal skin., *Toxicon : Official Journal of the International Society on Toxinology*. 53 (2009) 42–52.
- [31] C. Hine, E. Harputlugil, Y. Zhang, C. Ruckenstein, B.C. Lee, L. Brace, et al., Endogenous hydrogen sulfide production is essential for dietary restriction benefits., *Cell*. 160 (2015) 132–44.
- [32] G.D. Mendes, S. Bittencourt, C.F. Vespasiano, T. Babadópulos, T. Gagliano-Jucá, A.M.M.M. Arruda, et al., Propylthiouracil quantification in human plasma by high-performance liquid chromatography coupled with electrospray tandem mass spectrometry: application in a bioequivalence study., *Journal of Chromatography. B, Analytical Technologies in the Biomedical and Life Sciences*. 969 (2014) 19–28.
- [33] L. Li, M. Whiteman, Y.Y. Guan, K.L. Neo, Y. Cheng, S.W. Lee, et al.,

- Characterization of a novel, water-soluble hydrogen sulfide-releasing molecule (GYY4137): new insights into the biology of hydrogen sulfide., *Circulation*. 117 (2008) 2351–60.
- [34] A. Belcheva, M. Zhelyazkova, On the interaction between H1- and H2-histaminergic receptors in the histamine release and the realization of its vascular effects in rat skin., *Acta Physiologica et Pharmacologica Bulgarica*. 14 (1988) 49–54.
- [35] H. Jin, T. Koyama, Y. Hatanaka, S. Akiyama, F. Takayama, H. Kawasaki, Histamine-induced vasodilation and vasoconstriction in the mesenteric resistance artery of the rat., *European Journal of Pharmacology*. 529 (2006) 136–44.
- [36] R. Marks, M.W. Greaves, Vascular reactions to histamine and compound 48/80 in human skin: suppression by a histamine H2-receptor blocking agent., *British Journal of Clinical Pharmacology*. 4 (1977) 367–9.
- [37] M. Grossmann, M.J. Jamieson, W. Kirch, Histamine response and local cooling in the human skin: involvement of H1- and H2-receptors., *British Journal of Clinical Pharmacology*. 48 (1999) 216–22.
- [38] Y. Ji, Q.F. Pang, G. Xu, L. Wang, J.K. Wang, Y.M. Zeng, Exogenous hydrogen sulfide postconditioning protects isolated rat hearts against ischemia-reperfusion injury., *European Journal of Pharmacology*. 587 (2008) 1–7.
- [39] D. Johansen, K. Ytrehus, G.F. Baxter, Exogenous hydrogen sulfide (H<sub>2</sub>S) protects against regional myocardial ischemia-reperfusion injury--Evidence for a role of K ATP channels., *Basic Research in Cardiology*. 101 (2006) 53–60.
- [40] J.N. Sidhapuriwala, S.W. Ng, M. Bhatia, Effects of hydrogen sulfide on inflammation in caerulein-induced acute pancreatitis., *Journal of Inflammation (London, England)*. 6 (2009) 35.
- [41] M. Whiteman, L. Li, P. Rose, C.-H.H. Tan, D.B. Parkinson, P.K. Moore, The effect of hydrogen sulfide donors on lipopolysaccharide-induced formation of inflammatory mediators in macrophages., *Antioxidants & Redox Signaling*. 12 (2010) 1147–54.
- [42] J.W. Dear, S. Ghali, J.C. Foreman, Attenuation of human nasal airway responses to bradykinin and histamine by inhibitors of nitric oxide synthase., *British Journal of Pharmacology*. 118 (1996) 1177–82.
- [43] M.Y. Ali, C.Y. Ping, Y.-Y.Y. Mok, L. Ling, M. Whiteman, M. Bhatia, et al., Regulation of vascular nitric oxide in vitro and in vivo; a new role for endogenous hydrogen sulphide?, *British Journal of Pharmacology*. 149 (2006) 625–34.
- [44] M. Whiteman, L. Li, I. Kostetski, S.H. Chu, J.L. Siau, M. Bhatia, et al., Evidence for the formation of a novel nitrosothiol from the gaseous mediators nitric oxide and hydrogen sulphide., *Biochemical and Biophysical Research Communications*. 343 (2006) 303–10.
- [45] M.R. Filipovic, M. Eberhardt, V. Prokopovic, A. Mijuskovic, Z. Orescanin-Dusic, P. Reeh, et al., Beyond H<sub>2</sub>S and NO interplay: hydrogen sulfide and nitroprusside react directly to give nitroxyl (HNO). A new pharmacological source of HNO., *Journal of Medicinal Chemistry*. 56 (2013) 1499–508.
- [46] A. Saria, X. Hua, G. Skofitsch, J.M. Lundberg, Inhibition of compound 48/80--induced vascular protein leakage by pretreatment with capsaicin and a substance P antagonist., *Naunyn-Schmiedeberg's Archives of Pharmacology*. 328 (1984) 9–15.
- [47] W. Jiang, M.E. Kreis, C. Eastwood, A.J. Kirkup, P.P. Humphrey, D. Grundy, 5-HT(3) and histamine H(1) receptors mediate afferent nerve sensitivity to intestinal anaphylaxis in rats., *Gastroenterology*. 119 (2000) 1267–75.
- [48] R.C. Zanardo, V. Brancaleone, E. Distrutti, S. Fiorucci, G. Cirino, J.L. Wallace, Hydrogen sulfide is an endogenous modulator of leukocyte-mediated inflammation.,

- FASEB Journal : Official Publication of the Federation of American Societies for Experimental Biology. 20 (2006) 2118–20.
- [49] S. Fiorucci, E. Antonelli, E. Distrutti, G. Rizzo, A. Mencarelli, S. Orlandi, et al., Inhibition of hydrogen sulfide generation contributes to gastric injury caused by anti-inflammatory nonsteroidal drugs., *Gastroenterology*. 129 (2005) 1210–24.
- [50] Q. Guan, X. Wang, L. Gao, J. Chen, Y. Liu, C. Yu, et al., Hydrogen sulfide suppresses high glucose-induced expression of intercellular adhesion molecule-1 in endothelial cells., *Journal of Cardiovascular Pharmacology*. 62 (2013) 278–84.
- [51] W. Liang, J. Chen, L. Mo, X. Ke, W. Zhang, D. Zheng, et al., ATP-sensitive K<sup>+</sup> channels contribute to the protective effects of exogenous hydrogen sulfide against high glucose-induced injury in H9c2 cardiac cells., *Int. J. Mol. Med*. 37 (2016) 763–72.
- [52] J.K. Bell, D.S. McQueen, J.L. Rees, Involvement of histamine H4 and H1 receptors in scratching induced by histamine receptor agonists in Balb C mice., *British Journal of Pharmacology*. 142 (2004) 374–80.
- [53] P.J. Dunford, K.N. Williams, P.J. Desai, L. Karlsson, D. McQueen, R.L. Thurmond, Histamine H4 receptor antagonists are superior to traditional antihistamines in the attenuation of experimental pruritus., *The Journal of Allergy and Clinical Immunology*. 119 (2007) 176–83.
- [54] M. Schemann, E.M. Kugler, S. Buhner, C. Eastwood, J. Donovan, W. Jiang, et al., The mast cell degranulator compound 48/80 directly activates neurons., *PloS One*. 7 (2012) e52104.
- [55] D. Levy, V. Kainz, R. Burstein, A.M. Strassman, Mast cell degranulation distinctly activates trigemino-cervical and lumbosacral pain pathways and elicits widespread tactile pain hypersensitivity., *Brain, Behavior, and Immunity*. 26 (2012) 311–7.
- [56] J. Elies, J.L. Scragg, S. Huang, M.L. Dallas, D. Huang, D. MacDougall, et al., Hydrogen sulfide inhibits Cav3.2 T-type Ca<sup>2+</sup> channels., *FASEB Journal : Official Publication of the Federation of American Societies for Experimental Biology*. 28 (2015) 5376–87.