



Strathprints Institutional Repository

Uzelac, Marina and Kennedy, Alan R. and Hevia, Eva and Mulvey, Robert E. (2016) Transforming LiTMP lithiation of challenging diazines via gallium alkyl trans-metal-trapping. *Angewandte Chemie International Edition*, 55. pp. 13147-13150. ISSN 1433-7851 , <http://dx.doi.org/10.1002/anie.201607284>

This version is available at <http://strathprints.strath.ac.uk/57758/>

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<http://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to Strathprints administrator: strathprints@strath.ac.uk

Metalation

International Edition: DOI: 10.1002/anie.201607284
German Edition: DOI: 10.1002/ange.201607284

Transforming LiTMP Lithiation of Challenging Diazines through Gallium Alkyl Trans-Metal-Trapping


Marina Uzelac, Alan R. Kennedy, Eva Hevia,* and Robert E. Mulvey*

Abstract: This study establishes a new trans-metal-trapping (TMT) procedure based on a mixture of LiTMP (the base) and tris(trimethylsilylmethyl)gallium [$\text{Ga}(\text{CH}_2\text{SiMe}_3)_3$, GaR_3] (the trap) that, operating in a tandem manner, is effective for the regioselective deprotonation of sensitive diazines in hydrocarbon solution, as illustrated through reactions of pyrazine, pyridazine, and pyrimidine, as well as through the *N*-S heterocycle benzothiazole. The metallo-activated complexes of all of these compounds were isolated and structurally defined.

As one of the most used strategies in synthetic campaigns, metalation chemistry is currently enjoying a remarkable period of advancement.^[1] Lithium alkyls and amides remain front-runners for base candidates in routine C–H to C–metal transformations, but for non-routine, more formidable substrates, base selection is generally far from straightforward.^[2] One of nature's most important class of heterocycles utilized widely in agrochemicals, foodstuffs, pharmaceuticals, and many other commercial commodities, diazines are firmly in the formidable category, especially parent naked diazines devoid of substituents capable of assisting the direction of the metalation.^[3] To avoid competition from nucleophilic addition caused by a low-lying LUMO in the heteroaromatic ring, bulky LiTMP (TMP is 2,2,6,6-tetramethylpiperidide) is preferred to more basic alkyl lithium reagents as the base for these π -deficient diazines, as first noted by Quéguiner.^[4] However, lithiated diazine intermediates are generally unstable, so Knochel,^[5] Kondo,^[6] Mongin,^[7] and Hevia^[8] have employed coalitions of components typically but not exclusively based on the softer metal zinc to improve stabilities and to perform metalation under milder conditions. Though excellent progress has been made, these coalition approaches still have their limitations, as for example in the zincation of pyrazine using $[(\text{THF})\text{LiZn}(\text{TMP})\text{tBu}_2]$, where no stoichiometric control is possible because only 2,5-dizincation occurs

in a 1:1 base:pyrazine stoichiometry.^[8] Moreover, a lack of definitive structural information still impoverishes understanding of this area, which in the most extreme “black box” cases leads to a misidentification of the actual metalating base.^[9]

This paper reports a new trans-metal-trapping (TMT) procedure based on a mixture of LiTMP and tris(trimethylsilylmethyl)gallium [$\text{Ga}(\text{CH}_2\text{SiMe}_3)_3$, GaR_3] that is effective for regioselective diazine and benzothiazole deprotonation in hydrocarbon solution. Whereas alkyl lithium reactions are generally irreversible, LiTMP reactions tend to be $\text{p}K_{\text{a}}$ -dependent equilibria. In TMT, these equilibria are shifted towards the desired lithiated substrate product by its interception by a trapping agent (Scheme 1).^[10]



Scheme 1. Two-step lithiation and carbanion trapping process in equilibrium controlling Trans-Metal-Trapping (TMT).

We have previously employed the diorganoaluminum amide $i\text{Bu}_2\text{Al}(\text{TMP})$ as a trapping agent.^[10] For this challenging diazine work, we elected to test GaR_3 as a trap because its bulkiness compromises its ability to form a weakly basic ate complex with LiTMP and it has the potential to sedate the sensitive incipient carbanions on account of gallium's strong carbophilicity (stronger than that of aluminum). Moreover, this organometallic trap exhibits good hydrocarbon solubility and does not require low temperature procedures, giving it a decided advantage over salt traps (for example, MgCl_2 , ZnCl_2),^[11] which generally need the use of ethereal solvents and often require low temperatures to avoid competing salt metathesis reactions. This potential has been duly realized through isolation and characterization (crystallographic, elemental analyses, and NMR spectroscopic) of an unprecedented series of gallium diazide and related complexes. As well as launching the concept of gallium TMT, this study reports the first crystal structures of metallodiazine complexes made by metalation (C–H to C–metal) reactions for gallium and indeed, bar one exception for zinc,^[8] for any

[*] M. Uzelac, Dr. A. R. Kennedy, Prof. E. Hevia, Prof. R. E. Mulvey
WestCHEM, Department of Pure and Applied Chemistry
University of Strathclyde
Glasgow, G1 1XL (UK)
E-mail: eva.hevia@strath.ac.uk
r.e.mulvey@strath.ac.uk

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under <http://dx.doi.org/10.1002/anie.201607284>.

© 2016 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

metal. Furthermore, the study highlights that two-metal synergistic reactions are not confined to concerted, synchronized processes where the metals belong within the same reagent, but can be extended to tandem, stepwise processes involving two separately added reagents that do not form a co-complex.

The study first established through NMR spectroscopy that LiTMP and GaR₃ remain separate in benzene solution^[12] (Supporting Information). Such separation is essential to action the lithiation step of TMT because, whereas free LiTMP is a strong base, combining it with, for example, *i*Bu₃Al to form aluminate LiAl(TMP)(*i*Bu)₃ greatly diminishes deprotonating power.^[10a] TMT was then attempted on three classical naked diazines, pyrazine, pyridazine, and pyridimine, as well as the related nitrogen–sulfur ring compound benzothiazole.

Pyrazine previously required four molar equivalents of LiTMP in THF at –75 °C, but only to afford modest yields of 2-substituted derivatives (39–65 %, depending on the electrophile) mixed in some cases with 2,5-disubstituted species (16 %).^[4] When performed at room temperature in hexane solution, our new LiTMP-trialkylgallium approach in a 1:1:1 stoichiometry with pyrazine selectively afforded the 2-monogallated pyrazine manifested in the crystalline complex [1-(PMDETA)Li-3-(GaR₃)-C₄H₃N₂] **1** (isolated yield 61 %; note NMR monitoring of reaction showed **1** is obtained quantitatively, see the Supporting Information).

The role of PMDETA is to aid crystallization and stabilization of the sensitive metallo species by chelating to lithium, but it is added at a later stage to avoid undergoing a competing TMT deprotonation itself. Stoichiometric control was also evident when the base:trap:substrate ratio was doubled to 2:2:1, giving the 2,5-digallated pyrazine [1,4-((PMDETA)Li)₂-2,5-((GaR₃)₂C₄H₂N₂)] **2** cleanly as crystals isolated in a 44 % yield (Scheme 2). However, NMR monitoring of the reaction revealed **2** forms in a 55 % yield, though a second regioisomer is formed in 33 %, which appears to be the analogous product of 2,6-digallation. This stoichiometric control contrasts with the performance of zincate [(THF)LiZn(TMP)*i*Bu₂], which operates through a synchronized bimetallic synergy distinct to that of stepwise TMT, as it affords only the 2,5-disubstituted pyrazine even with a 1:1, base:substrate stoichiometry.^[8] Previously, excess LiTMP (1.5 equivalents) dispensed as 0.5 ZnCl₂·TMEDA/1.5 LiTMP in THF produced 59 % of the isolated 2-iodopyrazine that reportedly decomposes at room temperature, but in hexane a significant amount of coupled dimer product was also seen.^[7a] Note that a control reaction between pyrazine and gallate LiGaR₄^[13] did not produce any gallation, but only R group addition, with concomitant dearomatization of the heterocycle (Supporting Information).

It may seem surprising that sensitive pyrazinyl mono- and di-carbanions can be trapped in crystalline form at room temperature and structurally defined, but this is where the structures become informative as they show that the heterocyclic units of **1** (Figure 1) and **2** (Figure 2) are cooperatively stabilized through coordination by both the Li and Ga centers that tie up the lone pairs of the N and C atoms, respectively. Both structures are monomeric with aggregation blocked at




Figure 1. Molecular structure of [1-(PMDETA)Li-3-(GaR₃)-C₄H₃N₂] (**1**) with 50 % probability displacement ellipsoids. All H atoms except those in the C₄H₃N₂ ring have been omitted for clarity.




Figure 2. Molecular structure of [1,4-((PMDETA)Li)₂-2,5-((GaR₃)₂C₄H₂N₂)] (**2**) with 50 % probability displacement ellipsoids. All H atoms except those in the C₄H₂N₂ ring have been omitted for clarity.

Li by tridentate PMDETA, though centrosymmetric **2** is tetranuclear having two Ga and two Li centers. Notably, **2** is the more congested structure with its GaR₃ units having proximal dispositions to the (PMDETA)Li units; whereas in **1** these units have a 1,3-separation. The Ga–sp²C(diazide) bond lengths show little variation with each other or with the Ga–sp³C(R group) bonds (see the Supporting Information for full crystallographic details and supporting NMR characterization).

With its 1,2-placement of N atoms, pyridazine offers a choice of metalation sites. Site selectivity in its metalation is exceptionally challenging, as evident from previous work using excess LiTMP in 0.5 ZnCl₂·TMEDA/1.5 LiTMP/I₂, which in hexane at room temperature achieved only 27 % of the 3-iodo product mixed with the 4-iodo, and 3,6-diiodo derivatives as well as 54 % unreacted pyridazine (note yields determined from NMR data).^[7a] In THF, the yield of the 3-iodo product rises to 66 % but only under extreme reflux conditions. On its own, LiTMP (4 equivalents) in THF at –75 °C produced only 16–32 % yields of 3-substituted pyridazines following quenching with different electrophiles.^[4] When run in hexane solution at room temperature, our TMT reaction afforded a 51 % yield of the isolated product [2-(PMDETA)Li-3-(GaR₃)-C₄H₃N₂] **3**. Interestingly, ¹H NMR

monitoring of the reaction revealed an important effect of the order of metal reagent addition on reaction regioselectivity. Thus, when LiTMP is added as a solid to the hexane solution of GaR₃ and pyridazine, **3** is obtained in a 78% yield, along with small amounts of the C4-gallated regioisomer (16% yield). Contrastingly, if the substrate is added to the hexane suspension of LiTMP and GaR₃, the yield of **3** decreases to 50%, and more C4 metalated product is seen (36%). These contrasting results suggest an activating effect of the GaR₃ component, which perhaps can initially coordinate to the Lewis basic N atoms of the heterocycle, facilitating its lithiation at C3. The crystal structure of **3** (Figure 3) shows





Figure 3. Molecular structure of [2-(PMDETA)Li-3-(GaR₃)-C₄H₃N₂] (**3**) with 50% probability displacement ellipsoids. All H atoms except those in the C₄H₃N₂ ring have been omitted for clarity.


GaR₃ elects to sit at the most acidic 3-position adjacent to one N (confirmed in solution by NMR spectra; Supporting Information). A novel feature is the Li(PMDETA) unit bridging the two diazine N atoms [Li–N bond lengths 2.093(5) and 2.043(5) Å for the non-disordered molecule of the Z' = 2 structure] leading to a 1, “2.5”-separation. Consequently, the spiro Li center, connecting the 3- and 2 × 5-atom rings, has a 5-coordinate geometry.

Because pyrimidine was found to be totally inert to LiTMP from 0°C to reflux temperatures,^[4] it seemed the greatest challenge to TMT. However, as in the case of **1–3**, TMT was demonstrated tangibly through isolation and crystallographic characterization of a metalated derivative, here [1-(PMDETA)Li-6-(GaR₃)-(C₄H₃N₂)], **4**. This was made in a 27% isolated crystalline yield, though NMR monitoring of the reaction shows that under the conditions studied **4** forms in a higher 59% yield. Its structure (Figure S13) exhibits many of the features in **1–3** with the proximal 1,6-separation of its GaR₃ and (PMDETA)Li akin to that of those in dimetalated pyrazine **2**.

In view of the fact that the 2-lithiated derivative of the related nitrogen–sulfur heterocycle benzothiazole is known to exist simultaneously in ring-closed and ring-open forms, as best evidenced by Boche’s ¹³C NMR studies in D₈-THF at –75°C, we extended the TMT study to this fused heterocycle.^[14] Run at room temperature in hexane solution (Scheme 3), an equimolar TMT reaction produced the crystalline complex [2-(GaR₃)-3-{Li(PMDETA)}C₆H₄NCS]



Scheme 2. Contrasting reactions of pyridazine with LiTMP and different lithium–gallium combinations.



Scheme 3. Main products of reactions of TMT mixture with pyridazine, pyrimidine, and benzothiazole.

5 (Scheme 3) in a remarkably high isolated yield of 84%. Significantly, **5** is quantitative in solution with no ring-opened metallo(2-isocyano)thiophenolate isomer detected. Deprotonative gallation of the C2 center was evident from its downfield resonance at 209.5 ppm in the ¹³C NMR spectrum. In the crystal, **5** follows the pattern in the TMT diazine series with the GaR₃ and Li(PMDETA) units adjacent on deprotonated C and N atoms, respectively, with a Ga1–C13–N1–Li1 torsion angle of –13.9(4)° (Figure 4). This first Ga TMT reaction of a N–S heterocycle is competitive with Mongin’s LiTMP/CdCl₂·TMEDA in THF solution approach, which used excess (1.5) base equivalents for a 97% yield of 2-iodobenzothiazole after I₂ quenching, though no metallo intermediate was identified.^[7b,15]



For synthetic campaigns, it is important that the new C–Ga bonds in these systems are accessible to electrophiles. In preliminary NMR spectroscopic experiments (see the Supporting Information for full details), on subjecting the gallated benzothiazole **5** to methyl trifluoromethanesulfonate (methyl triflate, MeOTf) we obtained a yield of 62% of the desired methylated product and did not observe any opening of the azole ring. In contrast, though this strong electrophile did quench the monogallated pyridazine **1**, it proved too aggressive and decomposition ensued. Therefore, we turned to a second electrophile in trimethylsilyl chloride, which offers another potential pitfall in being too bulky. Reassur-

Communications



Metalation

M. Uzelac, A. R. Kennedy, E. Hevia,*

R. E. Mulvey* Transforming LiTMP Lithiation of
Challenging Diazines through Gallium
Alkyl Trans-Metal-Trapping

Gallium trumps lithium: While LiTMP cannot be used to metalate synthetically useful heterocycles under standard conditions owing to the instability of lithiated intermediates, adding a gallium alkyl trap in a two-step, trans-metal-trapping process permits metalation and allows unprecedented trapping and crystallization of five highly sensitive anions within bimetallic formulations.