



**Farooq, Abdulaziz and Parkinson, Kathryn N and Adamson, Ashley J and Pearce, Mark S and Reilly, Jessica K and Hughes, Adrienne R and Janssen, Xanne and Basterfield, Laura and Reilly, John J (2016) Timing of the decline in physical activity in childhood and adolescence : Gateshead Millennium Cohort Study. British Journal of Sports Medicine. pp. 1-19. ISSN 0306-3674 (In Press) , <http://dx.doi.org/10.1136/bjsports-2016-096933>**

This version is available at <https://strathprints.strath.ac.uk/59471/>

**Strathprints** is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<https://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to the Strathprints administrator: [strathprints@strath.ac.uk](mailto:strathprints@strath.ac.uk)

**Timing of the decline in physical activity in childhood and adolescence: Gateshead Millennium Cohort Study**

Abdulaziz Farooq<sup>1,2</sup>, Kathryn N Parkinson<sup>3</sup>, Ashley J Adamson<sup>3,4</sup>, Mark S Pearce<sup>3</sup>, Jessica K Reilly<sup>4</sup>, Adrienne R Hughes<sup>1</sup>, Xanne Janssen<sup>1</sup>, Laura Basterfield<sup>4</sup>, John J Reilly<sup>1</sup>

**Affiliations:**

<sup>1</sup> University of Strathclyde, Glasgow, Physical Activity for Health Group, School of Psychological Sciences & Health, Scotland, UK G1 1XQ;<sup>2</sup>Aspetar Orthopaedic and Sports Medicine Hospital, Doha, Qatar ;<sup>3</sup>Institute of Health & Society, Newcastle University, Newcastle upon Tyne, UK;<sup>4</sup>Human Nutrition Research Centre, Newcastle University, Newcastle upon Tyne, UK

**Corresponding author:** Prof John J Reilly, University of Strathclyde Glasgow, Physical Activity for Health Group, G1 1XQ, Scotland

[john.j.reilly@strath.ac.uk](mailto:john.j.reilly@strath.ac.uk)

**Competing interests:** None declared

**Word count:** 2949 words

**Keywords:** Physical activity; exercise; child; adolescent; longitudinal study

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, a worldwide licence (<http://www.bmj.com/sites/default/files/BMJ%20Author%20Licence%20March%202013.doc>) to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints, include within collections and create summaries, extracts and/or, abstracts of the Contribution and convert or allow conversion into any format including without limitation audio, iii) create any other derivative work(s) based in whole or part on the on the Contribution, iv) to exploit all subsidiary rights to exploit all subsidiary rights that currently exist or as may exist in the future in the Contribution, v) the inclusion of electronic links from the Contribution to third party material where-ever it may be located; and, vi) licence any third party to do any or all of the above. All research articles will be made available on an Open Access basis (with authors being asked to pay an open access fee—see<http://www.bmj.com/about-bmj/resources-authors/forms-policies-and-checklists/copyright-open-access-and-permission-reuse>).

The terms of such Open Access shall be governed by a [Creative Commons](#) licence—details as to which Creative Commons licence will apply to the research article are set out in our worldwide licence referred to above.

Transparency: The manuscript guarantor (John Reilly) confirms that the manuscript is an honest, accurate, and transparent account of the study, no important aspects were omitted, and no discrepancies are applicable.

Data sharing: The data on which the manuscript is based are available from the University of Strathclyde PURE/KnowledgeBase.

Sponsorship: The sponsors (Universities of Strathclyde and Newcastle) are responsible for research governance, but have had no role in manuscript content, or in the decision to publish, or where to publish.

## **Abstract**

**Background and Aim** There is a widely held and influential view that physical activity begins to decline at adolescence. This study aimed to identify the timing of changes in physical activity during childhood and adolescence.

**Methods** Longitudinal cohort study (Gateshead Millennium Study) with eight years of follow-up, from Northeast England. Cohort members comprise a socioeconomically representative sample studied at ages 7, 9, 12 and 15 years; 545 individuals provided physical activity data at two or more time points. Habitual total volume of physical activity and moderate-to-vigorous intensity physical activity (MVPA) were quantified objectively using the Actigraph accelerometer over 5-7 days at the four time points. Linear mixed models identified the timing of changes in physical activity across the 8 year period, and trajectory analysis was used to identify sub-groups with distinct patterns of age-related changes.

**Results** Four trajectories of change in total volume of physical activity were identified representing 100% of all participants: all trajectories declined from age 7. There was no evidence that physical activity decline began at adolescence, or that adolescent declines in physical activity were substantially greater than the declines during childhood, or greater in girls than boys. One group (19% of boys) had relatively high MVPA which remained stable between ages 7-15 years.

**Conclusions** Future policy and research efforts to promote physical activity should begin well before adolescence, and should include both boys and girls.

**What are the new findings ?**

- Physical activity declines markedly during childhood, and the age-related decline does not begin at adolescence
- The physical activity decline from childhood occurs in both boys and girls
- Physical activity decline is not uniform- a small but significant minority of boys maintained relatively high and stable levels of MVPA across childhood and adolescence

**How might the findings impact on clinical practice in future ?**

- Future efforts to arrest age-related decline in physical activity should begin well before adolescence
- We should abandon the concept of 'high risk groups' (e.g. adolescent girls) for low physical activity- most of the paediatric population is at high risk

## Introduction

Systematic reviews have established that daily physical activity, particularly moderate-to-vigorous intensity physical activity (MVPA), is important for both the current and future health of children and adolescents.<sup>1,2</sup> For many years the prevailing view among researchers, policymakers, and health and education practitioners, based largely on subjective measurement of physical activity, has been that levels of physical activity are adequate during childhood, but decline dramatically during adolescence, with the adolescent decline being much more marked in girls than boys.<sup>3-6</sup> A more marked decline in girls may be related to their advanced pubertal maturation compared to boys, though the relative importance of biological and environmental influences on child and adolescent physical activity remains unclear.<sup>7</sup> Much research and policy effort has been directed at adolescent girls, considered to be the 'high risk group' for low physical activity,<sup>3-6</sup> with the implicit or explicit assumption that children and boys are at low-risk. Many national and international physical activity surveillance programmes focus on adolescents,<sup>8</sup> also based on implicit or explicit assumptions that adolescence is the stage of the life-course when physical activity declines dramatically.

While levels of MVPA among adolescents are typically much lower than recommended<sup>8</sup> there is actually very little evidence that MVPA declines with the onset of adolescence, or that adolescent declines in MVPA are more marked in girls than boys. Dumith et al<sup>9</sup> published the most recent systematic review of changes in physical activity during adolescence, with 26 eligible longitudinal studies. Of these eligible studies only 2/26 included objective measures of the intensity of physical activity (accelerometry), most (16/26) were from the USA, and the representativeness of the samples in the eligible studies was usually unclear<sup>9</sup>. Moreover, since most studies completed data collection before 2000, they probably lack generalisability to contemporary populations given recent and dramatic societal changes: declines in active

commuting,<sup>10</sup> in the technology being used by children<sup>10</sup> (e.g. the ubiquitous nature of new screen-based media),<sup>11</sup> and in overweight and obesity (which appear to reduce physical activity).<sup>12</sup> The International Children's Accelerometry Database, though cross-sectional, suggests that objectively measured physical activity may be in decline in both sexes from around the age of school entry<sup>13</sup>. We are aware of only one single longitudinal study which measured physical activity across childhood and adolescence: in children born in Iowa in the early 1990's physical activity declined, in both sexes, from age 5 years in most study participants<sup>14</sup>. In summary, despite the well-established and influential concept that declines in physical activity begin at adolescence, an emerging body of evidence suggests that declines may occur earlier, but there is currently very little high quality longitudinal evidence on the timing of changes in physical activity across childhood and adolescence in contemporary cohorts. The primary aim of the present study was therefore to describe the timing of changes in physical activity across childhood and adolescence, to identify *when* changes take place.

## **Methods**

### **Participants, recruitment, consent, and study design.**

Participants were from the Gateshead Millennium Cohort Study (GMS). Details of this cohort study have been published previously,<sup>15</sup> but, in brief, the sample was socio-economically representative of the north-east of England.<sup>15</sup> This was a longitudinal study and the intention was to repeat measures in as many of the original cohort members as possible at subsequent time-points. Habitual physical activity measurements in the cohort began when participants were 7 years of age (October 2006 to December 2007), and were repeated at 9 years of age (October 2008 to September 2009), 12 years of age (October 2011 to December 2012) and 15 years of age (September 2014 to September 2015). The study was approved by the Gateshead and South Tyneside Local National Health Service Research Ethics Committee for data collection at age 7,

and by the Newcastle University Faculty of Medical Sciences Ethics Committee for subsequent measures. Informed written consent was obtained from the parent/main guardian of each child, and children and adolescents assented to participation.

### **Objective measurement of total volume of physical activity and MVPA**

Habitual total ‘volume’ of physical activity (i.e. light intensity physical activity combined with MVPA) and MVPA were measured using Actigraph GT1M (Actigraph, Pensacola, FL, USA) accelerometers at all four time points. The Actigraph is a small, unobtrusive and lightweight monitor with high validity (relative to reference measures of physical activity, direct observation, and energy expenditure), high reliability, and low reactivity in children,<sup>16</sup> and has been used successfully in many paediatric physical activity studies internationally.<sup>13</sup>

The methodology for measurement of physical activity in the GMS cohort is described in detail elsewhere.<sup>17-21</sup> In brief, participants were asked to wear the accelerometer during waking hours for 7 days, removing only for bed and water-based activities such as showering, bathing, and swimming. Families were also given a log sheet to note when the monitor had been worn and removed each day. The accelerometers were set to record data in 15 second sampling intervals (epochs) throughout the period the accelerometer was worn. Data were reduced manually using accelerometry output and log sheets, as described previously,<sup>17-21</sup> to delete periods of non-wear time (corresponding to times noted by the parent, child, or adolescent, as un-worn). Total volume of physical activity was measured using the accelerometer output or counts per minute (cpm). Measuring the intensity of physical activity, specifically MVPA, requires that an accelerometer output threshold - ‘cut-off point’ –is applied. Two evidence-based but distinct ‘cut-off points’ have been used widely in the literature, but in the present study the cut-point of 574 counts per 15s was used<sup>22</sup> and it has acceptable classification accuracy.<sup>23</sup> Since the Actigraph cut-off points



used in this study were derived from studies using Actigraph model 7164 and the more recent Actigraph GT1M model has a consistent bias of 9% relative to model 7164,<sup>24</sup> a +9% correction was applied to the raw accelerometry data before applying cut-off points. In UK children there are small seasonal variations in objectively measured habitual physical activity<sup>19,20,25</sup> and so accelerometry measurements at ages 7,9, and 12 years were made at the same time of year as described previously,<sup>17-21</sup> at age 15 years close matching for season was not possible in all cases, hence accelerometry data were adjusted for season (summer or winter) during analysis.

### **Anthropometric and descriptive characteristics of study participants**

Height was measured to 0.1cm with a Leicester portable height measure (Chasmors, London, UK) and weight measured to 0.1kg in light indoor clothing and % body fat estimated with a Tanita TBF300MA (Chasmors, London UK).<sup>17-21</sup> Body mass index (BMI, kg/m<sup>2</sup>) was calculated, and z-scores estimated to UK 1990 population reference data.<sup>26</sup> Socio-economic status was assessed using Townsend scores at birth,<sup>27</sup> an area-based measure derived from the UK census.

### **Statistical analysis**

The sample size for the present study was fixed by the size of the recruited cohort at age 7 years, and the extent of sample attrition over the eight-year period. However, the sample size was similar to or larger than most previous studies with two measures of physical activity.<sup>9,14</sup> Further, the accuracy of both the reported levels of PA and SB and the changes in levels over the two year period is increased by the use of accelerometry and the longitudinal design and analyses used by the study. Linear mixed models were used with SPSS v21.0 statistical software to describe age-related changes in physical activity. Gender, age groups and their interaction were fixed factors and participant ID was the clustering random variable using unstructured covariance structure for age group as repeated measurement. For the MVPA data at age 15 years, season (winter or

summer) was added as a factor to adjust for seasonal variation in physical activity. Posthoc pairwise differences were reported as absolute change with 95% confidence intervals (CI) after Bonferroni adjustment.

Group-based modelling of longitudinal data to identify distinct trajectories of age-related changes in physical activity was performed using the Traj module plug-in in STATA v11.0. This method, initially developed by Nagin et al,<sup>28</sup> has recently been applied to longitudinal physical activity data.<sup>14</sup> Group-based modelling to identify trajectories can identify patterns of change, incorporating multiple different time-trends and multiple different levels at baseline.<sup>28</sup> Given potential differences in trajectories of physical activity change between the sexes, they were modelled separately for boys and girls. The trajectories were modelled using either quadratic or linear trends. In a step by step manner, we increased the number of distinct trajectories that can be extracted from the data, first by evaluating the posterior probabilities for each model for tenability and by comparing fit statistics as defined by Bayesian Information Criterion (BIC) to be lowest. The process of specifying the number of distinct trajectories and selection of linear vs quadratic trend for each trajectory continued until there was no substantial improvement in model fit. Non-significant quadratic terms were replaced with linear terms irrespective of significance to ensure parsimony of the model.<sup>29</sup> Using a maximum probability assignment rule, each participant was classified under a trajectory for which posterior membership probability was the highest. Trajectories and the 95% confidence intervals (CI) surrounding each trajectory were plotted.

## **Results**

### **Characteristics of study participants**

A total of 545 individuals provided accelerometry data twice, and 217 at all four time points, with the sample size at each time point as follows: 431 (7 years), 428 (9 years), 385 (12 years),

and 278 (15 years). The study flow diagram is provided in supplementary Figure 1. Participants who provided data at least twice had similar physical activity and similar BMI-z scores at baseline compared to those who had missing or invalid accelerometer data (see supplementary Table 1). Girls with more than two missing/invalid data points had slightly higher BMI z-scores at baseline than those with complete data, but similar physical activity.

### **Summary data on physical activity at each of the four measurements**

Summary data on total volume of physical activity ([accelerometer count per minute, a measure of light intensity physical activity and MVPA combined](#)) and MVPA, at each time point between ages 7 - 15 years, is shown in Table 1. Mean accelerometry monitoring period ranged from 5.3 to 6.4 days and 667 to 725 minutes per day across the four time points.

### **Changes in physical activity analysed longitudinally using trajectory analysis**

The longitudinal trajectory analyses for changes in *total volume of physical activity* are shown in Figure 1 for boys and girls: in the boys three distinct trajectories were identified, but in all three sub-groups, representing 100% of the sample, the total volume of physical activity declined across the eight year period from age 7 years; in the girls a single trajectory of declining total physical activity across the eight year period from age 7 years characterised all study participants. None of the trajectories of total volume of physical activity showed initial high levels followed by sharp adolescent declines.

The longitudinal trajectories of change in MVPA for boys and girls are shown in Figure 2. In boys, four distinct sub-groups were identified: low and relatively stable/slowly declining MVPA from age 7 across the 8 years (3% of the sample); initially high at age 7, but rapidly declining

from age 7 (17% of the sample); gradually declining from age 7 (61% of the sample); relatively high and stable or increasing from age 7 years (19% of the sample).

The longitudinal trajectories of change in MVPA of girls are also shown in Figure 2. Three distinct trajectories were identified: low and relatively stable/ with modest declines (19%); gradually declining from age 7 years (62%); high initial MVPA with steeper declines from age 7 (19%). None of the trajectories of MVPA in either boys or girls were characterised by high and stable childhood values followed by sharp adolescent declines.

## **Discussion**

### **Main study findings and implications**

The present study suggests that total volume of physical activity is already declining by age 7 years in the UK. The present study found that 100% of boys and girls fitted into longitudinal trajectories which were inconsistent with the orthodox view that physical activity begins to decline at adolescence, declines much more rapidly at adolescence, and/or declines much more rapidly in adolescent girls than boys. The present study therefore suggests that much of the current UK and international policy and practice in this area, based on the orthodox view, is not evidence-based. The main implication of our findings for future research and public health policy in physical activity and health is the need for a much greater emphasis on childhood rather than adolescence, and on both boys and girls.

Trajectories of change in MVPA were more complex than trajectories of change in total volume of physical activity, but it is of note that in around 19% of boys MVPA was relatively high and fairly stable across the eight year period. These distinct trajectories of physical activity are new, and their determinants are unclear, but understanding why a minority of participants maintained a

relatively high and stable level of MVPA across childhood and adolescence would help inform future policy and clinical practice.

### **Consistency of evidence from present study with other studies**

The International Children's Accelerometry Database (ICAD), a global data-pooling initiative, has recently demonstrated that mean values for total volume of physical activity and MVPA both appear to decline steadily after the age of 5 years.<sup>13</sup> While this ICAD study was cross-sectional, and most samples not contemporary, ICAD data also provide no support for the view that physical activity decline either begins at adolescence, or increases markedly at adolescence. In the UK, recent nationally representative but cross-sectional accelerometry measures also suggest that total volume of physical activity begins to declines around the age of school-entry.<sup>30</sup>

Since the recent systematic review on changes in physical activity during adolescence by Dumith et al,<sup>9</sup> four English cohort studies have reported on objectively measured longitudinal changes in MVPA *during* adolescence, though these studies do not span **both** childhood and adolescence. Harding et al<sup>31</sup> (n 363; age 12-15 years) and Mitchell et al<sup>32</sup> (n 5436; 12-16 years) found negligible declines in objectively measured MVPA during adolescence. Corder et al<sup>6</sup> (n 480; 10-14 years) and Collings et al<sup>33</sup> (n 144; 15-17 years) found small declines in MVPA which were slightly greater in boys than girls. In summary, recent English longitudinal studies which used objective measures of physical activity are consistent with the present study in suggesting that physical activity does not decline especially markedly at adolescence, and does not decline more markedly in girls than boys during adolescence.

Data from the Iowa Bone Development Study was consistent with the present study in suggesting that total volume of physical activity declined either steadily or dramatically from age 5 years in

> 70% of participants<sup>14</sup>. Other evidence is also consistent with the findings of the present study in suggesting that lifestyles of English children become particularly ‘obesogenic’ during mid-childhood. In the ALSPAC cohort in England for example the incidence (appearance of new cases) of obesity across childhood and adolescence peaked around age 7 years,<sup>34</sup> and among non-obese individuals the rate of excessive weight gain also peaked at the same time.<sup>35</sup> The recent systematic review by Tanaka et al also found that accelerometer-measured daily sedentary time increased from around the age of school entry, and as sedentary time increased with age physical activity was displaced.<sup>36</sup>

### **Study strengths and weaknesses**

The main strengths of the present study were the longitudinal design, high quality objective measures of physical activity, socio-economically representative sample, and the contemporary nature of the cohort. One weakness may be generalisability to populations with substantially different physical, social, and/or policy environments. Further research will therefore be necessary to determine whether or not the findings of the present study are generalisable to other populations. The present study aimed to describe *when* changes in physical activity occurred. The issue of *why* any changes take place (e.g. the relative importance of biological versus environmental factors)<sup>7</sup> is important, but is a distinct question which was beyond the scope of the present study. Changes in physical activity arise as a result of a complex and currently poorly understood<sup>7</sup> combination of biological and environmental influences. An improved understanding of why changes in physical activity take place and why individuals follow one trajectory rather than another, would also be important to future public health applications, but would require measures which were not available in the present study (e.g. ideally direct measures of biological variables such as pubertal maturation)<sup>7</sup>. Trajectory analysis can also act as a tool to predict future MVPA based on current MVPA status, and future research could also identify how accurate such

predictions might be. The trajectory analyses identified relatively large sub-groups, with one exception: the 3% of boys with low and relatively stable/slowly declining MVPA over the 8 years. Exclusion of this sub-group would have reduced model fit as judged by the BIC and confidence intervals around the groups, and so it was retained. Future studies would be useful to confirm whether or not this small group exists in other populations. Finally, the periods between the four measurement periods in the present study were chosen so as to be adequate to detect longitudinal changes in physical activity between measurements (ie to permit real changes to be distinguishable from measurement imprecision). However, timing of measurements was determined largely by the availability of grant funding, and more frequent measurements might have provided more nuanced evidence on the timing of changes.

## **Conclusions**

The present study suggests that physical activity is in decline in most individuals of both sexes by age 6-7 years in the UK. The study questions the concept of the adolescent girl as a priority for research and policy effort in physical activity. Future research and public health policy should focus on (a) preventing the decline in physical activity which begins in childhood, not adolescence and (b) providing an improved understanding of the determinants of the different physical activity trajectories, including an understanding of the relative importance of biological and environmental influences<sup>7</sup>.

## **Acknowledgements and funding**

This study was supported by grants from the National Prevention Research Initiative reference (grant G0501306), and the Scottish Government Chief Scientist Office (grants CZH/4/484 and CZH/4/979) and the University of Strathclyde. The cohort was established with funding from the

Henry Smith Charity and Sport Aiding Medical Research in Kids (SPARKS) and followed up with grants from Gateshead NHS Trust R&D, Gateshead Council, Northern and Yorkshire NHS R&D, Northumberland, Tyne and Wear NHS Trust,. We appreciate the support of PCT, Gateshead Education Authority, local schools, the External Reference Group, the research team in conducting the study, and especially thank the families and children/adolescents for participating in the study. During this study period AJA was partly funded as member of Fuse, the Centre for Translational Research in Public Health, a UK Clinical Research Collaboration (UKCRC) Public Health Research Centre of Excellence. Funding for Fuse from the British Heart Foundation, Cancer Research UK, Economic and Social Research Council, Medical Research Council, and the National Institute for Health Research, under the auspices of the UKCRC, is gratefully acknowledged. AJA is currently funded by the National Institute of Health Research. Opinions expressed are not necessarily those of the funders.

### **Conflicts of interest**

The authors declare no conflicts of interest.

### **Author Contributions**

Conception/Design: KP,AA,MP, JR,AH,XJ, LB, JR; Data analysis/acquisition/interpretation: MF, JR, JR, LB, XJ.

Drafting/Revising critically for important content: all authors.

Final approval: all authors.

Accountability for study and manuscript: all authors.



## References Cited

1. Janssen I, Leblanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int J Behav Nutr Phys Act.* 2010;7:40.
2. Timmons BW, Leblanc AG, Carson V, et al. Systematic review of physical activity and health in the early years (aged 0-4 years). *Appl Physiol Nutr Metab.* 2012;37(4):773-792.
3. Physical Activity Review Group. *Five year review of 'Let's make Scotland more active: a strategy for physical activity': Update statement from the review group.* Edinburgh: NHS Health Scotland;2010.
4. SportsScotland. *Increasing demand for sport and physical activity for adolescent girls in Scotland: Exploring issues and suggesting solutions.* Edinburgh: sportsScotland;2006.
5. Physical Activity and Health Alliance. Physical Activity and Adolescent Girls – Knowledge Exchange Network. 2009; <http://www.paha.org.uk/Feature/physical-activity-and-adolescent-girls-knowledge-exchange-network>. Accessed Jul 2016.
6. Corder K, Sharp SJ, Atkin AJ, et al. Change in objectively measured physical activity during the transition to adolescence. *Br J Sports Med.* 2015;49(11):730-736.
7. Sherar LB, Cumming SP, Eisenmann JC, Baxter-Jones ADG, Malina RM. Adolescent biological maturity and physical activity: biology meets behavior. *Pediatr Exerc Sci* 2010; 22: 332-349.
8. Hallal PC, Andersen LB, Bull FC et al. Global physical activity levels: surveillance progress, pitfalls, and prospects. *Lancet.* 2012;380(9838):247-257.
9. Dumith SC, Gigante DP, Domingues MR et al. Physical activity change during adolescence: a systematic review and a pooled analysis. *Int J Epidemiol.* 2011;40(3):685-698.
10. Pratt M, Sarmiento OL, Montes F, et al. The implications of megatrends in information and communication technology and transportation for changes in global physical activity. *Lancet.* 2012;380(9838):282-293.
11. Christakis DA. Interactive media use at younger than the age of 2 years: time to rethink the American Academy of Pediatrics guideline? *JAMA Pediatr.* 2014;168(5):399-400.
12. Bauman AE, Reis RS, Sallis JF et al. Correlates of physical activity: why are some people physically active and others not? *Lancet.* 2012;380(9838):258-271.
13. Cooper AR, Goodman A, Page AS, et al. Objectively measured physical activity and sedentary time in youth: the International children's accelerometry database (ICAD). *Int J Behav Nutr Phys Act.* 2015;12:113
14. Kwon S, Janz KF, Letuchy EM et al. Developmental Trajectories of Physical Activity, Sports, and Television Viewing During Childhood to Young Adulthood: Iowa Bone Development Study. *JAMA Pediatr.* 2015;169(7):666-672.
15. Parkinson KN, Pearce MS, Dale A, et al. Cohort profile: the Gateshead Millennium Study. *Int J Epidemiol.* 2011;40(2):308-317.
16. Reilly JJ, Penpraze V, Hislop J et al. Objective measurement of physical activity and sedentary behaviour: review with new data. *Arch Dis Child.* 2008;93(7):614-619.
17. Basterfield L, Adamson AJ, Frary JK et al. Longitudinal study of physical activity and sedentary behavior in children. *Pediatrics.* 2011;127(1):e24-30.
18. Basterfield L, Pearce MS, Adamson AJ, et al. Physical activity, sedentary behavior, and adiposity in English children. *Am J Prev Med.* 2012;42(5):445-451.
19. King AC, Parkinson KN, Adamson AJ, et al. Correlates of objectively measured physical activity and sedentary behaviour in English children. *Eur J Public Health.* 2011;21(4):424-431.

20. Pearce MS, Basterfield L, Mann KD et al. Early predictors of objectively measured physical activity and sedentary behaviour in 8-10 year old children: the Gateshead Millennium Study. *PLoS One*. 2012;7(6):e37975.
21. Basterfield L, Adamson AJ, Pearce MS et al. Stability of habitual physical activity and sedentary behavior monitoring by accelerometry in 6- to 8-year-olds. *J Phys Act Health*. 2011;8(4):543-547.
22. Evenson KR, Catellier DJ, Gill K et al. Calibration of two objective measures of physical activity for children. *J Sports Sci*. 2008;26(14):1557-1565.
23. Trost SG, Loprinzi PD, Moore R, Pfeiffer KA. Comparison of accelerometer cutpoints for predicting activity intensity in youth. *Med Sci Sports Exerc* 2011; 43: 1360-1368.
24. Corder K, Brage S, Ramachandran A et al. Comparison of two Actigraph models for assessing free-living physical activity in Indian adolescents. *J Sports Sci*. 2007;25(14):1607-1611.
25. Carson V, Spence JC. Seasonal variation in physical activity among children and adolescents: a review. *Pediatr Exerc Sci*. 2010;22(1):81-92.
26. Cole TJ, Freeman JV, Preece MA. Body mass index reference curves for the UK, 1990. *Arch Dis Child*. 1995;73(1):25-29.
27. Townsend P, Phillimore P, Beattie A. *Health and Deprivation: Inequality in the North*. . London: Croom Helm, Limited; 1988.
28. Nagin D. *Group-Based Modeling of Development*. Cambridge: Harvard University Press; 2005.
29. Helgeson VS, Snyder P, Seltman H. Psychological and physical adjustment to breast cancer over 4 years: identifying distinct trajectories of change. *Health Psychol*. 2004;23(1):3-15.
30. Food Standards Scotland. National Diet and Nutrition Survey Rolling programme Results from Years 1-4 (combined) for Scotland (2008/09-2011/12). 2014; <http://www.foodstandards.gov.scot/national-diet-and-nutrition-survey-rolling-programme-results-years-1-4-combined-scotland-200809>. Accessed Jul 2016.
31. Harding SK, Page AS, Falconer C et al. Longitudinal changes in sedentary time and physical activity during adolescence. *Int J Behav Nutr Phys Act*. 2015;12:44.
32. Mitchell JA, Pate RR, Dowda M, et al. A prospective study of sedentary behavior in a large cohort of youth. *Med Sci Sports Exerc*. 2012;44(6):1081-1087.
33. Collings PJ, Wijndaele K, Corder K, et al. Magnitude and determinants of change in objectively-measured physical activity, sedentary time and sleep duration from ages 15 to 17.5y in UK adolescents: the ROOTS study. *Int J Behav Nutr Phys Act*. 2015;12:61.
34. Hughes AR, Sherriff A, Lawlor DA et al. Incidence of obesity during childhood and adolescence in a large contemporary cohort. *Prev Med*. 2011;52(5):300-304.
35. Hughes AR, Sherriff A, Lawlor DA et al. Timing of excess weight gain in the Avon Longitudinal Study of Parents and Children (ALSPAC). *Pediatrics*. 2011;127(3):e730-736.
36. Tanaka C, Reilly JJ, Huang WY. Longitudinal changes in objectively measured sedentary behaviour and their relationship with adiposity in children and adolescents: systematic review and evidence appraisal. *Obes Rev*. 2014;15(10):791-803.

### Figure legends

Figure 1: Change in total physical activity in boys (1A) and girls (1B) based on trajectory analysis.

Figure 2: Change in moderate-to-vigorous intensity physical activity in boys (2A) and girls (2B) based on trajectory analysis.

Table 1. Physical activity measurements and characteristics (Mean (Standard Deviation)) of study participants included in the analysis at the four time points

	Boys				Girls			
	Age 7 (n=217)	Age 9 (n=209)	Age 12 (n=176)	Age 15 (n=131)	Age 7 (n=214)	Age 9 (n=219)	Age 12 (n=209)	Age 15 (n=147)
Age (years)	7.5 (0.5)	9.3 (0.4)	12.5 (0.3)	15.2 (0.4)	7.5 (0.4)	9.3 (0.4)	12.5 (0.3)	15.2 (0.4)
<b>Anthropometry*</b>								
Height (cm)	125.5 (5.7)	135.9 (6.3)	154.2 (8.7)	171.6 (8.1)	124.7 (5.8)	135.4 (6.3)	155.2 (7.3)	163.0 (6.2)
Weight (kg)	26.5 (5.5)	33.0 (7.2)	48.3 (11.8)	63.4 (14.5)	26.7 (5.9)	33.8 (7.7)	51.0 (12.4)	61.8 (14.4)
Body mass index (kg/M <sup>2</sup> )	16.7 (2.4)	17.7 (2.8)	20.1 (3.7)	21.4 (4.0)	17.1 (2.6)	18.3 (3.0)	21.0 (4.0)	23.2 (4.6)
BMI z-score	0.4 (1.2)	0.5 (1.2) <sup>a</sup>	0.7 (1.2) <sup>ab</sup>	0.5 (1.3) <sup>c</sup>	0.4 (1.1)	0.6 (1.1) <sup>a</sup>	0.7 (1.1) <sup>ab</sup>	0.8 (1.1) <sup>ab</sup>
Fat mass (kg)	6.5 (3.2)	8.7 (5.0)	12.0 (8.4)	14.2 (10.6)	6.3 (3.6)	9.2 (5.0)	14.2 (8.6)	21.8 (11.5)
<b>Accelerometer data</b>								
Wear time (minutes/day)	672 (74.5)	673 (77.6)	712 (87.6)	725 (88.4)	667 (70.9)	671 (75.6)	718 (82.8)	720 (87.2)
Days worn	6.4 (1.0)	5.9 (1.3)	5.8 (1.4)	5.3 (1.6)	6.2 (1.2)	5.9 (1.3)	5.7 (1.5)	5.4 (1.7)
Total physical activity (cpm)†	780 (223) <sup>bcd</sup>	715 (192) <sup>cd</sup>	558 (189) <sup>d</sup>	482 (154)	746 (222) <sup>bcd</sup>	653 (193) <sup>cd</sup>	469 (191) <sup>d</sup>	400 (156)
MVPA (min/day)†	75.5 (25.5) <sup>bcd</sup>	70.0 (23.9) <sup>cd</sup>	60.3 (24.4) <sup>d</sup>	51.4 (22.4)	63.4 (20.9) <sup>bcd</sup>	56.4 (19.8) <sup>cd</sup>	47.4 (20.9) <sup>d</sup>	41.0 (16.8)

cpm: accelerometer counts per minute, a measure of the total amount of physical activity (light intensity physical activity and MVPA combined)

MVPA: Moderate-to-vigorous intensity physical activity, adjusted for season (Winter vs. Summer) at age 15 years

† Estimated marginal means derived from linear mixed model analysis.

a: significantly higher than age 7; b: significantly higher than age 9; c: significantly higher than age 12; d: significantly higher than age 15