

## **Drug-Free Platelets Can Act as Seeds for Aggregate Formation During Antiplatelet Therapy.**

Hoefer, T; Armstrong, PC; Finsterbusch, M; Chan, MV; Kirkby, NS; Warner, TD

For additional information about this publication click this link.

<http://qmro.qmul.ac.uk/jspui/handle/123456789/9199>

Information about this research object was correct at the time of download; we occasionally make corrections to records, please therefore check the published record when citing. For more information contact [scholarlycommunications@qmul.ac.uk](mailto:scholarlycommunications@qmul.ac.uk)

## Drug-Free Platelets Can Act as Seeds for Aggregate Formation During Antiplatelet Therapy

Thomas Hoefler, Paul C. Armstrong, Michaela Finsterbusch, Melissa V. Chan, Nicholas S. Kirkby, Timothy D. Warner

**Objective**—Reduced antiplatelet drug efficacy occurs in conditions of increased platelet turnover, associated with increased proportions of drug-free, that is, uninhibited, platelets. Here, we detail mechanisms by which drug-free platelets promote platelet aggregation in the face of standard antiplatelet therapy.

**Approach and Results**—To model standard antiplatelet therapy, platelets were treated in vitro with aspirin, the P2Y<sub>12</sub> receptor blocker prasugrel active metabolite, and aspirin plus prasugrel active metabolite. Different proportions of uninhibited platelets were then introduced. Light transmission aggregometry analysis demonstrated clear positive associations between proportions of drug-free platelets and percentage platelet aggregation in response to a range of platelet agonists. Using differential platelet labeling coupled with advanced flow cytometry and confocal imaging, we found aggregates formed in mixtures of aspirin-inhibited platelets and drug-free platelets were characterized by intermingled platelet populations. This distribution is in accordance with the ability of drug-free platelets to generate thromboxane A<sub>2</sub> and so drive secondary platelet activation. Conversely, aggregates formed in mixtures of prasugrel active metabolite–inhibited or aspirin plus prasugrel active metabolite–inhibited platelets together with drug-free platelets were characterized by distinct cores of drug-free platelets. This distribution is consistent with the ability of drug-free platelets to respond to the secondary activator ADP.

**Conclusions**—These experiments are the first to image the interactions of inhibited and uninhibited platelets in the formation of platelet aggregates. They demonstrate that a general population of platelets can contain subpopulations that respond strikingly differently to overall stimulation of the population and so act as the seed for platelet aggregation. (*Arterioscler Thromb Vasc Biol.* 2015;35:00-00. DOI: 10.1161/ATVBAHA.115.306219.)

**Key Words:** aspirin ■ flow cytometry ■ P2Y<sub>12</sub> receptor ■ prasugrel ■ thromboxane

Antiplatelet drugs are a cornerstone of preventive therapy for patients at risk of atherothrombotic events. Typically, these patients take low-dose aspirin (≈75–100 mg) once a day. Patients requiring further protection, such as following acute coronary syndrome, particularly with ST-segment elevation, or percutaneous coronary intervention, take a P2Y<sub>12</sub> receptor blocker, most commonly clopidogrel, in addition to aspirin.<sup>1–4</sup> This dosing regimen is commonly known as dual antiplatelet therapy (DAPT). Despite these therapeutic approaches, subsequent thrombotic events still occur and research has focused on ways to quantify individual patient risk. As a result, the concepts of antiplatelet drug resistance and high on-treatment platelet reactivity have developed.<sup>5–11</sup> Through this research, it has become apparent that patients who have conditions associated with increased platelet turnover also have particularly elevated risks.<sup>4,12–15</sup> Collectively, these observations lead to the

idea that increased risk while on antiplatelet therapy may be explained by the relationship between the pharmacokinetic and pharmacodynamic properties of aspirin and P2Y<sub>12</sub> inhibitors, and the dynamic changes associated with turnover of the platelet population.

Aspirin is rapidly absorbed from the stomach and small intestine reaching a plasma peak around 1 hour later.<sup>16</sup> It is then rapidly metabolized by esterases in the gut, liver, and circulation and has a circulating half-life of 13 to 19 minutes. Overall, this means that little active aspirin remains in the body 2 to 3 hours after consumption. However, as aspirin irreversibly acetylates and blocks the cyclooxygenase (COX)-1 enzyme in platelets, and as anucleated platelets are limited in their ability to generate new COX-1, the inhibitory effect of aspirin on platelets persists for their entire life span of 7 to 12 days.<sup>4,17</sup> The thienopyridine P2Y<sub>12</sub> blockers, clopidogrel

Received on: March 3, 2015; final version accepted on: July 30, 2015.

From The William Harvey Research Institute, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, Charterhouse Square, London, United Kingdom (T.H., P.C.A., M.F., M.V.C., T.D.W.); and National Heart and Lung Institute, Imperial College London, London, United Kingdom (N.S.K.).

The online-only Data Supplement is available with this article at <http://atvb.ahajournals.org/lookup/suppl/doi:10.1161/ATVBAHA.115.306219/-/DC1>.

Correspondence to Tim D. Warner, The William Harvey Research Institute, Barts and the London School of Medicine and Dentistry, Queen Mary London, Charterhouse Square, London EC1M 6BQ, United Kingdom. E-mail [t.d.warner@qmul.ac.uk](mailto:t.d.warner@qmul.ac.uk)

© 2015 The Authors. *Arteriosclerosis, Thrombosis, and Vascular Biology* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution, and reproduction in any medium, provided that the original work is properly cited.

*Arterioscler Thromb Vasc Biol* is available at <http://atvb.ahajournals.org>

DOI: 10.1161/ATVBAHA.115.306219

Nonstandard Abbreviations and Acronyms	
AA	arachidonic acid
COX	cyclooxygenase
DAPT	dual antiplatelet therapy
GP	glycoprotein
LTA	light transmission aggregometry
PAM	prasugrel active metabolite
TX	thromboxane

and prasugrel, are prodrugs with complex pharmacokinetics requiring metabolism to thiolactone active metabolites that are irreversible receptor antagonists of platelet P2Y<sub>12</sub> receptors.<sup>18–20</sup> Like aspirin, these active metabolites are short lived in the circulation but produce long-lasting effects because of irreversible inhibition of their target. Given these pharmacokinetic and pharmacodynamic profiles of aspirin and clopidogrel/prasugrel, we can reason that platelets that enter the circulation after the drugs have been cleared will not be inhibited and that the proportion of circulating drug-free platelets will increase in proportion to the rate of platelet turnover. Elevated rates of platelet production are associated with many conditions/procedures which feature increased thrombotic risk including patients undergoing coronary artery bypass surgery<sup>21</sup> or those with diabetes mellitus,<sup>22–27</sup> chronic kidney disease,<sup>28,29</sup> metabolic syndrome,<sup>30</sup> and essential thrombocythemia.<sup>31,32</sup> In these patient groups, increased circulating subpopulations of uninhibited platelets may be of crucial importance in providing the seed for the formation of occlusive platelet thrombi. Drug-free platelets could also drive the changes noted in *ex vivo* tests of platelet reactivity.

In the *in vitro* study reported here, we have assessed the impact of drug-free platelet subpopulations on overall platelet responses to antiplatelet therapy by modeling the entry of drug-free platelets into the circulation and used confocal microscopy and flow cytometric imaging to provide unique images of the interactions of inhibited and drug-free platelet subpopulations in the formation of platelet aggregates. These studies provide insight into potential mechanisms underlying the disease-associated incidence of thrombotic events and illustrate important concepts relevant to ideas of antiplatelet drug resistance.

## Materials and Methods

Materials and Methods are available in the online-only Data Supplement.

## Results

### Influence of Drug-Free Platelets on Light Transmission Aggregometry Responses

First, we modeled the effects of drug-free platelets on responses recorded by light transmission aggregometry (LTA) having previously established the effectiveness of antiplatelet drugs (Table I in the online-only Data Supplement). Increasing proportions of drug-free platelets were added to inhibited platelet populations, and the net aggregatory responses were recorded (Figure 1; Figure II in the online-only Data


Supplement). Aggregation to the COX-1 substrate arachidonic acid (AA; 1 mmol/L) in an aspirin-inhibited platelet population was returned to a full response by the inclusion of 30% drug-free platelets (Figure 1Ai). In general terms, increasing aggregation was seen with increasing platelet number in the absence of drug, that is, shifting from platelet-poor plasma to platelet-rich plasma (Figure III in the online-only Data Supplement). Measurement of thromboxane B<sub>2</sub> (TXB<sub>2</sub>) as a measure of the formation of the COX product TxA<sub>2</sub> demonstrated a linear increase of TXB<sub>2</sub> with increasing proportions of drug-free platelets ( $R^2=0.8867$ ), ranging from  $1.6\pm 0.4$  ng/mL TXB<sub>2</sub> when 0% platelets were aspirin-free to  $1220\pm 126$  ng/mL when 100% platelets were aspirin-free (Figure 1C). Similarly, aggregation to AA (1 mmol/L) in a prasugrel active metabolite (PAM)-inhibited platelet population was returned to a full response by the inclusion of 60% drug-free platelets (Figure 1Aii), and 80% drug-free platelets in the presence of an aspirin+PAM-inhibited platelet population (Figure 1Aiii). In each case, this relationship was nonlinear.

In contrast, for the P2Y<sub>12</sub> receptor ligand ADP, there was a linear increase in aggregation that followed the addition of drug-free platelets to platelets inhibited with either PAM or aspirin+PAM (Figure 1B and 1C). For instance, in PAM-inhibited platelets, addition of 20%, 40%, and 80% drug-free platelets increased aggregation from the control level of  $4\pm 3\%$  to  $13\pm 5\%$ ,  $29\pm 8\%$ , and  $53\pm 5\%$ , respectively. Conversely, the addition of drug-free platelets to PAM-treated platelets led to an inverse relationship with phosphorylation of vasodilator-stimulated phosphoprotein, a downstream measure of P2Y<sub>12</sub> inhibition, in the total platelet population (Figure IV in the online-only Data Supplement). Aspirin alone had little effect on aggregation induced by ADP and so responses of aspirin-inhibited platelets were minimally affected by the addition of drug-free platelets (Figure 1Ai). Similarly, aggregation in response to U46619 was linearly related to rising proportions of drug-free platelets mixed with PAM- or aspirin+PAM-inhibited platelets, but insensitive to aspirin-treatment (Figure II in the online-only Data Supplement).

Aspirin or PAM alone partially inhibited aggregations induced by the primary stimulus collagen or the glycoprotein (GP) VI agonist collagen-related peptide, which were returned to full responses by the addition of 40% or 60% to 80% drug-free platelets, respectively. The combination of aspirin+PAM required 80% drug-free platelets for the aggregation to collagen to return to a full response and 60% for collagen-related peptide (Figure II in the online-only Data Supplement).

Similar to the first series of experiments (Table I in the online-only Data Supplement), responses to ristocetin were unaffected by aspirin (Figure 1Ai), PAM (Figure 1Aii), or aspirin+PAM (Figure 1Aiii), consistent with the mechanism of action of ristocetin being platelet agglutination rather than platelet activation, and so independent of the generation of secondary mediators of aggregation.

In inhibited platelet populations, lumiaggregometry confirmed that the increases in platelet aggregations in response to AA or ADP following addition of increasing proportions of drug-free platelets correlated with matched increases in platelet release of ATP (Figure 1B). For example, aggregations



**Figure 1.** Light transmission aggregation, thromboxane A<sub>2</sub> (TXA<sub>2</sub>) formation, and ATP-release in inhibited platelet-rich plasma (PRP) containing rising proportions of drug-free platelets. **A**, Effects of adding drug-free PRP to PRP inhibited with aspirin (30 μmol/L; **Ai**), prasugrel active metabolite (PAM, 3 μmol/L; **Aii**), and aspirin+PAM (**Aiii**) on aggregatory responses to arachidonic acid (AA; 1 mmol/L), ADP (20 μmol/L), or ristocetin (2 mg/mL). Curves were constructed from multiple aggregatory responses using final aggregation values obtained by traditional light transmission aggregometry (LTA) after 5-minute aggregation. Percentage aggregation values are presented as mean±SEM of experiments using platelets prepared from 6 to 8 individuals. \*\*\**P*<0.001, \*\**P*<0.01, and \**P*<0.05 difference by paired ANOVA in aggregation from 0% drug-free platelets. **B**, Percentage aggregation and release of ATP, a measure of granule release, stimulated by AA or ADP in PRP pretreated with aspirin (30 μmol/L; **Bi**) or PAM (3 μmol/L), respectively, (**Bii**) mixed with increasing proportions of drug-free PRP. Curves were constructed from multiple aggregatory responses using aggregation values obtained by traditional LTA after 5 minutes and maximum ATP released measured by CHRONOLUME assay. Percentage aggregation values and ATP-release values are presented as mean±SEM of experiments using platelets prepared from 4 individuals for each comparative point. **C**, Concentrations of TXB<sub>2</sub>, a measure of TXA<sub>2</sub>, in unstimulated PRP or rising proportions of drug-free PRP mixed with aspirin-treated PRP on stimulation by AA (1 mmol/L). Values are presented as mean±SEM of experiments using platelets prepared from 4 individuals. \*\*\**P*<0.001, \*\**P*<0.01, and \**P*<0.05 difference by paired ANOVA in aggregation from 0% aspirin-free platelets.

induced by AA in the presence of 10%, 40%, and 80% aspirin-free platelets were 1±1%, 60±6%, and 74±3%, whereas the accompanying releases of ATP were 0, 3±0.1, and 4±1 nmol/10<sup>8</sup> platelets.

In order to mimic physiologically more relevant conditions, aggregation was also stimulated by the combination of subthreshold concentrations of the platelet agonists ADP, collagen, thrombin receptor activator peptide-6, and U46619. Aggregation in an inhibited platelet population was returned to a full response by the inclusion of 40% drug-free platelets in the presence of aspirin, 60% drug-free platelets in the presence of PAM, and 80% drug-free platelets in the presence of aspirin+PAM (Figure IID in the online-only Data Supplement).

### Aspirin-Free Platelets Intermingle With Aspirin-Inhibited Platelets, Whereas PAM-Free Platelets Form Clusters in the Center of Aggregates Covered in PAM-Inhibited Platelets

To elucidate the contribution of drug-free platelets to platelet aggregation, the morphologies of aggregates containing mixed platelet subpopulations were characterized by confocal imaging. Platelet aggregates comprising aspirin-inhibited platelets with a 20% subpopulation of drug-free platelets were characterized by intermingling of platelet populations (Figure 2A). Conversely, aggregates formed from combinations of 80% PAM-inhibited platelets and 20% drug-free platelets demonstrated clear clustering of drug-free platelets to the cores of


platelet aggregates on stimulation by ADP or AA. Aggregates formed by stimulation of the mixed platelet populations with ristocetin were always characterized by intermingled platelet populations (Figure 2B). Analysis of confocal images revealed 5× greater clustering of drug-free platelets in the presence of PAM after ADP stimulation than the clustering after stimulation by ristocetin.

In further studies, a full range of combinations of PAM-treated platelets or the clinically relevant combination of aspirin+PAM-treated platelets with drug-free platelets were examined. In these experiments, the central clustering of


drug-free platelets in aggregates increased in proportion to the fraction of drug-free platelets (Figure 3).

### GP IIb/IIIa Mediates Recruitment of PAM-Inhibited Platelets to the Aggregate Core Independently of TXA<sub>2</sub>

To assess the contribution of TXA<sub>2</sub> potentially formed by the drug-free core to the association of inhibited platelets to the platelet aggregates, mixed subpopulations containing drug-free and PAM-treated platelets were treated with aspirin (thereby inhibiting TXA<sub>2</sub> formation) or



**Figure 2.** Interaction between aspirin- or prasugrel active metabolite (PAM)-inhibited platelets and a drug-free subpopulation of platelets. Confocal images of platelet aggregates formed from combined platelet populations containing 80% aspirin-inhibited (red, **A**) or PAM-inhibited (red, **B**) platelets and 20% drug-free (green) platelets. Washed platelet aggregates were obtained at the end of 5-minute light transmission aggregometry (LTA) responses stimulated by arachidonic acid (AA; 250 μmol/L), ADP (20 μmol/L), or ristocetin (2 mg/mL). For experiments, platelet suspensions were pretreated with aspirin (30 μmol/L), PAM (3 μmol/L), or corresponding vehicle for 20 minutes, washed and labeled with either PKH67 (green) or PKH26 (red) before mixing and stimulation. Images were processed with Imaris software to show images of inhibited (red) and drug-free (green) platelets. **Bottom** row of both panel sets show 5-μm median focal sections. Scale bars indicate 20 μm. Each image is representative of images from platelets prepared from at least 4 different individuals.




**Figure 3.** Interaction between prasugrel active metabolite (PAM) or aspirin+PAM-inhibited platelets and a drug-free subpopulation of platelets. Three-dimensional (3D) reconstruction and median sections of confocal images of platelet aggregates formed from combined platelet populations containing 90% to 20% PAM-inhibited platelets (red, **left** panels) or aspirin+PAM-inhibited platelets (red, **right** panels) and 10% to 80% drug-free platelets (green). Washed platelet aggregates were obtained at the end of 5-minute light transmission aggregometry (LTA) responses stimulated by ADP (20  $\mu\text{mol/L}$ ). For experiments, platelet suspensions were pretreated with aspirin (30  $\mu\text{mol/L}$ ) plus PAM (3  $\mu\text{mol/L}$ ) or corresponding vehicle for 20 minutes, washed and labeled with either PKH67 (green) or PKH26 (red) before mixing and stimulation. Images were processed with Imaris software to show combined images of inhibited (red) and drug-free (green) platelets. Scale bars indicate 10  $\mu\text{m}$ .

corresponding vehicle before aggregation to ADP. Analysis of confocal images indicated no difference in the ratio of PAM-inhibited to drug-free platelet volume in formed aggregates in the absence ( $1.15 \pm 0.35$ ) or presence of aspirin ( $1.02 \pm 0.15$ ; Figure 4A). This was further supported by flow cytometry-based imaging analysis demonstrating no difference in measured area of PAM-inhibited platelets between

different treatments over a wide range of platelet proportions (Figure 4B).

Addition of the GP IIb/IIIa-inhibitor abciximab to PAM-inhibited platelets reduced their binding to the drug-free platelet aggregate core ( $0.48 \pm 0.11$  versus  $1.71 \pm 0.19$ ,  $P < 0.05$ ,  $n = 4$ ), indicating the association of PAM-inhibited platelets to the core was not an artifact of the imaging procedure (Figure 5A).




**Figure 4.** Aspirin has no effect on the association of inhibited platelets to prasugrel active metabolite (PAM)-free-platelet aggregates. **Ai**, Confocal images of platelet aggregates formed from combined platelet populations containing 80% PAM-inhibited or 80% aspirin+PAM-inhibited platelets (red) and 20% drug-free or aspirin-inhibited (green) platelets. Washed platelet aggregates were obtained at the end of 5-minute light transmission aggregometry (LTA) responses stimulated by ADP (20  $\mu\text{mol/L}$ ). For experiments, platelet suspensions were pretreated with PAM (3  $\mu\text{mol/L}$ ), PAM (3  $\mu\text{mol/L}$ ) plus aspirin (30  $\mu\text{mol/L}$ ), aspirin (30  $\mu\text{mol/L}$ ), or corresponding vehicle for 20 minutes, washed and labeled with either PKH67 (green) or PKH26 (red) before mixing and stimulation. Images were processed with Imaris software to show combined images of PAM-inhibited (red) and drug-free/aspirin-treated platelets (green). Scale bars indicate 10  $\mu\text{m}$ . Each image is representative of images from platelets prepared from at least 4 different individuals. **Aii**, Six confocal images per condition were analyzed for ratios between red and green platelet volume. Volumes of platelet subpopulations were calculated by Imaris software, and different treatments were compared by *t* test and found not to be significantly different. Scatter plots of combined platelet subpopulations consisting of 20% drug-free and 80% PAM-inhibited platelets in the absence (**Bi**) or presence of aspirin (**Bii**) post stimulation by ADP (20  $\mu\text{mol/L}$ ). **Biii**, Aggregates containing drug-free platelets were gated in region PAM-free positive and then analyzed by ImageStream. Ch1 shows the bright field image, Ch2 and Ch3 show the channel for PAM-free and PAM-inhibited platelets, respectively (scale bars indicate 14  $\mu\text{m}$ ). Washed platelet aggregates were obtained at the end of 5 minutes LTA responses stimulated by ADP (20  $\mu\text{mol/L}$ ). For experiments, platelet suspensions were pretreated with PAM (3  $\mu\text{mol/L}$ ), aspirin (30  $\mu\text{mol/L}$ ) plus PAM (3  $\mu\text{mol/L}$ ), aspirin (30  $\mu\text{mol/L}$ ), or corresponding vehicle for 20 minutes, washed and labeled with either PKH67 (green) or PKH26 (red) before mixing and stimulation. **Biv**, ImageStream data from 4 experiments were analyzed for the area of inhibited platelets associated with drug-free platelets over a range of platelet subpopulation proportions in the absence or presence of aspirin. Different treatments were compared by 2-way ANOVA and found not to be significantly different for all tested proportions.

The involvement of GP IIb/IIIa in the recruitment of PAM-inhibited platelets was confirmed by traditional flow cytometry and flow cytometric imaging analyses. Addition of abciximab to the inhibited platelet population reduced the measured platelet area of PAM-inhibited platelets associated with drug-free platelets (Figure 5B) and further caused a strong reduction in the incorporation of inhibited platelets into platelet aggregates as measured by a reduced number of aggregates containing both platelet subpopulations (Figure 5C).


To further elucidate the underlying mechanism causing the recruitment of PAM-treated platelets into the drug-free

platelet core, fibrinogen-binding studies were performed. The drug-free core was associated with the highest proportion of fibrinogen per volume of drug-free platelets ( $0.52 \pm 0.10$ ) followed by the PAM-treated shell ( $0.17 \pm 0.03$ ) and the control platelet sample ( $0.01 \pm 0.01$ ; Figure 6A). A similar pattern was found in the 60% drug-free proportion mixed with 40% PAM-treated platelets (fibrinogen/drug-free core,  $0.45 \pm 0.3$ ; fibrinogen/PAM-treated shell,  $0.1 \pm 0.06$ ; control,  $0.01 \pm 0.01$ ; data not shown). Flow cytometry analysis of fibrinogen binding to platelet subpopulations (Figure 6B) confirmed these findings (eg, 80%



**Figure 5.** Effect of abciximab pretreatment on the interaction of drug-free and P2Y<sub>12</sub>-inhibited platelet subpopulations. **Ai**, Confocal images of platelet aggregates formed from combined platelet populations containing 80% prasugrel active metabolite (PAM)- or PAM+abciximab-inhibited platelets (red) and 20% drug-free platelets (green). Washed platelet aggregates were obtained at the end of 5-minute light transmission aggregometry (LTA) responses stimulated by ADP (20 μmol/L). Images were processed with Imaris software. Scale bars indicate 10 μm. Each image is representative of images from platelets prepared from at least 4 different individuals. **Aii**, Six confocal images per experiment were analyzed for the ratios of red and green platelet volume. Volumes of platelet subpopulations were calculated by Imaris software and different treatments were compared by *t* test and determined as *P*=0.0013 between aggregates containing PAM-inhibited platelets mixed with drug-free platelets and aggregates containing PAM+abciximab-inhibited platelets and drug-free platelets. Data represent mean±SEM of 4 experiments. Scatter plots of combined platelet subpopulations consisting of 20% drug-free and 80% PAM-inhibited platelets (**Bi**) or 20% drug-free and 80% PAM+abciximab-inhibited platelets (**Bii**) post stimulation by ADP (20 μmol/L) obtained by ImageStream flow cytometry. For experiments, platelet suspensions were pretreated with PAM (3 μmol/L), PAM (3 μmol/L) plus abciximab (10 μmol/L), or corresponding vehicle for 20 minutes, washed and labeled with either PKH67 (green) or PKH26 (red) before mixing and stimulation. **Biii**, Flow cytometry-derived images of platelet aggregates formed from combined platelet populations containing 80% PAM-inhibited or PAM+abciximab-inhibited platelets (red) and 20% drug-free platelets (green; gated PAM-free positive in **Bi** or **Bii**). Ch1 shows the bright field images, Ch2 and Ch3 show the channels for drug-free and PAM- or PAM+abciximab-inhibited platelets, respectively (scale bars indicate 14 μm). For analyses, washed platelet aggregates were obtained at the end of 5-minute LTA responses stimulated by ADP (20 μmol/L). **Biv**, Images obtained during flow cytometry of 4 experiments were analyzed for inhibited platelets associated with drug-free platelets over a range of platelet combinations and treatments and compared by 2-way ANOVA. \*\*\**P*<0.001 and \**P*<0.05, *n*=4. **C**, Flow cytometry analysis was performed after 5-minute aggregation stimulated by ADP (20 μmol/L) in PAM- or PAM+abciximab-inhibited platelets in the presence of rising proportions of drug-free platelets. Events positive for PKH26 (inhibited platelets) and PKH67 (drug-free platelets) were identified as microaggregates. Data points represent mean±SEM of samples prepared from 5 individuals. \*\*\**P*<0.001 and \**P*<0.05 differences of microaggregates containing drug-free and PAM-treated platelets compared with microaggregates containing drug-free and PAM+abciximab-treated platelets compared with control responses by paired ANOVA; *n*=6, for each.





**Figure 6.** Fibrinogen is associated with both, the drug-free core and the P2Y<sub>12</sub>-inhibited shell. **Ai**, Confocal images of platelet aggregates formed from combined platelet populations containing 80% prasugrel active metabolite (PAM)-inhibited platelets (red) and 20% drug-free platelets (green) in the presence of AlexaFluor647-conjugated fibrinogen. Washed platelet aggregates were obtained at the end of 5-minute light transmission aggregometry (LTA) responses stimulated by ADP (20  $\mu$ mol/L). Images were processed with Imaris software. Scale bars indicate 10  $\mu$ m. Each image is representative of images from platelets prepared from at least 4 different individuals. **Aii**, Four confocal images per experiment were analyzed for the concentration of labeled fibrinogen associated with either red or green platelet volume. Volumes of platelet subpopulations were calculated with Imaris software, and fibrinogen concentrations per platelet volume were compared by *t* test and determined as  $P=0.0012$  or  $P=0.002$  between fibrinogen associated with unstimulated and ADP-stimulated PAM-treated platelets or drug-free platelets, respectively. **Bi**, Representative flow cytometric scatter plots of drug-free or PAM-treated platelet subpopulations against AlexaFluor647-conjugated fibrinogen fluorescence (x axis) following stimulation with saline or ADP (20  $\mu$ mol/L). Gated events were considered positive for fibrinogen binding and occurrence (%) calculated. **Bii**, Fibrinogen bound platelets (%) of drug-free or PAM-treated platelet populations across all tested proportions. Comparison by 1-way ANOVA found significantly higher fibrinogen binding in PAM-treated proportions compared with vehicle (saline) stimulated control. \*\*\* $P<0.001$ , \*\* $P<0.01$ , and \* $P<0.05$ . Data represent mean $\pm$ SEM of 4 to 5 experiments for all.

PAM-treated platelets: vehicle stimulated,  $2.4\pm 0.4\%$ ; ADP-stimulated,  $6.2\pm 0.6\%$ ;  $P<0.01$ ), indicating that GP IIb/IIIa, but not TXA<sub>2</sub>, is involved in the recruitment of PAM-inhibited platelets to the drug-free platelet core.


## Discussion

The confocal analyses we present here, together with quantitative data from LTA, indicate that overall aggregation responses in mixed populations of inhibited and drug-free platelets are


underpinned by distinct patterns of interaction that differ between inhibition of platelet COX by aspirin and blockade of platelet P2Y<sub>12</sub> receptors by thienopyridines. These findings define processes underlying in vitro platelet tests used to assess P2Y<sub>12</sub> receptor blocker and aspirin effectiveness<sup>5-11</sup> and provide insight to the potential interactions of platelet populations in vivo.

Others have previously reported that in tests of platelet reactivity conducted in vitro a relatively small population of

Aspirin-inhibited platelets containing a drug-free subpopulation:



Aspirin+PAM-inhibited platelets containing a drug-free subpopulation:



**Figure 7.** Schematic diagram of the platelet activation mechanisms resulting from presence of drug-free platelets in populations of aspirin-inhibited and dual antiplatelet-inhibited platelets. In aspirin-inhibited platelet populations mixed with drug-free platelets (**upper**), activation of cyclooxygenase-1 in drug-free platelets leads to thromboxane A<sub>2</sub> (TXA<sub>2</sub>) formation and release. Formed TXA<sub>2</sub> acts equally on both aspirin-inhibited and drug-free platelets, leading to activation and aggregate formation characterized by random intermingling of platelet subpopulations. In aspirin+PAM-inhibited platelets (modeling DAPT with an irreversible P2Y<sub>12</sub> receptor blocker) combined with drug-free platelets, exposure to ADP causes activation of the drug-free subpopulation and its clustering. This leads to formation of a distinct uninhibited core that promotes activation of the inhibited platelets. In this way, through different mechanisms, drug-free platelets act as the seed for platelet aggregate formation in the presence of both aspirin and DAPT. AA indicates arachidonic acid.

aspirin naïve platelets can support full platelet aggregation, subject to the stimulus being applied. For instance, di Minno et al<sup>15</sup> demonstrated that in LTA with platelet-rich plasma ≈10% drug-free platelets could support a full aggregation response to collagen (1 μg/mL) plus AA (1 mmol/L). More recent studies have associated reduced effectiveness of aspirin in vivo to increased platelet turnover as defined by the proportion of reticulated platelets in the circulation.<sup>33</sup> Similar analyses have indicated that increased proportions of reticulated platelets are associated with reduced effectiveness of clopidogrel in both rats<sup>34</sup> and humans,<sup>35</sup> in humans receiving DAPT of aspirin plus clopidogrel,<sup>36,37</sup> and most recently in humans receiving DAPT of aspirin plus prasugrel.<sup>38</sup> Studies of the duration of drug action after treatment withdrawal also indicate that the return of aggregatory responses is commensurate with the time for replenishment of circulating platelets.<sup>39</sup> In the first part of our studies, we modeled these effects in LTA using a panel of platelet agonists<sup>40</sup> and rising proportions of drug-free platelets against a background of standard antiplatelet therapy, aspirin and aspirin plus prasugrel (by the use of PAM).

We also studied PAM alone and provided aggregation curves from serially diluted platelet-rich plasma to assist in interpretation of data. Considering first the agonists most sensitive for testing the effects of aspirin, that is, AA and PAM, that is, ADP, we noted that whereas addition of 30% of drug-free platelets could return full responses to AA in the presence of aspirin, 80% of drug-free platelets were required to return full responses to ADP in the presence of PAM or aspirin+PAM. Aspirin and P2Y<sub>12</sub> receptor blockers have different targets on the platelet and inhibit with different functional modalities. Aspirin inhibits platelet COX-1, and so the ability of platelets to produce TXA<sub>2</sub>. This does not stop aspirin-inhibited platelets responding to TXA<sub>2</sub> produced by aspirin-free platelets, and so the hypothesis has developed that a minority of aspirin-free platelets are capable of generating sufficient TXA<sub>2</sub> to support a full aggregatory response.<sup>4,17,41</sup> While the in vivo proportion of aspirin uninhibited platelets required to provide a full response has been suggested to be as low as 5% or less,<sup>41</sup> the response of a population of platelets to a particular concentration of TXA<sub>2</sub> will depend on their existing state of activation

so there is unlikely to be a particular proportion relevant in all conditions; although clearly even a small subset of platelets can produce enough TXA<sub>2</sub> to aggregate a larger population.

PAM, by blocking P2Y<sub>12</sub> receptors, inhibits the ability of a platelet to respond to ADP released by other activated platelets. Unlike the effects of aspirin, the blockade of one platelet's P2Y<sub>12</sub> receptors cannot be compensated for by lack of blockade on another platelet; hence, we found a linear relationship between drug-free platelets and the response to ADP. For the other platelet agonists, we noted composite responses as expected from their relative dependencies on the TXA<sub>2</sub> and P2Y<sub>12</sub> receptor pathways. For example, collagen and collagen-related peptide activate both pathways; U46619 is sensitive to blockade of P2Y<sub>12</sub> receptors, because TXA<sub>2</sub> receptor activation leads to ADP release, but not to the effects of aspirin. Consistent with this, AA stimulates the generation of TXA<sub>2</sub> and so is sensitive both to aspirin, by inhibition of COX, and to PAM, because of inhibition of TXA<sub>2</sub> receptor amplification.<sup>42-44</sup> Overall, in the presence of aspirin+PAM, which models the effects of DAPT using aspirin and a thienopyridine, all agonists demonstrated increasing responses with increasing proportions of drug-free platelets. The exception was ristocetin, which we included as a control agent, because it causes platelet agglutination that does not require active platelet involvement.<sup>45</sup>

Our confocal and flow cytometric imaging studies permitted a unique and deeper exploration of the interactions between inhibited and drug-free platelet populations. Notably, drug-free platelets were distributed throughout aggregates of aspirin-inhibited platelets, whereas drug-free platelets formed clear cores within PAM- or aspirin+PAM-inhibited platelet aggregates. We conclude that the intermingled pattern of drug-free and aspirin-inhibited platelets is consistent with the drug-free subset driving the aggregatory response by producing TXA<sub>2</sub> throughout the platelet mixture. In this study, we have not examined the roles of COX-1 and COX-2 isoforms because we have used mature platelet populations, but in vivo newly formed platelets may express COX-2, which could provide a further level of complexity in the responsiveness to aspirin as could non-COX effects of aspirin.<sup>9,31,46</sup> Conversely, platelets with blocked P2Y<sub>12</sub> receptors cannot equally respond to the effects of ADP, do not become activated by ADP released as part of platelet amplification pathways, so are not part of platelet aggregate cores. It is interesting to contemplate that platelets with their P2Y<sub>12</sub> receptors blocked are entirely mixed with drug-free platelets in our system, yet after addition of a platelet activator the drug-free platelets bind to each other so assembling an activated core from the larger mixed platelet population.

Our imaging studies primarily concentrated on the effects of a population of 20% drug-free platelets. This ratio is what could be expected under a once-daily drug regimen in conditions, such as diabetes mellitus, in which circulating platelet lifetimes are ≈5 days.<sup>13,23,26</sup> For completeness, we also characterized a range of combinations between 10% drug-free and 80% drug-free platelets. These studies demonstrated in PAM-inhibited or aspirin+PAM-inhibited platelets that the size of the drug-free platelet aggregate core grew with the proportion of drug-free platelets, whereas inhibited platelets were

only ever associated with the outer portion. Furthermore, we found that addition of aspirin did not reduce the association of inhibited platelets, indicating that this was not dependent on the formation of TXA<sub>2</sub>. However, the recruitment of P2Y<sub>12</sub>-inhibited platelets to the forming aggregate was the result of active aggregation because these inhibited platelets bound fibrinogen more than unstimulated control platelets, and their association to the drug-free core was reduced by the GP IIb/IIIa blocker abciximab.

In summary, our studies shed new light on the responses recorded in many studies of ex vivo platelet reactivity and their association to antiplatelet drug therapy. Although it has previously been hypothesized that subpopulations of aspirin-free platelets act as individual generators of TXA<sub>2</sub>, we demonstrate for the first time a mechanism by which this is achieved; aspirin-free platelets are distributed throughout aspirin-inhibited platelet aggregates allowing generated TXA<sub>2</sub> to activate a larger proportion of platelets (Figure 7). In contrast, platelets free of inhibition by irreversible P2Y<sub>12</sub> receptor blockers, such as clopidogrel and prasugrel, form the core of platelet aggregates and act as the nexus for the formation of larger aggregates because P2Y<sub>12</sub> receptor inhibited platelets are drawn in via other activation pathways. For DAPT, this leads to a complicated interaction between the ability of a minority of aspirin uninhibited platelets to drive a full TXA<sub>2</sub>-dependent response and a linear relationship between P2Y<sub>12</sub> receptor blockade and platelet aggregation. Aggregatory responses, measured by LTA, have been associated with thrombotic risk and these have been linked to particular patient groups in which increased platelet turnover occurs.<sup>4,12-15</sup> Our in vitro studies clearly indicate that such an increased ratio of drug-free platelets may be potentially critical to modulating thrombotic risk in conditions that are associated with increased platelet turnover, such as diabetes mellitus, chronic kidney disease, metabolic syndrome, and essential thrombocythemia. This is consistent with twice a day administration of aspirin providing an improved antiplatelet effect compared with standard once a day therapy in such patient groups.<sup>13,26,27,32,47</sup> In essence, our studies indicate that in a patient receiving antiplatelet therapy in the form of either aspirin or DAPT consisting of aspirin plus a thienopyridine, a drug-free subpopulation can exist that will respond differently to platelet activators. This differential interaction and its potential to drive thrombosis are important both to consideration of individualized therapies and to the development of antiplatelet strategies.

### Acknowledgments

We are grateful to Professor Sussan Nourshargh for use of confocal microscopes.

### Sources of Funding

This work was supported by grants from the Medical Research Council, the British Heart Foundation (PG-12-68-29779), the Wellcome Trust (101604/Z/13/Z), and the William Harvey Research Foundation. T.D. Warner has received research grant funding and consultancy fees from Astra Zeneca.

## Disclosures

None.

## References

- Jneid H, Anderson JL, Wright RS, Adams CD, Bridges CR, Casey DE, Jr, Ettinger SM, Fesmire FM, Ganiats TG, Lincoff AM, Peterson ED, Philippides GJ, Theroux P, Wenger NK, Zidar JP. 2012 ACCF/AHA focused update of the guideline for the management of patients with unstable angina/non-ST-elevation myocardial infarction (updating the 2007 guideline and replacing the 2011 focused update): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2012;60:645–681. doi: 10.1016/j.jacc.2012.06.004.
- Baigent C, Blackwell L, Collins R, Emberson J, Godwin J, Peto R, Buring J, Hennekens C, Kearney P, Meade T, Patrono C, Roncaglioni MC, Zanchetti A. Aspirin in the primary and secondary prevention of vascular disease: collaborative meta-analysis of individual participant data from randomised trials. *Lancet*. 2009;373:1849–1860
- Antiplatelet Trialists' Collaboration. Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. *Bmj*. 2002;324:71–86
- Patrono C. Low-dose aspirin in primary prevention: cardioprotection, chemoprevention, both, or neither? *Eur Heart J*. 2013;34:3403–3411. doi: 10.1093/eurheartj/ehs058.
- Harrison P, Frelinger AL 3rd, Furman MI, Michelson AD. Measuring antiplatelet drug effects in the laboratory. *Thromb Res*. 2007;120:323–336. doi: 10.1016/j.thromres.2006.11.012.
- Bonello L, Tantry US, Marcucci R, et al; Working Group on High On-Treatment Platelet Reactivity. Consensus and future directions on the definition of high on-treatment platelet reactivity to adenosine diphosphate. *J Am Coll Cardiol*. 2010;56:919–933. doi: 10.1016/j.jacc.2010.04.047.
- Gurbel PA, Tantry US. Clopidogrel response variability and the advent of personalised antiplatelet therapy. A bench to bedside journey. *Thromb Haemost*. 2011;106:265–271. doi: 10.1160/TH11-03-0167.
- Pinto Slotow TL, Bonello L, Gavini R, et al. Prevalence of aspirin and clopidogrel resistance among patients with and without drug-eluting stent thrombosis. *Am J Cardiol*. 2009;104:525–530. doi: 10.1016/j.amjcard.2009.04.015.
- Undas A, Brummel-Ziedins KE, Mann KG. Antithrombotic properties of aspirin and resistance to aspirin: beyond strictly antiplatelet actions. *Blood*. 2007;109:2285–2292. doi: 10.1182/blood-2006-01-010645.
- Tantry US, Bonello L, Aradi D, et al; Working Group on On-Treatment Platelet Reactivity. Consensus and update on the definition of on-treatment platelet reactivity to adenosine diphosphate associated with ischemia and bleeding. *J Am Coll Cardiol*. 2013;62:2261–2273. doi: 10.1016/j.jacc.2013.07.101.
- Patrono C, Rocca B. Drug insight: aspirin resistance—fact or fashion? *Nat Clin Pract Cardiovasc Med*. 2007;4:42–50. doi: 10.1038/ncpcardio0728.
- Ferreiro JL, Angiolillo DJ. Diabetes and antiplatelet therapy in acute coronary syndrome. *Circulation*. 2011;123:798–813. doi: 10.1161/CIRCULATIONAHA.109.913376.
- Rocca B, Santilli F, Pitocco D, et al. The recovery of platelet cyclooxygenase activity explains interindividual variability in responsiveness to low-dose aspirin in patients with and without diabetes. *J Thromb Haemost*. 2012;10:1220–1230. doi: 10.1111/j.1538-7836.2012.04723.x.
- Rocca B, Dragani A, Pagliaccia F. Identifying determinants of variability to tailor aspirin therapy. *Expert Rev Cardiovasc Ther*. 2013;11:365–379. doi: 10.1586/erc.12.144.
- Di Minno G, Silver MJ, Murphy S. Monitoring the entry of new platelets into the circulation after ingestion of aspirin. *Blood*. 1983;61:1081–1085.
- Casado-Arroyo R, Sostres C, Lanas A. Optimizing the use of aspirin for cardiovascular prevention. *Drugs*. 2013;73:803–814. doi: 10.1007/s40265-013-0061-z.
- Davì G, Patrono C. Platelet activation and atherothrombosis. *N Engl J Med*. 2007;357:2482–2494. doi: 10.1056/NEJMra071014.
- Cattaneo M. New P2Y<sub>12</sub> blockers. *J Thromb Haemost*. 2009;7 Suppl 1:262–265. doi: 10.1111/j.1538-7836.2009.03382.x.
- Cattaneo M. The platelet P2Y<sub>12</sub> receptor for adenosine diphosphate: congenital and drug-induced defects. *Blood*. 2011;117:2102–2112. doi: 10.1182/blood-2010-08-263111.
- Floyd CN, Passacuale G, Ferro A. Comparative pharmacokinetics and pharmacodynamics of platelet adenosine diphosphate receptor antagonists and their clinical implications. *Clin Pharmacokinet*. 2012;51:429–442. doi: 10.2165/11630740-000000000-00000.
- Zimmermann N, Wenk A, Kim U, Kienzle P, Weber AA, Gams E, Schrör K, Hohlfeld T. Functional and biochemical evaluation of platelet aspirin resistance after coronary artery bypass surgery. *Circulation*. 2003;108:542–547. doi: 10.1161/01.CIR.0000081770.51929.5A.
- DiChiara J, Bliden KP, Tantry US, Hamed MS, Antonino MJ, Suarez TA, Bailon O, Singla A, Gurbel PA. The effect of aspirin dosing on platelet function in diabetic and nondiabetic patients: an analysis from the aspirin-induced platelet effect (ASPECT) study. *Diabetes*. 2007;56:3014–3019. doi: 10.2337/db07-0707.
- Patrono C. Platelet survival in diabetes mellitus using an aspirin-labelling technique. *Thromb Res*. 1979;15:793–802.
- Dassin E, Najean Y, Poirier O, Passa P, Bensoussan D. *In vivo* platelet kinetics in 31 diabetic patients. Correlation with the degree of vascular impairment. *Thromb Haemost*. 1978;40:83–88.
- Ferguson JC, Mackay N, Philip JA, Sumner DJ. Platelet and fibrinogen kinetic studies in diabetes mellitus. *Br J Haematol*. 1973;25:545.
- Spectre G, Arnetz L, Östenson CG, Brismar K, Li N, Hjemdahl P. Twice daily dosing of aspirin improves platelet inhibition in whole blood in patients with type 2 diabetes mellitus and micro- or macrovascular complications. *Thromb Haemost*. 2011;106:491–499. doi: 10.1160/TH11-04-0216.
- Capodanno D, Patel A, Dharmashankar K, Ferreiro JL, Ueno M, Kodali M, Tomasello SD, Capranzano P, Seecheran N, Darlington A, Tello-Montoliu A, Desai B, Bass TA, Angiolillo DJ. Pharmacodynamic effects of different aspirin dosing regimens in type 2 diabetes mellitus patients with coronary artery disease. *Circ Cardiovasc Interv*. 2011;4:180–187. doi: 10.1161/CIRCINTERVENTIONS.110.960187.
- Abrahamsen AF. Platelet survival studies in man. With special reference to thrombosis and atherosclerosis. *Scand J Haematol Suppl*. 1968;3:1–53.
- Htun P, Fateh-Moghadam S, Bischofs C, Banya N, Müller K, Bigalke B, Stellos K, May AE, Flather M, Gawaz M, Geisler T. Low responsiveness to clopidogrel increases risk among CKD patients undergoing coronary intervention. *J Am Soc Nephrol*. 2011;22:627–633. doi: 10.1681/ASN.2010020220.
- Smith JP, Haddad EV, Taylor MB, Oram D, Blakemore D, Chen Q, Boutaud O, Oates JA. Suboptimal inhibition of platelet cyclooxygenase-1 by aspirin in metabolic syndrome. *Hypertension*. 2012;59:719–725. doi: 10.1161/HYPERTENSIONAHA.111.181404.
- Dragani A, Pascale S, Recchiti A, Mattosco D, Lattanzio S, Petrucci G, Mucci L, Ferrante E, Habib A, Ranelletti FO, Ciabattoni G, Davì G, Patrono C, Rocca B. The contribution of cyclooxygenase-1 and -2 to persistent thromboxane biosynthesis in aspirin-treated essential thrombocythemia: implications for antiplatelet therapy. *Blood*. 2010;115:1054–1061. doi: 10.1182/blood-2009-08-236679.
- Pascale S, Petrucci G, Dragani A, Habib A, Zaccardi F, Pagliaccia F, Pocaterra D, Ragazzoni E, Rolandi G, Rocca B, Patrono C. Aspirin-insensitive thromboxane biosynthesis in essential thrombocythemia is explained by accelerated renewal of the drug target. *Blood*. 2012;119:3595–3603. doi: 10.1182/blood-2011-06-359224.
- Guthikonda S, Lev EI, Patel R, DeLao T, Bergeron AL, Dong JF, Kleiman NS. Reticulated platelets and uninhibited COX-1 and COX-2 decrease the antiplatelet effects of aspirin. *J Thromb Haemost*. 2007;5:490–496. doi: 10.1111/j.1538-7836.2007.02387.x.
- Kuijpers MJ, Megens RT, Nikoohesal E, Feijge MA, De Mey JG, oude Egbrink MG, van Giezen JJ, Heemskerk JW. Role of newly formed platelets in thrombus formation in rat after clopidogrel treatment: comparison to the reversible binding P2Y<sub>12</sub> antagonist ticagrelor. *Thromb Haemost*. 2011;106:1179–1188. doi: 10.1160/TH11-04-0252.
- Ibrahim H, Nadipalli S, DeLao T, Guthikonda S, Kleiman NS. Immature platelet fraction (IPF) determined with an automated method predicts clopidogrel hyporesponsiveness. *J Thromb Thrombolysis*. 2012;33:137–142. doi: 10.1007/s11239-011-0665-7.
- Guthikonda S, Alviar CL, Vaduganathan M, Arikian M, Tellez A, DeLao T, Granada JF, Dong JF, Kleiman NS, Lev EI. Role of reticulated platelets and platelet size heterogeneity on platelet activity after dual antiplatelet therapy with aspirin and clopidogrel in patients with stable coronary artery disease. *J Am Coll Cardiol*. 2008;52:743–749. doi: 10.1016/j.jacc.2008.05.031.
- Cesari F, Marcucci R, Caporale R, Panizza R, Romano E, Gensini GF, Abbate R, Gori AM. Relationship between high platelet turnover and platelet function in high-risk patients with coronary artery disease on dual

- antiplatelet therapy. *Thromb Haemost.* 2008;99:930–935. doi: 10.1160/TH08-01-0002.
38. Perl L, Lerman-Shivek H, Rechavia E, Vaduganathan M, Leshem-Lev D, Zemer-Wassercug N, Dadush O, Codner P, Bental T, Battler A, Kornowski R, Lev EI. Response to prasugrel and levels of circulating reticulated platelets in patients with ST-segment elevation myocardial infarction. *J Am Coll Cardiol.* 2014;63:513–517. doi: 10.1016/j.jacc.2013.07.110.
  39. Gurbel PA, Bliden KP, Butler K, et al. Randomized double-blind assessment of the ONSET and OFFSET of the antiplatelet effects of ticagrelor versus clopidogrel in patients with stable coronary artery disease: the ONSET/OFFSET study. *Circulation.* 2009;120:2577–2585. doi: 10.1161/CIRCULATIONAHA.109.912550.
  40. Cattaneo M, Cerletti C, Harrison P, Hayward CP, Kenny D, Nugent D, Nurden P, Rao AK, Schmaier AH, Watson SP, Lussana F, Pugliano MT, Michelson AD. Recommendations for the standardization of light transmission aggregometry: A consensus of the working party from the platelet physiology subcommittee of ssc/isth. *J Thromb Haemost.* 2013. doi: 10.1111/jth.12231.
  41. Reilly IA, FitzGerald GA. Inhibition of thromboxane formation *in vivo* and *ex vivo*: implications for therapy with platelet inhibitory drugs. *Blood.* 1987;69:180–186.
  42. Armstrong PC, Dhanji AR, Truss NJ, Zain ZN, Tucker AT, Mitchell JA, Warner TD. Utility of 96-well plate aggregometry and measurement of thrombi adhesion to determine aspirin and clopidogrel effectiveness. *Thromb Haemost.* 2009;102:772–778. doi: 10.1160/TH09-04-0215.
  43. Armstrong PC, Dhanji AR, Tucker AT, Mitchell JA, Warner TD. Reduction of platelet thromboxane A2 production *ex vivo* and *in vivo* by clopidogrel therapy. *J Thromb Haemost.* 2010;8:613–615. doi: 10.1111/j.1538-7836.2009.03714.x.
  44. Armstrong PC, Leadbeater PD, Chan MV, Kirkby NS, Jakubowski JA, Mitchell JA, Warner TD. In the presence of strong P2Y12 receptor blockade, aspirin provides little additional inhibition of platelet aggregation. *J Thromb Haemost.* 2011;9:552–561. doi: 10.1111/j.1538-7836.2010.04160.x.
  45. Scott JP, Montgomery RR, Retzinger GS. Dimeric ristocetin flocculates proteins, binds to platelets, and mediates von Willebrand factor-dependent agglutination of platelets. *J Biol Chem.* 1991;266:8149–8155.
  46. Rocca B, Secchiero P, Ciabattoni G, Ranalletti FO, Catani L, Guidotti L, Melloni E, Maggiano N, Zauli G, Patrono C. Cyclooxygenase-2 expression is induced during human megakaryopoiesis and characterizes newly formed platelets. *Proc Natl Acad Sci U S A.* 2002;99:7634–7639. doi: 10.1073/pnas.112202999.
  47. Grove EL, Hvas AM, Mortensen SB, Larsen SB, Kristensen SD. Effect of platelet turnover on whole blood platelet aggregation in patients with coronary artery disease. *J Thromb Haemost.* 2011;9:185–191. doi: 10.1111/j.1538-7836.2010.04115.x.



### Significance

Many individuals at risk of heart attacks and strokes receive prophylactic antiplatelet therapy, aspirin often with the addition of a thienopyridine P2Y<sub>12</sub> receptor blocker. Because of the short half lives of these drugs within the body and the continual release into the circulation of newly formed platelets, it has been hypothesized that significant populations of uninhibited platelets can emerge during standard once a day dosing. Here by modeling these events *in vitro* and imaging the interactions of different platelet populations, we have for the first time provided analysis of the potential underlying mechanisms and demonstrated that differently inhibited platelet subpopulations respond very differently to platelet activators. Understanding these mechanisms is central to modulating thrombotic risk in conditions that are associated with increased platelet turnover, such as diabetes mellitus, chronic kidney disease, metabolic syndrome, and essential thrombocythemia.


# Arteriosclerosis, Thrombosis, and Vascular Biology

## SUPPLEMENTAL MATERIAL


### TABLES

aggregatory stimulus	% of control aggregation (vehicle-treated)		
	aspirin	PAM	aspirin + PAM
AA	2±2*	27±12*	4±2*
ADP	76±6	7±4*	5±2*
collagen	22±5*	56±15*	9±2*
CRP-XL	53±10*	47±10*	37±10*
ristocetin	85±8	95±7	115±9
U46619	113±5	25±11*	24±7*

**Table S1. *In vitro* effects of aspirin, PAM and aspirin+PAM on aggregatory responses of platelets.** Percentage values of control vs aspirin (30 µmol/L), PAM (3 µmol/L) and aspirin+PAM aggregation in response to stimulation by AA (1 mmol/L), ADP (20 µmol/L), collagen (1 µg/ml), CRP-XL (0.1 µg/ml), ristocetin (2 mg/ml) and U46619 (1 µmol/L). are presented as mean±SEM of experiments using platelets prepared from 6 individuals; \* p<0.05 compared to control responses.




**Figure S1: Schematic diagram summarising the experimental setup used for imaging platelet aggregates.** 1 - platelet suspensions obtained from PRP; 2 – platelets labelled and treated with anti-platelet drugs; 3 – differently treated and labelled platelets combined in rising proportions; 4 – aggregation stimulated by a range of agonists; 5 – formed aggregates transferred to a microscope slide; 6 – aggregates imaged by confocal microscopy; 7 – images processed and analysed by Imaris software.




**Figure SII: Light transmission aggregation in inhibited PRP containing rising proportions of drug-free platelets.** Aggregatory responses to collagen (1  $\mu\text{g/ml}$ ), CRP-XL (0.1  $\mu\text{g/ml}$ ), U46619 (1  $\mu\text{mol/L}$ ) or ADP (5  $\mu\text{mol/L}$ ) after adding drug-free PRP to PRP inhibited with (A) aspirin (30  $\mu\text{mol/L}$ ), (B) PAM (3  $\mu\text{mol/L}$ ) and (C) aspirin+PAM. (D) The aggregatory responses to the combination of ADP (0.1  $\mu\text{mol/L}$ ), collagen (0.1  $\mu\text{g/ml}$ ), TRAP-6 (0.1  $\mu\text{mol/L}$ ) and U46619 (0.1  $\mu\text{mol/L}$ ) in drug-free PRP with (Di) aspirin, (Dii) PAM or (Diii) aspirin+PAM-inhibited PRP. Curves were constructed from multiple aggregatory responses using final aggregation values obtained by traditional LTA after five minutes aggregation. Percentage aggregation values are presented as mean $\pm$ SEM of experiments using platelets prepared from 5-8 individuals. \*\*\*  $p < 0.001$ , \*\*  $p < 0.01$  and \*  $p < 0.05$  difference by paired ANOVA in aggregation from 100% drug-free platelets.





**Figure SIII: Light transmission aggregation in PRP containing variable platelet concentrations.** Effects of variable platelet concentrations, obtained by centrifuging whole blood at different speeds, on aggregatory responses to A) AA (1 mmol/L), or B) ADP (20 μmol/L). Data is presented as percentage of platelets obtained after centrifugation at 175 x g. Curves were constructed from multiple aggregatory responses using final aggregation values obtained by traditional LTA after five minutes aggregation. Percentage aggregation values and % platelets are presented as mean±SEM of experiments using platelets prepared from 3 individuals.



**Figure SIV: Phosphorylation of vasodilator stimulated phosphoprotein (VASP) in PRP containing rising proportions of drug-free platelets.** (A) Geometric mean fluorescence intensity of phospho(Ser<sup>239</sup>)-VASP, measured by flow cytometry after adding drug-free PRP to PRP inhibited with (PAM (3  $\mu\text{mol/L}$ ) and stimulated with ADP (20  $\mu\text{mol/L}$ ). B) Representative phospho(Ser<sup>239</sup>)-VASP fluorescence histograms of each PRP-proportion stimulated with ADP (20  $\mu\text{mol/L}$ ). Data are presented as mean $\pm$ SEM of experiments using platelets prepared from 4 individuals.

# Arteriosclerosis, Thrombosis, and Vascular Biology



JOURNAL OF THE AMERICAN HEART ASSOCIATION

## Drug-Free Platelets Can Act as Seeds for Aggregate Formation During Antiplatelet Therapy

Thomas Hoefler, Paul C. Armstrong, Michaela Finsterbusch, Melissa V. Chan, Nicholas S. Kirkby and Timothy D. Warner

*Arterioscler Thromb Vasc Biol.* published online August 13, 2015;  
*Arteriosclerosis, Thrombosis, and Vascular Biology* is published by the American Heart Association, 7272  
Greenville Avenue, Dallas, TX 75231

Copyright © 2015 American Heart Association, Inc. All rights reserved.  
Print ISSN: 1079-5642. Online ISSN: 1524-4636

The online version of this article, along with updated information and services, is located on the  
World Wide Web at:

<http://atvb.ahajournals.org/content/early/2015/08/13/ATVBAHA.115.306219>

Free via Open Access

Data Supplement (unedited) at:

<http://atvb.ahajournals.org/content/suppl/2015/08/13/ATVBAHA.115.306219.DC1.html>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Arteriosclerosis, Thrombosis, and Vascular Biology* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:

<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Arteriosclerosis, Thrombosis, and Vascular Biology* is online at:

<http://atvb.ahajournals.org/subscriptions/>