



## **Invasive crayfish as drivers of fine sediment dynamics in rivers: Field and laboratory evidence**

Harvey, GL; Henshaw, AJ; Moorhouse, TP; Clifford, NJ; Holah, H; Grey, J; Macdonald, DW

“The final publication is available at <http://onlinelibrary.wiley.com/doi/10.1002/esp.3486/full>”

For additional information about this publication click this link.

<http://qmro.qmul.ac.uk/xmlui/handle/123456789/11788>

Information about this research object was correct at the time of download; we occasionally make corrections to records, please therefore check the published record when citing. For more information contact [scholarlycommunications@qmul.ac.uk](mailto:scholarlycommunications@qmul.ac.uk)

1 This is the final pre-publication version of:  
2 Harvey, G. L., Henshaw, A.J., Moorhouse, T.P., Clifford, N. J., Holah, H., Grey, J.  
3 and Macdonald, D. W. (2014) Invasive crayfish as drivers of fine sediment dynamics  
4 in rivers: field and laboratory evidence. *Earth Surface Processes and Landforms* 39:  
5 259-271.  
6

7 **Invasive crayfish as drivers of fine sediment dynamics in rivers:**  
8 **field and laboratory evidence**

9 Harvey, G. L.\*, Henshaw, A.J., Moorhouse, T.P., Clifford, N. J., Holah, H., Grey, J.  
10 and Macdonald, D. W.

11

12 \*corresponding author (g.l.harvey@qmul.ac.uk)

13

14 **ABSTRACT**

15 Despite increasing recognition of the potential of aquatic biota to act as 'geomorphic  
16 agents', key knowledge gaps exist in relation to biotic drivers of fine sediment  
17 dynamics at microscales and particularly the role of invasive species. This paper  
18 explores the impacts of invasive signal crayfish on suspended sediment dynamics at  
19 the patch scale through laboratory and field study. Three hypotheses are presented  
20 and tested: (1) that signal crayfish generate pulses of fine sediment mobilisation  
21 through burrowing and movement that are detectable in the flow field; (2) that such  
22 pulses may be more frequent during nocturnal periods when signal crayfish are  
23 known to be most active; and (3) that cumulatively the pulses would be sufficient to  
24 drive an overall increase in turbidity. Laboratory mesocosm experiments were used  
25 to explore crayfish impacts on suspended sediment concentrations for two  
26 treatments: clay banks and clay bed substrate. For the field study, high frequency  
27 near-bed and mid-flow turbidity time series from a lowland river with known high  
28 densities of signal crayfish were examined. Laboratory data demonstrate the direct

1 influence of signal crayfish on mobilisation of pulses of fine sediment through  
2 burrowing into banks and fine bed material, with evidence of enhanced activity levels  
3 around the mid-point of the nocturnal period. Similar patterns of pulsed fine  
4 sediment mobilisation identified under field conditions follow a clear nocturnal trend  
5 and appear capable of driving an increase in ambient turbidity levels. The findings  
6 indicate that signal crayfish have the potential to influence suspended sediment  
7 yields, with implications for morphological change, physical habitat quality and the  
8 transfer of nutrients and contaminants. This is particularly important given the spread  
9 of signal crayfish across Europe and their presence in extremely high densities in  
10 many catchments. Further process-based studies are required to develop a full  
11 understanding of impacts across a range of river styles.

12 **KEY WORDS:** invasive species; *Pacifastacus leniusculus*; suspended sediment  
13 dynamics; turbidity; river management  
14

## 15 **INTRODUCTION**

16 Fine sediment, comprising sand, silt and clay, accounts for the majority of total fluvial  
17 sediment flux, and has important implications for catchment water resource  
18 management, particularly since changes in fine sediment dynamics can modify  
19 sediment yields and initiate morphological adjustments downstream (Walling and  
20 Collins, 2008). Such changes may have implications for channel morphology and  
21 conveyance capacity, aquatic ecosystem health and chemical water quality. For  
22 instance, aggradation of fine sediments can degrade aquatic habitats, reduce  
23 survival rates of fish and aquatic invertebrates (Marks and Rutt, 1997; Heywood and  
24 Walling, 2006), alter community structure (Wood and Armitage, 1997) and  
25 undermine river restoration efforts (Kondolf, 1998) as well as reduce flow

1 conveyance, with the potential to increase flood risks (Singer et al., 2008). In  
2 addition, fine sediments play a significant role in the transport of both nutrients and  
3 pollutants within fluvial environments (Bowes et al., 2003; House, 2003; Carter et al.,  
4 2006), with implications for water quality.

5

6 Fine sediment dynamics can be classified into three domains with associated modes  
7 of explanation (Naden, 2010). At the catchment scale, fine sediment flux is  
8 determined primarily by the geomorphological characteristics of the fluvial system  
9 and the history of erosion and sediment transfer (e.g. Trimble, 1983). Time-scales of  
10 interest in determining catchment fine sediment fluxes may range from years to  
11 centuries, and catchment sediment budgets may be created from estimates of fine  
12 sediment storage and the apportionment of sediment sources (e.g. using  
13 fingerprinting techniques; Oldfield et al., 1979). At the reach-scale of river systems,  
14 fine sediment flux is usually explored through measurement of suspended sediment  
15 concentrations (and/or water turbidity) and discharge. The amount of fine sediment  
16 transported by rivers typically follows a power law relationship with discharge (e.g.  
17 Bogardi, 1974), but often with large amounts of scatter in the relationship resulting  
18 from factors such as seasonal variability (e.g. associated with the growth and  
19 senescence of aquatic vegetation; Cotton et al., 2006) and hysteresis effects over  
20 the course of an individual hydrological event (Walling 1977; Williams, 1989). At  
21 small spatio-temporal scales in river channels (e.g. channel cross section; seconds  
22 to minutes), the physical factors that determine fine sediment dynamics are the  
23 physical properties of fine sediments and the high-frequency properties of the flow.  
24 Examination of such characteristics requires high-frequency sampling of water  
25 turbidity and flow velocities, and such studies have illustrated the role of turbulent

1 flow structures in initiating short duration, high magnitude bursts of fine sediment  
2 transport (Lapointe, 1992; 1996; Clifford et al., 1996).

3

4 Recent literature has also given increasing attention to the complexity introduced into  
5 sediment entrainment and transport processes by the presence of aquatic organisms  
6 acting as 'geomorphic agents' or 'ecosystem engineers' (Jones et al., 1994). This  
7 includes aquatic and riparian vegetation (e.g. Gurnell and Petts, 2006; Bertoldi et al.,  
8 2009; Gurnell et al., 2012), aquatic invertebrates (Statzner et al., 2003; Wharton et  
9 al., 2006; Johnson et al., 2009), fish, particularly through spawning activities (e.g.  
10 salmonids; Hassan et al., 2008; 2011) and mammals (e.g. Wright et al., 2002).  
11 Within this context, the influence of aquatic organisms on fluvial sediment dynamics  
12 may be particularly pronounced for invasive species; these may be present at high  
13 densities, and represent a disturbance to which the river system may not be resilient  
14 (Harvey et al., 2011). The susceptibility of natural communities to further invasions  
15 is likely to increase, exacerbated by the interacting stressors of habitat loss and  
16 climate change (Macdonald and Burnham, 2010). A sound understanding of the  
17 ways in which invasive aquatic biota influence fine sediment dynamics from the  
18 patch to the catchment scale of river systems is vital for catchment sediment  
19 budgeting and management efforts, but represents a new and under-developed  
20 research area.

21

22 This paper focuses on the signal crayfish (*Pacifastacus leniusculus*), a species  
23 endemic to North America, that has been introduced to over 20 countries in Europe  
24 since the 1960s (Lewis, 2002). It has rapidly colonised river systems in Britain since  
25 the mid-1970s (Holdich, 2000). Much of the research on signal crayfish to date has

1 focused on biotic interactions. As relatively large bodied omnivores they are typically  
2 detrimental to other invertebrates and fish in the recipient community via predation,  
3 competition, or habitat alteration via consumption of aquatic macrophytes (Guan and  
4 Wiles, 1997; 1998; Crawford et al., 2006). Native white-clawed crayfish  
5 (*Austropotamobius pallipes*) populations in the UK have been decimated by the  
6 arrival and rapid spread of signals over the last few decades (Alderman et al., 1990;  
7 Holdich, 2003; Bubb et al., 2005, Haddaway et al., 2012). Signal crayfish are known  
8 to burrow extensively (Lewis, 2002), can achieve extremely high densities - up to 20  
9 individuals m<sup>2</sup> (Abrahamsson and Goldman, 1970; Bubb, 2004) and are known to  
10 inhabit a range of environments including stillwaters (Ruokonen et al., 2013).

11

12 Several species of crayfish, including signal crayfish, are known to act as  
13 'geomorphic agents', impacting directly upon the physical environment of river  
14 systems (Newton, 2010). This includes bioturbation of sediments through  
15 movements such as walking and tail flips (Statzner and Sagnes, 2008); alteration of  
16 bedform roughness and modifying bed material particle size distributions and particle  
17 consolidation (Statzner et al., 2000; 2003); changes in the composition of fine  
18 sediments associated with the processing of organic matter (Usio, 2000); and  
19 extensive burrowing into soft river banks which may increase bank erosion and  
20 associated increased inputs of fine sediments (Lewis, 2002). Much of this detailed  
21 research on sediment mobilisation, however, focuses on experimental laboratory  
22 streams and on coarser (gravel-sized) bed material. Reaches characterised by fine  
23 bed and bank material may be more susceptible to these geomorphic impacts, and  
24 hence there is a pressing need for improved process understanding at local scales in  
25 order to inform up-scaling and assess the nature and extent of potential implications

1 for catchment-level sediment management, water quality and physical habitat. Such  
2 insights are particularly important given that currently there are no effective  
3 measures for control or mitigation available for signal crayfish (Holdich et al., 1999).  
4 Harvey et al. (2011) review the existing knowledge base for the impacts of signal  
5 crayfish on fine sediment dynamics through their interactions with bed and bank  
6 material through a range of activities. They suggest that patch scale impacts on fine  
7 sediment dynamics through burrowing, movement and feeding activities may lead to  
8 changes in bed material size and structural arrangement, microtopography, bank  
9 erosion, flow resistance and hydrodynamics. Such patch scale impacts may in turn  
10 influence sediment yields, morphology, conveyance capacity and transport of  
11 sediment-associated nutrients and contaminants at larger spatio-temporal scales,  
12 and hence it is recommended that further research is urgently required at the  
13 interface between freshwater ecology, fluvial geomorphology and hydraulics at  
14 various spatiotemporal scales in order to quantify the significance of different  
15 impacts (Harvey et al., 2011).

16

17 This paper presents findings from combined field and laboratory work which aims to  
18 explore signal crayfish impacts on fine sediment dynamics at the patch scale. In  
19 particular, three hypotheses are tested:

20

21 1. Signal crayfish generate pulses of fine sediment mobilisation through their  
22 interactions with river bed and bank material (burrowing and movement) that are  
23 detectable in the flow.

24

1 2. The frequency of pulsed sediment mobilisation events is greater during nocturnal  
2 periods when crayfish in general, and signal crayfish in particular are known to be  
3 most active (Hill and Lodge, 1994; Lozan, 2000; Styris have et al., 2007; Grey and  
4 Jackson, 2012).

5

6 3. The cumulative influence of sediment pulses is sufficient to result in a change to  
7 ambient turbidity levels during nocturnal periods.

8

## 9 **RESEARCH DESIGN AND METHODS**

### 10 *Research design*

11 A combination of laboratory and field study was employed to test the research  
12 hypotheses. Laboratory mesocosm experiments focused on interactions with fine  
13 bed sediment and artificially created banks. Laboratory mesocosms provided  
14 controlled conditions under which detectable changes in suspended sediment  
15 concentration could be directly related to crayfish interacting with artificial banks and  
16 substrates constructed from fine sediments. This was designed to provide 'proof of  
17 concept' for impacts on fine sediment mobilisation. The field study then addressed  
18 whether similar patterns in turbidity were identifiable under field conditions, providing  
19 an indication of the real-world relevance of the process. Given the episodic nature of  
20 bank erosion processes (and burrowing activity) and hence the complexities of  
21 exploring bank erosion processes in the field, field experiments focused on sediment  
22 mobilisation in the near-bed region.

23

24 In both laboratory and field studies, turbidity sensors were used to indicate  
25 suspended sediment concentrations. Turbidity is a measure of the degree to which



1 light passing through a water body is scattered and/or absorbed by particulate matter  
2 held in suspension and has been used to indicate suspended sediment  
3 concentrations in a range of field applications (e.g. Clifford et al., 1995; Gippel, 1995;  
4 Brasington and Richards, 2000; Harvey and Clifford, 2010). The focus on  
5 continuous measurement of turbidity rather than time-integrated suspended  
6 sediment sampling is appropriate in this application, since it allowed continuous high  
7 frequency sampling (Lapointe, 1992; Clifford and French, 1996) and minimised the  
8 disturbance to flow and fine sediments at sampling locations. Turbidity was recorded  
9 using Partech IR40 and/or IR100 infrared turbidity sensors connected to a Campbell  
10 Scientific CR10X data logger, recording data at a frequency of 5Hz. The 0-5 mA  
11 signal produced by the turbidity sensors is converted to a voltage by placing a  
12 resistor in the current loop (Clifford et al., 1995). The IR40 and IR100 have path  
13 lengths of 40 mm and 100 mm, and suspended sediment concentration detection  
14 ranges of 0-200 mg $l^{-1}$  and 0-1500 mg $l^{-1}$ , respectively.

15

16 While the relationship between turbidity and suspended sediment concentration is  
17 affected by particle size, shape and colour, it can be calibrated with respect to either  
18 *in situ* particles (Clifford et al., 1995) or Formazin Turbidity Units (FTU) using a  
19 formazin solution (HMSO, 1984). Calibration curves are generally S-shaped and can  
20 be described by linear relationships for a large part of the range. For the laboratory  
21 experiments, turbidity was converted to sediment concentration (mg $l^{-1}$ ) by calibration  
22 with suspended sediment samples collected across a range of concentrations during  
23 the experiments. Samples were filtered through 2.5 $\mu$ m papers and dried at 100°C to  
24 obtain dry weight. Regression curves were fitted to the relationship between voltage  
25 output and suspended sediment concentration ( $R^2 = 0.99$ ). Results for the

1 laboratory study are reported as suspended sediment concentration ( $\text{mg l}^{-1}$ ). For the  
2 field study, turbidity was converted to FTU through calibration with formazin solution  
3 across a range of concentrations (0-1200 FTU). Regression curves were fitted to the  
4 relationship between voltage output and FTU ( $R^2 = 0.99$ ). Results for the field study  
5 are reported as turbidity (FTU). Suspended sediment concentration was not directly  
6 measured in the field as this would have necessitated disturbing the water column to  
7 obtain water samples, potentially compromising the experiment.

8

### 9 *Laboratory mesocosms*

10 Laboratory experiments were conducted within replicate treatment and control  
11 mesocosm tanks (0.24 x 0.45 x 0.3m) filled with 10l of tap water. There was no flow  
12 through the tanks so both treatment and control tanks were aerated using standard  
13 aquarium air stones and tubing and an AirBlow 100 pump unit. The tanks were  
14 maintained at 15°C under a fixed light regime of 12 hours light (07:00-19:00)  
15 followed by 12 hours of darkness (19:00-07:00). Experiments comprised two  
16 treatments: artificial clay banks and artificial clay substrates, both prepared using  
17 bentonite clay. Each treatment was replicated three times, giving a total of six  
18 experimental runs. For the bank experiments, replicate artificial banks were created  
19 by moulding damp sediment and placing it under weights for several days to create  
20 solid clay blocks of a consistent volume (0.08 x 0.10 x 20.5 m) and shear strength  
21 ( $0.05\text{kg/cm}^3$ ; measured using a shear vane). Banks were held in place at each side  
22 of the tank using wooden panels. For bed substrate experiments, the base of each  
23 tank was covered with bentonite to a depth of 0.03 m and compressed under weights  
24 in the same ways as artificial banks. This was necessary to minimise mobilisation of  
25 substrate during filling of tanks with water. The substrate was also covered with

1 plastic sheeting as the tanks were filled to prevent disturbance of the sediment.  
2 Signal crayfish were collected and identified from field sites within the Greater  
3 London area. One medium-sized crayfish (carapace length 22.5 – 36 mm) was  
4 placed into each treatment tank and no crayfish were present within the control  
5 tanks. Crayfish were not sexed but had both claws intact. Crayfish were stored  
6 within holding tanks with food for several days prior to the experimental runs.  
7 Turbidity sensors were positioned 10 cm above the bed in each of the control and  
8 treatment tanks. Figure 1a illustrates the experimental design.

9

#### 10 *Field study*

11 High-frequency turbidity records were obtained from a site on the eastern arm of the  
12 River Windrush (UK grid reference SP 401 050; Figure 2) on 1 and 2 July 2009. The  
13 Windrush drains an area of 362.6 km<sup>2</sup> from its source in Gloucestershire to its  
14 confluence with the River Thames at Newbridge. The majority of the catchment is  
15 underlain by pervious Oolitic limestone and land use is predominantly agricultural,  
16 although extensive gravel workings and the town of Witney (population 22,265) are  
17 prominent features of the lower valley. The mean annual rainfall in the catchment is  
18 743 mm (1961-90) and the mean daily flow of the Windrush at Newbridge is 3.35  
19 m<sup>3</sup>s<sup>-1</sup> (1950-2009). The Windrush has a large population of signal crayfish. The  
20 native white-clawed crayfish are no longer present in the majority of the catchment  
21 and are confined to its headwaters which have yet to be invaded by the signal  
22 crayfish. The field site forms part of a wider project by the Wildlife Conservation  
23 Research Unit, University of Oxford, researching signal crayfish movements, growth  
24 rates and removal strategies (Moorhouse and Macdonald 2011a, b, c) which  
25 provides quantitative data on signal crayfish densities in the study reach. Channel

1 width at the study location was approximately 5.5 m. Signal crayfish densities at the  
2 time of the study were extremely high. From the 500 m section surrounding the  
3 turbidity monitoring station 6158 individual crayfish were captured from 35 traps set  
4 for a total of 24 nights between April and September 2009 (for further information  
5 see Moorhouse and Macdonald, 2010). Accurate estimation of crayfish densities is  
6 impossible from these trapping data, due to low recapture rates, small nightly home-  
7 ranges of crayfish (modal nightly movement size in these rivers 0-5 m; Moorhouse  
8 and Macdonald, 2011) and biases towards the capture of the largest individuals  
9 (Moorhouse and Macdonald 2010). Both of these latter issues mean that the majority  
10 of the population would likely remain untrapped with traps employed at the spacing  
11 used in this study. The captured individuals, therefore, only represent a small fraction  
12 of the actual population present during the study period. An approximate calculation  
13 of Catch Per Unit Effort (catch per trap per night) reveals > 7 individuals per trap.  
14 Assuming that each trap captured and retained every individual within a 2.5 m radius  
15 this would equate to a minimal density of one crayfish per 3 m<sup>2</sup> of river bed. Given  
16 the known trap bias towards capturing only the largest individuals (those with a  
17 carapace of 50 mm or greater; Moorhouse and Macdonald 2010), and the likelihood  
18 that the majority of even large bodied crayfish within this radius will remain  
19 untrapped or escape (e.g. Holdich et al., 1999) this minimal estimate almost certainly  
20 excludes a large number of crayfish.

21

22 Turbidity monitoring equipment was deployed for a period of 16 hours,  
23 encompassing approximately nine hours of daylight and seven hours of darkness  
24 between 5pm on 1st July 2009 and 9am on 2nd July 2009, incorporating the  
25 nocturnal period when signal crayfish are typically most active (Styrishave et al.,

1 2007). Crayfish activity is also known to be greatest during warmer summer periods  
2 (Lozan, 2000; Johnson et al., 2010). Due to uncertainties associated with the  
3 expected field range of suspended sediment concentrations, two types of turbidity  
4 sensor with differing ranges (IR40C and IR100C) were mounted as a pair at 0.8 of  
5 the water depth from the surface (flow depth of 0.14 m above the bed) in order to  
6 capture-near bed turbidity variations. A further IR40C sensor was mounted at 0.6 of  
7 the water depth from the surface (flow depth of 0.28 m above the bed) to record mid-  
8 flow turbidity levels. The monitoring equipment was positioned within a 'glide'  
9 physical biotope unit (Bisson et al., 1981; Newson and Newson, 2000), characterised  
10 by homogeneous smooth boundary turbulent surface flow conditions and water  
11 depths, and a bed substrate consisting of sand and silt (grain size < 2mm diameter).  
12 Turbidity sensors were installed at a point in the channel cross section that was  
13 expected to be associated with high levels of crayfish movement: towards the bank,  
14 adjacent to overhanging marginal vegetation that provides cover (Jowett et al.,  
15 2008). Figure 1b illustrates the experimental design.

16

17 Hydrological conditions were stable for the duration of the monitoring period:  
18 discharge and water depth at the monitoring location were  $0.46 \text{ m}^3\text{s}^{-1}$  and 0.7 m  
19 respectively and remained constant. Flow stage was low, reflecting the  
20 meteorological conditions encountered at the time of the study: maximum daytime air  
21 temperatures exceeded  $28^\circ\text{C}$  and no rainfall was detected at Met Office operated  
22 rain gauges in the surrounding catchment for at least four days prior to the  
23 experimental period. Discharge records from a downstream gauging station at  
24 Newbridge, below the confluence of the eastern and western forks of the Windrush,  
25 confirm stationarity in flow conditions throughout the monitoring period (Figure 3).

1 High frequency flow properties were sampled over the period of record on an  
2 approximately hourly basis in order to identify changes in turbulent bursting and  
3 event-driven boundary layer organisation (McQuivey, 1973; Clifford et al., 1993;  
4 Ashworth et al., 1996) which can result in short-term sediment suspension (Lapointe,  
5 1992; 1996). High frequency flow measurements were taken immediately adjacent  
6 to turbidity sensors, in the near-bed region (0.8 of the flow depth from the surface),  
7 using a Sontek field 10 MHz 3D Acoustic Doppler Velocimeter (ADV) sampling  
8 streamwise (U), cross stream (V) and vertical (W) velocities at a frequency of 16Hz  
9 for a sampling period of 5 minutes.

10

#### 11 *Data analysis*

12 The first stage of analysis involved visual examination of calibrated turbidity traces to  
13 check for data quality issues and anomalies, and to identify scales of variation the  
14 data set and detrending requirements. For both field and laboratory data sets,  
15 shorter-duration pulsed turbidity events superimposed on longer-term trends were  
16 visually identifiable on time series plots. Pulses were isolated from low amplitude  
17 trends in ambient suspended sediment concentration/turbidity in order to estimate  
18 the number of pulses, their duration and the maximum turbidity value achieved for  
19 each pulse. For geophysical time series characterised by a variety of higher  
20 frequency fluctuations superimposed onto low amplitude, low frequency variations,  
21 detrending is required, and commonly takes the form of a low-order polynomial  
22 regression (e.g. Gordon, 1974; Clifford and French, 1993; French et al., 1993). This  
23 is achieved through visual inspection of model fit, since the nature of high magnitude  
24 fluctuations in such time series will limit the utility of traditional measures of fit such  
25 as  $R^2$ . Following detrending, a conservative analytical regime was adopted based

1 on identification of an appropriate threshold value to delimit sediment pulses and to  
2 focus analysis on the more pronounced suspension events. For laboratory data,  
3 visual inspection of fitted regression curves confirmed that a moving average filter  
4 was most effective in detrending the time series and this was used to detrend the  
5 series. A threshold value of 1.5 was applied to residuals based on visual inspection  
6 of remaining variance in ambient suspended sediment concentration. Pulses were  
7 less discrete for laboratory time series as a result of the non-circulatory nature of the  
8 tanks, and hence pulses separated by less than 2 seconds were combined into a  
9 single event. For field data the low amplitude trend was more subtle relative to the  
10 magnitude of discrete turbidity pulses and detrending involved downweighting of high  
11 magnitude values to within the ambient turbidity range (as identified visually from  
12 timeplots; see Chatfield, 2004). Visual inspection of fitted regression curves  
13 revealed that 4<sup>th</sup> order polynomial regressions best described the low amplitude  
14 trend in field data and these were used to detrend the time series.

15

16 Turbidity residuals were derived from the low magnitude trends and downweighted  
17 values were reinstated within the field time series to allow analysis of turbidity pulse  
18 characteristics. The same threshold value of 1.5 was applied to delimit pulses in the  
19 field time series in order to remove remaining variance in ambient turbidity as  
20 identified by visual inspection. The threshold value chosen will necessarily influence  
21 the estimated absolute number of events and their duration but allows comparability  
22 across data sets. Non-parametric Mann-Whitney U tests were used to identify  
23 statistically significant differences between control and treatment laboratory time  
24 series, and between darkness and daylight hours in the field ambient turbidity data  
25 set.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

For the field data set only, the character of pulsed turbidity events was explored further by classifying pulses separately by shape and magnitude through application of Principal Components Analysis (PCA) and Hierarchical Cluster Analysis (HCA). The approach has been used by Hannah et al. (2000) to classify hydrographs. In order to classify pulses according to event 'shape', a varimax PCA was applied to a data matrix of N columns of pulses by n rows of sequential turbidity values. In order to achieve a matrix of turbidity pulses with an equal number of rows for each pulse, events were artificially extended to the length of the longest duration pulse in the time series using the ambient turbidity threshold value. In this case, the process of applying PCA reduces the matrix of empirical turbidity response curves into a smaller number of generalised curves representing pulses of similar character. PC loadings, therefore, describe the correlation between empirical turbidity response curves and the derived generalised curves. Nine Principal Components (PCs) with eigenvalues greater than 1 cumulatively explained 94% of the variance in the IR40C turbidity series. These represent 9 generalised turbidity response curves. HCA (average linkage method) was then applied to the PC loadings for each pulse, hence grouping pulses with similar loadings on the nine PCs (i.e. those with similar shape; Hannah et al., 2000). The number of clusters within the data set was identified from breaks in the dendrogram. The second process of classifying pulses according to their magnitude involved performing a separate HCA (Ward's method) on a set of variables used to describe the 'magnitude' characteristics for each event: mean turbidity; turbidity range; standard deviation of turbidity; and time to peak. The two HCA outputs were combined to yield a series of shape-magnitude clusters (Hannah et al., 2000). Following the identification and analysis of sediment pulses in the field



1 data, the individual pulses were removed from the time series in order to explore the  
2 presence of a nocturnal trend in ambient turbidity levels outside of pulsed events.

3

## 4 **RESULTS**

### 5 *Visual inspection of laboratory and field time series*

6 Replicate time series for the bank and substrate laboratory experiments are  
7 presented in Figure 4. For each of the two experiments, there was a clear distinction  
8 between control and treatment time series, with evidence of considerable  
9 mobilisation of sediment within the treatment tanks in the form of both shorter  
10 duration, higher magnitude 'pulses' of sediment mobilisation and/or extended  
11 duration lower magnitude increases, superimposed on an overall trend of increasing  
12 suspended sediment concentration following the start of the experimental period.  
13 For the bank treatment, control traces did show some fluctuations over time which  
14 reflects localised bank collapse as a result of saturation. This is negligible in relation  
15 to the treatment tanks, however, and Mann-Whitney U tests confirm statistically  
16 significant differences in suspended sediment concentration between control and  
17 treatment pairs for both bank and bed experiments ( $p < 0.001$ ). Further analysis  
18 focuses on treatment time series only.

19

20 Bank treatment time series were characterised by higher overall suspended  
21 sediment concentrations, greater prominence of shorter-duration fluctuations and  
22 greater similarity between runs compared to substrate time series. No clear  
23 nocturnal trend was apparent for either treatment, but time series data suggest a  
24 tendency towards a reduction in suspended sediment concentration following the  
25 end of the night-time period (Figure 4). For the substrate treatment, there was a

1 considerable difference in the character of sediment traces between runs. A large  
2 number of high magnitude, short duration pulses occur on one of the runs while the  
3 remaining two were characterised by a series of more prolonged, lower magnitude  
4 increases in suspended sediment concentration.

5

6 These trends in suspended sediment concentration correspond with visual  
7 observation of crayfish behaviour in the two treatment tanks. Example still  
8 photographs and video clips are provided in Figure 5 and in the supplementary video  
9 material respectively. For substrate treatments, crayfish dug pits in corners of the  
10 tank in which they positioned themselves for periods of time (Figure 5a and b; and  
11 see supplementary video material). For bank treatments, crayfish created and  
12 inhabited burrows in the artificial banks (Figure 5c and d) but the displacement of  
13 sediment through burrowing was combined with associated periodic, localised  
14 collapse of bank material which also mobilised sediment (Figure 5e and f),  
15 contributing to suspended sediment concentrations. Crayfish also mobilised  
16 sediment by walking across the sediment (see supplementary video material).

17

18 Field turbidity time series derived from the near-bed and mid-flow sensors are  
19 presented in Figure 6 and reveal two key features. First, a large number of high  
20 magnitude turbidity pulses were detected in the near-bed region by both sensors. In  
21 contrast, the mid-flow sensor fails to detect these with the exception of one pulse  
22 which appears to have been detected by all three sensors. Second, there was a  
23 clear increase in the frequency of near-bed turbidity pulses between approximately  
24 21:00 and 06:00, corresponding closely with the period of darkness between sunset  
25 (21:20) and sunrise (04:48). High frequency flow data presented in Figure 7

1 demonstrate stationarity in U V and W velocity components throughout the sampling  
2 period in terms of both average velocity values and variation around the mean. This  
3 suggests that no significant changes in flow variability occurred during the sampling  
4 period, at either low or high frequencies, that would indicate a turbulence-induced  
5 change in pulsed turbidity events during the monitoring period.

6

### 7 *Pulsed turbidity events*

8 Discrete suspended sediment pulses occurring within the laboratory data sets are  
9 explored in Figure 8. The number of pulses differed considerably between  
10 experimental runs for the same treatment, suggesting that crayfish behaviour varied  
11 between runs and exerted a considerable influence on the character of turbidity time  
12 series. Overall, the total number of pulses was higher for the bank treatments  
13 (between 743 and 1941 pulses; with a cumulative duration of between 7 and 10  
14 hours of the record) relative to the substrate treatment (between 398 and 737 pulses;  
15 with a cumulative duration between 1 and 7 hours of the record). A small amount of  
16 bank collapse occurred under the control treatment suggesting that saturation of  
17 artificial banks played a small role, but bank collapse was greatly increased by  
18 crayfish burrowing. Two of the substrate time series showed less evidence of high  
19 magnitude pulsed events, but instead were characterised lower magnitude, longer  
20 duration periods of elevated suspended sediment associated with crayfish activity  
21 occurring further from the sensor (in corners of the tanks; see Figure 5 and  
22 supplementary video material).

23

24 The number of pulses occurring in each hour of the monitoring period varied  
25 between runs and ranged from zero to 300. There was no clear nocturnal trend in

1 the number of pulsed events as identified previously from time plots, although both  
2 bank and substrate experiments showed evidence of an enhanced number of pulses  
3 between 00:00 and 03:00 which then declines. For the substrate treatment there  
4 was considerable variability between runs, but a total of 421 pulses were recorded  
5 across the three runs between 00:00 and 03:00 (mean 35 pulses per hour; standard  
6 error 6.4) compared to 245 pulses during the period of light between 07:00 and  
7 10:00 (mean 26 per hour; standard error 6.0). For the bank treatment a total of 917  
8 pulses were recorded across the three runs between 0:00 and 03:00 (mean 76  
9 pulses per hour; standard error 10.8) compared to 331 between 07:00 and 10:00  
10 (mean 28 per hour; standard error 7.3). The vast majority of pulses were short in  
11 duration (across bank runs median = 3.75s and upper quartile = 8.75s; for substrate  
12 runs median = 3.25s and upper quartile = 10.75s), and with maximum suspended  
13 sediment concentration values up to 5.4 mg $l^{-1}$  (for bank runs median 3.75 mg $l^{-1}$  and  
14 upper quartile = 5.14 mg $l^{-1}$ ; for substrate runs median = 3.95 mg $l^{-1}$  and upper quartile  
15 = 5.40 mg $l^{-1}$ ). A smaller number of pulses endured for over 1 minute, up to a  
16 maximum of 20 minutes, reflecting periods of more sustained crayfish activity.  
17 Likewise a small number of short duration, high magnitude pulses with peak  
18 suspended sediment concentration values exceeding 100mg $l^{-1}$  occurred in the  
19 record for both treatments, indicating sediment mobilisation events closer to the  
20 turbidity sensor. Since the pulse characteristics for the laboratory time series will be  
21 strongly controlled by tank hydraulics, and hence may be considerably different in  
22 magnitude and duration to crayfish-induced sediment mobilisation events occurring  
23 under field conditions, pulse characteristics for laboratory data were not analysed in  
24 further detail.

25

1 Figure 9 plots characteristics for each pulsed turbidity event identified on the IR40  
2 near-bed sensor for the field turbidity series. Since several high magnitude turbidity  
3 pulses detected by the IR100C sensor resulted in the readings outside of the sensor  
4 range, and only one pulse was detected on the mid-flow IR40C sensor, turbidity  
5 pulse analysis focused on the near-bed IR40C sensor only for field data. In total, 88  
6 events were identified for the IR40C near-bed sensor, an order of magnitude below  
7 the values identified in laboratory data. This was expected given the small surface  
8 area of the laboratory tanks and the absence of flow to diffuse and advect sediment  
9 plumes. One pulse was removed from the time series for the IR40C since it was  
10 over 15 minutes in length (compared to <40 s for all other events) and of similar  
11 magnitude throughout. This was considered to represent a temporary sensor  
12 obstruction such as a piece of organic debris, rather than a discrete sediment  
13 suspension event.

14

15 Figure 9a illustrates a clear nocturnal trend in pulse frequency for the field data set:  
16 no pulses were detected between 17:00 and 19:00 and the number of pulsed events  
17 per hour then increased from 1 between 19:00 and 19:59 to a peak of 23 between  
18 23:00 and 23:59, and subsequently decreased to zero by 07:00 the following day.  
19 Pulse duration and maximum turbidity revealed a large degree of scatter through the  
20 period of record (Figure 9c and d), although there appeared to be a tendency  
21 towards higher variability in pulse duration/ magnitude towards the mid-point of the  
22 nocturnal period (particularly around 24:00 and 03:00). The relationship between  
23 pulse duration and magnitude (Figure 9b) shows a tendency for higher peak turbidity  
24 values to be associated with longer duration pulses, but there is considerable scatter  
25 observed for the longest duration pulses. The majority of pulses were associated

1 with durations between 1s and 15s and maximum turbidity values up to 1500 FTU,  
2 with a smaller cluster of much longer duration events up to a maximum of 37s and  
3 with a maximum turbidity up to 2600 FTU.

4

5 Figure 10 plots turbidity time series for individual pulses classified according to the  
6 shape-magnitude clusters derived from PCA and HCA. Each plot in Figure 10  
7 corresponds to a shape cluster and uses symbology to differentiate between  
8 magnitude clusters for each. In total, eight pulse 'shape' clusters (A-H) were  
9 identified by applying HCA to the PC loadings that described correlation between  
10 empirical turbidity response curves and the generalised curves derived from the  
11 PCA. In a separate process, three pulse 'magnitude' clusters (1-3 describing low,  
12 intermediate and high magnitudes respectively) were identified based on pulse  
13 magnitude characteristics (mean turbidity; turbidity range; standard deviation of  
14 turbidity; and time to peak). Clusters A and B (Figure 10a and b) contained the  
15 lowest magnitude and shortest duration pulses; these pulse shapes were associated  
16 only with the lowest 'magnitude' cluster. The pulses in cluster B have a longer lag  
17 time, higher overall event magnitude and lower level of negative skewing relative to  
18 cluster A. The shapes of pulses in clusters C, D and E (Figure 10c,d,e) were more  
19 complex than A and B, and demonstrate intermediate durations and variable  
20 magnitudes. Most pulses falling within these clusters show a lag in peak turbidity of  
21 up to 7 seconds from the onset of the event. Clusters F, G, and H described a small  
22 number of pulse events (3, 1 and 2 pulses respectively) and are therefore plotted  
23 together for brevity in Figure 10f. Together, they represent longer duration pulses  
24 with long lags and multiple peaks. Clusters E, F and G all contained pulses in which  
25 a secondary peak occurred following a delay of between 15 and 30 s from the onset

1 of the pulse. These secondary peaks could represent a new, potentially unrelated  
2 turbidity event but since there is not a complete return to ambient turbidity, these  
3 pulses were treated as an individual event for the purposes of this analysis. Given  
4 their close proximity in time and similarity in shape, it is possible that these pulses  
5 may represent a repetition of the same phenomenon occurring in the same spatial  
6 location. The occurrence of the different shape-magnitude pulse clusters varied  
7 throughout the period of record. Events belonging to cluster A were distributed  
8 throughout the period between 19:00 and 06:00 but the period of greatest variability  
9 in cluster type occurred between 22:00 and 03:00, and almost all of the events  
10 associated with clusters B, C, D and E, G and H fell within this period.

11

#### 12 *Nocturnal trend in ambient turbidity for field data*

13 Figure 11 plots average hourly turbidity values for the low magnitude trend in  
14 ambient turbidity in field data (i.e. following the removal of discrete high magnitude  
15 turbidity pulses) for all three sensors. An abrupt increase in turbidity at around 19:50  
16 which lasted for approximately 50 minutes on the IR100C sensor only was  
17 considered to represent an anomalous event associated with a piece of debris  
18 becoming trapped around the sensor and was therefore excluded from the ambient  
19 data set. For the near-bed sensors there was a clear nocturnal trend associated  
20 with an increase in ambient turbidity from approximately 21:00 up to around 02:00,  
21 which then receded to approximately the original level by around 08:00. This trend  
22 was more pronounced for the IR100C which is more sensitive at the lower part of the  
23 turbidity range, thereby affording greater resolution (Henshaw, 2009). The mid-flow  
24 IR40C sensor curve demonstrated a slightly different character from the near-bed  
25 sensor curves, rising from around 21:00 until reaching a peak around 04:00, but

1 failing to return to the original level during the period of record and remaining  
2 approximately 5 FTU higher than the starting turbidity value. In order to test the  
3 statistical significance of these nocturnal variations, Mann Whitney U tests were  
4 performed on light and darkness groupings of minute-averaged ambient turbidity  
5 values for each of the three turbidity sensors according to the timing of sunset and  
6 sunrise. This confirmed statistically significant differences in the mean daylight and  
7 darkness ambient turbidity values for each of the three sensors ( $p < 0.001$ ).

8

## 9 **DISCUSSION**

10 The analysis of high frequency suspended sediment and turbidity time series from  
11 laboratory and field studies revealed a number of interesting features that support  
12 the research hypotheses. First, signal crayfish were able to mobilise fine sediment  
13 through different types of movement, creating pulsed sediment suspension events of  
14 varying magnitude and duration under controlled laboratory conditions. These  
15 pulsed events were sufficient to drive a statistically significant increase in suspended  
16 sediment concentrations in still-water tanks. A higher total number of pulses were  
17 recorded for the bank treatment and this reflects a combination of factors. First,  
18 there was greater potential for burrowing into the artificial banks relative to the bed  
19 substrate due to the greater depth and width of consolidated material. For bank  
20 treatments, crayfish were observed to burrow extensively into artificial clay banks,  
21 consistent with field observations of dense networks of burrows in invaded habitats  
22 (Guan, 1994; Holdich et al., 1999). In addition, the occurrence of sediment pulses in  
23 the bank treatment reflects the combined contribution of 'direct' impacts of burrow  
24 excavation activity transferring sediment into water column and the 'indirect' impacts  
25 associated with periodic partial bank collapse as a result of burrowing. This was



1 combined with crayfish movements across destabilised material. For substrate  
2 treatments, pulsed suspension events were associated with crayfish movements  
3 across the bed and the digging of pits in the corner of the tanks. Other species of  
4 crayfish have also been observed to mobilise fine sediment during walking, foraging  
5 and swimming (e.g. *Paranephrops planifrons*: Parastacidae; Parkyn et al., 1997) and  
6 signal crayfish are known to mobilise larger gravel-sized sediment particles (Johnson  
7 et al., 2010). Signal crayfish have also been shown to create pit and mound features  
8 in coarse sediment beds under laboratory conditions (Johnson et al., 2010), similar  
9 to those observed in fine sediment beds for this study. This may partly reflect a lack  
10 of cover in substrate tanks, leading to the crayfish adopting a defensive position.  
11 While there is no clear nocturnal trend in laboratory data, the number of pulses was  
12 enhanced between 00:00 and 03:00 and declined into the subsequent period of light  
13 at the end of the night-time period (see Figure 4 and Figure 8). Across the three  
14 experimental runs, the number of pulses in the period between 00:00 and 03:00 was  
15 almost three times higher than the period of light between 07:00 and 10:00 for the  
16 bank treatment, and almost two times higher for the substrate treatment.

17

18 The laboratory experiments provide clear evidence of the ability of signal crayfish to  
19 mobilise fine sediment under controlled conditions. However, relationships between  
20 crayfish behaviour and suspended sediment concentrations for laboratory  
21 experiments will be greatly influenced by both the hydraulic conditions of the tanks  
22 and the altered behaviour of the crayfish as a result of their placement in a different  
23 environment. Tanks were not recirculating, meaning that pulsed suspension events  
24 were not subject to advection and diffusion processes that would be significant under  
25 field conditions (Rutherford, 1994). This will increase overall suspended sediment

1 levels and limits the potential for detailed analysis and interpretation of sediment  
2 pulse characteristics for laboratory time series. The artificial lighting and  
3 temperature control as well as the lack of cover, food sources, predation, competition  
4 and other biotic interactions will have influenced crayfish behaviour during the  
5 experimental period. For example, crayfish are known to cease or curtail various  
6 behaviours in the presence of conspecific alarm odours (Hazlett, 2003), and in the  
7 absence of such odours (as would be the case in laboratory conditions) it is likely  
8 that variations signal crayfish activity may not conform to a nocturnal pattern (e.g.  
9 Lozan, 2000).

10

11 The field data set reveals pulsed turbidity events similar to those identified in  
12 laboratory data. These occur in the near-bed region and are detected by two  
13 separate sensors, but similar pulses are not detected in the mid-flow region at the  
14 same location. Fewer pulses are identified in the field time series relative to  
15 laboratory data. This was expected, given the small surface area of laboratory  
16 mesocosms and the influence of advection and diffusion processes under field  
17 conditions. Importantly, turbidity pulses identified on field time series appear tightly  
18 constrained to the period of darkness between sunset and sunrise. The pulsed  
19 events vary in duration, magnitude and 'shape' but can be grouped statistically into  
20 distinct clusters of similar magnitude and shape characteristics. There is a notable  
21 lack of short duration (<5 s) high magnitude observations characteristic of  
22 turbulence-induced sediment suspension events (McQuivey 1973; French and  
23 Clifford, 1992; Lapointe, 1996; Roy et al., 1996; Buffin-Belanger and Roy, 2005), and  
24 few extended duration (>15 s) events. The turbidity pulses are most frequent and  
25 most varied in their characteristics towards the mid-point of nocturnal period. These

1 temporal trends correlate with known nocturnal increases in signal crayfish activity  
2 including movement, burrowing, fighting and feeding (Flint, 1977; Guan and Wiles,  
3 1998; Lozan, 2000; Styris have et al., 2007).

4

5 Observational data on crayfish activity during the monitoring period are not available  
6 for the field study, limiting direct process inference. However, the known behavioural  
7 characteristics of signal crayfish, combined with their presence in extremely high  
8 densities at the field site (Moorhouse and Macdonald 2011a, b, c), and the similar  
9 features identified in laboratory data sets, implies they are likely to be responsible for  
10 the observed trends in field data. Importantly, the turbidity increases are not  
11 explained by temporal variations in high frequency velocity characteristics, and seem  
12 independent of discharge variations or possible enhanced supply from catchment  
13 run-off events in the immediate past. Additionally, the turbidity increases are more  
14 marked near to the river bed, indicating a local origin, and the nocturnal pattern  
15 observed in field data, with clear onset and cessation aligned with sunset and  
16 sunrise implies a causal or generating mechanism which was biological rather than  
17 physical in origin. The data also complement previous research that links  
18 bioturbation by the red swamp crayfish (*Procambarus clarkii*) with increased  
19 suspended solids and dissolved nutrients in a wetland enclosure study (Angeler et  
20 al., 2001).

21

22 The occurrence of pulsed turbidity events within the nocturnal period of the field time  
23 series corresponds with a statistically significant increase in ambient turbidity values  
24 (i.e. excluding pulses) in the near-bed region for the same period. Furthermore, a  
25 statistically significant increase in ambient turbidity is also recorded in the mid-flow

1 region which appears time-lagged relative to the near-bed trends. These data  
2 suggest a combined effect of localised turbidity events occurring both within the  
3 immediate vicinity of the sensors and upstream that was sufficient to drive a  
4 statistically significant increase in ambient turbidity both at the bed and also in the  
5 mid-flow region. The difference between the near-bed and mid-flow trends could  
6 reflect the cumulative effects of crayfish activity upstream and the diffusion and  
7 downstream advection of mobilised sediments which may have a time-lagged  
8 influence in the mid-flow region.

9

10 The combined field and laboratory approach taken in this paper was designed to  
11 overcome some of the extreme practical difficulties associated with process-based  
12 research at the geomorphology-ecology interface. For this study, laboratory  
13 experimentation provides a controlled environment for a proof of concept study that  
14 identifies detectable pulsed sediment suspension events of varying character that  
15 can be directly attributed to crayfish interacting with bed and bank material through  
16 movement and burrowing. However, crayfish behaviour will be influenced by the  
17 artificial surroundings of the laboratory, and the non-recirculatory nature of the tanks  
18 does not provide an accurate description of the character and significance of the  
19 pulsed mobilisation events under field conditions. Observational evidence is much  
20 more difficult to secure in the field as a result of a number of practical constraints,  
21 notably the need to isolate hydrological and hydraulic influences, maintenance,  
22 power and data storage constraints of field equipment, and the nocturnal behaviour  
23 of signal crayfish. In this case the field data set complements the laboratory data set  
24 by identifying similar pulsed sediment suspension features in a reach with extremely  
25 high densities of crayfish. This suggests real world significance, with the potential to

1 drive an overall increase in turbidity and hence influence sediment dynamics at  
2 larger spatio-temporal scales (Harvey et al., 2011). While the signal crayfish is of  
3 particular interest due to its relatively large body size and aggressive nature, other  
4 invasive species of crayfish in the UK are also known to burrow into river banks (e.g.  
5 red swamp crayfish *Procambarus clarkii* and spiny-cheeked crayfish *Orconectes*  
6 *limosus*) and hence may have similar impacts upon mobilisation of fine sediment  
7 from the river bed and banks.

8

9 The results presented in this paper suggest that further research into the impact of  
10 signal crayfish on fine sediment dynamics is warranted. Future studies may  
11 incorporate in situ enclosure/ exclosure field experiments (removing animals in order  
12 to help identify the extent of impacts (e.g. localised versus significant downstream  
13 advection)) across sites with different fine sediment bed material characteristics and  
14 for different hydrological conditions. Opportunities for paired control-impact studies  
15 may also be identified at invasion fronts, allowing a comparison of fine sediment  
16 dynamics across reaches with similar physical characteristics in the presence and  
17 absence of a crayfish population. Observational data may be collected alongside  
18 suspended sediment monitoring through Passive Integrated Transponder (PIT)  
19 technology in order to detect animal movement patterns at the reach scale (e.g. see  
20 Bubb et al., 2006) and/or underwater videography in order to directly observe  
21 movements at a local scale. This may allow relationships between rates of  
22 movement across the bed and suspended sediment concentrations to be explored,  
23 and different types of activity (e.g. movement, feeding, fighting) to be linked to  
24 different styles of sediment mobilisation.

25

1 **CONCLUSION**

2 This paper presents findings from combined field and laboratory studies that  
3 demonstrate the potential for signal crayfish to act as drivers of fine sediment  
4 dynamics at the patch scale of river systems. Results support the three research  
5 hypotheses presented. Signal crayfish generate pulses of fine sediment mobilisation  
6 through interactions with river bed and bank material (burrowing and movement) that  
7 are detectable in the flow. Pulsed sediment mobilisation events are tightly  
8 constrained to the nocturnal period in which signal crayfish are known to be most  
9 active under field conditions, and show evidence of enhanced frequencies in the  
10 nocturnal period under laboratory conditions. For both data sets, pulsed events  
11 appear sufficient in their cumulative magnitude and frequency to drive a change to  
12 ambient suspended sediment/ turbidity levels. Whilst similar activities and  
13 associated sediment disturbances could be associated with native species, the  
14 larger body size and more aggressive nature of the signal crayfish, and its presence  
15 in many catchments in extremely high densities, may lead to more significant  
16 sediment disturbance compared to that which may be expected for native species.  
17 Furthermore, signal crayfish are known to burrow much more extensively in invaded  
18 environments relative to both their native range and to other species such as the  
19 UK's native white-clawed crayfish.

20

21 If the impacts of signal crayfish on fine sediment mobilisation and transport become  
22 significant in specific reaches and/or for certain periods of time, downstream impacts  
23 could include morphological change, increased turbidity levels, and the mobilisation  
24 and transport of sediments and sediment-associated nutrients and contaminants.  
25 Such impacts could have detrimental effects for the ecological status of water

1 bodies, and for flood risk through changes in conveyance capacity. Further process-  
2 based studies are required to develop a full understanding of relationships between  
3 fine sediment mobilisation and transport associated with the various mechanistic  
4 abilities of crayfish; relationships between impact and body size/sex; interspecific  
5 interactions; the influence of additional habitat-related variables such as food  
6 sources and refugia; temporal variations in activity levels including seasonal  
7 (temperature) as well as diurnal; and the nature and scale of potential impacts  
8 across a range of river styles.

9

## 10 **ACKNOWLEDGEMENTS**

11 The research was supported by a *British Society for Geomorphology* Research  
12 Grant ('Field evidence for signal crayfish impacts on fine sediment dynamics in a  
13 lowland river'). Tom Moorhouse was funded by the Esmée Fairbairn Foundation.  
14 The authors would like to thank Matej Faller for helpful comments on the manuscript.

15

## 16 **REFERENCES**

17 Abrahamsson SAA and Goldman CR (1970) Distribution, density and production of  
18 the crayfish *Pacifastacus leniusculus* Dana in Lake Tahoe, California – Nevada.

19 *Oikos* **21**: 83–91.

20

21 Alderman DJ, Holdich DM, and Reeve I. (1990) Signal crayfish as vectors in crayfish  
22 plague in Britain. *Aquaculture* 86: 3–6.

23

24 Ashworth PJ, Bennett SJ, Best JL and McLelland SJ. (1996) Coherent flow  
25 structures in open channels. John Wiley and Sons Ltd, Chichester.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

Bertoldi W, Gurnell AM and Drake NA (2011) The topographic signature of vegetation development along a braided river: results of a combined analysis of airborne lidar, colour air photographs and ground measurements. *Water Resources Research* **47**: W06525

Bisson PA, Nielson JL, Palson RA, Grove LE. (1981) A system of naming habitat in small streams, with examples of habitat utilization by salmonids during low streamflow. In Acquisition and Utilization of Aquatic Habitat Inventory Information: Proceedings of a Symposium. American Fisheries Society Western Division: Bethesda, MD.

Bogardi J. (1974) Sediment Transport in Alluvial Streams. Akadémiai Kiadó, Budapest

Bowes MJ, House, WA, and Hodgkinson, RA. (2003) Phosphorous dynamics along a river continuum. *Science of the Total Environment* **313**: 199-212.

Brasington J and Richards KS. (2000) Turbidity and suspended sediment dynamics in the Nepal Middle Hills. *Hydrological Processes* **14**: 2259-2274.

Bubb DH, Thom TJ, and Lucas MJ (2004) Movement and dispersal of the invasive signal crayfish *Pacifastacus leniusculus* in upland rivers. *Freshwater Biology* **49**: 357–368.



- 1   Bubb DH, Thom TJ, and Lucas MC (2005) The within catchment invasion of the non-  
2   indigenous signal crayfish (*Dana*) in upland rivers. *Bulletin Francais de Peche et de*  
3   *Pisciculture* **376–377**: 665–673.
- 4
- 5   Bubb DH, Thom TJ, Lucas MC. (2006) Movement patterns of the invasive signal  
6   crayfish determined by PIT telemetry. *Canadian Journal of Zoology* **84**: 1202–1209.
- 7
- 8   Buffin-Belanger T and Roy AG. (2005) 1 min in the life of a river: selecting the  
9   optimal record length for the measurement of turbulence in fluvial boundary layers.  
10  *Geomorphology* **68**: 77-94.
- 11
- 12  Carter J, Walling DE, Owens PN and Leeks GJL. (2006) Spatial and temporal  
13  variability in the concentration and speciation of metals in suspended sediment  
14  transported by the River Aire, Yorkshire, UK. *Hydrological Processes* **20**: 3007-  
15  3027.
- 16
- 17  Chatfield C. (2004) *The Analysis of Time Series: An Introduction*. CRC Press LLC:  
18  Boca Raton, Florida.
- 19
- 20  Clifford NJ, Richards KS, Brown RA and Lane SN. (1995) Scales of variation of  
21  suspended sediment concentration and turbidity in a glacial meltwater stream.  
22  *Geografiska Annaler* **77A**(1-2), 45-65.
- 23
- 24  Clifford NJ, French JR and Hardisty J. (1993) *Turbulence: Perspectives on Flow and*  
25  *Sediment Transport*. Chichester, John Wiley and Sons Ltd.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

Clifford NJ, Richards KS, Brown RA, Lane SN. (1995) Laboratory and field assessment of an infrared turbidity probe and its response to particle size and variation in suspended sediment concentration. *Hydrological Sciences Journal* **40(6)**: 771–791.

Clifford NJ and French JR. (1996) Criteria for suspended sediment monitoring in river environments: field evaluation of contrasting turbidity sensor types using simulated sediment transport events. In Maxwell WHC, Preul HC and Stout GE (eds.) *Proceedings Rivertech96: 1st International Conference on New/emerging Concepts for Rivers*: 754-761.

Cotton JA, Wharton G, Bass JAB, Heppell CM and Wotton RS. (2006) Plant-water-sediment interactions in lowland permeable streams: investigating the effect of seasonal changes in vegetation cover on flow patterns and sediment accumulation. *Geomorphology* **77**: 320-334.

Crawford L, Yeomans WE, and Adams CE (2006) The impact of introduced signal crayfish *Pacifastacus leniusculus* on stream invertebrate communities. *Aquatic Conservation: Marine and Freshwater Ecosystems* **16**: 611–621.

Flint RW. (1977) Seasonal activity, migration and distribution of crayfish, *Pacifastacus leniusculus*, in Lake Tahoe. *American Midland Naturalist* **97**: 280-292.

1 French JR and Clifford NJ. (1992) Characteristics and "Event-Structure" of near-bed  
2 turbulence in a macrotidal saltmarsh environment. *Estuarine, Coastal and Shelf*  
3 *Science* **34**: 49-69.

4

5 Gippel CJ (1995) Potential of turbidity monitoring for measuring the transport of  
6 suspended solids in streams. *Hydrological Processes* **9**: 83-97.

7

8 Grey J and Jackson MC. (2012) 'Leaves and eats shoots': direct terrestrial feeding  
9 can supplement invasive red swamp crayfish in times of need. *PLoS ONE* **7(8)**:  
10 e42575.

11

12 Guan RZ and Wiles PR. (1997) Ecological impact of introduced crayfish on benthic  
13 fishes in a British lowland river. *Conservation Biology* **11**: 641–647.

14

15 Guan RZ and Wiles PR. (1998) Feeding ecology of the signal crayfish *Pacifastacus*  
16 *leniusculus* in a British lowland river. *Aquaculture* **169**: 177-163.

17

18 Gurnell AM and Petts GE. (2006) Trees as riparian engineers: the Tagliamento  
19 River, Italy. *Earth Surface Processes and Landforms* **31**: 1558-1574.

20

21 Gurnell AM, Bertoldi W and Corenblit D. (2012) Changing river channels: the roles of  
22 hydrological processes, plants and pioneer fluvial landforms. *Earth Science Reviews*  
23 **111**: 129-141.

24

1 Haddaway NR, Wilcox RH, Heptonstall REA, Griffiths HM, Mortimer RJG, Christmas  
2 M, and Dunnet AMI. (2012) Predatory functional response and prey choice identify  
3 predation differences between native/invasive and parasitised/unparasitised crayfish.  
4 *PLoS ONE* **7(2)**: e32229.

5

6 Hannah DM, Smith BPG, Gurnell AM and McGregor GR. (2000) An approach to  
7 hydrograph classification. *Hydrological Processes* **14**: 317-338.

8

9 Harvey GL and Clifford NJ. (2009) Microscale hydrodynamics and coherent flow  
10 structures in rivers: Implications for the characterization of physical habitat. *River*  
11 *Research and Applications* **25**: 160-180.

12

13 Harvey GL and Clifford NJ. (2010) Experimental field assessment of suspended  
14 sediment pathways for characterising hydraulic habitat. *Earth Surface Processes*  
15 *and Landforms* **35**: 600–610.

16

17 Harvey GL, Moorhouse TP, Clifford NJ, Henshaw AJ, Johnson MF, Macdonald DW,  
18 Reid I and Rice SP. (2011) Evaluating the role of invasive aquatic species as drivers  
19 of fine sediment-related river management problems: the case of the signal crayfish  
20 (*Pacifastacus leniusculus*). *Progress in Physical Geography* **35**: 517–533.

21

22 Hassan MA, Peticrew EL, Montgomery DR, Gottesfeld AS and Rex JF. (2011)  
23 Salmon as biogeomorphic agents in gravel bed rivers: the effect of fish on sediment  
24 mobility and spawning habitat. *Geophysical Monograph Series* 194: 337-352.

25

1 Hassan MA, Gottesfeld AS, Montgomery DR, Tunnicliffe JF, Clarke GKC, Wynn G  
2 Jones-Cox H, Poirier R, Maclsaac E, Herunter H, and Mcdonald SJ. (2008) Salmon-  
3 driven bedload transport and bed morphology in mountain streams. *Geophysical*  
4 *Research Letters* 35: L04405.

5  
6 Hazlett BA. (2003) The effects of starvation on crayfish responses to alarm odor.  
7 *Ethology* **109**: 587-592.

8  
9 Henshaw AJ. (2009) Impacts of land use changes and land management practices  
10 on upland catchment sediment dynamics: Pontbren, mid-Wales. Unpublished PhD  
11 thesis. University of Nottingham.

12  
13 Heywood MJT and Walling DE. (2006) The sedimentation of salmonid spawning  
14 gravels in the Hampshire Avon catchment, UK: implications for the dissolved oxygen  
15 content of intragravel water and embryo survival. *Hydrological Processes* **21**: 770-  
16 788.

17  
18 Hill AM and Lodge DM. (1994) Diel changes in resource demand - competition and  
19 predation in species replacement among crayfishes. *Ecology* **75**: 2118-2126.

20  
21 Holdich DM, Rogers WD and Reynolds JD. (1999) Native and alien crayfish in the  
22 British Isles. In: Gherardi F and Holdich DM.(Eds.) *Crayfish in Europe as alien*  
23 *species: how to make the best of a bad situation?* 221–236. Balkema, Rotterdam.

24

- 1 Holdich DM. (2000) The Development of Ecological Requirements to inform the  
2 Production of Conservation Objectives for white-clawed crayfish. English Nature,  
3 Peterborough.
- 4
- 5 House WA. (2003) Geochemical cycling of phosphorous in rivers. *Applied*  
6 *Geochemistry* 18: 739-748.
- 7
- 8 HMSO (1984) Methods for the examination of waters and associated materials:  
9 Colour and turbidity of waters – 1981 Tentative methods. HMSO, London.
- 10
- 11 Johnson MF, Reid I, Rice SP and Wood PJ (2009) Stabilization of fine-gravels by  
12 net-spinning caddisfly larvae. *Earth Surface Processes and Landforms* 34: 413 -  
13 423.
- 14 Johnson MF, Rice SP and Reid I (2010) Topographic disturbance of subaqueous  
15 gravels by signal crayfish (*Pacifastacus leniusculus*). *Geomorphology* 123: 269-278.
- 16
- 17 Jones CG, Lawton JH, and Shachak M (1994) Organisms as ecosystem engineers.  
18 *Oikos* 69: 373–386.
- 19
- 20 Jowett IG, Parkyn SM and Richardson J. (2008) Habitat characteristics of crayfish  
21 (*Paranephrops planifrons*) in New Zealand streams using generalized additive  
22 models (GAMs). *Hydrobiologia* 596: 353 – 365.
- 23
- 24 Kondolf GM (1998) Lessons learned from river restoration projects in California.  
25 *Aquatic Conservation: Marine and Freshwater Ecosystems* 8:39-52.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

Lane SN, Tayefi V Reid SC, Yu D, Hardy RJ. (2007) Interactions between sediment delivery, channel change, climate change and flood risk in a temperate upland environment. *Earth Surface Processes and Landforms* **32**: 429-446

Lapointe M. (1992) Burst-like sediment suspension events in a sand bed river. *Earth Surface Processes and Landforms* **17**:253-2970.

Lapointe MF. (1996) Frequency spectra and intermittency of the turbulent suspension process in a sand- bed river. *Sedimentology* **43**: 439–449.

Lewis SD. (2002) *Pacifastacus* Biology of freshwater crayfish. In: Holdich DM. (Ed.) *Biology of Freshwater Crayfish*: 511–540. Blackwell, Oxford.

Lozan JL. (2000) On the threat to the European crayfish: A contribution with the study of the activity behaviour of four crayfish species (Decapoda: Astacidae). *Limnologica* **30**: 156-161.

Macdonald D and Burnham D. (2010) *The state of Britain’s mammals: a focus on invasive species*. People's Trust for Endangered Species: London. ISBN 0-9545711-5-0.

Marks SD and Rutt GP. (1997) Fluvial sediment inputs to upland gravel bed rivers draining forested catchments: potential ecological impacts. *Hydrology and Earth System Sciences* **1**: 499-508.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

McQuivey RS. (1973) Summary of turbulence data from rivers, conveyance channels and laboratory flumes. United States Geological Survey Professional Paper, 802B.

Moorhouse, T.P., Macdonald, D.W. (2011a). The effect of manual removal on movement distances in populations of signal crayfish (*Pacifastacus leniusculus*). *Freshwater Biology* **56**: 2370-2377.

Moorhouse, T.P., Macdonald, D.W. (2011b). The effect of removal by trapping on body condition in populations of signal crayfish. *Biological Conservation* **144**: 1826-1831.

Moorhouse, T.P., Macdonald, D.W., (2011c). Immigration rates of signal crayfish (*Pacifastacus leniusculus*) in response to manual control measures. *Freshwater Biology* **56**: 993-1001.

Naden PS. (2010) The fine-sediment cascade. In: Burt T and Allison R. (eds) *Sediment Cascades*. Wiley-Blackwell, Chicester.

Newson MD and Newson CL. (2000) Geomorphology, ecology and river channel habitat; mesoscale approaches to basin-scale challenges. *Progress in Physical Geography* **24(2)**: 195–217.

Newton A. (2010) Geomorphology: Crayfish at work. *Nature Geoscience* **3**: 592.



1 Oldfield F, Rummary TA, Thompson R and Walling DE. (1979) Identification of  
2 suspended sediment sources by means of magnetic measurements: some  
3 preliminary results. *Water Resources Research* **15**: 211-218.  
4

5 Parkyn SM, Rabeni CF, and Collier KJ (1997) Effects of crayfish (*Paraneohrops*  
6 *planifrons*: Parastacidae) on in-stream processes and benthic faunas: A density  
7 manipulation experiment. *New Zealand Journal of Marine and Freshwater Research*  
8 **31**: 685–692.  
9

10 Roy AG, Buffin-Bélanger T and Deland S. (1996) Scales of turbulent coherent flow  
11 structures in a gravel-bed river. In: Ashworth P J, Bennett SJ, Best JL and  
12 McLelland SJ (eds.) *Coherent Flow Structures in Open Channels*, Chichester, John  
13 Wiley and Sons Ltd pp. 147-164.  
14

15 Ruokonen TJ, Kiljunen M, Karjalainen J and Hämäläinen H. (2012) Invasive  
16 crayfish increase habitat connectivity: a case study in a large boreal lake.  
17 *Knowledge and Management of Aquatic Ecosystems* **407**, 08.  
18

19 Rutherford JC. (1994) *River Mixing*. John Wiley & Sons: Chichester.  
20

21 Singer, MB, Aalto, R and James, LA. (2008) Status of the Lower Sacramento Valley  
22 flood-control system within the context of its natural geomorphic setting. *Natural*  
23 *Hazards Review*, **9**: 104-115.  
24

- 1 Statzner B, Fievet E, Champagne JY, Morel R and Herouin E. (2000) Crayfish as  
2 geomorphic agents and ecosystem engineers: Biological behavior affects sand and  
3 gravel erosion in experimental streams. *Limnology and Oceanography* **45**: 1030-  
4 1040.
- 5
- 6 Statzner B, Peltret O and Tomanova S. (2003) Crayfish as geomorphic agents and  
7 ecosystem engineers: effect of a biomass gradient on baseflow and flood-induced  
8 transport of gravel and sand in experimental streams. *Freshwater Biology* **48**: 147-  
9 163.
- 10
- 11 Statzner B and Sagnes P. (2008) Crayfish and fish as bioturbators of streambed  
12 sediments: Assessing joint effects of species with different mechanistic abilities.  
13 *Geomorphology* **93**: 267-287.
- 14
- 15 Styrrishave B, Bojsen BH, Witthoff H and Andersen O. (2007) Diurnal variations in  
16 physiology and behaviour of the noble crayfish *Astacus astacus* and the signal  
17 crayfish *Pacifastacus leniusculus*. *Marine and Freshwater Behaviour and Physiology*  
18 **40**: 63-77.
- 19
- 20 Trimble SW. (1983) A sediment budget for Coon Creek Basin in the Driftless area,  
21 Wisconsin, 1853-1977. *American Journal of Science* **283**: 454-474.
- 22
- 23 Usio N (2000) Effects of crayfish on leaf processing and invertebrate colonisation of  
24 leaves in a headwater stream: decoupling of a trophic cascade. *Oecologia* **124**: 608-  
25 617.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

Walling DE. (1977) Limitation of the rating curve technique for estimating suspended sediment loads, with particular reference to British rivers. In: Erosion and Solid Matter Transport in Inland Waters (Proc. Paris Symp., July 1977), IAHS Publ. no. 122, 34-48.


Walling DE and Collins AL. (2008) The catchment sediment budget as a management tool. *Environmental Science and Policy* **11**: 136-143.

Wharton G, Cotton JA, Wotton RS, Bass JAB, Heppell CM, Trimmer M, Sanders IA and Warren L. (2006) Macrophytes and suspension-feeding invertebrates modify flows and fine sediments in the Frome and Piddle catchments, Dorset (UK). *Journal of Hydrology* **330**: 171-184.


Williams GP. (1989) Sediment concentration versus water discharge during single hydrologic events in rivers. *Journal of Hydrology* **111**: 89-106.

Wright JP, Jones CG and Flecker AS. (2002) An ecosystem engineer, the beaver, increases species richness at the landscape scale. *Oecologia* **132**: 96-101.


1 **FIGURES**




2  
3 Figure 1 Experimental design for laboratory mesocosm experiments and field study:  
4 (a) mesocosm tanks with artificial clay banks; (b) mesocosm tanks with artificial clay  
5 substrate; (c) control and treatment tanks; and (d) photography of monitoring location  
6 at the field site on the River Windrush with diagram to show the equipment set-up.  
7




8  
9 Figure 2 Map showing the location of the field study site on the River Windrush,  
10 Oxfordshire, UK.  
11  
12




1  
 2 Figure 3 Time series of 15-minute discharge data for the River Windrush at  
 3 Newbridge during the monitoring period, showing long-term flow percentiles.  
 4 Source: Environment Agency.  
 5  
 6  
 7




8  
 9 Figure 4 Suspended sediment concentration time series for laboratory mesocosm  
 10 experiments, showing each of the three runs for (a) bank treatments; and (b)  
 11 substrate treatments.




1  
 2 Figure 5 Photographs illustrating key crayfish interactions with bank and bed material  
 3 observed during the laboratory mesocosm experiments: (a) crayfish digging a pit in  
 4 the left corner of the tank; (b) crayfish digging a pit in the right corner of the tank, with  
 5 a previously created pit visible in the left corner; (c) burrow visible in the artificial  
 6 bank; (d) partial collapse of bank following extensive burrowing. (e) and (f) show the  
 7 artificial bank treatment before the start of the experiment, and following removal of  
 8 the crayfish at the end of the monitoring period respectively.  
 9



10  
 11 Figure 6 Turbidity time series for: (a) IR40C near-bed; (b) IR100C near-bed; and (c)  
 12 IR40C mid-flow. The dotted line in (b) represents the upper limit of the IR100C  
 13 range – turbidity values which meet the line indicate turbidity levels beyond the range  
 14 of the sensor. All field readings were within range for the IR40C.




1  
 2 Figure 7 High frequency velocity data for the field site on the River Windrush. Box-  
 3 plots show median, interquartile range (boxes), 1.5 times the interquartile range  
 4 (whiskers) and outliers (points) for streamwise (U), cross stream (V) and vertical (W)  
 5 velocities.  
 6  
 7



1  
 2 Figure 8 Characteristics of suspended sediment pulses identified in the laboratory  
 3 time series, showing: (a) number of pulses per hour for bank treatments (mean,  
 4 maximum, minimum; n=3 runs); (b) number of pulses per hour for bank treatments  
 5 (mean, maximum, minimum; n=3 runs); (c) duration and maximum turbidity of  
 6 individual pulses for bank treatments (across the 3 runs); (d) duration and maximum




1 turbidity of individual pulses for bed treatments (across the 3 runs).



2  
3 Figure 9 Characteristics of turbidity pulses identified in the field turbidity time series  
4 for the IR40C near-bed sensor, showing: (a) number of pulses per hour of the  
5 record; (b) duration and (c) maximum turbidity of pulses according to their position in  
6 the record; and (d) the relationship between pulse duration and maximum turbidity.  
7

1



2

3

4

5

6


7

8


9

Figure 10 Event time-plots presented in shape-magnitude clusters. For (a) to (e), each plot represents a ‘shape’ cluster (A – E respectively), and within each plot solid lines represent low magnitude, dashed lines intermediate magnitude and dotted lines high magnitude pulses (identified from the magnitude HCA). For (f), a series of clusters that describe a small number of pulse events are plotted together for brevity, with a separate legend.


(a) IR40 near-bed



(b) IR100 near-bed



(c) IR40 mid-flow



1  
2  
3  
4  
5

Figure 11 Trends in ambient turbidity, following removal of short-term pulsed events for: (a) IR40C near-bed sensor; (b) IR100C near-bed sensor; and (c) IR40C mid-flow sensor. Vertical dotted lines mark the timing of sunset and sunrise.