

Density functional theory study of the oligomerization of carboxylic acids.

Di Tommaso, D; Watson, KL

“The final publication is available at <http://pubs.acs.org/doi/abs/10.1021/jp509100u>”

For additional information about this publication click this link.

<http://qmro.qmul.ac.uk/jspui/handle/123456789/7510>

Information about this research object was correct at the time of download; we occasionally make corrections to records, please therefore check the published record when citing. For more information contact scholarlycommunications@qmul.ac.uk

Density Functional Theory Study of the Oligomerization of Carboxylic Acids

Devis Di Tommaso, and Ken Leo Watson

J. Phys. Chem. A, **Just Accepted Manuscript** • DOI: 10.1021/jp509100u • Publication Date (Web): 30 Oct 2014

Downloaded from <http://pubs.acs.org> on November 10, 2014

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



1
2
3 **Density Functional Theory Study of the Oligomerization of**
4
5
6 **Carboxylic Acids**
7
8

9
10 Devis Di Tommaso* and Ken L. Watson
11

12
13 School of Biological and Chemical Sciences, Queen Mary University of London, 20 Gordon
14
15 Street, London WC1 H0AJ, United Kingdom
16
17

18
19
20
21 30 October 2014
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 Corresponding author: Dr. Devis DI TOMMASO, School of Biological and Chemical Sciences,
52
53 Queen Mary University of London, 20 Gordon Street, London WC1 H0AJ, United Kingdom.
54

55
56 Email: d.ditommaso@qmul.ac.uk. Phone: +44 (0)20 7882 6226
57
58

Abstract

We present a density functional theory [M06-2X/6-31+G(d,p)] study of the structures and free energies of formation of oligomers of four carboxylic acids (formic acid, acetic acid, tetrolic acid and benzoic acid) in water, chloroform and carbon tetrachloride. Solvation effects were treated using the SMD continuum solvation model. The low-lying energy structures of molecular complexes were located by adopting an efficient search procedure to probe the potential energy surfaces of the oligomers of carboxylic acids $(CA)_n$ ($n = 2-6$). The free energies of the isomers of $(CA)_n$ in solution were determined as the sum of the electronic energy, vibrational-rotational-translational gas-phase contribution and solvation free energy. The assessment of the computational protocol adopted in this study with respect to the dimerization of acetic acid, $(AA)_2$, and formic acid, $(FA)_2$, locates new isomers of $(AA)_2$ and $(FA)_2$ and gave dimerization constants in good agreement with the experimental values. The calculation of the self-association of acetic acid, tetrolic acid and benzoic acid shows the following: (i) Classic carboxylic dimers are the most stable isomer of $(CA)_2$ in both the gas phase and solution; (ii) Trimers of carboxylic acid are stable in apolar aprotic solvents; (iii) Molecular clusters consisting of two interacting classic carboxylic dimers $(D+D)_T$ are the most stable type of tetramers but their formation from the self-association of classic carboxylic dimers is highly unfavourable; (iv) For acetic acid and tetrolic acid the reactions $(CA)_2 + 2CA \rightarrow (D+D)_T$ and $(CA)_3 + CA \rightarrow (D+D)_T$ are exoergonic but these aggregation pathways go through unstable clusters that could hinder the formation of tetrameric species; (v) For tetrolic acid the pre-nucleation species that are more likely to form in solution are dimeric and trimeric structures that have encoded structural motifs resembling the α - and β -solid forms of tetrolic acid; (vi) Stable tetramers of benzoic acid could form in carbon tetrachloride from the aggregation of trimers and monomers; (vii) Higher order clusters such as acetic acid pentamers and tetrolic acid hexamers are highly unstable in all solvation environments.

Keywords: Molecular self-association; Free-energy computations; Quantum mechanical continuum solvation models;; Carboxylic oligomers.

1. Introduction

Thermodynamically stable pre-nucleation solvated clusters can have encoded structural motifs resembling the solid state form crystallizing from solution.¹⁻³ Consequently, the determination of the structure and stability of molecular complexes is of importance to understand and predict the process of polymorph selection during crystallization from solution, that is, the ability of a molecule to crystallize in one specific solid state form depending on the solution composition (nature of the solvent and/or presence of additives in the solution).⁴⁻⁵ However, in order to determine which molecular clusters are thermodynamically stable in solution, a method for the quantification of their free energy of formation is required.

Computational studies of the aggregation of organic molecules have largely relied on *unbiased* molecular dynamics, by means of which the structural information of molecular aggregates containing between two (dimer) and five (pentamer) units could be attained.⁵⁻⁸ However, it is not possible using this technique to determine the free energy of formation for molecular clusters in solution. Free energy profiles or free energy surfaces associated with the process of molecular aggregation could be obtained by means of sampling techniques such as umbrella sampling^{9,10} and metadynamics,^{11,12} but the application of these methods require the identification of a suitable set of collective variables capable of describing the mechanism of self-association of the solute molecules.^{9,13,14} However, even for simple aromatic molecules such as benzoic acid, which could interact during the process of self-association *via* OH/O and OH/ π hydrogen (H) bonding as well as π - π interactions, the choice of collective variables is not obvious.

1
2
3 An alternative computational approach that can be adopted for the calculation of free
4 energies of molecules and clusters in solution is based on quantum mechanical continuum
5 solvation models,¹⁵⁻¹⁷ in which the solute is described at a quantum mechanical level (*ab*
6 *initio* or DFT) and the solvent is treated as a polarizable continuum. In particular, the
7 quantum mechanical treatment of molecular associates using modern density functionals
8 specifically developed for non-bonding interactions^{18,19} can give an accurate description of
9 H-bonding and van der Waals forces that are important in weakly interacting systems at an
10 accessible computational cost. Moreover, polarizable continuum models represent a practical
11 approach to simulate the solvation environment and determine the effect of the medium to the
12 structure and stability of solutes in solution. This approach has been extensively applied to
13 model the dimerization of organic molecules,^{15,20-24} but it has been hardly ever used to
14 quantify the thermodynamic stability in solution of higher order molecular clusters such as
15 trimers, tetramers, pentamers or hexamers. In fact, one of the main challenges in modelling
16 the formation of weakly interacting molecular aggregates is the complexity of their potential
17 energy surfaces (PES), which can be characterized by a large number of local minima
18 separated by low-energy saddle points. Consequently, to obtain an accurate evaluation of the
19 thermodynamic stability of molecular oligomers in solution, the most stable isomers on the
20 PES of these species should be located.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 Here we report an approach to calculate the low-lying energy structures and free
45 energies in solution of molecular clusters, where the solute is described at the DFT [M06-
46 2X/6-31+G(d,p)] level of theory,^{18,25} the solvent is treated using the SMD continuum
47 solvation model,²⁶ and an efficient approach is employed to locate the low-lying energy
48 structures on the PES of molecular complexes. We have applied this method to a selected set
49 of molecular systems characterized by a systematic increase in the complexity of the
50
51
52
53
54
55
56
57
58
59
60

1
2
3 molecular structure and inter-molecular interactions: formic acid, acetic acid, tetrolic acid and
4
5 benzoic acid (see Table 1). We have chosen formic acid and acetic acid because previous
6
7 theoretical^{13,23,24,27} and experimental²⁸ investigations of the dimerization of these two
8
9 molecules can be used as a benchmark to verify the accuracy of the methodology employed
10
11 in the present work. Tetrolic acid is an example of a molecule that crystallizes in different
12
13 polymorphic forms, and that exhibits selective crystallization depending on the solvent: the α -
14
15 form, based on a classic carboxylic motif, crystallizes from apolar aprotic solvents (CCl₄ and
16
17 CHCl₃) and the β -form, based on a catameric chain structure, crystallizes from aqueous and
18
19 alcoholic solutions.⁴ Benzoic acid is a model system that can form complex intermolecular
20
21 interactions, i.e. OH/O and OH/ π hydrogen bonds, all of which are important generally in
22
23 polymorph formation.
24
25
26
27

28 29 30 **2. Methods**

31 32 **2.1 Electronic structure calculations**

33
34
35 All calculations were carried out with the NWChem (version 6.3)²⁹ and Gaussian09³⁰ codes
36
37 using the Minnesota 06 global hybrid density functional with 54% HF exchange (M06-2X)
38
39 along with the 6-31+G(d,p) basis set. We have chosen M06-2X because its assessment
40
41 against representative databases showed that this method is one of the most accurate density
42
43 functionals for a combination of main-group thermochemistry, kinetics and noncovalent
44
45 interaction.^{18,25,31} We have used the 6-31+G(d,p) basis set because when evaluating gas-phase
46
47 binding energies the mere addition of diffuse functions on nonhydrogenionic atoms to a
48
49 doubly polarized valence double zeta basis set provides a good compromise between
50
51 accuracy and computational cost.^{18,32} Thermal contributions were calculated using the gas-
52
53 phase harmonic frequencies at the optimised geometries with the temperature and pressure
54
55
56
57

1
2
3 fixed at the values of 298 K and 1 atm, respectively. The frequencies were scaled by a factor
4
5 of 0.97 in order to account for systematic errors in the density functional and for
6
7 anharmonicity.^{15,33} Free energies of solvation were calculated using the SMD solvation
8
9 model²⁶ and the optimized gas-phase geometries. The SMD model was originally tested over
10
11 a set of 2892 solvation free energies and transfer free energies for neutral solutes and ions in
12
13 water and non-aqueous solutions, and the mean unsigned error (MUE) over 26 combinations
14
15 of various basis sets and density functionals was 0.8 kcal mol⁻¹ for neutral solutes, and 4.3
16
17 kcal mol⁻¹ for ions. In particular, the MUE in calculated solvation free energies of carboxylic
18
19 acids was only 0.25 kcal mol⁻¹ in water and 0.55 kcal mol⁻¹ in organic solvents using the
20
21 M06-2X/6-31+G(d,p) level of theory.²⁶ The SMD model together with the M06-2X density
22
23 functional was also applied to predict the free energies of aqueous solvation for 61 drug-like
24
25 molecules in the SAMPL1 test set and the authors reported a MUE of 2.0 kcal mol⁻¹.³⁴ The
26
27 SMD/M06-2X level of theory was also used by Ribeiro and co-workers to compute the
28
29 partition coefficients of nucleobases between chloroform and water with a MUE of 0.8 kcal
30
31 mol⁻¹.¹⁵ We have also verified the re-optimization of the carboxylic acid molecules and
32
33 clusters in the SMD solvation model but the effect on the free energies of association was
34
35 minor. For example, the Boltzmann average values of the trimerization constants of benzoic
36
37 acid in water, chloroform and carbon tetrachloride computed according to the aggregation
38
39 reactions $3\text{BA} \rightarrow (\text{BA})_3$ and $(\text{BA})_2 + \text{BA} \rightarrow (\text{BA})_3$ changed by only 0.1 pK units when the
40
41 structures of the monomer, dimers and trimers were optimized in the solution.
42
43
44
45
46
47
48

49 **2.2 Free energies of association in solution**

50
51
52 The free energy of formation of carboxylic acid clusters were computed according to the
53
54 following equation:
55
56
57
58
59
60

$$\Delta G_{ass}^* = G_{AB}^* - G_A^* - G_B^* \quad (1)$$

where G_X^* is the total Gibbs free energy of the species X ($X = AB, A$ or B) in the liquid at 298 K, which was determined (see equation 2) by the addition of

$$G_X^* = E_{e,gas} + \delta G_{VRT,gas}^{\circ} + \Delta G_{solv}^* + RT \ln[\tilde{R}T] \quad (2)$$

the gas-phase total electronic energy of the solute ($E_{e,gas}$), the vibrational-rotational-translational contribution to the to the gas-phase Gibbs free energy ($\delta G_{VRT,gas}^{\circ}$) at $T = 298$ K under a standard-state partial pressure of 1 atm, the solvation free energy of the solute corresponding to transfer from an ideal gas at a concentration of 1 M to an ideal solution at a liquid-phase concentration of 1 M (ΔG_{solv}^*), and the free energy change of 1 mol of an ideal gas from 1 atm to 1 mol L⁻¹ ($RT \ln[\tilde{R}T] = 1.89$ kcal mol⁻¹ at 298.15 K where $\tilde{R} = 0.082052$ K⁻¹).^{15,35,36} For multiple stationary points of a molecular cluster the free energy was determined as the Boltzmann ensemble average

$$\langle G(X) \rangle = \sum_{i=1}^N f_i G(X_i) \quad (3)$$

where f_i is the Boltzmann factor corresponding to the i^{th} isomer, $G(X_i)$ is the corresponding free energy and N is the number of low-lying energy structures located on the PES of X . The Boltzmann factor was determined according to

$$f_i = \frac{e^{-G(X_i)/RT}}{\sum_j e^{-G(X_j)/RT}} \quad (4)$$

1
2
3 where R is the universal gas constant, T is the absolute temperature ($T = 298$ K) and the index
4
5 j runs over all isomers.
6
7

8 9 2.3 Localisation of stationary points on the PES of carboxylic acid clusters

10
11
12 The potential energy surfaces of weakly interacting carboxylic acid complexes, $(CA)_n$, can be
13
14 characterized by a very large number of minima. However, according to eqns. 3 and 4 only
15
16 the low-lying energy structures of $(CA)_n$ will have a significant contribution to the Boltzmann
17
18 average value of the Gibbs free energy. To locate the most stable isomers of a molecular
19
20 cluster we have adopted the following computational strategy:
21
22

- 23
24 i. For each carboxylic acid cluster $(CA)_n$ we generated several thousands of candidate
25
26 structures using the Granada software,³⁷ which has been designed to randomly
27
28 distribute one or more molecules around a central unit (a monomer, dimer, trimer etc.)
29
30 placed at the centre of a cube of defined side length, and extensively applied to study
31
32 the microhydration and noncovalent interaction of molecular complexes.³⁸⁻⁴³ For
33
34 example, to generate candidate structures for the trimer $(CA)_3$, we considered the
35
36 random distribution of one CA molecule around the most stable carboxylic acid dimer
37
38 (the cyclic dimeric structure), and from the random distribution of two CA molecules
39
40 around a central carboxylic acid molecule. To sample the potential energy surface of
41
42 the homologous clusters of the smaller carboxylic acids, formic acid and acetic acid,
43
44 we simply generated a very large number of configurations (between 7500 and
45
46 10000). For tetrolic acid and benzoic acid instead, we initially generated, using the
47
48 Granada code, 80'000 configurations in a cube of approximately 10 Å, and then
49
50 selected only those configurations such that the minimum distance between the
51
52 central molecular unit (a monomer, dimer, trimer, etc.) and the randomly distributed
53
54
55
56
57
58
59
60

1
2
3 molecule(s) was less than 4.0 Å. To increase the sampling we rounded in excess the
4
5 number of configurations by gradually increasing the cutoff distance used. The
6
7 molecular units used to generate the candidate structures were fully optimized at the
8
9 M06-2X/6-31+G(d,p) level of theory.

- 10
11
12 ii. The energies of the candidate structures were evaluated at the M06-2X/6-31+G(d,p)
13
14 level of theory using the electronic structure code NWChem and the Boltzmann factor
15
16 f_i corresponding to the i^{th} configuration was determined as

$$f_i = \frac{e^{-(E_i - E_0)/RT}}{\sum_j e^{-(E_j - E_0)/RT}} \quad (5)$$

17
18
19
20
21
22
23
24
25 where E_i was the energy of the i^{th} candidate structure and E_0 was the energy of the
26
27 most stable candidate structure.

- 28
29
30 iii. We considered the candidate structures with a Boltzmann factor $f_i \geq 0.01$ and, to
31
32 increase our sampling, ten to fifteen randomly selected structure such that $3 \leq E_i -$
33
34 $E_0 \leq 15 \text{ kcal mol}^{-1}$. These systems were subject to full geometry optimization using
35
36 the Gaussian 09 code. The thermochemical properties and solvation energies of the
37
38 optimized gas-phase geometries were also evaluated using Gaussian 09.
39
40
41

42
43 The gas phase energies, thermal corrections to the Gibbs free energy, energies in the
44
45 solvation model (including the SMD non-electrostatic terms), and the geometries of the 772
46
47 low-lying energy structures of the carboxylic acids clusters located in this study are listed in
48
49 Supporting Information.

3 Results and Discussion

3.1 Assessment of the method

In this section we report the structures and energetics of formation of the dimers of acetic acid, (AA)₂, and formic acid, (FA)₂, in both polar (water) and apolar (chloroform and carbon tetrachloride) solvents. The values of absolute dimerization constants of (AA)₂ and (FA)₂ have been obtained experimentally,²⁸ and the potential energy surfaces of these two dimers have been subject to extensive theoretical investigation.^{13,23,24,27} The aim of this section is therefore to assess the searching procedure described in section 2.3 to locate the stationary points on the PES of (CA)_n complexes, and the SMD/M06-2X/6-31+G(d,p) level of theory to evaluate the thermodynamic properties in solution of carboxylic acid clusters.

Figure 1 displays the optimized structures and relative energies in the gas-phase of acetic acid dimers. The analysis of the vibrational spectrum of these structures confirmed that they are minima on the PES of (AA)₂. Chochouloušová and co-workers previously applied a molecular dynamics / quenching procedure to investigate the PES of the acetic acid dimer and located six low-lying energy isomers of (AA)₂ (**I** to **V** and **IX** in Figure 1).²³ Using our computational protocol we could locate isomers **I-V** but not **IX**, which structure was taken from reference 23. On the other hand, we have found three new isomers of (AA)₂ (**VI**, **VII** and **VIII** in Figure 1) that are lower in energy than **IX**. The relative energy of **IX** with respect to the global minima **I** is 14 kcal mol⁻¹ and we have not locate this isomer probably because only some of the candidate structures with a relative energy lower than 15 kcal mol⁻¹ were subject to geometry optimization.

The structures and gas-phase energies of the low-lying structures of the formic acid dimer are shown in Figure 2. We located five of the six isomers of formic acid dimers

previously reported by Chochoulošová et al.^{27,44} (**I-V**) but we could not determine isomer **VII** ($\Delta E = 13 \text{ kcal mol}^{-1}$), which structure was taken from reference 27. However, the contribution of **VII** to the Boltzmann average value $\langle G(\text{FA})_2 \rangle$ is negligible. Figure 2 also reports a new isomer of $(\text{FA})_2$ (structure **VI**) that is lower in energy than **VII**. We can therefore conclude that the strategy adopted in this study to probe the potential energy surfaces of molecular clusters is capable of locating the most stable isomers of the oligomers of carboxylic acid.

In Table 2 are compared the values of dimerization constant (pK_D) of acetic acid and formic acid computed at the SMD/M06-2X/6-31+G(d,p) level of theory with those obtained experimentally⁴⁵⁻⁴⁷, and by means of free energy (umbrella sampling) molecular dynamics simulations of the formation of $(\text{AA})_2$ and $(\text{FA})_2$ in explicit water.¹³ In this work the dimerization constants in solution were computed *via* $K_D = -\log[\exp(\Delta G_{ass}^*/RT)]$, where ΔG_{ass}^* is the free energy of dimerization computed according to eqn. 1 using the Boltzmann averaged free energies of $(\text{AA})_2$ and $(\text{FA})_2$. In water both SMD/M06-2X/6-31+G(d,p) and potential mean force (PMF) methods give values of pK_D that are 0.2 – 1.2 units lower than the experiment, and in chloroform the dimerization constant of acetic acid computed at the M06-2X/6-31+G(d,p) level is 0.5 units lower than the experimental value. Using the methodology employed in this work, the uncertainty in the evaluation of the free energy of association (ΔG_{ass}^*) can be written as the sum of

$$\varepsilon(\Delta G_{ass}^*) = \varepsilon(\Delta E_{e, \text{gas}}) + \varepsilon(\Delta \delta G_{\text{VRT}, \text{gas}}) + \varepsilon(\Delta \Delta G_{\text{solv}}^*) \quad (6)$$

the error in the evaluation of the gas phase interaction energy in the gas phase ($\Delta E_{e, \text{gas}}$), the error in the vibrational-rotational-translational contribution to the gas-phase reaction free

energy ($\Delta \delta G_{\text{VRT,gas}}^{\circ}$), and the difference between the error in the evaluation of the solvation free energies of the product and reactants ($\Delta \Delta G_{\text{solv}}^*$). In particular, for the dimerization process the error in the solvation free energy contribution to the reaction free energy is

$$\varepsilon(\Delta \Delta G_{\text{solv}}^*) = \varepsilon[\Delta G_{\text{solv}}^*(\text{D})] - 2\varepsilon[\Delta G_{\text{solv}}^*(\text{M})] \quad (7)$$

where $\varepsilon[\Delta G_{\text{solv}}^*(\text{D})]$ and $\varepsilon[\Delta G_{\text{solv}}^*(\text{M})]$ are the mean unsigned errors in the calculated solvation free energies of carboxylic acids, which, as discussed in Section 2.2, have been found to be 0.25 kcal mol⁻¹ in water and 0.55 kcal mol⁻¹ in organic solvents. Therefore, we expect the uncertainty associated with the SMD/M06-2X/6-31+G(d,p) solvation model to be in the range of 0 – 0.75 kcal mol⁻¹. The differences between the calculated and experimental values of the free energy of dimerization of acetic acid (AA)₂ and formic acid (FA)₂ (0.3 – 1.9 kcal mol⁻¹ in Table 2) are therefore consistent with the errors in the evaluation of the free energies of solvation of the monomers and dimers of formic acid and acetic acid. Table 2 also reports the values of pK_D obtained with the counterpoise correction (CPC) of Boys and Bernardi to account for the basis set superposition error (BSSE)⁴⁸ during geometry optimization. The constants obtained with (“cpc”) and without (“no-cpc”) the counterpoise method differ by less than 0.2 units. As the cost of calculating the CPC is typically greater than that of calculating the original gas-phase interaction energy,⁴⁹ we have not included this term in the evaluation of the gas-phase energy of carboxylic acid clusters. These results indicate that the SMD/M06-2X/6-31+G(d,p) method slightly over-stabilizes the formation of (AA)₂ and (FA)₂ in solution but is still sufficiently accurate to provide a quite accurate prediction of the thermodynamic stability in solution of carboxylic acid clusters.

3.2 Acetic acid

The free energies of formation in the gas-phase (ΔG_{ass}°) and in polar (water) and non-polar (chloroform and carbon tetrachloride) solvents (ΔG_{ass}^*) of acetic acid (AA) clusters consisting of two (dimer), three (trimer), four (tetramer) and five (pentamer) molecular units are listed in Table 3.

The cyclic structure **I**, which is reported in Figure 1, is the most stable dimeric species in both the gas phase and solution. When chloroform and carbon tetrachloride are used the formation of the cyclic dimer is highly exoergonic ($\Delta G_{ass}^* = -3.20$ kcal mol⁻¹ in CHCl₃ and $\Delta G_{ass}^* = -5.87$ kcal mol⁻¹ in CCl₄). In water the free energy of reaction is -0.18 kcal mol⁻¹ but as discussed in section 3.1 our calculation slightly overestimates the interaction between acetic acid molecules. Isomer **VIII** of (AA)₂ could also be stable in carbon tetrachloride ($\Delta G_{ass}^* = -0.33$ kcal) but all other types of dimers have positive values of the free energy of dimerization (see Table 3). Therefore, according to our simulations, and in agreement with previous studies,^{23,24,35} in apolar solvents acetic acid molecules tend to form stable cyclic carboxylic dimers **I**.

To locate the low-lying energy structures of the acetic acid trimer we have generated 7500 candidate (AA)₃ structures by considering the aggregation of three monomers [trimers of type (AA)_{3,(3M)}] and of the cyclic dimer **I** with one acetic acid molecule [trimers of type (AA)_{3,(D+M)}]. The gas-phase energies of these two types of structures, which are reported in Figure 3, are scattered in two different regions and the trimers of type (AA)_{3,(D+M)} are approximately 20 kcal mol⁻¹ lower than the trimers of type (AA)_{3,(3M)}. From the trimeric structures of type (AA)_{3,(D+M)} we have distinguished eight possible low-lying isomers of (AA)₃, which all consist of an acetic acid molecule interacting *via* H-bonding with the

1
2
3 carboxylic or methyl groups of the cyclic dimeric unit (see Figure 4). In Table 3 we have
4
5 listed the energetics of formation of the acetic acid trimers **I-VIII** computed according to the
6
7 trimerization reactions (i) $(AA)_2 + AA \rightarrow (AA)_{3,(D+M)}$ and (ii) $3AA \rightarrow (AA)_{3,(D+M)}$. In
8
9 chloroform and carbon tetrachloride, acetic acid trimers are stable with respect to non-
10
11 associated monomers [reaction (ii)] and the lowest energy isomers (structures **I** and **V** in
12
13 Figure 4) correspond to the favourite aggregation modes for $(AA)_3$ found during molecular
14
15 dynamics simulations of acetic acid in explicit carbon tetrachloride solvent molecules.⁶
16
17 Moreover, the thermodynamic stability of acetic acid trimers in CCl_4 supports the findings of
18
19 Bulmer and Shurvell,⁵⁰ who proposed that species larger than dimers should be present in
20
21 appreciable concentration in solution in order to account for infrared data on the acetic acid –
22
23 carbon tetrachloride system.
24
25
26
27

28
29 For tetramers we located the isomers of $(AA)_4$ by evaluating the energies of 10000
30
31 candidate structures of type $(AA)_{4,(D+D)}$, where D is the cyclic dimer **I** in Figure 1 [structures
32
33 of type $(AA)_{4,(D+D)}$ were found to be considerably lower in energy than those of type
34
35 $(AA)_{4,(Tr+M)}$ and $(AA)_{4,(D+2M)}$]. As the number of units in a molecular cluster increases, it
36
37 becomes more difficult to classify an optimized structure to a specific type of isomer, as
38
39 previously done for the dimeric and trimeric structures of acetic acid. Consequently, the
40
41 energetics of formation of $(AA)_4$ in Table 3 are the Boltzmann average values of the energies
42
43 and free energies of 47 optimized tetramers, which have been computed according to the
44
45 following reactions: (i) $(AA)_2 + (AA)_2 \rightarrow (AA)_{4,(D+D)}$; (ii) $(AA)_3 + AA \rightarrow (AA)_{4,(D+D)}$; $(AA)_2$
46
47 + $2AA \rightarrow (AA)_{4,(D+D)}$. Representative low-lying structures of type $(AA)_{4,(D+D)}$ of acetic acid
48
49 tetramers are reported in Figure 5. The free energy of self-association of two cyclic dimers **I**
50
51 [reaction (i)] is endoergonic ($\Delta G > 0$) in all media but reactions (ii) ($\Delta G_{ass}^* = -1.36 \text{ kcal mol}^{-1}$)
52
53 and (iii) ($\Delta G_{ass}^* = -0.56 \text{ kcal mol}^{-1}$) are exoergonic ($\Delta G < 0$) in carbon tetrachloride, which
54
55
56
57
58
59
60

would suggest that acetic acid tetramers exist in apolar solution. However, the aggregation pathways $(AA)_3 + AA \rightarrow (AA)_{4,(D+D)}$ and $(AA)_2 + 2 AA \rightarrow (AA)_{4,(D+D)}$ do not take into account that in solutions containing monomers, dimers and trimers we could have first the formation of intermediate tetrameric structures of type $(AA)_{4,(D+2M)}$ or $(AA)_{4,(Tr+M)}$, followed by the process of reorganization of the molecules in the clusters to give the lower free energy species $(AA)_{4,(D+D)}$. The free energy differences between the tetramers of type $(AA)_{4,(Tr+M)}$, $(AA)_{4,(D+2M)}$ and $(AA)_{4,(D+D)}$ are reported in Figure 6, where the values were obtained from the Boltzmann average free energies of the isomers of $(AA)_{4,(Tr+M)}$, $(AA)_{4,(D+2M)}$ and $(AA)_{4,(D+D)}$, respectively. Figure 6 shows that the formation of tetramers of type $(AA)_{4,(D+2M)}$ ($\Delta G_{ass}^* = 2.65 \text{ kcal mol}^{-1}$ in CCl_4) and $(AA)_{4,(Tr+M)}$ ($\Delta G_{ass}^* = 1.17 \text{ kcal mol}^{-1}$ in CCl_4) is not favourable, and, consequently, the formation of the tetrameric structures $(AA)_{4,(D+D)}$ could be hindered by the thermodynamic instability of the intermediate species $(AA)_{4,(D+2M)}$ and $(AA)_{4,(Tr+M)}$. Moreover, there could also be an activation barrier associated with the reorganization of acetic acid molecules from the molecular arrangement in the tetramers $(AA)_{4,(D+2M)}$ or $(AA)_{4,(Tr+M)}$ to the arrangement observed in the $(AA)_{4,(D+D)}$ species (see Figure 5).

The energetics of formation of pentamers in Table 3, which representative most stable structures are reported in Figure 7, is highly endoergonic in all solvation environments and this higher order species is unlikely to be found in solution. To probe the PES and locate the low-lying isomers of $(AA)_5$ we evaluated the energies of 10000 candidate structures of type $(AA)_{5,(Tr+D)}$, and the energetics in Table 3 are the Boltzmann average values of the energies of 46 optimized pentameric structures.

3.3 Tetrolic acid

The free energies of formation of dimers, trimers, tetramers and hexamers of tetrolic acid (TTA) are listed in Table 4. We located nine isomers on the PES of the tetrolic acid dimer, which structures are reported in Figure 8. The cyclic structure **I** is the only dimeric species that is stable in chloroform ($\Delta G_{ass}^* = -1.80 \text{ kcal mol}^{-1}$) and carbon tetrachloride ($\Delta G_{ass}^* = -4.62 \text{ kcal mol}^{-1}$) but not in water ($\Delta G_{ass}^* = 0.85 \text{ kcal mol}^{-1}$). Isomer **I** corresponds to the structural synthon found in the polymorph of tetrolic acid that crystallizes from chloroform and carbon tetrachloride (α -TTA).^{4,51} The formation of all other dimers is endoergonic in both the gas-phase and solution but it is interesting to notice that the second most stable isomer of (TTA)₂ (structure **II** in Figure 8) corresponds to the catemeric unit found in the β form of TTA.⁵¹

The energetics of formation of the trimers in Table 4 represent the Boltzmann average of the energies and free energies of 40 low-lying structures, which were selected from 6000 candidate structures of type (3M)_{Tr} and 3000 candidate structures of type (D+M)_{Tr}, where D and M are the optimised geometries of the cyclic dimer **I** and of the tetrolic acid molecule, respectively. The most stable structures of (TTA)₃, which are reported in Figure 9, show that the isomers of the tetrolic acid trimer consist of a cyclic dimer **I** interacting with a tetrolic acid molecule and, similarly to what found for acetic acid, in the most stable isomer of (AA)₃ (structure **I** in Figure 9) the carboxylic acid molecule forms both an O-H...O and an C-H...O bond to the dimer. Notice also that species **I** corresponds to the structural unit found in the β form of TTA.⁵¹

Tetramers of tetrolic acid could form from the aggregation of two cyclic dimer **I**, (TTA)_{4,(D+D)}, a cyclic dimer **I** with two tetrolic acid molecules (TTA)_{4,(D+2M)}, and the trimer **I** with one tetrolic acid molecule (TTA)_{4,(Tr+M)}. However, Figure 10 shows that tetramers of

1
2
3 type $(TTA)_{4,(D+D)}$ are significantly lower in energy than tetramers of type $(TTA)_{4,(D+2M)}$ and
4
5 $(TTA)_{4,(Tr+M)}$. Some representative $(TTA)_{4,(D+D)}$ structures are reported in Figure 11 where we
6
7 have highlighted some of the H-bonding between the cyclic dimeric units that stabilize the
8
9 clusters. The energetics of formation of $(TTA)_{4,(D+D)}$ in Table 4, which have been obtained
10
11 from the Boltzmann average of the energies of 70 minima on the PES of $(TTA)_{4,(D+D)}$,
12
13 indicate that the self-association of tetrolic acid dimers [reaction $(TTA)_2 + (TTA)_2 \rightarrow (TTA)_4$
14
15 in Table 4] is endoergonic, but the formation of tetramers could occur in carbon tetrachloride
16
17 only along the aggregation pathway $(TTA)_3 + (TTA) \rightarrow (TTA)_4$ ($\Delta G_{ass}^* = -1.35 \text{ kcal mol}^{-1}$).
18
19 However, this reaction does not take into account the energetics required for the formation of
20
21 intermediate clusters of type $(TTA)_{4,(Tr+M)}$, which could form when a trimer **I** interacts with a
22
23 tetrolic acid molecule. In Figure 12 the free energy profiles along the reaction pathways (i)
24
25 $(TTA)_3 + (TTA) \rightarrow (TTA)_4$ and (ii) $(TTA)_2 + 2 (TTA) \rightarrow (TTA)_4$ indicate that species of
26
27 type $(TTA)_{4,(Tr+M)}$ and $(TTA)_{4,(D+2M)}$ are unstable in solution and they could therefore hinder
28
29 the formation of the thermodynamically stable $(TTA)_{4,(D+D)}$ tetramers.
30
31
32
33
34
35

36 Table 4 also reports the Boltzmann average values of the energetics of formation of
37
38 hexamers, which low lying energy structures have been located by considering 8000
39
40 structures of type $(TTA)_{6,(3D)}$, where D was the cyclic dimer **I**. The values in Table 4 indicate
41
42 that the formation of $(TTA)_6$ species is highly endoergonic in all solvation environments.
43
44
45

46 Based on our simulations we therefore suggest that in homogeneous tetrolic acid
47
48 aqueous solutions the monomers of TTA dominate, whereas in apolar aprotic solvents, like
49
50 carbon tetrachloride or chloroform, tetrolic acid molecules form stable dimeric (**I** in Figure 8)
51
52 and trimeric (**I** in Figure 9) species that have structural features resembling the α and β
53
54 polymorphic forms of TTA.
55
56
57
58
59
60

3.4 Benzoic acid

The energetics of formation of the dimers, trimers and tetramers of benzoic acid (BA) in the gas-phase and in solution are reported in Table 5. Molecules of benzoic acid have the ability to form H-bonds, π - π and OH/ π interactions and therefore it would have been difficult to locate the most stable isomers of $(BA)_n$ clusters only by means of chemical intuition.

Using the computational protocol adopted in this work we located eighteen isomers on the PES of $(BA)_2$ and the structure of the six most stable benzoic acid dimers are displayed in Figure 13. The energetics of the dimerization reaction indicate that in chloroform ($\Delta G_{ass}^* = -2.97 \text{ kcal mol}^{-1}$) and carbon tetrachloride ($\Delta G_{ass}^* = -7.77 \text{ kcal mol}^{-1}$) the classic carboxylic structure **I** is thermodynamically very stable. Isomers **III** ($\Delta G_{ass}^* = -0.75 \text{ kcal mol}^{-1}$) and **IV** ($\Delta G_{ass}^* = -0.56 \text{ kcal mol}^{-1}$), which consist of two molecules interacting *via* a single H-bonding, could also form in carbon tetrachloride. For all other dimeric structures, where the benzoic acid molecules interact *via* weaker π - π and OH/ π interactions (**II**, **V** and **VI** in Figure 13), the energetics of dimerization is positive in both the gas phase and solution (see Table 5).

To locate the low-lying energy structures of benzoic acid trimers we generated and computed the energies of 10000 candidate structures of type $(3M)_{Tr}$ and $(D+M)_{Tr}$. The energetics of trimerization reaction in Table 5 represents the Boltzmann average values of 36 low-lying $(BA)_3$ isomers. The aggregation of the cyclic dimer **I** with a benzoic acid molecule $[(BA)_2 + BA \rightarrow (BA)_3]$ is not thermodynamically favourable, but in carbon tetrachloride the free energy change for the tetramerization reaction $3(BA) \rightarrow (BA)_3$ is highly exoergonic ($-5.20 \text{ kcal mol}^{-1}$). This result agrees with experiments of apolar (benzene) solutions of benzoic acids by Goralski and Taniewska, who found that the presence of monomers, dimers as well

1
2
3 as trimers were required to account for osmometric data.⁵² We have verified the stability of
4
5 benzoic acid trimers in benzene by computing the energetics of the benzoic acid – benzene
6
7 system, and the resulting Boltzmann average free energy of trimerization reaction $3(\text{BA}) \rightarrow$
8
9 $(\text{BA})_3$ is negative ($-1.94 \text{ kcal mol}^{-1}$). On the other hand, recent density functional theory
10
11 [PCM/B3P86/6-31G(d,p)] calculations of the trimerization of benzoic acid in benzene did not
12
13 support the formation of substantial amount of the trimeric phase.⁵³ However, in this study
14
15 the authors considered a single $(\text{BA})_3$ structure, which gas-phase stabilization energy ($\Delta E_{e,gas}$
16
17 $= -22.58 \text{ kcal mol}^{-1}$)⁵³ is lower than the Boltzmann average value reported in Table 5 ($\Delta E_{e,gas}$
18
19 $= -28.20 \text{ kcal mol}^{-1}$). In fact, several $(\text{BA})_3$ isomers located in this work are significantly
20
21 more stable than the structure reported by Pham and co-workers,⁵³ where three benzoic
22
23 molecules interacting *via* the three carboxylic groups, such as isomer **V** in Figure 14, which
24
25 consist of a benzoic acid molecule interacting *via* H-bonding as well as π - π interaction with
26
27 the cyclic dimer **I**. We would like also to mention that using our searching technique we
28
29 determined several isomers with a gas-phase stabilization energy in the range $[-15.6, -23.9]$
30
31 kcal mol^{-1} [the structures and energetics of the trimers of type $(\text{BA})_{3,(D+M)}$ and $(\text{BA})_{3,(3M)}$
32
33 reported in Supporting Information]. Moreover, it is also important to stress that when using
34
35 DFT methods to compute relative energies of weakly bounded complexes such as $(\text{CA})_n$ the
36
37 inclusion of diffuse functions on non-hydrogenic atoms is crucial,^{32,54} and the 6-31G(d,p)
38
39 basis set employed by Pham and co-workers to evaluate the gas-phase reaction energies could
40
41 have not provided a sufficient accuracy for the evaluation of the energetics of dimerization
42
43 and trimerization reactions of benzoic acid.
44
45
46
47
48
49

50
51 For the tetramers we determined the structure and free energies of 22 $(\text{BA})_4$ low-lying
52
53 energy structures, which were selected from 5000 candidate structures of type $(\text{BA})_{4,(D+D)}$,
54
55 where D is the benzoic acid cyclic dimer **I** in Figure 13. The isomers of $(\text{BA})_4$ consist of two
56
57
58
59
60

1
2
3 dimeric units arranged in a stacked configuration, which allows the formation of both H-
4
5 bonding and/or π - π interactions between the molecules of benzoic acid (see Figure 15). In
6
7 chloroform and carbon tetrachloride, where benzoic acid molecules form stable classic
8
9 carboxylic species, the aggregation of two dimers $[(BA)_2 + (BA)_2 \rightarrow (BA)_{4,(D+D)}]$ is
10
11 endoergonic ($\Delta G_{ass,CHCl_3}^\circ = 1.65 \text{ kcal mol}^{-1}$ and $\Delta G_{ass,CCl_4}^\circ = 3.04 \text{ kcal mol}^{-1}$), but according to
12
13 the values reported in Table 5, the $(BA)_{4,(D+D)}$ tetramers could form along the reaction
14
15 pathways $(BA)_2 + 2 BA \rightarrow (BA)_4$ and $(BA)_3 + BA \rightarrow (BA)_4$. Figure 16 reports the relative
16
17 free energies of formation of the molecular aggregates $(BA)_{4,(D+D)}$, $(BA)_{4,(D+2M)}$ and
18
19 $(BA)_{4,(Tr+2M)}$, where the values have been obtained from the Boltzmann average of the free
20
21 energies of the molecular clusters of type $(BA)_{4,(D+D)}$ (tetramers consisting of two benzoic acid
22
23 cyclic dimer **I** units), $(BA)_{4,(Tr+2M)}$ (tetramers containing the most stable trimer in CCl_4 ,
24
25 isomer **I** in Figure 14), and $(BA)_{4,(D+2M)}$. Structures of type $(BA)_{4,(D+D)}$ represent the lowest
26
27 energy molecular aggregates but stable intermediate clusters of type $(BA)_{4,(Tr+2M)}$ could also
28
29 form from the aggregation of trimers and monomers $[(BA)_3 + BA \rightarrow (BA)_4]$ in apolar
30
31 solvents. On the other hand, the formation of the tetramers $(BA)_{4,(D+D)}$ along the reaction
32
33 pathway $(BA)_2 + 2 BA \rightarrow (BA)_4$ is thermodynamically hindered by the instability of species
34
35 of type $(BA)_{4,(D+2M)}$.
36
37
38
39
40
41
42
43
44

45 **4. Summary and Conclusions**

46
47 Density functional theory [M06-2X/6-31+G(d,p)] continuum solvation (SMD) calculations
48
49 were used to determine the structure, thermodynamic stability and free energy pathways for
50
51 the formation of carboxylic acid clusters: formic acid $(FA)_n$ ($n = 2$), acetic acid $(AA)_n$ ($n = 2$ -
52
53 5), tetrolic acid $(TTA)_n$ ($n = 2$ -6) and benzoic acid $(BA)_n$ ($n = 2$ -4). The low-lying energy
54
55 structures of weakly interacting molecular complexes were located by adopting a search
56
57
58
59
60

1
2
3 procedure to probe the potential energy surface of $(CA)_n$. The free energies of the low-lying
4 energy isomers of $(CA)_n$ in solution were determined as the sum of the electronic energy,
5 vibrational-rotational-translational gas-phase contribution and solvation free energy. Based
6
7
8
9 on the results we conclude the following:

- 10
11
12
- 13 i. The application of the computational approach adopted in the present study to
14 compute the dimerization of acetic acid and formic acid resulted in the location of
15 three new low lying isomers of $(AA)_2$ and one new isomer of $(FA)_2$. The Boltzmann
16 averaged dimerization constants (pK_D) of $(AA)_2$ and $(FA)_2$ were in good agreement
17 with the values of pK_D obtained experimentally.
18
19
 - 20 ii. The most stable isomer of the carboxylic acid dimer, in both the gas phase and
21 solution, is the classic carboxylic species, which formation is endoergonic in apolar
22 aprotic solvents like chloroform and carbon tetrachloride.
23
24
 - 25 iii. For all carboxylic acids we found that the formation of trimers according to reaction
26 $3CA \rightarrow (CA)_3$ is thermodynamically spontaneous in chloroform and carbon
27 tetrachloride. The stability of $(CA)_3$ species in solution support experimental data of
28 acetic acid – carbon tetrachloride⁵⁰ and benzoic acid – benzene benzoic⁵² systems,
29 which suggested that trimers are present in appreciable concentrations in apolar
30 aprotic solvents.
31
32
 - 33 iv. The structure of the most stable trimer of tetrolic acid consist of the cyclic dimeric
34 unit, corresponding to the structural synthon found in the α polymorph of tetrolic acid,
35 which forms both an O-H \cdots O and an C-H \cdots O bond to a tetrolic acid molecule and that
36 corresponds to the catameric unit found in the β -TTA.²⁸
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 v. For all carboxylic acid, the most stable tetrameric structures consist of two interacting
4 cyclic dimers, $(CA)_{4,(D+D)}$, but the self-association reaction $(CA)_2 + (CA)_2 \rightarrow$
5 $(CA)_{4,(D+D)}$ is endoergonic in all solvation environments.
6
7
8
9
10 vi. Tetramers of type $(CA)_{4,(D+D)}$ could form in carbon tetrachloride and chloroform along
11 the reaction pathways $(CA)_2 + 2 (CA) \rightarrow (CA)_4$ and $(CA)_3 + CA \rightarrow (CA)_4$. For acetic
12 acid and tetrolic acid these aggregation reactions involve the formation of
13 intermediate clusters of type $(CA)_{4,(D+2M)}$ and $(CA)_{4,(Tr+M)}$ that are unstable in solution.
14 For benzoic acid tetrameric structures could also form from the aggregation of trimers
15 and monomers in carbon tetrachloride and chloroform $[(BA)_3 + BA \rightarrow (BA)_4]$.
16
17
18
19
20
21
22
23
24

25 The free energy pathways for the formation of the dimers, trimers and tetramers of acetic
26 acid, tetrolic acid and benzoic acid as computed at the SMD/M06-2X/6-31+G(d,p) level of
27 theory are summarized in Figure 17.
28
29
30
31

- 32
33 vii. Higher order clusters such as acetic acid pentamers and tetrolic acid hexamers are
34 highly unstable in all solvation environments.
35
36
37

38 In conclusion, this work proposes the application of a computational approach for the
39 determination of the structure and stability in solution of molecular clusters. We conducted
40 all calculations using an accurate, but computationally demanding, hybrid density functional
41 method (M06-2X). As all calculation of the thermodynamic properties of weakly interacting
42 carboxylic acid molecular clusters should consider the location of the low-lying isomers other
43 than the deepest minimum structures, the application of this method to large and/or
44 conformationally flexible molecules would require prohibiting computational resources.
45 However, a hybrid methodology could be adopted where the gas-phase contributions to the
46 free energy ($E_{e,gas} + \delta G_{VRT,gas}$ in eqn. 2) is computed using a forcefield model, which
47
48
49
50
51
52
53
54
55
56
57
58
59
60

accuracy could be assessed with respect to the structures and energetics reported in this study, and the solvation free energy term is evaluated using the accurate SMD model (ΔG_{solv}^* in eqn. 3).

Acknowledgements

D.D.T. thanks the U.K.'s Royal Society for the award of a Royal Society Industry Fellowship. This research utilised Queen Mary's MidPlus computational facilities, supported by QMUL Research-IT and funded by EPSRC grant EP/K000128/1. Via our membership of the UK's HPC Materials Chemistry Consortium, which is funded by EPSRC (EP/L000202), this work made use of the facilities of HECToR and ARCHER the UK's national high-performance computing service, which is funded by the Office of Science and Technology through EPSRC's High End Computing Programme. The authors would like to thank Prof. Luis A. Montero-Cabrera and Dr. Cercis Morera-Boado (Universidad de La Habana, Havana) for providing us with a copy of the Granada code.

Supporting Information Available

The gas phase energies, thermal corrections to the Gibbs free energy, energies in the solvation model (including the SMD non-electrostatic terms), and the geometries of the 772 low-lying energy structures of the carboxylic acids clusters located in this study are listed in Supporting Information. This information is free of charge via the Internet at

<http://pubs.acs.org>

References

- (1) Gebauer, D.; Kellermeier, M.; Gale, J. D.; Bergstrom, L.; Colfen, H. Pre-nucleation Clusters as Solute Precursors in Crystallisation. *Chem. Soc. Rev.* **2014**, *43*, 2348-2371

- 1
2
3 (2) Davey, R. J.; Blagden, N.; Righini, S.; Alison, H.; Quayle, M. J.; Fuller, S. Crystal
4 Polymorphism as a Probe for Molecular Self-assembly During Nucleation from Solutions:
5 The Case of 2,6-dihydroxybenzoic Acid. *Cryst Growth Des* **2001**, *1*, 59-65.
6
7 (3) Davey, R. J.; Schroeder, S. L. M.; ter Horst, J. H. Nucleation of Organic Crystals - A
8 Molecular Perspective. *Angew. Chem. Int. Edit.* **2013**, *52*, 2-16.
9
10 (4) Parveen, S.; Davey, R. J.; Dent, G.; Pritchard, R. G. Linking Solution Chemistry to
11 Crystal Nucleation: The Case of Tetrolic acid. *Chem. Commun.* **2005**, 1531-1533.
12
13 (5) Hamad, S.; Moon, C.; Catlow, C. R. A.; Hulme, A. T.; Price, S. L. Kinetic Insights
14 into the Role of the Solvent in the Polymorphism of 5-fluorouracil from Molecular Dynamics
15 Simulations. *J. Phys. Chem. B* **2006**, *110*, 3323-3329.
16
17 (6) Gavezzotti, A. Molecular Aggregation of Acetic Acid in a Carbon Tetrachloride
18 Solution: A Molecular Dynamics Study with a View to Crystal Nucleation. *Chem. Eur. J.*
19 **1999**, *5*, 567-576.
20
21 (7) Hamad, S.; Hughes, C. E.; Catlow, C. R. A.; Harris, K. D. M. Clustering of Glycine
22 Molecules in Aqueous Solution Studied by Molecular Dynamics Simulation. *J. Phys. Chem.*
23 *B* **2008**, *112*, 7280-7288.
24
25 (8) Gavezzotti, A.; Filippini, G.; Kroon, J.; vanEijck, B. P.; Klewinghaus, P., The Crystal
26 Polymorphism of Tetrolic Acid: A Molecular Dynamics Study of Precursors in Solution, and
27 a Crystal Structure Generation. *Chem. Eur. J.* **1997**, *3*, 893-899.
28
29 (9) Chen, J.; Trout, B. L., Computational Study of Solvent Effects on the Molecular Self-
30 assembly of Tetrolic Acid in Solution and Implications for the Polymorph Formed from
31 Crystallization. *J. Phys. Chem. B* **2008**, *112*, 7794-7802.
32
33 (10) Kästner, J., Umbrella Sampling. *Wiley Interdisciplinary Reviews: Computational*
34 *Molecular Science* **2011**, *1*, 932-942.
35
36 (11) Laio, A.; Parrinello M. Escaping Free-energy Minima. *Proc. Natl. Acad. Sci. USA*
37 **2002**, *99*, 12562-12566.
38
39 (12) Laio, A.; Gervasio, F. L. Metadynamics: a Method to Simulate Rare Events and
40 Reconstruct the Free Energy in Biophysics, Chemistry and Material Science. *Rep. Prog.*
41 *Phys.* **2008**, *71*, 126601.
42
43 (13) Chen, J. H.; Brooks, C. L.; Scheraga, H. A. Revisiting the Carboxylic Acid Dimers in
44 Aqueous Solution: Interplay of Hydrogen Bonding, Hydrophobic Interactions, and Entropy.
45 *J. Phys. Chem. B* **2008**, *112*, 242-249.
46
47 (14) Salvalaglio, M.; Vetter, T.; Giberti, F.; Mazzotti, M.; Parrinello, M. Uncovering
48 Molecular Details of Urea Crystal Growth in the Presence of Additives. *J. Am. Chem. Soc.*
49 **2012**, *134*, 17221-17233.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 (15) Ribeiro, R. F.; Marenich, A. V.; Cramer, C. J.; Truhlar, D. G., The Solvation,
4 Partitioning, Hydrogen Bonding, and Dimerization of Nucleotide Bases: a Multifaceted
5 Challenge for Quantum Chemistry. *Phys. Chem. Chem. Phys.* **2011**, *13*, 10908-10922.
6
7 (16) Tomasi, J.; Mennucci, B.; Cammi, R. Quantum Mechanical Continuum Solvation
8 Models. *Chem. Rev.* **2005**, *105*, 2999-3093.
9
10 (17) Cramer, C. J.; Truhlar, D. G., Implicit Solvation Models: Equilibria, Structure,
11 Spectra, and Dynamics. *Chem. Rev.* **1999**, *99*, 2161-2200.
12
13 (18) Zhao, Y.; Truhlar, D. G. The M06 Suite of Density Functionals for Main Group
14 Thermochemistry, Thermochemical Kinetics, Noncovalent Interactions, Excited States, and
15 Transition Elements: Two New Functionals and Systematic Testing of Four M06-class
16 Functionals and 12 Other Functionals. *Theoretical Chemistry Accounts* **2008**, *120*, 215-241.
17
18 (19) Klimeš, J.; Michaelides, A., Perspective: Advances and Challenges in Treating van
19 der Waals Dispersion Forces in Density Functional Theory. *J Chem Phys* **2012**, *137*, 12.
20
21 (20) Friant-Michel, P.; Ruiz-Lopez, M. F., Glycine Dimers: Structure, Stability, and
22 Medium Effects. *ChemPhysChem* **2010**, *11*, 3499-3504.
23
24 (21) Ramondo, F.; Bencivenni, L.; Caminiti, R.; Pieretti, A.; Gontrani, L. Dimerisation of
25 Urea in Water Solution: a Quantum Mechanical Investigation. *Phys. Chem. Chem. Phys.*
26 **2007**, *9*, 2206-2215.
27
28 (22) Di Tommaso, D. The Molecular Self-association of Carboxylic Acids in Solution:
29 Testing the Validity of the Link Hypothesis Using a Quantum Mechanical Continuum
30 Solvation Approach. *CrystEngComm* **2013**, *15*, 6564-6577.
31
32 (23) Chocholousova, J.; Vacek, J.; Hobza, P. Acetic Acid Dimer in the Gas Phase,
33 Nonpolar Solvent, Microhydrated Environment, and Dilute and Concentrated Acetic Acid:
34 Ab Initio Quantum Chemical and Molecular Dynamics Simulations. *J. Phys. Chem. A* **2003**,
35 *107*, 3086-3092.
36
37 (24) Pasalic, H.; Aquino, A. J. A.; Tunega, D.; Haberhauer, G.; Gerzabek, M. H.; Georg,
38 H. C.; Moraes, T. F.; Coutinho, K.; Canuto, S.; Lischka, H. Thermodynamic Stability of
39 Hydrogen-Bonded Systems in Polar and Nonpolar Environments. *J. Comput. Chem.* **2010**,
40 *31*, 2046-2055.
41
42 (25) Zhao, Y.; Truhlar, D. G., Exploring the Limit of Accuracy of the Global Hybrid Meta
43 Density Functional for Main-Group Thermochemistry, Kinetics, and Noncovalent
44 Interactions. *J. Chem. Theory Comput.* **2008**, *4*, 1849-1868.
45
46 (26) Marenich, A. V.; Cramer, C. J.; Truhlar, D. G. Universal Solvation Model Based on
47 Solute Electron Density and on a Continuum Model of the Solvent Defined by the Bulk
48 Dielectric Constant and Atomic Surface Tensions. *J. Phys. Chem. B* **2009**, *113*, 6378-6396.
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 (27) Chocholousova, J.; Vacek, J.; Hobza, P. Potential Energy and Free Energy Surfaces of
4 the Formic Acid Dimer: Correlated Ab Initio Calculations and Molecular Dynamics
5 Simulations. *Phys. Chem. Chem. Phys.* **2002**, *4*, 2119-2122.
6
7 (28) Schrier, E. E.; Robinson, R. A. Study of Free Energy Relationships in Some Amino
8 Acid-Sodium Chloride-Water Systems. *J. Biol. Chem.* **1971**, *246*, 2870-2874.
9
10 (29) Valiev, M.; Bylaska, E. J.; Govind, N.; Kowalski, K.; Straatsma, T. P.; Van Dam, H.
11 J. J.; Wang, D.; Nieplocha, J.; Apra, E.; Windus, T. L. et al. NWChem: A Comprehensive
12 and Scalable Open-source Solution for Large Scale Molecular Simulations. *Comput. Phys.*
13 *Commun.* **2010**, *181*, 1477-1489.
14
15 (30) Gaussian 09, Revision C.01, Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria,
16 G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G.
17 A. et al. Gaussian, Inc., Wallingford CT, **2009**.
18
19 (31) Riley, K. E.; Pitonak, M.; Jurecka, P.; Hobza, P. Stabilization and Structure
20 Calculations for Noncovalent Interactions in Extended Molecular Systems Based on Wave
21 Function and Density Functional Theories. *Chem. Rev.* **2010**, *110*, 5023-5063.
22
23 (32) Lynch, B. J.; Zhao, Y.; Truhlar, D. G. Effectiveness of Diffuse Basis Functions for
24 Calculating Relative Energies by Density Functional Theory. *J. Phys. Chem. A* **2003**, *107*,
25 1384-1388.
26
27 (33) Alecu, I. M.; Zheng, J. J.; Zhao, Y.; Truhlar, D. G. Computational Thermochemistry:
28 Scale Factor Databases and Scale Factors for Vibrational Frequencies Obtained from
29 Electronic Model Chemistries. *J. Chem. Theory Comput.* **2010**, *6*, 2872-2887.
30
31 (34) Marenich, A. V.; Cramer, C. J.; Truhlar, D. G. Performance of SM6, SM8, and SMD
32 on the SAMPL1 Test Set for the Prediction of Small-Molecule Solvation Free Energies. *J.*
33 *Phys. Chem. B* **2009**, *113*, 4538-4543.
34
35 (35) Colominas, C.; Teixido, J.; Cemeli, J.; Luque, F. J.; Orozco, M. Dimerization of
36 Carboxylic Acids: Reliability of Theoretical Calculations and the Effect of Solvent. *J. Phys.*
37 *Chem. B* **1998**, *102*, 2269-2276.
38
39 (36) Tang, E.; Di Tommaso, D.; de Leeuw, N. H. Accuracy of the Microsolvation-
40 Continuum Approach in Computing the pK(a) and the Free Energies of Formation of
41 Phosphate Species in Aqueous Solution. *Phys Chem Chem Phys* **2010**, *12*, 13804-13815.
42
43 (37) Montero, L. A. GRANADA and Q programs for PC computers. 1996.
44
45 (38) Codorniu-Hernandez, E.; Mesa-Ibirico, A.; Montero-Cabrera, L. A.; Martinez-
46 Luzardo, F.; Borrmann, T.; Stohrer, W. D. Theoretical Study of Flavonoids and Proline
47 Interactions. Aqueous and Gas Phases. *J. Mol. Struc. (Theochem)* **2003**, *623*, 63-73.
48
49 (39) Montero, L. A.; Molina, J.; Fabian, J. Multiple Minima Hypersurfaces of Water
50 Clusters for Calculations of Association Energy. *Int. J. Quantum. Chem.* **2000**, *79*, 8-16.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 (40) Sanchez-Garcia, E.; Mardyukov, A.; Studentkowski, M.; Montero, L. A.; Sander, W.
4 Furan-Formic Acid Dimers: An Ab Initio and Matrix Isolation Study. *J. Phys. Chem. A* **2006**,
5 *110*, 13775-13785.
6
7 (41) Sanchez-Garcia, E.; Mardyukov, A.; Tekin, A.; Crespo-Otero, R.; Montero, L. A.;
8 Sander, W.; Jansen, G. Ab Initio and Matrix Isolation Study of the Acetylene-Furan Dimer.
9 *Chem Phys* **2008**, *343*, 168-185.
10
11 (42) Sanchez-Garcia, E.; Montero, L. A.; Sander, W. Computational Study of Noncovalent
12 Complexes Between Formamide and Formic Acid. *J. Phys. Chem. A* **2006**, *110*, 12613-
13 12622.
14
15 (43) Sanchez-Garcia, E.; Studentkowski, M.; Montero, L. A.; Sander, W. Noncovalent
16 complexes between dimethyl ether and formic acid - An ab initio and matrix isolation study.
17 *ChemPhysChem* **2005**, *6*, 618-624.
18
19 (44) Madeja, F.; Havenith, M.; Nauta, K.; Miller, R. E.; Chocholousova, J.; Hobza, P.,
20 Polar Isomer of Formic Acid Dimers Formed in Helium Nanodroplets. *J. Chem. Phys.* **2004**,
21 *120*, 10554-10560.
22
23 (45) Pistolis, G.; Paleos, C. M.; Malliaris, A. Molecular Recognition in Organic-Solvents -
24 the Importance of Excimer Fluorescence Spectroscopy. *J. Phys. Chem.* **1995**, *99*, 8896-8902.
25
26 (46) Sellini, H.; Maurizot, J. C.; Dimicoli, J. L.; Helene, C. Hydrogen-Bonding of Amino-
27 Acid Side Chains to Nucleic-Acid Bases. *Febs. Lett.* **1973**, *30*, 219-224.
28
29 (47) Schrier, E. E.; Pottle, M.; Scheraga, H. A. The Influence of Hydrogen and
30 Hydrophobic Bonds on the Stability of the Carboxylic Acid Dimers in Aqueous Solution. *J.*
31 *Am. Chem. Soc.* **1964**, *86*, 3444-3449.
32
33 (48) Boys, S. F.; Bernardi, F. Calculation of Small Molecular Interactions by Differences
34 of Separate Total Energies - Some Procedures with Reduced Errors. *Mol. Phys.* **1970**, *19*,
35 553-566.
36
37 (49) Deng, J.; Gilbert, A. T. B.; Gill, P. M. W. Communication: Efficient Counterpoise
38 Corrections by a Perturbative Approach. *J. Chem. Phys.* **2011**, *135*, 081105.
39
40 (50) Bulmer, J. T.; Shurvell, H. F. Factor-Analysis as a Complement to Band Resolution
41 Techniques .1. Method and Its Application to Self-Association of Acetic-Acid. *J. Phys.*
42 *Chem.* **1973**, *77*, 256-262.
43
44 (51) Benghiat, V.; Leiserow, L. Molecular Packing Modes. 6. Crystal and Molecular
45 Structures of 2 Modifications of Tetrolic Acid. *J. Chem. Soc. Perk. Trans 2* **1972**, 1763-1768.
46
47 (52) Goralski, P.; Taniewskaosinska, S. On the Analysis of Calorimetric Data in the Study
48 of Dimerization of Carboxylic-Acids in Non-Polar Solvents. *Can. J. Chem.* **1983**, *61*, 2209-
49 2213.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 (53) Pham, H. H.; Taylor, C. D.; Henson, N. J. First-Principles Prediction of the Effects of
4 Temperature and Solvent Selection on the Dimerization of Benzoic Acid. *J. Phys. Chem. B*
5 **2013**, *117*, 868-876.
6

7 (54) Di Tommaso, D.; de Leeuw, N. H. The Onset of Calcium Carbonate Nucleation: A
8 Density Functional Theory Molecular Dynamics and Hybrid Microsolvation/Continuum
9 Study. *J. Phys. Chem. B* **2008**, *112*, 6965-6975.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Tables

Table 1. Simulation systems investigated in the present work.





Formic acid (FA)	
Acetic acid (AA)	
Tetrolic acid (TTA)	
Benzoic acid (BA)	

Table 2. Calculated and experimental values of the dimerization constants (pK_D) of acetic acid (AA)₂ and formic acid (FA)₂.

	solvent	M06-2X (no-cpc) ^a	M06-2X (cpc) ^b	PMF ^c	exptl
(AA) ₂	H ₂ O	-0.25	-0.32	0.038 ± 0.01	0.0-1.2 ^d
	CHCl ₃	-2.33	-2.52		-1.8 ^e
	CCl ₄	-4.27	-4.46		
(FA) ₂	H ₂ O	0.79	0.76	0.25 ± 0.02	1.2-2.1 ^f
	CHCl ₃	-1.13	-1.14		
	CCl ₄	-3.18	-3.19		

^aWe use “no-cpc” to denote calculations without counterpoise correction for the basis set superposition error (BSSE). ^bWe use “cpc” to denote geometry optimization with counterpoise correction for BSSE. ^cTaken from Ref [13]. ^dTaken from Ref [45]. ^eTaken from Ref [46]. ^fTaken from Ref [47].

Table 3. Energetics [kcal mol⁻¹] of formation of acetic acid (AA) clusters as computed at the M06-2X/6-31+G(d,p) level of theory: $\Delta E_{e,gas}$ is the gas phase interaction energy; ΔG_{ass}° is the standard state (1 atm) gas-phase free energy of association at 298 K; ΔG_{ass}^* is the standard state (1 mol L⁻¹) free energies of reaction in the liquid-phase.

species	reaction	type	$\Delta E_{e,gas}$	ΔG_{ass}°	ΔG_{ass}^*		
					H ₂ O	CHCl ₃	CCl ₄
dimer	AA + AA → (AA) ₂	I	-17.70	-6.56	-0.18	-3.20	-5.87
		II	-9.19	1.06	4.69	3.00	1.13
		III	-9.96	0.43	3.81	2.30	0.71
		IV	-7.16	3.03	5.09	4.14	3.00
		V	-7.16	1.97	2.93	2.14	1.26
		VI	-10.05	0.63	3.79	2.30	0.91
		VII	-6.36	4.76	5.37	5.20	4.77
		VIII	-10.07	-0.10	2.61	1.22	-0.33
		IX	-4.66	5.08	5.97	5.66	5.20
trimer	(AA) ₂ + AA → (AA) ₃ ^a	I	-9.36	0.59	3.02	2.02	0.75
		II	-9.06	2.63	4.71	2.88	2.72
		III	-5.73	4.21	4.52	2.69	3.52
		IV	-7.78	3.14	4.66	2.83	2.87
		V	-7.99	2.60	5.17	3.34	2.55
		VI	-8.81	4.56	5.03	3.21	5.04
		VII	-7.99	3.11	5.24	3.41	3.21
	3 AA → (AA) ₃ ^a	I	-27.07	-5.91	2.90	-1.13	-5.07
		II	-26.77	-3.88	4.58	-0.27	-3.10
		III	-23.44	-2.29	4.40	-0.45	-2.29
		IV	-25.49	-3.37	4.53	-0.32	-2.95
		V	-25.69	-3.91	5.05	0.20	-3.27
		VI	-25.51	-1.95	4.91	0.06	-0.78
		VII	-25.69	-3.40	5.12	0.27	-3.16
tetramer	(AA) ₂ + (AA) ₂ → (AA) ₄ ^b		-10.40	4.70	3.69	4.78	5.26
	(AA) ₃ + AA → (AA) ₄ ^b		-18.78	-2.12	0.46	-0.46	-1.36
	(AA) ₂ + 2 AA → (AA) ₄ ^b		-28.10	0.00	3.57	1.63	-0.56
pentamer	(AA) ₃ + (AA) ₂ → (AA) ₅ ^c		-11.82	5.50	4.32	5.46	6.12
	2 (AA) ₂ + AA → (AA) ₄ ^c		-21.14	6.13	7.43	7.55	6.93

^aTrimers generated using as building units the gas phase geometries of the acetic acid molecule and the cyclic dimer **I**. ^bTetramers generated using as building units the cyclic dimer **I**. ^cPentamers generated using as building units the cyclic dimer **I** and the trimeric structure **I**.

Table 4. Energetics [kcal mol⁻¹] of formation of tetrolic acid (TTA) clusters as computed at the M06-2X/6-31+G(d,p) level of theory: $\Delta E_{e,gas}$ is the gas phase interaction energy; ΔG_{ass}° is the standard state (1 atm) gas-phase free energy of association at 298 K; ΔG_{ass}^* is the standard state (1 mol L⁻¹) free energies of reaction in the liquid-phase.

species	reaction	type	$\Delta E_{e,gas}$	ΔG_{ass}°	ΔG_{ass}^*		
					H ₂ O	CHCl ₃	CCl ₄
dimer	TTA + TTA → (TTA) ₂	I	-17.56	-5.38	0.85	-1.80	-4.62
		II	-9.61	2.03	5.13	4.19	2.79
		III	-9.33	2.21	5.34	4.45	2.99
		IV	-7.06	5.25	7.23	7.27	6.47
		V	-6.79	4.00	6.41	6.03	4.52
		VI	-6.17	5.22	5.98	4.42	4.60
		VII	-5.04	5.96	6.30	6.78	6.37
		VIII	-5.42	9.30	8.73	9.28	9.11
		IX	-1.02	11.03	11.14	10.65	10.26
trimer	(TTA) ₂ + TTA → (TTA) ₃ ^a		-8.90	1.48	3.50	3.97	2.37
			-26.47	-3.90	4.30	2.19	-2.23
tetramer	(TTA) ₂ + (TTA) ₂ → (TTA) ₄ ^b		-8.11	3.72	3.30	5.76	5.61
			-16.53	-1.54	-0.30	-0.05	-1.35
			-25.68	0.22	4.09	3.83	0.84
hexamer	3 (TTA) ₂ → (TTA) ₆ ^c		-20.23	10.28	3.77	10.61	11.81

^a Trimers generated using the gas phase geometries of the tetrolic acid molecule and the cyclic dimer **I**. ^bTetramers generated using two acetic acids cyclic dimers **I**. ^cHexamers generated using three acetic acid cyclic dimers.


Table 5. Energetics [kcal mol⁻¹] of formation of benzoic acid (BA) clusters as computed at the M06-2X/6-31+G(d,p) level of theory: $\Delta E_{e,gas}$ is the gas phase interaction energy; ΔG_{ass}° is the standard state (1 atm) gas-phase free energy of association at 298 K; ΔG_{ass}^* is the standard state (1 mol L⁻¹) free energies of reaction in the liquid-phase.

species	reaction	structure	$\Delta E_{e,gas}$	ΔG_{ass}°	ΔG_{ass}^*		
					H ₂ O	CHCl ₃	CCl ₄
dimer	BA + BA → (BA) ₂	I	-18.51	-6.48	0.21	-2.97	-7.77
		II	-9.46	2.44	2.40	3.63	1.15
		III	-9.23	1.33	2.64	2.25	-0.75
		IV	-9.20	1.34	4.50	3.30	-0.56
		V	-7.98	3.68	2.33	4.26	2.14
		VI	-7.041	4.24	1.26	4.09	2.43
		VII	-7.36	4.96	2.51	4.80	3.12
		VIII	-6.90	3.98	1.06	3.90	2.21
		IX	-6.83	4.53	1.54	4.44	2.80
		X	-6.55	4.72	1.77	4.56	2.93
		XI	-6.54	5.89	2.63	5.56	3.96
		XII	-6.47	5.70	3.40	5.56	3.90
		XIII	-5.73	6.10	2.33	5.58	3.09
		XIV	-5.56	5.34	2.42	4.66	4.03
		XV	-5.29	6.06	3.03	5.54	3.88
		XVI	-4.48	5.97	4.96	5.89	3.45
		XVII	-4.21	5.10	4.71	5.15	2.29
		XVIII	-3.48	7.33	5.41	6.93	4.93
trimer	(BA) ₂ + BA → (BA) ₃ ^a		-9.73	3.44	0.31	3.61	2.67
	3 (BA) → (BA) ₃ ^a		-28.20	-3.17	0.43	0.55	-5.20
tetramer	(BA) ₂ + (BA) ₂ → (BA) ₄ ^b		-14.14	1.19	-4.07	1.65	3.04
	(BA) ₂ + 2 (BA) → (BA) ₄ ^b		-32.65	-5.38	-3.55	-1.41	-4.83
	(BA) ₃ + (BA) → (BA) ₄ ^b		-22.96	-8.78	-4.07	-5.03	-7.50


^aTrimers generated using the gas phase geometries of the benzoic acid molecule and the cyclic dimer **I**. ^bTetramer generated using two benzoic acids cyclic dimers **I**.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Figures



1
2
3
4
5 **Figure 1.** Structures of acetic acid dimers. In bold are the structures that have been located in this study. In parenthesis are given the energy
6 differences with respect to structure **I**. Distances in Å and energies in kcal mol⁻¹.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49



1
2
3
4
5 **Figure 2.** Structures of formic acid dimers. In bold are the structures that have been located in this study. In parenthesis are reported the energy
6 differences with respect to structure **I**. Distances in Å and energies in kcal mol⁻¹.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49





Figure 3. Relative energies of candidate structures of acetic acid trimers generated from the aggregation of three acetic acid molecules, (3M)_T, and of the cyclic dimer **I** with one acetic acid molecule, (D+M)_T. Values computed at the M06-2X/6-31+G(d,p) level of theory.



1
2
3
4
5 **Figure 4.** Structures of acetic acid trimers. In parenthesis are given the energy differences with respect to structure **I**. Distances in Å and
6 energies in kcal mol⁻¹.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49




Figure 5. Structures of acetic acid tetramers. Distances in Å.




Figure 6. Calculated reaction free energy of acetic acid tetramers.




Figure 7. Structures of the acetic acid pentamers. Distances in Å.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 8. Structures of tetrolic acid dimers. In parenthesis are given the energy differences with respect to structure I. Distances in Å.




Figure 9. Structures of tetrolic acid trimers. The gas phase interaction energies and the standard state (1 mol L⁻¹) free energies of formation in water, chloroform and carbon tetrachloride (values in parenthesis) computed according to the association reaction 3 BA → (BA)₃ are also reported. Distances in Angstrom and energies in kcal mol⁻¹.




Figure 10. Relative energies of the candidate structures of the tetrolic acid tetramer generated using as building units two cyclic dimers **I**, (D+D)_T, the cyclic dimer **I** with two tetrolic acid molecules, (D+2M)_T, and the most stable trimer **I** with one tetrolic acid molecule, (Tr+M)_T. Values computed at the M06-2X/6-31+G(d,p) level of theory.




Figure 11. Structures of the tetrolic acid tetramers. Distances in Å.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49




Figure 12. Calculated reaction free energy of tetrolic acid tetramers.




Figure 13. Structures of the six most stable benzoic acid dimers. In parenthesis are given the energy differences with respect to structure I. Distances in Å and energies in kcal mol⁻¹.




Figure 14. Structures and gas phase interaction energies ($\Delta E_{e,gas}$) of the most stable benzoic acid trimers. In parenthesis are reported the standard state (1 mol L^{-1}) free energies of reaction in carbon tetrachloride (ΔG_{ass}^*) computed according to the association reaction $3 \text{ BA} \rightarrow (\text{BA})_3$. Distances in Angstrom and energies in kcal mol^{-1} .




Figure 15. Structures of representative benzoic acid tetramers. Distances in Angstrom.




Figure 16. Calculated reaction free energy of benzoic acid tetramers.




Figure 17. Free energy pathways of the formation in carbon tetrachloride of dimers, trimers and tetramers of acetic acid (AA), tetrolic acid (TTA) and benzoic acid (BA). Values in kcal mol⁻¹.

Table of Contents - Graphic

