



Permutations and the combinatorics of gauge invariants for general N

Ramgoolam, S

<http://arxiv.org/abs/1605.00843>

For additional information about this publication click this link.

<http://qmro.qmul.ac.uk/xmlui/handle/123456789/12488>

Information about this research object was correct at the time of download; we occasionally make corrections to records, please therefore check the published record when citing. For more information contact scholarlycommunications@qmul.ac.uk

Permutations and the combinatorics of gauge invariants for general N

Sanjaye Ramgoolam *

*Centre for Research in string theory,
School of Physics and Astronomy,
Queen Mary, University of London
E-mail: s.ramgoolam@qmul.ac.uk*

Group algebras of permutations have proved highly useful in solving a number of problems in large N gauge theories. I review the use of permutations in classifying gauge invariants in one-matrix and multi-matrix models and computing their correlators. These methods are also applicable to tensor models and have revealed a link between tensor models and the counting of branched covers. The key idea is to parametrize $U(N)$ gauge invariants using permutations, subject to equivalences. Correlators are related to group theoretic properties of these equivalence classes. Fourier transformation on symmetric groups by means of representation theory offers nice bases of functions on these equivalence classes. This has applications in AdS/CFT in identifying CFT duals of giant gravitons and their perturbations. It has also lead to general results on quiver gauge theory correlators, uncovering links to two dimensional topological field theory and the combinatorics of trace monoids.

Proceedings of the Corfu Summer Institute 2015 "School and Workshops on Elementary Particle Physics and Gravity"

1-27 September 2015

Corfu, Greece

*Speaker.

1. Introduction

The AdS/CFT correspondence [1] relates ten dimensional string theory to four dimensional conformal quantum field theory. In CFT, the operator-state correspondence relates quantum states to local operators, which are gauge invariant composites built from the elementary fields. The enumeration of these composites and the computation of their correlators is an algebraic problem. The emergence from these algebraic structures of the extra dimensions of the string background, of the strings and branes in these backgrounds, which we expect from the duality is an example of geometry emerging from algebra. The idea that space-time geometry is to be reconstructed from algebras underlies work in non-commutative geometry, a subject of interest to many in the audience at the Corfu2015 conference. In this talk, I will be describing some ways in which geometrical structures emerge from algebraic data in the context of CFT correlators.

The physical context of what I will describe is $\mathcal{N} = 4$ SYM, the canonical example of gauge string duality, where the dual is string theory in $AdS_5 \times S^5$. The gauge group of the CFT4 is $U(N)$. The fields transform in the adjoint. In addition to gauge fields and fermions, we have six hermitian adjoint scalars X_1, \dots, X_6 . They transform in the vector representation of an $SO(6)$ global symmetry, which forms part of the $PSU(2, 2|4)$ superconformal algebra.

Another algebra, more precisely a sequence of algebras, will play a crucial role in this talk. This is the sequence of group algebras of symmetric groups $\mathbb{C}(S_n)$, for natural numbers n . Aside from the obvious Lagrangian symmetries, the direct sum of symmetric group algebras,

$$\mathbb{C}(S_\infty) \equiv \bigoplus_{n=0}^{\infty} \mathbb{C}(S_n) \tag{1.1}$$

will be found to play an important role in the description of the state space of CFT. Permutation algebras are useful in organizing the multiplicities of representations of $PSU(2, 2|4)$. In the simplest sector of interest, the maximally supersymmetric half-BPS sector, the multiplicity of representations will be related to problems of invariant theory of one matrix, and will be organized by the conjugacy classes of S_n . For the less supersymmetric quarter and eighth-BPS sectors, we encounter a multi-matrix problem and a generalization of conjugacy class algebras will play a role.

A very useful tool for thinking about the matrix invariant theory problems we encounter, and the role of permutations in these problems, will be tensor spaces and linear operators acting on these tensor spaces. Diagrams for representing linear operators, their products and traces, provide a powerful way to uncover the hidden simplicity of multi-index tensor manipulations. These tools will be illustrated as we proceed from the simplest example of 1-matrix to more complex multi-matrix problems encountered in AdS/CFT and to problems of tensor invariants encountered in tensor models.

2. Half-BPS sector in $\mathcal{N} = 4$ SYM: 1-complex matrix model

Using two of the six hermitian scalars in $N = 4$ SYM, define $Z = X_1 + iX_2$. Gauge invariant operators such as

$$\text{tr}Z^3, \text{tr}Z^2\text{tr}Z, (\text{tr}Z)^3 \tag{2.1}$$

which are holomorphic gauge-invariant polynomials in the matrix elements Z_j^i are annihilated by half of the 32 supercharges in the super-algebra $PSU(2,2|4)$. Acting with the generators of the super-algebra produces an ultra-short representation. All ultra-short representations are generated in this way. Their correlators have non-renormalization properties (see [2] and refs. therein to earlier work) which allows comparison between free conformal field theory computations and supergravity. These holomorphic gauge invariant operators satisfy the BPS condition $\Delta = J$ relating the dimension to a $U(1)$ charge inside the $SO(6)$ R-symmetry. The enumeration of these BPS operators using a Young diagram basis, the computation of correlators in this basis and the identification of Young diagram states with giant gravitons was developed in [3].

For a fixed $\Delta = n$, and restricting to $n < N$, the number of gauge invariant polynomials is equal to the number of partitions of n , i.e. the number of ways of writing n as a sum of positive integers. This is also the number of conjugacy classes of the symmetric group and this is no accident. Gauge invariants can be constructed from permutations $\sigma \in S_n$ which determine the order in which the lower indices are contracted with the upper indices

$$\mathcal{O}_\sigma \equiv Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n)}}^{i_n} \quad (2.2)$$

It is useful to think of Z as an operator acting on an N -dimensional vector space V_N and $Z^{\otimes n}$ as an operator on the tensor product $V_N^{\otimes n}$. There is a standard action of $\sigma \in S_n$ on the n -fold tensor product. Choosing a set of basis vectors $\{e_i, 1 \leq i \leq N\}$ for V_N , permutations act as

$$\sigma (e_{i_1} \otimes e_{i_2} \cdots e_{i_n}) = e_{i_{\sigma(1)}} \otimes \cdots \otimes e_{i_{\sigma(n)}} \quad (2.3)$$

The tensor product of Z -operators acts in the standard way as

$$Z^{\otimes n}(e_{i_1} \otimes e_{i_2} \cdots e_{i_n}) = (Z_{i_1}^{j_1} Z_{i_2}^{j_2} \cdots Z_{i_n}^{j_n}) e_{j_1} \otimes e_{j_2} \cdots e_{j_n} \quad (2.4)$$

One checks that

$$\begin{aligned} \mathcal{O}_\sigma(Z) &= \text{tr}_{V_N^{\otimes n}}(Z^{\otimes n} \sigma) \\ &= \langle e^{i_1} \otimes e^{i_2} \otimes \cdots \otimes e^{i_n} | Z^{\otimes n} \sigma | e_{i_1} \otimes e_{i_2} \cdots \otimes e_{i_n} \rangle \end{aligned} \quad (2.5)$$

Using the standard connection between linear operators and diagrams, such as used in knot theory for example, we can represent this diagrammatically as in Figure 1. In this picture a line represents a state in $V_N^{\otimes n}$, a box is a linear operator and a contraction is an identification of an upper line with a lower line. On the left of the picture, this is represented by joining top and bottom. On the right picture the upper index line ends on a horizontal line, the lower index line ends on another horizontal line and the two horizontal lines are understood to be identified. The diagrammatic way of thinking about operators and correlators is explained further in [4, 5].

It is plausible and very easy to check that, for any permutation $\gamma \in S_n$,

$$\gamma Z^{\otimes n} \gamma^{-1} = Z^{\otimes n} \quad (2.6)$$

This is an equality of operators acting on $V_N^{\otimes n}$. It follows that

$$\mathcal{O}_{\gamma\sigma\gamma^{-1}}(Z) = \text{tr}_{V_N^{\otimes n}}(Z^{\otimes n} \gamma\sigma\gamma^{-1}) = \text{tr}_{V_N^{\otimes n}}(\gamma^{-1} Z^{\otimes n} \gamma\sigma) = \mathcal{O}_\sigma(Z) \quad (2.7)$$

The enumeration of gauge invariants is the same as enumerating conjugacy classes in S_n . Thus

$$\mathbb{C}(S_\infty) = \bigoplus_n \mathbb{C}(S_n) \tag{2.8}$$

is a hidden algebra which organizes gauge invariants.

It turns out that permutations are important, not just for enumerating gauge invariants, but also for computing their correlators. Consider a two point function involving a holomorphic gauge invariant polynomial of fixed degree and an anti-holomorphic polynomial of fixed degree in the free field limit. This is non-zero if the two polynomials have the same degree. The non-vanishing two-point functions are thus of the form

$$\langle \mathcal{O}_{\sigma_1}(Z)(x_1)(\mathcal{O}_{\sigma_2}(Z)(x_2))^\dagger \rangle \tag{2.9}$$

where $\sigma_1, \sigma_2 \in S_n$. The correlator is calculated using the basic correlator

$$\langle Z_{j_1}^{i_1}(x_1)(Z^\dagger_{j_2}{}^{i_2}(x_2)) \rangle = \frac{\delta_{j_2}^{i_1} \delta_{j_1}^{i_2}}{(x_1 - x_2)^2} \tag{2.10}$$

and Wick's theorem, which implies that

$$\begin{aligned} \langle Z_{j_1}^{i_1} \dots Z_{j_n}^{i_n} Z_{l_1}^{\dagger k_1} \dots Z_{l_n}^{\dagger k_n} \rangle &= \delta_{j_{\gamma(1)}}^{k_1} \delta_{l_1}^{i_{\gamma(1)}} \dots \delta_{j_{\gamma(n)}}^{k_n} \delta_{l_n}^{i_{\gamma(n)}} \\ &= \delta_{j_{\gamma(1)}}^{k_1} \delta_{l_{\gamma^{-1}(1)}}^{i_1} \dots \delta_{j_{\gamma(n)}}^{k_n} \delta_{l_{\gamma^{-1}(n)}}^{i_n} \end{aligned} \tag{2.11}$$

We have dropped the space-time dependence which is trivial. Performing the free field path integral amounts to relating the Z, Z^\dagger operators in tensor space by permutation operators. This can be expressed diagrammatically in Figure 2.




Figure 1: Operator as trace in tensor space

The correlator can be computed by putting together (2.2) and (2.11). The calculation can also be understood diagrammatically as in Figure 3. The result is

$$(x_1 - x_2)^{2n} \langle \mathcal{O}_{\sigma_1}(Z(x_1))(\mathcal{O}_{\sigma_2}(Z(x_2)))^\dagger \rangle = \sum_{\gamma \in S_n} \text{tr}_{V_N^{\otimes n}}(\sigma_1 \gamma \sigma_2^{-1} \gamma^{-1})$$




Figure 2: Wick contraction sum as operator in tensor space




Figure 3: Correlator as trace in tensor space

$$\begin{aligned}
 &= \sum_{\gamma \in S_n} N^{C_{\sigma_1 \gamma \sigma_2^{-1} \gamma^{-1}}} \\
 &= \sum_{\gamma, \sigma_3 \in S_n} N^{C_{\sigma_3}} \delta(\sigma_1 \gamma \sigma_2 \gamma^{-1} \sigma_3) \\
 &= \frac{n!}{|T_1| |T_2|} \sum_{\sigma'_1 \in T_1, \sigma'_2 \in T_2, \sigma_3 \in S_n} \delta(\sigma'_1 \sigma'_2 \sigma_3) N^{C_{\sigma_3}} \quad (2.12)
 \end{aligned}$$

The first step comes from the diagrammatic manipulation. In the second we use that, the trace in $V_N^{\otimes n}$ of a permutation is given in terms of the number of cycles in σ , denoted by C_σ

$$\text{tr}_{V_N^{\otimes n}}(\sigma) = N^{C_\sigma} \quad (2.13)$$

In the third line, we have introduced an extra permutation and a delta function on the permutation group to write a neater formula. $\delta(\sigma)$ is defined to be 1 if σ is the identity permutation and 0 otherwise. Finally since \mathcal{O}_σ only depends on the conjugacy class, we have replaced σ_1, σ_2 by sums of permutations in their respective conjugacy classes T_1, T_2 , dividing out by the sizes of these conjugacy classes $|T_1|, |T_2|$. With these sums in place, the γ sum can be replaced by $n!$ since the sums commute with γ .

This shows that the introduction of permutations in S_n as a way to parametrize the gauge invariant operators is extremely useful, since correlators can be given neatly in terms of group theory of S_n .

The final formula has an interpretation in terms of counting of branched covering maps from Riemann surfaces (worldsheets Σ) to the sphere \mathbb{P}^1 branched over three points on the sphere. These

branched covers are holomorphic maps

$$f : \Sigma \rightarrow \mathbb{P}^1 \tag{2.14}$$

with vanishing holomorphic derivatives at points on Σ which are in the inverse image of three points on the \mathbb{P}^1 , which can be taken as $\{0, 1, \infty\}$ [6, 7]. These types of holomorphic maps are called Belyi maps and have deep implications for number theory, in connection with the absolute Galois group [8].

It is rather amazing that correlators in the distinguished half-BPS sector of $\mathcal{N} = 4$ SYM, equivalently in a distinguished $AdS_5 \times S^5$ background of string theory, are related directly to Belyi maps, of central importance in number theory, a subject which Gauss famously called the Queen of Mathematics.

The power of the permutation approach to gauge-invariants becomes particularly manifest when we consider finite N effects. While the $\mathcal{O}_\sigma(Z)$, with conjugation equivalence, form a good basis of operators when $n \leq N$ this is not the case for $n > N$. In this case, there are relations among traces due to the Cayley Hamilton theorem which implies that $tr Z^{N+1}$ can be written in terms of multi-traces. An elegant way to give a basis at finite N is to use the Fourier transform relation from group theory, which relates conjugacy classes to irreducible representations

Irreducible representations R of S_n , correspond to Young diagrams with n boxes. We can define operators corresponding to Young diagrams as

$$\mathcal{O}_R(Z) = \frac{1}{n!} \sum_{\sigma \in S_n} \chi_R(\sigma) \mathcal{O}_\sigma(Z) \tag{2.15}$$

where $\chi_R(\sigma)$ is the character of the S_n group element σ in the irrep R , i.e. the trace of the matrix representing σ in irrep R . It is shown that [3] these operators have diagonal 2-point function.

$$\langle \mathcal{O}_R(Z) (\mathcal{O}_S(Z))^\dagger \rangle = \delta^{RS} \frac{n! \text{Dim} R}{d_R} \tag{2.16}$$

Failure of diagonality of the trace basis had been used [9] to argue that giant gravitons are not traces. Single giants which are large in the S^5 were identified with subdeterminant operators. With the Young diagram basis in hand, there are natural candidates for CFT duals of single and multi-giant graviton states [3].

3. Multi-matrix model

The above approach to matrix invariants, based on permutations and diagrammatic tensor space methods, extends to multi-matrix models. This should be contrasted with many other approaches to 1-matrix models, e.g. reduction to eigenvalue dynamics, which do not admit multi-matrix generalization. The enumeration and correlators of multi-matrix invariants has direct applications to quarter and eighth BPS sectors of SYM. I will explain how permutation methods described above generalize to give restricted Schur operators. These were introduced in the study of open string excitations of giant gravitons [10, 11, 12, 13], subsequently shown to diagonalize the 2-point functions in the 2-matrix sector [14] and understood in terms of enhanced symmetries in gauge theory [15].

Consider the two-matrix problem, of enumerating gauge invariants built from two matrices. In the context of the quarter-BPS sector of SYM, we are interested in holomorphic polynomial gauge invariants built from two matrices Z, Y , where the gauge symmetry acts as

$$(Z, Y) \rightarrow (UZU^\dagger, UYU^\dagger) \quad (3.1)$$

As in the 1-matrix case, we observe that gauge invariants can be parametrized by permutations which determine how the lower indices are contracted with the upper indices

$$\mathcal{O}_\sigma(Z, Y) = \text{tr}_{V_N^{\otimes m+n}} (Z^{\otimes m} \otimes Y^{\otimes n} \sigma) = Z_{i_{\sigma(1)}}^{i_1} Z_{i_{\sigma(2)}}^{i_2} \cdots Z_{i_{\sigma(m)}}^{i_m} Y_{i_{\sigma(m+1)}}^{i_{m+1}} \cdots Y_{i_{\sigma(m+n)}}^{i_{m+n}} \quad (3.2)$$

Here the permutation σ is in the symmetric group S_{m+n} , where m is the number of Z and n is the number of Y in the gauge-invariant operator. In this two-matrix case, the equivalence takes the form

$$\mathcal{O}_\sigma(Z, Y) = \mathcal{O}_{\gamma\sigma\gamma^{-1}}(Z, Y) \quad (3.3)$$

for $\gamma \in S_m \times S_n$. In the large N limit, i.e. when $m+n < N$, these are the only equivalences. Counting gauge invariant operators is equivalent to enumerating orbits of $S_m \times S_n$ permutations γ acting on permutations $\sigma \in S_{m+n}$ by conjugation :

$$\sigma \sim \gamma\sigma\gamma^{-1} \quad \text{for} \quad \gamma \in S_m \times S_n \quad (3.4)$$

From the Burnside Lemma, we know that the number of orbits is equal to the number of fixed points, so that

$$\text{Number of operators with } m, n \text{ copies of } Z, Y = \frac{1}{m!n!} \sum_{\gamma \in S_m \times S_n} \sum_{\sigma \in S_{m+n}} \delta(\gamma\sigma\gamma^{-1}\sigma^{-1})$$

This formula can be used to obtain the generating function

$$\begin{aligned} \sum_{m,n=0}^{\infty} z^m y^n (\text{Number of operators with } m, n \text{ copies of } Z, Y) \\ = \prod_{i=1}^{\infty} \frac{1}{(1-z^i - y^i)} \end{aligned}$$

The expression also has an interpretation as a partition function for topological lattice gauge theory with S_{m+n} gauge group on a two-torus, with a defect inserted on a circle, which constrains the holonomy to be in the $S_m \times S_n$ subgroup. The free field 2-point function is

$$\begin{aligned} \langle \mathcal{O}_{\sigma_1}(Z, Y) (\mathcal{O}_{\sigma_2}(Z, Y))^\dagger \rangle &= \sum_{\gamma} \text{tr}_{V_N^{\otimes m+n}} (\sigma_1 \gamma \sigma_2 \gamma^{-1}) \\ &= \sum_{\gamma \in S_m \times S_n} \sum_{\beta \in S_{m+n}} \delta(\sigma_1 \gamma \sigma_2^{-1} \gamma^{-1} \beta) N^{C_\beta} \end{aligned} \quad (3.5)$$

As in the 1-matrix case, finite N counting and correlators can be obtained by going to a Fourier basis using matrix elements of the permutation $\sigma \in S_{m+n}$ in irreps R . There are linear operators $D^R(\sigma)$ which give functions $D_{IJ}^R(\sigma)$ where I, J can be taken to run over an orthonormal basis for

the irrep R . Since the functions of interest are invariant under $S_m \times S_n$, it is useful to use group theoretic data associated with reduction of the irrep R of S_{m+n} into irreps of $S_m \times S_n$. These can be parametrized by a pair of Young diagram R_1, R_2 with m, n boxes respectively. A representation R of S_{m+n} can be decomposed into a direct sum of representations R_1, R_2 , which appear with multiplicities $g(R_1, R_2, R)$ equal to the Littlewood-Richardson coefficients. Thus there is a subgroup-adapted basis in V_R , with states of the form

$$|R_1, R_2, m_1, m_2; \nu_1\rangle \quad (3.6)$$

where m_1, m_2 are state labels for the irreps R_1, R_2 and ν_1 is a multiplicity label taking values between 1 and $g(R_1, R_2, R)$. we may write the relevant decomposition as

$$V_R^{(S_{m+n})} = \bigoplus_{R_1, R_2} V_{R_1}^{(S_m)} \otimes V_{R_2}^{(S_n)} \otimes V_{R_1, R_2}^R \quad (3.7)$$

where the dimension of the multiplicity space V_{R_1, R_2}^R is $g(R_1, R_2, R)$. The subgroup basis states can be expanded in terms of a general orthonormal basis

$$|R_1, R_2, m_1, m_2; \nu_1\rangle = \sum_I |R, I\rangle \langle R, I | R_1, R_2, m_1, m_2; \nu_1\rangle \quad (3.8)$$

The branching coefficients $\langle R, I | R_1, R_2, m_1, m_2; \nu_1\rangle$ can be used to define functions $\chi_{R_1, R_2, \nu_1, \nu_2}^R(\sigma)$, called restricted characters, labelled by three Young diagrams and two multiplicity indices. These functions are invariant under conjugation by $\gamma \in S_m \times S_n$. These restricted characters are used to define restricted Schur Polynomials

$$\chi_{R_1, R_2, \nu_1, \nu_2}^R(Z, Y) = \sum_{\sigma \in S_{m+n}} \chi_{R_1, R_2, \nu_1, \nu_2}^R(\sigma) \mathcal{O}_\sigma(Z, Y) \quad (3.9)$$

In this Fourier basis, finite N constraints are easy to implement and they follow from Schur-Weyl duality [16, 17]. They amount to restricting the Young diagram R to have no more than N rows. Thus the finite N counting of 2-matrix invariants is given by

$$\begin{aligned} & \text{Finite } N \text{ counting of 2-matrix gauge invariants} \\ &= \sum_{\substack{R \text{ with } m+n \text{ boxes} \\ R \text{ has no more than } N \text{ rows}}} \sum_{R_1 \text{ with } m \text{ boxes}} \sum_{R_2 \text{ with } n \text{ boxes}} (g(R_1, R_2, R))^2 \end{aligned} \quad (3.10)$$

This counting formula can be obtained directly [18] from the group integral formula for counting gauge invariants which was introduced by Sundborg [19]. Another formula [20] for the counting in terms of $U(2)$ representations Λ is

$$\begin{aligned} & \text{Finite } N \text{ counting of 2-matrix gauge invariants} \\ &= \sum_{\substack{R \text{ with } m+n \text{ boxes} \\ R \text{ has no more than } N \text{ rows}}} \sum_{\Lambda \text{ with } m+n \text{ boxes}} C(R, R, \Lambda) M(\Lambda; m, n) \end{aligned}$$

$$(3.11)$$

$C(R, R, \Lambda)$ is the S_n Kronecker coefficient (Clebsch-series multiplicity) for Λ in $R \otimes R$, and $M(\Lambda, m, n)$ is the multiplicity of the irrep of the symmetric representations $[m] \otimes [n]$ of $S_m \times S_n$ in the decomposition of Λ into irreps of $S_m \times S_n$. The equivalence of the two counting formulae was shown in [21].

The restricted Schur operators have diagonal 2-point function in the free field limit [14]

$$\langle \chi_{R_1, R_2, v_1, v_2}^R(Z, Y) (\chi_{S_1, S_2, \mu_1, \mu_2}^S(Z, Y))^\dagger \rangle \propto \delta_{R, S} \delta_{v_1, v_2} \delta_{\mu_1, \mu_2} \quad (3.12)$$

Diagonal bases in the 2-matrix problem, with well-defined $U(2)$ global symmetry quantum numbers have also been constructed [20, 22]. Brauer algebras provide another useful free field basis [23]. The operators in these different bases diagonalise commuting sets of Casimirs for enhanced symmetries in the free field limit [15]. For the two-matrix case, we have a $U(N) \times U(N)$ symmetry of left multiplication by unitary matrices

$$Z, Y \rightarrow UZ, VY \quad (3.13)$$

We can also multiply on the right. For each matrix there is in fact a $U(N^2)$ symmetry. The Noether charges for these different symmetries can be used to construct Casimirs which measure the representation theory labels used to construct the different bases. Finding minimal sets of charges determining the labels of the restricted Schur basis leads to the study of the structure of permutation centralizer algebras [24]. The starting point of this discussion is that the averaging over the equivalence classes in (3.3) and using the product in $\mathbb{C}(S_{m+n})$ to define a non-commutative associative algebra $\mathcal{A}(m, n)$ on these equivalence classes. The restricted characters in 3.9 lead to the Matrix (Wedderburn-Artin) decomposition of the algebra $\mathcal{A}(m, n)$ in terms of the expressions

$$Q_{R_1, R_2, v_1, v_2}^R = \sum_{\sigma \in S_{m+n}} \chi_{R_1, R_2, v_1, v_2}^R(\sigma) \sigma \quad (3.14)$$

This gives gives the intrinsic permutation meaning of the restricted characters, without reference to matrices. The structure of these algebras is closely related to Littlewood-Richardson coefficients, and can be viewed as a sort of categorification thereof. The relations between the centre and an appropriately defined Cartan (the diagonals of the Wedderburn Artin decomposition) play an important role in the definition of minimal sets of charges.

3.1 Quivers

Recently these results have been generalized to the counting and correlators of quiver gauge theories, where the gauge group is a product of unitary groups and the matter is in bifundamentals or fundamentals [18, 25, 26]. Links to two dimensional topological field theory have been described and some surprising connections to trace monoids which have applications in computer science have been described. This arises because the counting function for large ranks of the Unitary groups is an infinite product of inverse determinants. The inverse determinant counts words constructed from an alphabet consisting of letters corresponding to simple loops, with partial commutation relations between the letters.

4. Tensor models

Consider a complex 3-index tensor $\Phi_{i,j,k}$ which transforms as $V_N \otimes V_N \otimes V_N$ and its conjugate $\bar{\Phi}^{i,j,k}$, which transforms as $\bar{V}_N \otimes \bar{V}_N \otimes \bar{V}_N$. The three ranks can, more generally, be chosen to be unequal. Polynomial invariants are constructed by contracting the upper indices with the lower indices [27].

Permutation methods, of the type above which have been useful in enumerating multi-matrix invariants as outlined above, were used to approach the tensor invariants in [28]. Explicit generating functions for the counting, new connections with branched covers, as well as group theoretic expressions for correlators were found.

Choose a positive integer n which determines the number of Φ and $\bar{\Phi}$ used to construct the invariants. These invariants can be parametrized by a triple of permutations $(\sigma_1, \sigma_2, \sigma_3)$ which contract the i -indices, the j -indices and the k -indices.

$$\mathcal{O}_{\sigma_1, \sigma_2, \sigma_3}(\Phi, \bar{\Phi}) = \bar{\Phi}^{i_1, j_1, k_1} \dots \bar{\Phi}^{i_n, j_n, k_n} \Phi_{i_{\sigma_1(1)}, j_{\sigma_2(1)}, k_{\sigma_3(1)}} \dots \Phi_{i_{\sigma_1(n)}, j_{\sigma_2(n)}, k_{\sigma_3(n)}} \quad (4.1)$$

There is an S_n symmetry of permuting the Φ and an S_n symmetry of permuting the $\bar{\Phi}$'s. Thus the counting of 3-index tensor invariants of degree n is equivalent to counting permutation triples $(\sigma_1, \sigma_2, \sigma_3)$ subject to equivalences generated by $\gamma_1, \gamma_2 \in S_n$.

$$(\sigma_1, \sigma_2, \sigma_3) \sim (\gamma_1 \sigma_1 \gamma_2, \gamma_1 \sigma_2 \gamma_2, \gamma_1 \sigma_3 \gamma_2) \quad (4.2)$$

Using the Burnside Lemma, the number of these invariants is

$$\frac{1}{n!^2} \sum_{\gamma_1, \gamma_2 \in S_n} \sum_{\sigma_1, \sigma_2, \sigma_3} \delta(\gamma_1 \sigma_1 \gamma_2 \sigma_1^{-1}) \delta(\gamma_1 \sigma_2 \gamma_2 \sigma_2^{-1}) \delta(\gamma_1 \sigma_3 \gamma_2 \sigma_3^{-1}) \quad (4.3)$$

This expression can be easily simplified. The first step is to solve for one of the γ 's. The upshot is that we can write the counting as

$$\frac{1}{n!} \sum_{\gamma \in S_n} \sum_{\tau_1, \tau_2 \in S_n} \delta(\gamma \tau_1 \gamma^{-1} \tau_1^{-1}) \delta(\gamma \tau_2 \gamma^{-1} \tau_2^{-1}) \quad (4.4)$$

This is the counting of equivalence classes of pairs (τ_1, τ_2) , where the equivalence is generated by simultaneous conjugation by a permutation γ . This in turn is the same as counting branched covers of degree n from Riemann surfaces to a sphere with three branch points. After choosing a base point on the sphere, labelling the inverse images as $\{1, \dots, n\}$, and lifting paths going round the three branch points, we recover permutations $(\tau_1, \tau_2, (\tau_1 \tau_2)^{-1})$. The product of these three permutations is the identity, as it should be, since the path surrounding all three branch points is homotopic to the trivial path, with trivial monodromy. Generalizing the above arguments shows that counting of tensor invariants of rank d is equivalent to counting branched covers with d branch points. This is used to construct explicit generating functions in [28]. As in the use of permutations for multi-matrix theories, the application to tensor models also shows that finite N effects can be captured by performing the Fourier transform to representations.

Averaging over the equivalence classes defined by (4.2) and taking the products in $\mathbb{C}(S_n) \otimes \mathbb{C}(S_n) \otimes \mathbb{C}(S_n)$ leads to another important example of the permutation centralizer algebras defined in [24] (see concluding section there). The structure of this algebra is closely related to Kronecker product coefficients in S_n , forming a sort of categorification thereof.

5. Conclusions and Outlook

I have described a body of techniques based on permutations and diagrams to approach problems of enumerating gauge invariants and computing correlators in 1-matrix models, multi-matrix models, quiver gauge theories and tensor models. They are also useful in computing the spectrum of loop-corrected dilatation operators, particularly in the sector of perturbations around half-BPS giant gravitons (see [29, 30, 31, 32, 33] and references therein). Some progress has been made in finding the general quarter or eighth BPS ground states, which are ground states of the 1-loop dilatation operator acting on holomorphic invariants built from two or three complex matrices [34, 35]. The general solution at finite N remains elusive. One would like to construct the BPS operators in correspondence with states of a harmonic oscillator in two dimensions, as expected from quantization of giant graviton moduli spaces [36, 37]. Generalizations beyond unitary groups have been studied [38, 39].

It is noteworthy that the algebra $\mathbb{C}(S_n)$, in particular its centre, controls the n -dependent combinatorics of the orbifold theories. The $\mathcal{N} = (4, 4)$ superconformal theory on the orbifold $(T^4)^n/S_n$ is dual to $AdS_3 \times S^3$ (some papers on the relevant combinatorics are [1, 40, 41, 42, 43, 44]). This gives another instance where $\mathbb{C}(S_\infty)$ knows about the interactions of gravitons in AdS space, via holography. Elsewhere it has also been used in the classification of brane-tilings [45, 46, 47], the counting of Feynman graphs [48, 49], the systematic classification of light-cone string diagrams [50] and large N 2d Yang Mills theory [51, 52].

The algebra $\mathbb{C}(S_\infty)$, along with its representation theory and relations to Lie groups via Schur-Weyl duality, contains a lot of information about holography and quantum field theory. Further examples and a deeper understanding of this phenomenon would be fascinating.

Acknowledgments

I thank the organizers of the Corfu2015 workshop on Non-commutative Field Theory and Gravity for the opportunity to present this talk in September 2015 and the generous deadline extension allowed to complete the proceedings contribution. I thank the audience for stimulating questions and feedback. I have taken the liberty to add a few small points in this article which draw on papers that have appeared since, but which fit well with the content of the talk. I am also grateful to all my collaborators on the research presented here: Joseph Ben Geloun, Tom Brown, Stefan Cordes, Steve Corley, David Garner, Laurent Freidel, Amihay Hanany, Yang Hui He, Paul Heslop, Robert de Mello Koch, Antal Jevicki, Vishnu Jejjala, Yusuke Kimura, Paolo Mattioli, Mihail Mihailescu, Greg Moore, Jurgis Pasukonis, Diego Rodriguez-Gomez, Rak Kyeong Seong. My research is supported by STFC consolidated grant ST/L000415/1 “String Theory, Gauge Theory & Duality”

References

- [1] J. M. Maldacena, “The Large N limit of superconformal field theories and supergravity,” *Int. J. Theor. Phys.* **38** (1999) 1113 [*Adv. Theor. Math. Phys.* **2** (1998) 231] doi:10.1023/A:1026654312961 [hep-th/9711200].

- [2] M. Baggio, J. de Boer and K. Papadodimas, “A non-renormalization theorem for chiral primary 3-point functions,” JHEP **1207** (2012) 137 doi:10.1007/JHEP07(2012)137 [arXiv:1203.1036 [hep-th]].
- [3] S. Corley, A. Jevicki and S. Ramgoolam, “Exact correlators of giant gravitons from dual N=4 SYM theory,” Adv. Theor. Math. Phys. **5** (2002) 809 [hep-th/0111222].
- [4] S. Corley and S. Ramgoolam, “Finite factorization equations and sum rules for BPS correlators in N=4 SYM theory,” Nucl. Phys. B **641** (2002) 131 doi:10.1016/S0550-3213(02)00573-4 [hep-th/0205221].
- [5] J. Pasukonis and S. Ramgoolam, “Quivers as Calculators: Counting, Correlators and Riemann Surfaces,” JHEP **1304** (2013) 094 doi:10.1007/JHEP04(2013)094 [arXiv:1301.1980 [hep-th]].
- [6] R. d. M. Koch and S. Ramgoolam, “From Matrix Models and Quantum Fields to Hurwitz Space and the absolute Galois Group,” arXiv:1002.1634 [hep-th].
- [7] T. W. Brown, “Complex matrix model duality,” Phys. Rev. D **83** (2011) 085002 doi:10.1103/PhysRevD.83.085002 [arXiv:1009.0674 [hep-th]].
- [8] S. K. Lando, A. K. Zvonkin, "Graphs on Surfaces and Their Applications," Springer 2010.
- [9] V. Balasubramanian, M. Berkooz, A. Naqvi and M. J. Strassler, “Giant gravitons in conformal field theory,” JHEP **0204** (2002) 034 doi:10.1088/1126-6708/2002/04/034 [hep-th/0107119].
- [10] V. Balasubramanian, D. Berenstein, B. Feng and M. x. Huang, “D-branes in Yang-Mills theory and emergent gauge symmetry,” JHEP **0503** (2005) 006 doi:10.1088/1126-6708/2005/03/006 [hep-th/0411205].
- [11] R. de Mello Koch, J. Smolic and M. Smolic, “Giant Gravitons - with Strings Attached (I),” JHEP **0706** (2007) 074 doi:10.1088/1126-6708/2007/06/074 [hep-th/0701066].
- [12] R. de Mello Koch, J. Smolic and M. Smolic, “Giant Gravitons - with Strings Attached (II),” JHEP **0709** (2007) 049 doi:10.1088/1126-6708/2007/09/049 [hep-th/0701067].
- [13] D. Bekker, R. de Mello Koch and M. Stephanou, “Giant Gravitons - with Strings Attached. III.,” JHEP **0802** (2008) 029 doi:10.1088/1126-6708/2008/02/029 [arXiv:0710.5372 [hep-th]].
- [14] R. Bhattacharyya, S. Collins and R. d. M. Koch, “Exact Multi-Matrix Correlators,” JHEP **0803** (2008) 044 doi:10.1088/1126-6708/2008/03/044 [arXiv:0801.2061 [hep-th]].
- [15] Y. Kimura and S. Ramgoolam, “Enhanced symmetries of gauge theory and resolving the spectrum of local operators,” Phys. Rev. D **78** (2008) 126003 doi:10.1103/PhysRevD.78.126003 [arXiv:0807.3696 [hep-th]].
- [16] W. Fulton and J. Harris, "Representation theory: A First Course," Springer, 1991.
- [17] S. Ramgoolam, “Schur-Weyl duality as an instrument of Gauge-String duality,” AIP Conf. Proc. **1031** (2008) 255 doi:10.1063/1.2972012 [arXiv:0804.2764 [hep-th]].
- [18] J. Pasukonis and S. Ramgoolam, “Quivers as Calculators: Counting, Correlators and Riemann Surfaces,” JHEP **1304** (2013) 094 doi:10.1007/JHEP04(2013)094 [arXiv:1301.1980 [hep-th]].
- [19] B. Sundborg, “The Hagedorn transition, deconfinement and N=4 SYM theory,” Nucl. Phys. B **573** (2000) 349 doi:10.1016/S0550-3213(00)00044-4 [hep-th/9908001].
- [20] T. W. Brown, P. J. Heslop and S. Ramgoolam, “Diagonal free field matrix correlators, global symmetries and giant gravitons,” JHEP **0904** (2009) 089 doi:10.1088/1126-6708/2009/04/089 [arXiv:0806.1911 [hep-th]].

- [21] S. Collins, “Restricted Schur Polynomials and Finite N Counting,” *Phys. Rev. D* **79** (2009) 026002 doi:10.1103/PhysRevD.79.026002 [arXiv:0810.4217 [hep-th]].
- [22] T. W. Brown, P. J. Heslop and S. Ramgoolam, “Diagonal multi-matrix correlators and BPS operators in N=4 SYM,” *JHEP* **0802** (2008) 030 doi:10.1088/1126-6708/2008/02/030 [arXiv:0711.0176 [hep-th]].
- [23] Y. Kimura and S. Ramgoolam, “Branes, anti-branes and brauer algebras in gauge-gravity duality,” *JHEP* **0711** (2007) 078 doi:10.1088/1126-6708/2007/11/078 [arXiv:0709.2158 [hep-th]].
- [24] P. Mattioli and S. Ramgoolam, “Permutation Centralizer Algebras and Multi-Matrix Invariants,” *Phys. Rev. D* **93** (2016) no.6, 065040 doi:10.1103/PhysRevD.93.065040 [arXiv:1601.06086 [hep-th]].
- [25] P. Mattioli and S. Ramgoolam, “Quivers, Words and Fundamentals,” *JHEP* **1503** (2015) 105 doi:10.1007/JHEP03(2015)105 [arXiv:1412.5991 [hep-th]].
- [26] P. Mattioli and S. Ramgoolam, “Gauge Invariants and Correlators in Flavoured Quiver Gauge Theories,” arXiv:1603.04369 [hep-th].
- [27] R. Gurau and V. Rivasseau, “The 1/N expansion of colored tensor models in arbitrary dimension,” *Europhys. Lett.* **95** (2011) 50004 doi:10.1209/0295-5075/95/50004 [arXiv:1101.4182 [gr-qc]].
- [28] J. Ben Geloun and S. Ramgoolam, “Counting Tensor Model Observables and Branched Covers of the 2-Sphere,” arXiv:1307.6490 [hep-th].
- [29] V. De Comarmond, R. de Mello Koch and K. Jefferies, “Surprisingly Simple Spectra,” *JHEP* **1102** (2011) 006 doi:10.1007/JHEP02(2011)006 [arXiv:1012.3884 [hep-th]].
- [30] R. d. M. Koch, G. Mashile and N. Park, “Emergent Threebrane Lattices,” *Phys. Rev. D* **81** (2010) 106009 doi:10.1103/PhysRevD.81.106009 [arXiv:1004.1108 [hep-th]].
- [31] R. d. M. Koch, M. Dessein, D. Giataganas and C. Mathwin, “Giant Graviton Oscillators,” *JHEP* **1110** (2011) 009 doi:10.1007/JHEP10(2011)009 [arXiv:1108.2761 [hep-th]].
- [32] R. de Mello Koch and S. Ramgoolam, “A double coset ansatz for integrability in AdS/CFT,” *JHEP* **1206** (2012) 083 doi:10.1007/JHEP06(2012)083 [arXiv:1204.2153 [hep-th]].
- [33] N. Bornman, R. d. M. Koch and L. Tribelhorn, “Rotating Restricted Schur Polynomials,” arXiv:1602.05675 [hep-th].
- [34] J. Pasukonis and S. Ramgoolam, “From counting to construction of BPS states in N=4 SYM,” *JHEP* **1102** (2011) 078 doi:10.1007/JHEP02(2011)078 [arXiv:1010.1683 [hep-th]].
- [35] Y. Kimura, “Quarter BPS classified by Brauer algebra,” *JHEP* **1005** (2010) 103 doi:10.1007/JHEP05(2010)103 [arXiv:1002.2424 [hep-th]].
- [36] A. Mikhailov, “Giant gravitons from holomorphic surfaces,” *JHEP* **0011** (2000) 027 doi:10.1088/1126-6708/2000/11/027 [hep-th/0010206].
- [37] I. Biswas, D. Gaiotto, S. Lahiri and S. Minwalla, “Supersymmetric states of N=4 Yang-Mills from giant gravitons,” *JHEP* **0712** (2007) 006 doi:10.1088/1126-6708/2007/12/006 [hep-th/0606087].
- [38] P. Caputa, R. de Mello Koch and P. Diaz, “A basis for large operators in N=4 SYM with orthogonal gauge group,” *JHEP* **1303** (2013) 041 doi:10.1007/JHEP03(2013)041 [arXiv:1301.1560 [hep-th]].
- [39] P. Caputa, R. d. M. Koch and P. Diaz, *JHEP* **1306** (2013) 018 doi:10.1007/JHEP06(2013)018 [arXiv:1303.7252 [hep-th]].

- [40] J. de Boer, “Six-dimensional supergravity on $S^3 \times \text{AdS}(3)$ and 2-D conformal field theory,” Nucl. Phys. B **548** (1999) 139 doi:10.1016/S0550-3213(99)00160-1 [hep-th/9806104].
- [41] F. Larsen and E. J. Martinec, “U(1) charges and moduli in the D1 - D5 system,” JHEP **9906** (1999) 019 doi:10.1088/1126-6708/1999/06/019 [hep-th/9905064].
- [42] A. Jevicki, M. Mihailescu and S. Ramgoolam, “Gravity from CFT on $S^2 \times N(X)$: Symmetries and interactions,” Nucl. Phys. B **577** (2000) 47 doi:10.1016/S0550-3213(00)00147-4 [hep-th/9907144].
- [43] O. Lunin and S. D. Mathur, “Correlation functions for $M^N/S(N)$ orbifolds,” Commun. Math. Phys. **219** (2001) 399 doi:10.1007/s002200100431 [hep-th/0006196].
- [44] A. Pakman, L. Rastelli and S. S. Razamat, Phys. Rev. D **80** (2009) 086009 doi:10.1103/PhysRevD.80.086009 [arXiv:0905.3451 [hep-th]].
- [45] V. Jejjala, S. Ramgoolam and D. Rodriguez-Gomez, “Toric CFTs, Permutation Triples and Belyi Pairs,” JHEP **1103** (2011) 065 doi:10.1007/JHEP03(2011)065 [arXiv:1012.2351 [hep-th]].
- [46] A. Hanany, V. Jejjala, S. Ramgoolam and R. K. Seong, “Consistency and Derangements in Brane Tilings,” arXiv:1512.09013 [hep-th].
- [47] A. Hanany, Y. H. He, V. Jejjala, J. Pasukonis, S. Ramgoolam and D. Rodriguez-Gomez, JHEP **1106** (2011) 056 doi:10.1007/JHEP06(2011)056 [arXiv:1104.5490 [hep-th]].
- [48] R. de Mello Koch and S. Ramgoolam, “Strings from Feynman Graph counting : without large N,” Phys. Rev. D **85** (2012) 026007 doi:10.1103/PhysRevD.85.026007 [arXiv:1110.4858 [hep-th]].
- [49] R. de Mello Koch, S. Ramgoolam and C. Wen, “On the refined counting of graphs on surfaces,” Nucl. Phys. B **870** (2013) 530 doi:10.1016/j.nuclphysb.2013.01.023 [arXiv:1209.0334 [hep-th]].
- [50] L. Freidel, D. Garner and S. Ramgoolam, “Permutation combinatorics of worldsheet moduli space,” Phys. Rev. D **91** (2015) no.12, 126001 doi:10.1103/PhysRevD.91.126001 [arXiv:1412.3979 [hep-th]].
- [51] D. J. Gross and W. Taylor, “Two-dimensional QCD is a string theory,” Nucl. Phys. B **400** (1993) 181 doi:10.1016/0550-3213(93)90403-C [hep-th/9301068].
- [52] S. Cordes, G. W. Moore and S. Ramgoolam, “Large N 2-D Yang-Mills theory and topological string theory,” Commun. Math. Phys. **185** (1997) 543 doi:10.1007/s002200050102 [hep-th/9402107].