



A compactness theorem for complete Ricci shrinkers

MUELLER, R; Halshofer, R

© Springer Basel AG 2011

This is a pre-copyedited, author-produced PDF of an article accepted for publication in Geometric and Functional Analysis following peer review. The version of record is available <http://link.springer.com/article/10.1007/s00039-011-0137-4>

For additional information about this publication click this link.

<http://qmro.qmul.ac.uk/xmlui/handle/123456789/16128>

Information about this research object was correct at the time of download; we occasionally make corrections to records, please therefore check the published record when citing. For more information contact scholarlycommunications@qmul.ac.uk

A compactness theorem for complete Ricci shrinkers

Robert Haslhofer, Reto Müller

Abstract

We prove precompactness in an orbifold Cheeger-Gromov sense of complete gradient Ricci shrinkers with a lower bound on their entropy and a local integral Riemann bound. We do not need any pointwise curvature assumptions, volume or diameter bounds. In dimension four, under a technical assumption, we can replace the local integral Riemann bound by an upper bound for the Euler characteristic. The proof relies on a Gauss-Bonnet with cutoff argument.

1 Introduction

Let us start with some background: The classical Cheeger-Gromov theorem says that every sequence of closed Riemannian manifolds with uniformly bounded curvatures, volume bounded below, and diameter bounded above has a $C^{1,\alpha}$ -convergent subsequence [12, 21, 20]. The convergence is in the sense of Cheeger-Gromov, meaning $C^{1,\alpha}$ -convergence of the Riemannian metrics after pulling back by suitable diffeomorphisms. Without diameter bounds, the global volume bound should be replaced by a local volume non-collapsing assumption [14], and the appropriate notion of convergence is convergence in the pointed Cheeger-Gromov sense. If one can also control all the derivatives of the curvatures, e.g. in the presence of an elliptic or parabolic equation, the convergence is smooth [23]. To remind the reader about the precise definition, a sequence of complete smooth Riemannian manifolds with basepoints (M_i^n, g_i, p_i) converges to $(M_\infty^n, g_\infty, p_\infty)$ in the *pointed smooth Cheeger-Gromov sense* if there exist an exhaustion of M_∞ by open sets U_i containing p_∞ and smooth embeddings $\phi_i : U_i \rightarrow M_i$ with $\phi_i(p_\infty) = p_i$ such that the pulled back metrics $\phi_i^* g_i$ converge to g_∞ in C_{loc}^∞ .

Now, let us describe the problem under consideration: Hamilton's Ricci flow in higher dimensions without curvature assumptions leads to the formation of intriguingly complex singularities [22, 24]. The specific question we are concerned with is about the compactness properties of the corresponding space of singularity models. Namely, given a sequence of *gradient shrinkers*, i.e. a sequence of smooth, connected, complete Riemannian manifolds (M_i^n, g_i) satisfying

$$\text{Rc}_{g_i} + \text{Hess}_{g_i} f_i = \frac{1}{2} g_i \tag{1.1}$$

for some smooth function $f_i : M \rightarrow \mathbb{R}$ (called the potential), under what assumptions can we find a convergent subsequence? In the *compact case*, this problem was first studied by Cao-Sesum [8], see also Zhang [41], and Weber succeeded in removing their pointwise Ricci bounds [39]. We have profited from these previous works and the papers by Anderson, Bando, Kasue, Nakajima and Tian about the Einstein case [2, 29, 5, 33], as well as from the papers [3, 34, 35, 36, 37].

In this article, we generalize the shrinker orbifold compactness result to the case of *noncompact manifolds*. The obvious motivation for doing this is the fact that most interesting singularity models are noncompact, the cylinder being the most basic example. We manage to remove all volume and diameter assumptions, and we do not need any positivity assumptions for the curvatures nor pointwise curvature bounds (as the blow-down shrinker shows [18], even the Ricci curvature can have both signs). In fact, if the curvature is uniformly bounded below, it is easy to pass to a smooth limit (Theorem 2.5). The general case without positivity assumptions is much harder.

Having removed all other assumptions, we prove a precompactness theorem for complete Ricci shrinkers, assuming only a lower bound for the Perelman entropy and local $L^{n/2}$ bounds for the Riemann tensor (Theorem 1.1). The assumptions allow orbifold singularities to occur (these are isolated singularities modelled on \mathbb{R}^n/Γ for some finite subgroup $\Gamma \subset O(n)$), and the convergence is in the *pointed orbifold Cheeger-Gromov sense*. In particular, this means that the sequence converges in the pointed Gromov-Hausdorff sense (this is the natural notion of convergence for complete metric spaces), and that the convergence is in the smooth Cheeger-Gromov sense away from the isolated point singularities (see Section 3 for the precise definitions).

Our results are most striking in dimension four. In this case, the local L^2 Riemann bound is *not* an a priori assumption, but we prove it modulo a technical assumption on the soliton potential (Theorem 1.2). Our proof is based on a 4d-Chern-Gauss-Bonnet with cutoff argument (see Section 4). In particular, the key estimate of the cubic boundary term (Lemma 4.4) is based on a delicate use of partial integrations and soliton identities.

Before stating our main results, let us explain a few facts about gradient shrinkers, see Section 2 and Appendix A for proofs and further references. Associated to every gradient shrinker (M^n, g, f) , there is a family of Riemannian metrics $g(t)$, $t \in (-\infty, 1)$, evolving by Hamilton's Ricci flow $\frac{\partial}{\partial t}g(t) = -2\text{Rc}_{g(t)}$ with $g(0) = g$, which is self-similarly shrinking, i.e. $g(t) = (1-t)\phi_t^*g$ for the family of diffeomorphisms ϕ_t generated by $\frac{1}{1-t}\nabla f$, see [7, 42]. In this article however, we focus on the elliptic point of view. Gradient shrinkers always come with a natural basepoint, a point $p \in M$ where the potential f attains its minimum (such a minimum always exists and the distance between two minimum points is bounded by a constant depending only on

the dimension). The potential grows like one-quarter distance squared, so $2\sqrt{f}$ can be thought of as distance from the basepoint. Moreover, the volume growth is at most Euclidean, hence it is always possible to normalize f (by adding a constant) such that

$$\int_M (4\pi)^{-n/2} e^{-f} dV_g = 1. \quad (1.2)$$

Then the gradient shrinker has a well defined *entropy*,

$$\mu(g) = \mathcal{W}(g, f) = \int_M (|\nabla f|_g^2 + R_g + f - n)(4\pi)^{-n/2} e^{-f} dV_g > -\infty. \quad (1.3)$$

The entropy was introduced by Perelman in his famous paper [30] to solve the long standing problem of ruling out collapsing with bounded curvature (see [4, 9, 25, 28] for detailed expositions of Perelman's work). For general Ricci flows, the entropy is time-dependent, but on gradient shrinkers it is constant and finite (even without curvature assumptions). Assuming a lower bound for the entropy is natural, because it is non-decreasing along the Ricci flow in the compact case or under some mild technical assumptions. Under a local scalar curvature bound, a lower bound on the entropy gives a local volume non-collapsing bound.

The main results of this article are the following two theorems.

Theorem 1.1

Let (M_i^n, g_i, f_i) be a sequence of gradient shrinkers (with normalization and basepoint p_i as above) with entropy uniformly bounded below, $\mu(g_i) \geq \underline{\mu} > -\infty$, and uniform local energy bounds,

$$\int_{B_r(p_i)} |\text{Rm}_{g_i}|_{g_i}^{n/2} dV_{g_i} \leq E(r) < \infty, \quad \forall i, r. \quad (1.4)$$

Then a subsequence of (M_i^n, g_i, f_i, p_i) converges to an orbifold gradient shrinker in the pointed orbifold Cheeger-Gromov sense.

Here is a cute way to rephrase this theorem: *The space of Ricci flow singularity models with bounded entropy and locally bounded energy is orbifold compact.*

In the case $n = 4$, we obtain a particularly strong compactness result under a technical assumption on the potential.

Theorem 1.2

Let (M_i^4, g_i, f_i) be a sequence of four-dimensional gradient shrinkers (with normalization and basepoint p_i as above) with entropy uniformly bounded below, $\mu(g_i) \geq \underline{\mu} > -\infty$, Euler characteristic bounded above, $\chi(M_i) \leq \bar{\chi} < \infty$, and the technical assumption that the potentials do not have critical points at large distances, more precisely

$$|\nabla f_i|(x) \geq c > 0 \quad \text{if } d(x, p_i) \geq r_0, \quad (1.5)$$

for some constant $r_0 < \infty$. Then we have the weighted L^2 estimate

$$\int_{M_i} |\text{Rm}_{g_i}|_{g_i}^2 e^{-f_i} dV_{g_i} \leq C(\underline{\mu}, \bar{\chi}, c, r_0) < \infty. \quad (1.6)$$

In particular, the energy condition (1.4) is satisfied and by Theorem 1.1 a subsequence converges in the pointed orbifold Cheeger-Gromov sense.

As explained above, to appreciate our theorems it is most important to think about the assumptions that we do *not* make.

Remark. The technical assumption (1.5) is satisfied in particular if the scalar curvature satisfies

$$R_{g_i}(x) \leq \alpha d(x, p_i)^2 + C \quad (1.7)$$

for some $\alpha < \frac{1}{4}$. The scalar curvature grows at most like one-quarter distance squared and the average scalar curvature on $2\sqrt{f}$ -balls is bounded by $n/2$ (see Section 2 and Appendix A), so the technical assumption is rather mild. However, it would be very desirable to remove (or prove) it. Of course, (1.5) would also follow from a diameter bound.

In this article, the potentials of the gradient shrinkers play a central role in many proofs. In particular, we can view (a perturbation of) f as a Morse function, use e^{-f} as weight or cutoff function and use balls defined by the distance $2\sqrt{f}$ instead of the Riemannian distance. This has the great advantage, that we have a formula for the second fundamental form in the Gauss-Bonnet with boundary argument.

There are very deep and interesting other methods that yield comparable results, in particular the techniques developed by Cheeger-Colding-Tian in their work on the structure of spaces with Ricci curvature bounded below (see [13] for a nice survey) and the nested blowup and contradiction arguments of Chen-Wang [15]. Finally, let us mention the very interesting recent paper by Song-Weinkove [32].

This article is organized as follows. In Section 2, we collect and prove some properties of gradient shrinkers. In Section 3, we prove Theorem 1.1. Finally, we prove Theorem 1.2 in Section 4 using the Chern-Gauss-Bonnet theorem for manifolds with boundary and carefully estimating the boundary terms. We would like to point out that the Sections 3 and 4 are completely independent of each other and can be read in any order.

Acknowledgments: We greatly thank Tom Ilmanen for suggesting this problem. We also thank him and Carlo Mantegazza for very interesting discussions, and the anonymous referee for useful suggestions that greatly helped to improve the exposition. The first author was partially supported by the Swiss National Science Foundation, the research of the second author was supported by the Italian FIRB Ideas “Analysis and Beyond” and by The Leverhulme Trust.

2 Some properties of gradient shrinkers

Let us start by collecting some basic facts about gradient shrinkers (for a recent survey about Ricci solitons, see [6]). Tracing the soliton equation,

$$R_{ij} + \nabla_i \nabla_j f = \frac{1}{2} g_{ij}, \quad (2.1)$$

gives

$$R + \Delta f = \frac{n}{2}. \quad (2.2)$$

Using the contracted second Bianchi identity, inserting the soliton equation (2.1), and commuting the derivatives, we compute

$$\frac{1}{2} \nabla_i R = \nabla_i R - \nabla_j R_{ij} = -\nabla_i \nabla_j \nabla_j f + \nabla_j \nabla_i \nabla_j f = R_{ik} \nabla_k f. \quad (2.3)$$

As observed by Hamilton, from this formula and equation (2.1), it follows that

$$C_1(g) := R + |\nabla f|^2 - f \quad (2.4)$$

is constant (note that we always assume that our manifold is connected). By (2.1), the Hessian of f is uniquely determined by g . Thus, the potential has the form $f(x, y) = \tilde{f}(x) + \frac{1}{4}|y - y_0|^2$ after splitting $M \cong \tilde{M} \times \mathbb{R}^k$ isometrically. Note that the constant $C_1(g)$ and also the normalization (1.2) do not depend on the point $y_0 \in \mathbb{R}^k$. It follows that \tilde{f} is completely determined by fixing the normalization (1.2), and that $C_1(g)$ is independent of f after fixing this normalization. Gradient shrinkers always have nonnegative scalar curvature,

$$R \geq 0. \quad (2.5)$$

This follows from the elliptic equation

$$R + \langle \nabla f, \nabla R \rangle = \Delta R + 2|\text{Rc}|^2 \quad (2.6)$$

by the maximum principle, see [42] for a proof in the noncompact case without curvature assumptions. Equation (2.6) is the shrinker version of the evolution equation $\frac{\partial}{\partial t} R = \Delta R + 2|\text{Rc}|^2$ under Ricci flow.

The following two lemmas show, that the shrinker potential f grows like one-quarter distance squared and that gradient shrinkers have at most Euclidean volume growth.

Lemma 2.1 (Growth of the potential)

Let (M^n, g, f) be a gradient shrinker with $C_1 = C_1(g)$ as in (2.4). Then there exists a point $p \in M$ where f attains its infimum and f satisfies the quadratic growth estimate

$$\frac{1}{4}(d(x, p) - 5n)_+^2 \leq f(x) + C_1 \leq \frac{1}{4}(d(x, p) + \sqrt{2n})^2 \quad (2.7)$$

for all $x \in M$, where $a_+ := \max\{0, a\}$. If p_1 and p_2 are two minimum points, then their distance is bounded by

$$d(p_1, p_2) \leq 5n + \sqrt{2n}. \quad (2.8)$$

Lemma 2.2 (Volume growth)

There exists a constant $C_2 = C_2(n) < \infty$ such that every gradient shrinker (M^n, g, f) with $p \in M$ as in Lemma 2.1 satisfies the volume growth estimate

$$\text{Vol } B_r(p) \leq C_2 r^n. \quad (2.9)$$

The proofs are small but crucial improvements of the proofs by Cao-Zhou and Munteanu [10, 26]. In fact, their results are not strong enough for our purpose for which it is necessary, in particular, to remove the dependence on the geometry on a unit ball in Theorem 1.1 of [10] and to show that the constant in the volume growth estimate can be chosen uniformly for a sequence of shrinkers. In order to keep this section compact, we moved the proofs of both lemmas to Appendix A.

From now on, we fix a point $p \in M$ where f attains its minimum.

By Lemma 2.1 and Lemma 2.2, any function φ that satisfies the growth estimate

$$|\varphi(x)| \leq C e^{\alpha d(x,p)^2} \quad \text{for some } \alpha < \frac{1}{4} \quad (2.10)$$

is integrable with respect to the measure $e^{-f} dV$. In particular, the integral $\int_M e^{-f} dV$ is finite and f can be normalized (by adding a constant if necessary) to satisfy the normalization constraint (1.2).

From now on, we will fix the normalization (1.2).

Let us now explain the logarithmic Sobolev inequality, compare with Carrillo-Ni [11]. Any polynomial in $R, f, |\nabla f|, \Delta f$ is integrable with respect to the measure $e^{-f} dV$. Indeed, using one after another (2.5), $0 \leq |\nabla f|^2$, (2.4), and Lemma 2.1, we compute

$$0 \leq R(x) \leq R(x) + |\nabla f|^2(x) = f(x) + C_1 \leq \frac{1}{4}(d(x,p) + \sqrt{2n})^2, \quad (2.11)$$

and using furthermore (2.2) this implies

$$-\frac{n}{2} \leq -\Delta f(x) \leq -\frac{n}{2} + \frac{1}{4}(d(x,p) + \sqrt{2n})^2. \quad (2.12)$$

So, any polynomial in $R, f, |\nabla f|, \Delta f$ has at most polynomial growth and thus in particular satisfies the growth estimate (2.10). It follows that the entropy

$$\mu(g) := \mathcal{W}(g, f) = \int_M (|\nabla f|^2 + R + f - n)(4\pi)^{-n/2} e^{-f} dV \quad (2.13)$$

is well defined. To obtain another expression for the entropy, we will use the partial integration formula

$$\int_M \Delta f e^{-f} dV = \int_M |\nabla f|^2 e^{-f} dV, \quad (2.14)$$

which is justified as follows: Let $\eta_r(x) := \eta(d(x,p)/r)$, where $0 \leq \eta \leq 1$ is a cutoff function such that $\eta(s) = 1$ for $s \leq 1/2$ and $\eta(s) = 0$ for $s \geq 1$. Then

$$\int_M \eta_r \Delta f e^{-f} dV = \int_M \eta_r |\nabla f|^2 e^{-f} dV - \int_M \langle \nabla \eta_r, \nabla f \rangle e^{-f} dV.$$

Now, using the estimates for $f, |\nabla f|$ and the volume growth, we see that

$$\int_M |\nabla \eta_r| |\nabla f| e^{-f} dV \leq C \int_{B_r(p) \setminus B_{r/2}(p)} \frac{1}{r} d(x,p) e^{-\frac{(d(x,p)-5n)^2}{4}} dV \leq C r^n e^{-\frac{(r/2-5n)^2}{4}}$$

converges to zero for $r \rightarrow \infty$, and (2.14) follows from the dominated convergence theorem. Moreover, note that (2.2) and (2.4) imply the formula

$$2\Delta f - |\nabla f|^2 + R + f - n = -C_1. \quad (2.15)$$

Putting everything together, we conclude that

$$\mu(g) = \int_M (2\Delta f - |\nabla f|^2 + R + f - n) (4\pi)^{-n/2} e^{-f} dV = -C_1(g), \quad (2.16)$$

where we also used the normalization (1.2) in the last step. In other words, the *auxiliary constant* $C_1(g)$ of the gradient shrinker is minus the *Perelman entropy*. Carrillo-Ni made the wonderful observation that Perelman's logarithmic Sobolev inequality holds even for noncompact shrinkers without curvature assumptions [11, Thm. 1.1], i.e.

$$\inf \mathcal{W}(g, \tilde{f}) \geq \mu(g), \quad (2.17)$$

where the infimum is taken over all $\tilde{f} : M \rightarrow \mathbb{R} \cup \{+\infty\}$ such that $\tilde{u} = e^{-\tilde{f}/2}$ is smooth with compact support and $\int_M \tilde{u}^2 dV = (4\pi)^{n/2}$. Essentially, this follows from $\text{Rc}_f = \text{Rc} + \text{Hess } f \geq 1/2$ and the Bakry-Emery theorem [38, Thm. 21.2].

Remark. In fact, equality holds in (2.17), which can be seen as follows. First observe that, as a function of g and \tilde{u} ,

$$\mathcal{W}(g, \tilde{u}) = (4\pi)^{-n/2} \int_M (4|\nabla \tilde{u}|^2 + (R-n)\tilde{u}^2 - \tilde{u}^2 \log \tilde{u}^2) dV, \quad (2.18)$$

and that one can take the infimum over all properly normalized Lipschitz functions \tilde{u} with compact support. Now, the equality follows by approximating $u = e^{-f/2}$ by $\tilde{u}_r := C_r \eta_r u$, with η_r as above and with constants

$$C_r = \sqrt{\frac{(4\pi)^{n/2}}{\int_M \eta_r^2 u^2 dV}} \searrow 1 \quad (2.19)$$

to preserve the normalization. Indeed, arguing as before we see that

$$\begin{aligned} \int_M (R-n)\eta_r^2 u^2 &\rightarrow \int_M (R-n)u^2, & \int_M C_r^2 \eta_r^2 u^2 \log u^2 &\rightarrow \int_M u^2 \log u^2, \\ \int_M |\nabla(\eta_r u)|^2 &\rightarrow \int_M |\nabla u|^2, & \int_M C_r^2 \eta_r^2 \log(C_r^2 \eta_r^2) u^2 &\rightarrow 0, \end{aligned} \quad (2.20)$$

which together yields $\mathcal{W}(g, \tilde{u}_r) \rightarrow \mathcal{W}(g, u)$.

From Perelman's logarithmic Sobolev inequality (2.17) and the local bounds for the scalar curvature (2.11), we get the following non-collapsing lemma.

Lemma 2.3 (Non-collapsing)

There exists a function $\kappa(r) = \kappa(r, n, \underline{\mu}) > 0$ such that for every gradient shrinker (M^n, g, f) (with basepoint p and normalization as before) with entropy bounded below, $\mu(g) \geq \underline{\mu}$, we have the lower volume bound $\text{Vol } B_\delta(x) \geq \kappa(r)\delta^n$ for every ball $B_\delta(x) \subset B_r(p)$, $0 < \delta \leq 1$.

The proof is strongly related to Perelman's proof for finite-time Ricci flow singularities (see Kleiner-Lott [25, Sec. 13] for a nice and detailed exposition), and can be found in Appendix A. Given a lower bound $\mu(g) \geq \underline{\mu}$, we also get an upper bound $\mu(g) \leq \bar{\mu} = \bar{\mu}(\underline{\mu}, n)$ using $\tilde{u} = c^{-1/2}\eta(d(x, p))$ as test function. Of course, the conjecture is $\mu(g) \leq 0$ even for noncompact shrinkers without curvature assumptions.

Equipped with the above lemmas, we can now easily prove the non-collapsed pointed Gromov-Hausdorff convergence in the general case, and the pointed smooth Cheeger-Gromov convergence in the case where the curvature is uniformly bounded below.

Theorem 2.4 (Non-collapsed Gromov-Hausdorff convergence)

Let (M_i^n, g_i, f_i) be a sequence of gradient shrinkers with entropy uniformly bounded below, $\mu(g_i) \geq \underline{\mu} > -\infty$, and with basepoint p_i and normalization as before. Then the sequence is volume non-collapsed at finite distances from the basepoint and a subsequence (M_i, d_i, p_i) converges to a complete metric space in the pointed Gromov-Hausdorff sense.

Proof. The first part is Lemma 2.3. For the second part, to find a subsequence that converges in the pointed Gromov-Hausdorff sense, it suffices to find uniform bounds $N(\delta, r)$ for the number of disjoint δ -balls that fit within an r -ball centered at the basepoint [21, Prop. 5.2]. Assume $\delta \leq 1$ without loss of generality. By Lemma 2.2 the ball $B_r(p)$ has volume at most $C_2 r^n$, while by Lemma 2.3 each ball $B_\delta(x) \subset B_r(p)$ has volume at least $\kappa\delta^n$. Thus, there can be at most $N(\delta, r) = C_2 r^n / \kappa\delta^n$ disjoint δ -balls in $B_r(p)$. \square

Remark. Alternatively, the Gromov-Hausdorff convergence also follows from the volume comparison theorem of Wei-Wylie for the Bakry-Emery Ricci tensor [40], using the estimates for the soliton potential from this section. This holds even without entropy and energy bounds, but in that case the limit can be collapsed and very singular.

Theorem 2.5 (Smooth convergence in the curvature bounded below case)

Let (M_i^n, g_i, f_i) be a sequence of gradient shrinkers (with basepoint p_i and normalization as before) with entropy uniformly bounded below, $\mu(g_i) \geq \underline{\mu} > -\infty$, and curvature uniformly bounded below, $\text{Rm}_{g_i} \geq \underline{K} > -\infty$. Then a subsequence (M_i, g_i, f_i, p_i)

converges to a gradient shrinker $(M_\infty, g_\infty, f_\infty, p_\infty)$ in the pointed smooth Cheeger-Gromov sense (i.e. there exist an exhaustion of M_∞ by open sets U_i containing p_∞ and smooth embeddings $\phi_i : U_i \rightarrow M_i$ with $\phi_i(p_\infty) = p_i$ such that $(\phi_i^* g_i, \phi_i^* f_i)$ converges to (g_∞, f_∞) in C_{loc}^∞).

Proof. Recall the following Cheeger-Gromov compactness theorem from the very beginning of the introduction: For every sequence (M_i^n, g_i, p_i) of complete Riemannian manifolds with uniform local bounds for the curvatures and all its derivatives,

$$\sup_{B_r(p_i)} |\nabla^k \text{Rm}_{g_i}| \leq C_k(r), \quad (2.21)$$

and with a uniform local volume-noncollapsing bound around the basepoint,

$$\text{Vol}_{g_i} B_1(p_i) \geq \kappa, \quad (2.22)$$

we can find a subsequence that converges to a limit $(M_\infty, g_\infty, p_\infty)$ in the pointed smooth Cheeger-Gromov sense.

Moreover, if we have also uniform local bounds for the shrinker potential and all its derivatives,

$$\sup_{B_r(p_i)} |\nabla^k f_i| \leq C_k(r), \quad (2.23)$$

then by passing to another subsequence if necessary the functions $\phi_i^* f_i$ will also converge to some function f_∞ in C_{loc}^∞ (the embeddings $\phi_i : U_i \rightarrow M_i$ come from the pointed Cheeger-Gromov convergence). From the very definition of smooth convergence it is clear that the shrinker equation will pass to the limit, i.e. that $(M_\infty, g_\infty, f_\infty)$ will be a gradient shrinker. Thus, it remains to verify (2.21), (2.22), and (2.23) for our sequence of shrinkers.

By Lemma 2.3, we have uniform local volume non-collapsing, in particular condition (2.22) is satisfied. From (2.11) we have uniform local bounds for the scalar curvature, and together with the assumption $\text{Rm}_{g_i} \geq \underline{K}$ this gives uniform local Riemann bounds,

$$\sup_{B_r(p_i)} |\text{Rm}_{g_i}| \leq C_0(r). \quad (2.24)$$

From (2.11) and the bounds $\underline{\mu} \leq -C_1(g_i) \leq \bar{\mu}$, we get local C^1 bounds for f_i ,

$$\sup_{B_r(p_i)} |f_i| \leq C_0(r), \quad \sup_{B_r(p_i)} |\nabla f_i| \leq C_1(r). \quad (2.25)$$

Finally, by some very well known arguments, we can bootstrap the elliptic system

$$\begin{aligned} \Delta \text{Rm} &= \nabla f * \nabla \text{Rm} + \text{Rm} + \text{Rm} * \text{Rm}, \\ \Delta f &= \frac{n}{2} - R, \end{aligned} \quad (2.26)$$

starting from (2.24) and (2.25) to arrive at (2.21) and (2.23). Here, the second equation is just the traced soliton equation (2.2), while the first equation is obtained

from the soliton equation (2.1) and the Bianchi identity as follows:

$$\begin{aligned}
\nabla_p \nabla_p R_{ijkl} &= -\nabla_p \nabla_k R_{ijlp} - \nabla_p \nabla_\ell R_{ijpk} \\
&= -\nabla_k \nabla_p R_{ijlp} - \nabla_\ell \nabla_p R_{ijpk} + (\text{Rm} * \text{Rm})_{ijkl} \\
&= \nabla_k (\nabla_i R_{jl} - \nabla_j R_{il}) + \nabla_\ell (\nabla_j R_{ik} - \nabla_i R_{jk}) + (\text{Rm} * \text{Rm})_{ijkl} \\
&= \nabla_k (R_{jilp} \nabla_p f) + \nabla_\ell (R_{ijkp} \nabla_p f) + (\text{Rm} * \text{Rm})_{ijkl} \\
&= (\nabla f * \nabla \text{Rm} + \text{Rm} + \text{Rm} * \text{Rm})_{ijkl}.
\end{aligned} \tag{2.27}$$

Here, we used the Bianchi identity and the commutator rule in the first three lines and in the fourth and fifth line we used the soliton equation. This finishes the proof of the theorem. \square

Remark. The more interesting case without positivity assumptions is treated in Section 3. A related simple and well known example for singularity formation is the following. Consider the Eguchi-Hanson metric g_{EH} [17], a Ricci-flat metric on TS^2 which is asymptotic to $\mathbb{R}^4/\mathbb{Z}_2$ (remember that the unit tangent bundle of the 2-sphere is homeomorphic to S^3/\mathbb{Z}_2). Then $g_i := \frac{1}{i} g_{\text{EH}}$ is a sequence of Ricci-flat metrics, that converges to $\mathbb{R}^4/\mathbb{Z}_2$ in the orbifold Cheeger-Gromov sense. In particular, an orbifold singularity develops as the central 2-sphere (i.e. the zero section) shrinks to a point. For the positive Kähler-Einstein case, see Tian [33], in particular Theorem 7.1.

Remark. For a sequence of gradient shrinkers with entropy uniformly bounded below, by Lemma 2.1 and Lemma 2.2, $(4\pi)^{-n/2} e^{-f_i} dV_{g_i}$ is a sequence of uniformly tight probability measures. Thus, a subsequence of $(M_i, d_i, e^{-f_i} dV_{g_i}, p_i)$ converges to a pointed measured complete metric space $(M_\infty, d_\infty, \nu_\infty, p_\infty)$ in the pointed *measured* Gromov-Hausdorff sense. By (2.11), Lemma 2.1 and a Gromov-Hausdorff version of the Arzela-Ascoli theorem, there exists a continuous limit function $f_\infty : M_\infty \rightarrow \mathbb{R}$. It is an interesting question, if ν_∞ equals e^{-f_∞} times the Hausdorff measure.

Remark. It follows from the recent work of Lott-Villani and Sturm that the condition $\text{Rc}_f \geq 1/2$ is preserved in a weak sense [38].

3 Proof of orbifold Cheeger-Gromov convergence

In this section, we prove Theorem 1.1. For convenience of the reader, we will also explain some steps that are based on well known compactness techniques.

The structure of the proof is the following: First, we show that we have a uniform estimate for the local Sobolev constant (Lemma 3.2). Using this, we prove the ε -regularity Lemma 3.3, which says that we get uniform bounds for the curvatures on balls with small energy. We can then pass to a smooth limit away from locally finitely many singular points using in particular the energy assumption (1.4) and the ε -regularity lemma. This limit can be completed as a metric space by adding locally finitely many points. Finally, we prove that the singular points are of smooth

orbifold type.

We start by giving a precise definition of the convergence.

Definition 3.1 (Orbifold Cheeger-Gromov convergence)

A sequence of gradient shrinkers (M_i^n, g_i, f_i, p_i) converges to an orbifold gradient shrinker $(M_\infty^n, g_\infty, f_\infty, p_\infty)$ in the pointed orbifold Cheeger-Gromov sense, if the following properties hold.

1. There exist a locally finite set $S \subset M_\infty$, an exhaustion of $M_\infty \setminus S$ by open sets U_i and smooth embeddings $\phi_i : U_i \rightarrow M_i$, such that $(\phi_i^* g_i, \phi_i^* f_i)$ converges to (g_∞, f_∞) in C_{loc}^∞ on $M_\infty \setminus S$.
2. The maps ϕ_i can be extended to pointed Gromov-Hausdorff approximations yielding a convergence $(M_i, d_i, p_i) \rightarrow (M_\infty, d_\infty, p_\infty)$ in the pointed Gromov-Hausdorff sense.

Here, an orbifold gradient shrinker is a complete metric space that is a smooth gradient shrinker away from locally finitely many singular points. At a singular point q , M_∞ is modeled on \mathbb{R}^n/Γ for some finite subgroup $\Gamma \subset O(n)$ and there is an associated covering $\mathbb{R}^n \supset B_\rho(0) \setminus \{0\} \xrightarrow{\pi} U \setminus \{q\}$ of some neighborhood $U \subset M_\infty$ of q such that $(\pi^* g_\infty, \pi^* f_\infty)$ can be extended smoothly to a gradient shrinker over the origin.

For the arguments that follow, it will be very important to have a uniform local Sobolev constant that works simultaneously for all shrinkers in our sequence.

Lemma 3.2 (Estimate for the local Sobolev constant)

There exist $C_S(r) = C_S(r, n, \underline{\mu}) < \infty$ and $\delta_0(r) = \delta_0(r, n, \underline{\mu}) > 0$ such that for every gradient shrinker (M^n, g, f) (with basepoint p and normalization as before) with $\mu(g) \geq \underline{\mu}$, we have

$$\|\varphi\|_{L^{\frac{2n}{n-2}}} \leq C_S(r) \|\nabla \varphi\|_{L^2} \quad (3.1)$$

for all balls $B_\delta(x) \subset B_r(p)$, $0 < \delta \leq \delta_0(r)$ and all functions $\varphi \in C_c^1(B_\delta(x))$.

Proof. The main point is that the estimate for the local Sobolev constant will follow from the noncollapsing and the volume comparison for the Bakry-Emery Ricci tensor. The detailed argument goes as follows:

The first reduction is that it suffices to control the optimal constant $C_1(B)$ in the L^1 -Sobolev inequality,

$$\|\psi\|_{L^{\frac{n}{n-1}}(B)} \leq C_1(B) \|\nabla \psi\|_{L^1(B)}, \quad (3.2)$$

for all $\psi \in C_c^1(B)$, where in our case B will always be an open ball in a Riemannian manifold. Indeed, applying (3.2) for $\psi = \varphi^{(2n-2)/(n-2)}$ and using Hölder's inequality

we can compute

$$\begin{aligned} \left(\|\varphi\|_{L^{\frac{2n}{n-2}}(B)} \right)^{\frac{2n-2}{n-2}} &\leq \frac{2n-2}{n-2} C_1(B) \|\varphi^{n/(n-2)} \nabla \varphi\|_{L^1(B)} \\ &\leq \frac{2n-2}{n-2} C_1(B) \left(\|\varphi\|_{L^{\frac{2n}{n-2}}(B)} \right)^{\frac{n}{n-2}} \|\nabla \varphi\|_{L^2(B)}, \end{aligned}$$

so the L^1 -Sobolov inequality (3.2) implies the L^2 -Sobolov inequality

$$\|\varphi\|_{L^{\frac{2n}{n-2}}(B)} \leq C_2(B) \|\nabla \varphi\|_{L^2(B)}, \quad (3.3)$$

with $C_2(B) = \frac{2n-2}{n-2} C_1(B)$. Next, it is a classical fact, known under the name Federer-Fleming theorem, that the optimal constant $C_1(B)$ in (3.2) is equal to the optimal constant $C_I(B)$ in the isoperimetric inequality,

$$|\Omega|^{\frac{n-1}{n}} \leq C_I(B) |\partial\Omega|, \quad (3.4)$$

for all regions $\Omega \Subset B$ with C^1 -boundary. Third, by a theorem of Croke [16, Thm. 11], the isoperimetric constant can be estimated by

$$C_I(B) \leq C(n) \omega(B)^{-\frac{n+1}{n}}, \quad (3.5)$$

where $C(n) < \infty$ is an explicit constant whose value we do not need and $\omega(B)$ is the visibility angle defined as

$$\omega(B) = \inf_{y \in B} |U_y| / |S^{n-1}|, \quad (3.6)$$

where $U_y = \{v \in T_y B ; |v| = 1, \text{ the geodesic } \gamma_v \text{ is minimizing up to } \partial B\}$.

Putting everything together, to finish the proof of our lemma it suffices to find a lower bound for the visibility angle (3.6) for $B = B_\delta(x) \subset B_r(p)$ inside a shrinker for $\delta \leq \delta_0(r)$, where $\delta_0(r)$ will be chosen sufficiently small later. To find such a lower bound, we will use the volume comparison theorem for the Bakry-Emery Ricci tensor due to Wei-Wylie [40] which we now explain:

Fix $y \in M$, use exponential polar coordinates around y and write $dV = \mathcal{A}(r, \theta) dr \wedge d\theta$ for the volume element, where $d\theta$ is the standard volume element on the unit sphere S^{n-1} . Let $\mathcal{A}_f(r, \theta) = \mathcal{A}(r, \theta) e^{-f}$. Note that $\text{Rc}_f = \text{Rc} + \nabla^2 f \geq 0$ by the soliton equation. The angular version of the volume comparison theorem for the Bakry-Emery Ricci tensor [40, Thm 1.2a] says that if in addition $|\nabla f| \leq a$ on $B_R(y)$ then

$$\frac{\int_0^R \mathcal{A}_f(s, \theta) ds}{\int_0^r \mathcal{A}_f(s, \theta) ds} \leq e^{aR} \left(\frac{R}{r} \right)^n. \quad (3.7)$$

for $0 < r \leq R$. If we have moreover $\max_{B_R(y)} f \leq \min_{B_R(y)} f + b$ then this implies

$$\int_0^R \mathcal{A}(s, \theta) ds \leq e^{aR+b} \left(\frac{R}{r} \right)^n \int_0^r \mathcal{A}(s, \theta) ds, \quad (3.8)$$

and finally by sending r to zero we obtain the form of the volume comparison estimate that we will actually use, namely

$$\int_0^R \mathcal{A}(s, \theta) ds \leq \frac{1}{n} e^{aR+b} R^n. \quad (3.9)$$

In our application everything will stay inside a ball $B_{r+1}(p)$ around the basepoint of the soliton, so by (2.11) we can take $a := \frac{1}{2}(r+1 + \sqrt{2n})$ and $b := a^2$.

Now, coming back to actually estimating the visibility angle of $B_\delta(x) \subset B_r(p)$, we let $y \in B_\delta(x)$ and apply the above ideas. Since the volume is computed using exponential polar coordinates around y , we have the estimate

$$|B_1(x)| - |B_\delta(x)| \leq \int_{U_y} \int_0^{1+\delta} \mathcal{A}(s, \theta) ds d\theta, \quad (3.10)$$

where U_y denotes the set of unit tangent vectors in whose direction the geodesics are minimizing up to the boundary of $B_\delta(x)$. Using (3.9), we can estimate this by

$$|B_1(x)| - |B_\delta(x)| \leq \frac{1}{n} e^{2a+b} |U_y| (1+\delta)^n, \quad (3.11)$$

and minimizing over $y \in B_\delta(x)$ we obtain the inequality

$$|B_1(x)| - |B_\delta(x)| \leq C \omega(B_\delta(x)) (1+\delta)^n \quad (3.12)$$

with $C = C(r, n) = \frac{1}{n} e^{2a+b} |S^{n-1}|$. Moreover, using (3.9) again, we obtain the upper bound

$$|B_\delta(x)| \leq \int_{S^{n-1}} \int_0^\delta \mathcal{A}(s, \theta) ds d\theta \leq C \delta^n. \quad (3.13)$$

Finally, we have the lower volume bound

$$|B_1(x)| \geq \kappa, \quad (3.14)$$

for $\kappa = \kappa(r+1, n, \underline{\mu})$ from Lemma 2.3. If we now choose $\delta_0 = \delta_0(r, n, \underline{\mu}) = (\kappa/2C)^{1/n}$, then putting together (3.12), (3.13) and (3.14) gives the lower bound

$$\omega(B_\delta(x)) \geq \frac{\kappa}{2^{n+1}C} \quad (3.15)$$

for the visibility angle for $\delta \leq \delta_0$, and this finishes the proof of the lemma. \square

Using the uniform estimate for the local Sobolev constant, we obtain the following ε -regularity lemma.

Lemma 3.3 (ε -regularity)

There exist $\varepsilon_1(r) = \varepsilon_1(r, n, \underline{\mu}) > 0$ and $K_\ell(r) = K_\ell(r, n, \underline{\mu}) < \infty$ such that for every gradient shrinker (M^n, g, f) (with basepoint p and normalization as before) with $\mu(g) \geq \underline{\mu}$ and for every ball $B_\delta(x) \subset B_r(p)$, $0 < \delta \leq \delta_0(r)$, we have the implication

$$\|\text{Rm}\|_{L^{n/2}(B_\delta(x))} \leq \varepsilon_1(r) \Rightarrow \sup_{B_{\delta/4}(x)} |\nabla^\ell \text{Rm}| \leq \frac{K_\ell(r)}{\delta^{2+\ell}} \|\text{Rm}\|_{L^{n/2}(B_\delta(x))}. \quad (3.16)$$

Proof. The gradient shrinker version of the evolution equation of the Riemann tensor under Ricci flow, $\frac{\partial}{\partial t} \text{Rm} = \Delta \text{Rm} + Q(\text{Rm})$, is the elliptic equation

$$\Delta \text{Rm} = \nabla f * \nabla \text{Rm} + \text{Rm} + \text{Rm} * \text{Rm}. \quad (3.17)$$

Here, we used (2.1) to eliminate $\nabla^2 f$ in $L_{\nabla f} \text{Rm} = \nabla f * \nabla \text{Rm} + \nabla^2 f * \text{Rm}$. Alternatively, we have given another derivation of the elliptic equation (3.17) in (2.27). Now, we set $u := |\text{Rm}|$ and compute

$$\begin{aligned} -u\Delta u &= -\frac{1}{2}\Delta u^2 + |\nabla u|^2 \\ &= -\frac{1}{2}\Delta |\text{Rm}|^2 + |\nabla |\text{Rm}||^2 \\ &= -\langle \text{Rm}, \Delta \text{Rm} \rangle - |\nabla \text{Rm}|^2 + |\nabla |\text{Rm}||^2. \end{aligned} \quad (3.18)$$

By equation (3.17) and Young's inequality, we can estimate

$$\begin{aligned} -\langle \text{Rm}, \Delta \text{Rm} \rangle &\leq C_3 (|\text{Rm}| |\nabla f| |\nabla \text{Rm}| + |\text{Rm}|^2 + |\text{Rm}|^3) \\ &\leq \frac{1}{10} |\nabla \text{Rm}|^2 + (1 + \frac{10}{4} C_3 |\nabla f|^2) C_3 |\text{Rm}|^2 + C_3 |\text{Rm}|^3, \end{aligned} \quad (3.19)$$

for some constant $C_3 = C_3(n) < \infty$ depending only on the dimension. Finally we use Kato's inequality $|\nabla |\text{Rm}|| \leq |\nabla \text{Rm}|$, and the estimate (2.11) for $|\nabla f|$. Putting everything together, we obtain the elliptic inequality

$$-u\Delta u \leq \frac{1}{10} |\nabla u|^2 + C_4 u^2 + C_3 u^3 \quad (3.20)$$

on $B_r(p)$, where $C_4 = C_4(r, n) := (1 + \frac{5}{8} C_3 (r + \sqrt{2n})^2) C_3$. Given an elliptic inequality like (3.20) it is well known to PDE-experts that if the $L^{n/2}$ -norm of u is sufficiently small on a ball, then one gets L^∞ -bounds on a smaller ball, more precisely

$$\|u\|_{L^{n/2}(B_\delta(x))} \leq \varepsilon \Rightarrow \sup_{B_{\delta/2}(x)} |u| \leq \frac{K}{\delta^2} \|u\|_{L^{n/2}(B_\delta(x))}, \quad (3.21)$$

for some constants $\varepsilon > 0$ and $K < \infty$. For convenience of the reader, we sketch the necessary Moser-iteration argument here: To keep the notation reasonably concise let us assume $\delta = 1$ and $n = 4$, the general case works similarly. Choose a cutoff-function $0 \leq \eta \leq 1$ that is 1 on $B_{3/4}(x)$, has support in $B_1(x)$, and satisfies $|\nabla \eta| \leq 8$. Multiplying (3.20) by η^2 and integrating by parts we obtain

$$\frac{9}{10} \int_M \eta^2 |\nabla u|^2 dV \leq 2 \int_M \eta |\nabla \eta| u |\nabla u| dV + \int_M (C_4 \eta^2 u^2 + C_3 \eta^2 u^3) dV. \quad (3.22)$$

Dealing with the first term on the right hand side by Young's inequality and absorption, this gives the estimate

$$\frac{1}{2} \int_M \eta^2 |\nabla u|^2 dV \leq (C_4 + 160) \int_{B_1} u^2 dV + C_3 \int_M \eta^2 u^3 dV. \quad (3.23)$$

For the last term, using Hölder's inequality, the assumption that the energy on B_1 is less than ε , and the Sobolev-inequality, we get

$$\begin{aligned} \int_M \eta^2 u^3 dV &\leq \left(\int_{B_1} u^2 dV \right)^{1/2} \left(\int_M (\eta u)^4 dV \right)^{1/2} \\ &\leq \varepsilon C_S^2 \int_M |\nabla(\eta u)|^2 dV \\ &\leq 2\varepsilon C_S^2 \int_M \eta^2 |\nabla u|^2 dV + 50\varepsilon C_S^2 \int_{B_1} u^2 dV, \end{aligned} \quad (3.24)$$

where $C_S < \infty$ is the local Sobolev constant on B_1 . The main idea is that if we choose ε so small that $2\varepsilon C_S^2 C_3 \leq \frac{1}{4}$ then the $\int \eta^2 |\nabla u|^2$ term can be absorbed, giving

$$\frac{1}{4} \int_M \eta^2 |\nabla u|^2 dV \leq (C_4 + 200) \int_{B_1} u^2 dV, \quad (3.25)$$

and using the Sobolev inequality we arrive at the L^4 -estimate

$$\|u\|_{L^4(B_{3/4})} \leq 2C_S \sqrt{C_4 + 200} \|u\|_{L^2(B_1)}. \quad (3.26)$$

Now we choose a sequence of radii $r_k = \frac{1}{2} + \frac{1}{2^k}$ interpolating between $r_1 = 1$ and $r_\infty = \frac{1}{2}$. We multiply (3.20) by $\eta_k^2 u^{p_k}$, where $p_k = 2^k - 2$, and $0 \leq \eta_k \leq 1$ is a cutoff function that equals 1 on $B_{r_{k+1}}$, has support in B_{r_k} , and satisfies $|\nabla \eta_k| \leq 2/(r_k - r_{k+1})$. Carrying out similar steps as above we obtain the iterative estimates

$$\|u\|_{L^{2^{k+1}}(B_{r_{k+1}})} \leq C_k \|u\|_{L^{2^k}(B_{r_k})}. \quad (3.27)$$

The product of the constants C_k converges and sending $k \rightarrow \infty$ gives the desired estimate

$$\|u\|_{L^\infty(B_{1/2})} \leq K \|u\|_{L^2(B_1)}. \quad (3.28)$$

Note that the estimate (3.21) is of course only useful if we can get uniform constants $\varepsilon > 0$ and $K < \infty$ for our sequence of shrinkers. This crucial point is taken care of by Lemma 3.2, so we indeed get constants $\varepsilon_1(r) = \varepsilon_1(r, n, \underline{\mu}) > 0$ and $K_0(r) = K_0(r, n, \underline{\mu}) < \infty$, such that

$$\|\text{Rm}\|_{L^{n/2}(B_\delta(x))} \leq \varepsilon_1(r) \Rightarrow \sup_{B_{\delta/2}(x)} |\text{Rm}| \leq \frac{K_0(r)}{\delta^2} \|\text{Rm}\|_{L^{n/2}(B_\delta(x))}, \quad (3.29)$$

for every ball $B_\delta(x) \subset B_r(p)$, $0 < \delta \leq \delta_0(r)$. Once one has L^∞ control, the hard work is done and it is standard to bootstrap the elliptic equation (3.17) to get C^∞ bounds on the ball $B_{\delta/4}(x)$. The only slightly subtle point is that higher derivatives of f appear when differentiating (3.17), but one can get rid of them again immediately using the soliton equation (2.1). This finishes the proof of the ε -regularity lemma. \square

Let us now explain how to finish the proof of Theorem 1.1. Let (M_i^n, g_i, f_i) be a sequence of gradient shrinkers satisfying the assumptions of Theorem 1.1. By Theorem 2.4, we can assume (after passing to a subsequence) that the sequence converges in the pointed Gromov-Hausdorff sense. By passing to another subsequence, we can also assume that the auxiliary constants converge.

Let $r < \infty$ large and $0 < \delta \leq \delta_1(r, E(r), n, \underline{\mu})$ small enough. The assumption (1.4) gives a uniform bound $E(r)$ for the energy contained in $B_r(p_i)$. So there can be at most $\frac{E_1(r)}{\varepsilon_1(r)}$ disjoint δ -balls in $B_r(p_i)$ that contain energy more than $\varepsilon_1(r)$, and away from those bad balls we get C^∞ -estimates for the curvatures using the ε -regularity lemma. Recall that we also have volume-noncollapsing by Lemma 2.3, and that we get C_{loc}^∞ bounds for f_i in regions with bounded curvature, using the elliptic equation

$$\Delta f = \frac{n}{2} - R. \quad (3.30)$$

Thus, putting everything together (and playing around with the parameters r and δ a bit), we can find on any (M_i, g_i) suitable balls $B_\delta(x_i^k(\delta))$, $1 \leq k \leq L_i(r) \leq L(r) = L(r, E(r), n, \underline{\mu})$, such that on

$$X_i := B_r(p_i) \setminus \bigcup_{k=1}^{L_i(r)} B_\delta(x_i^k(\delta)) \subset M_i, \quad (3.31)$$

we have the estimates

$$\begin{aligned} \sup_{X_i} |\nabla^\ell \text{Rm}_{g_i}| &\leq C_\ell(\delta, r, n, \underline{\mu}) \\ \sup_{X_i} |\nabla^\ell f_i| &\leq C_\ell(\delta, r, n, \underline{\mu}). \end{aligned} \quad (3.32)$$

Together with the volume-noncollapsing, this is exactly what we need to pass to a smooth limit. Thus, sending $r \rightarrow \infty$ and $\delta \rightarrow 0$ suitably and passing to a diagonal subsequence, we obtain a (possibly incomplete) smooth limit gradient shrinker. Since we already know, that the manifolds M_i converge in the pointed Gromov-Hausdorff sense, this limit can be completed as a metric space by adding locally finitely many points and the convergence is in the sense of Definition 3.1. We have thus proved Theorem 1.1 up to the statement that the isolated singular points are of orbifold shrinker type.

This claimed orbifold structure at the singular points is a local statement, so we can essentially refer to [8, 41]. Nevertheless, let us sketch the main steps, following Tian [33, Sec. 3 and 4] closely (see also [2, 5] for similar proofs).

Step 1 (C^0 -multifold): The idea is that blowing up around a singular point will show that the tangent cone is a union of finitely many flat cones over spherical space forms. Improving this a bit, one also gets C^0 -control over g , and thus the structure of a so called C^0 -multifold (“multi” and not yet “orbi”, since we have to wait until the

next step to see that “a union of finitely many flat cones over spherical space forms” can actually be replaced by “a single flat cone over a spherical space form”). The precise argument goes as follows: Near an added point $q \in S \subset M_\infty$, we have

$$|\nabla^\ell \text{Rm}_{g_\infty}|_{g_\infty}(x) \leq \frac{\varepsilon(\varrho(x))}{\varrho(x)^{2+\ell}}, \quad (3.33)$$

where $\varrho(x) = d_\infty(x, q)$. Here and in the following, $\varepsilon(\varrho)$ denotes a quantity that tends to zero for $\varrho \rightarrow 0$ and we always assume that ϱ is small enough. With $\varepsilon(\varrho) \rightarrow 0$ in (3.33), together with the Bakry-Emery volume comparison and the non-collapsing, it follows that the tangent cone at q is a finite union of flat cones over spherical space forms S^{n-1}/Γ_β . The tangent cone is unique and by a simple volume argument we get an explicit bound (depending on $r, n, \underline{\mu}$) for the order of the orbifold groups Γ_β and the number of components $\#\{\beta\}$. As in [33, Lemma 3.6, Eq. (4.1)] there exist a neighborhood $U \subset M_\infty$ of q and for every component U_β of $U \setminus \{q\}$ an associated covering $\pi_\beta : B_\varrho^* = B_\varrho(0) \setminus \{0\} \rightarrow U_\beta$ such that $g^\beta := \pi_\beta^* g_\infty$ can be extended to a C^0 -metric over the origin with the estimates

$$\begin{aligned} \sup_{B_\varrho^*} |g^\beta - g_E|_{g_E} &\leq \varepsilon(\varrho), \\ |D^I g^\beta|_{g_E}(x) &\leq \frac{\varepsilon(\varrho(x))}{\varrho(x)^{|I|}}, \quad x \in B_\varrho^*, \quad 1 \leq |I| \leq 100, \end{aligned} \quad (3.34)$$

where g_E is the Euclidean metric, D the Euclidean derivative and I a multiindex.

Step 2 (C^0 -orbifold): The idea is that if $U \setminus \{q\}$ had two or more components, than all geodesics in an approximating sequence would pass through a very small neck, but this yields a contradiction to the volume comparison theorem. For the precise argument, let $q \in S \subset M_\infty$ be an added point and choose points $x_i \in M_i$ converging to q . By the non-collapsing and the Bakry-Emery volume comparison with the bounds for f_i and $|\nabla f_i|$, there exists a constant $C < \infty$ such that for ϱ small enough any two points in $\partial B_\varrho(x_i)$ can be connected by a curve in $B_\varrho(x_i) \setminus B_{\varrho/C}(x_i)$ of length less than $C\varrho$. This follows by slightly modifying the proof of [3, Lemma 1.2] and [1, Lemma 1.4]. Since the convergence is smooth away from q , it follows that $U \setminus \{q\}$ is connected. In particular, the tangent cone at q consists of a single flat cone over a spherical space form S^{n-1}/Γ .

Step 3 (C^∞ -orbifold): The final step is to get C^∞ bounds in suitable coordinates. Let g^1 be the metric on B_ϱ^* from Step 1 and 2. In the case $n = 4$, using Uhlenbeck’s method for removing finite energy point singularities in the Yang-Mills field [37], we get an improved curvature decay

$$|\text{Rm}_{g^1}|_{g^1}(x) \leq \frac{1}{|x|^\delta} \quad (3.35)$$

for $\delta > 0$ as small as we want on a small enough punctured ball B_ϱ^* . The proof goes through almost verbatim as in [33, Lemma 4.3]. The only difference is that instead

of the Yang-Mills equation we use

$$\nabla_i R_{ijkl} = \nabla_k R_{lj} - \nabla_l R_{kj} = -\nabla_k \nabla_l \nabla_j f + \nabla_l \nabla_k \nabla_j f = R_{\ell k j p} \nabla_p f \quad (3.36)$$

and the estimates for $|\nabla f|$ from Section 2. The point is that the Yang-Mills equation $\nabla_i R_{ijkl} = 0$ is satisfied up to some lower order term that can be dealt with easily. The case $n \geq 5$ is more elementary, and Sibner's test function [31] gives L^∞ bounds for the curvature, in particular (3.35) is also satisfied in this case.

Due to the improved curvature decay, there exists a diffeomorphism $\psi : B_{\varrho/2}^* \rightarrow \psi(B_{\varrho/2}^*) \subset B_\varrho^*$ that extends to a homeomorphism over the origin such that $\psi^* g^1$ extends to a $C^{1,\alpha}$ metric over the origin (for any $\alpha < 1 - \delta$). By composing with another diffeomorphism (denoting the composition by φ), we can assume that the standard coordinates on $B_{\varrho/4}$ are harmonic coordinates for $\varphi^* g^1$. Finally, let $\pi := \pi_1 \circ \varphi$, $g := \pi^* g_\infty$ and $f := \pi^* f_\infty$. Then, for (g, f) we have the elliptic system

$$\begin{aligned} \Delta_g f &= |\nabla f|_g^2 - f + \frac{n}{2} - C, \\ \text{Rc}_g &= \frac{1}{2}g - \text{Hess}_g f. \end{aligned} \quad (3.37)$$

This is indeed elliptic, since $R_{ij}(g) = -\frac{1}{2} \sum_k \partial_k \partial_k g_{ij} + Q_{ij}(g, \partial g)$ in harmonic coordinates. It is now standard to bootstrap (3.37) starting with the $C^{1,\alpha}$ -bound for g and the $C^{0,1}$ -bound for f to obtain C^∞ -bounds for (g, f) and to conclude that (g, f) can be extended to a smooth gradient shrinker over the origin. This finishes the proof of Theorem 1.1.

Remark. Every added point is a singular point. Indeed, suppose Γ is trivial and $K_i = |\text{Rm}_{g_i}|_{g_i}(x_i) = \max_{B_\varrho(x_i)} |\text{Rm}_{g_i}|_{g_i} \rightarrow \infty$, $x_i \rightarrow q$ for some subsequence. Then, a subsequence of $(M_i, K_i g_i, x_i)$ converges to a non-flat, Ricci-flat manifold with the same volume ratios as in Euclidean space, a contradiction.

Remark. As discovered by Anderson [2], one can use the following two observations to rule out or limit the formation of singularities a priori: For n odd, $\mathbb{R}P^{n-1}$ is the only nontrivial spherical space form and it does not bound a smooth compact manifold. For $n = 4$, every nontrivial orientable Ricci-flat ALE manifold has nonzero second Betti number.

4 A local Chern-Gauss-Bonnet argument

In this section, we prove Theorem 1.2. To explain and motivate the Gauss-Bonnet with cutoff argument, we will first prove a weaker version (Proposition 4.3).

Recall from Section 2 that f grows like one-quarter distance squared, that R and $|\nabla f|^2$ grow at most quadratically, and that the volume growth is at most Euclidean. These growth estimates will be used frequently in the following.

The next lemma, first observed by Munteanu-Sesum [27], will be very useful in the following. To keep this section self-contained, we give a quick proof.

Lemma 4.1 (Weighted L^2 estimate for Ricci)

For $\lambda > 0$ and $\underline{\mu} > -\infty$ there exist constants $C(\lambda) = C(\lambda, \underline{\mu}, n) < \infty$ such that for every gradient shrinker (M^n, g, f) with $\mu(g) \geq \underline{\mu}$ and normalization as before,

$$\int_M |\text{Rc}|^2 e^{-\lambda f} dV \leq C(\lambda) < \infty. \quad (4.1)$$

Proof. Take a cutoff function η as in Section 2 and set $\eta_r(x) = \eta(d(x, p)/r)$. Note that $\text{div}(e^{-f} \text{Rc}) = 0$ by (2.3). Using this, the soliton equation, a partial integration and the inequality $ab \leq a^2/4 + b^2$, we compute

$$\begin{aligned} \int_M \eta_r^2 |\text{Rc}|^2 e^{-\lambda f} dV &= \int_M \eta_r^2 \langle \tfrac{1}{2}g - \nabla^2 f, \text{Rc} \rangle e^{-\lambda f} dV \\ &= \int_M \left(\tfrac{1}{2} \eta_r^2 R + (1 - \lambda) \eta_r^2 \text{Rc}(\nabla f, \nabla f) + 2\eta_r \text{Rc}(\nabla \eta_r, \nabla f) \right) e^{-\lambda f} dV \\ &\leq \tfrac{1}{2} \int_M \eta_r^2 |\text{Rc}|^2 e^{-\lambda f} dV + \int_M \eta_r^2 \left(\tfrac{1}{2} R + (1 - \lambda)^2 |\nabla f|^4 \right) e^{-\lambda f} dV \\ &\quad + 4 \int_M |\nabla \eta_r|^2 |\nabla f|^2 e^{-\lambda f} dV. \end{aligned}$$

The first term can be absorbed. The second term is uniformly bounded and the last term converges to zero as $r \rightarrow \infty$ by the growth estimates from Section 2. \square

As a consequence of Lemma 4.1, we can replace the Riemann energy bound in Theorem 1.1 by a Weyl energy bound in dimension four.

Corollary 4.2 (Weyl implies Riemann energy condition)

Every sequence of 4-dimensional gradient shrinkers (M_i, g_i, f_i) (with normalization and basepoint as usual) with entropy bounded below, $\mu(g_i) \geq \underline{\mu}$, and a local Weyl energy bound

$$\int_{B_r(p_i)} |W_{g_i}|_{g_i}^2 dV_{g_i} \leq C(r) < \infty, \quad \forall i, r \quad (4.2)$$

satisfies the energy condition (1.4).

Remark. As a consistency check, note that in dimension $n = 3$, Rm is determined by Rc and thus only a lower bound for the entropy is needed and the limit is smooth. Of course, the existence of a smooth limit also follows from Theorem 2.5 and the fact that $\text{Rm} \geq 0$ on gradient shrinkers for $n = 3$. All this is not surprising, since the only 3-dimensional gradient shrinkers are the Gaussian soliton, the cylinder, the sphere and quotients thereof [6].

In the following, the goal is to get local energy bounds from 4d-Gauss-Bonnet with boundary. For a 4-manifold N with boundary ∂N , the Chern-Gauss-Bonnet formula says (see e.g. [19])

$$32\pi^2\chi(N) = \int_N (|\text{Rm}|^2 - 4|\text{Rc}|^2 + R^2)dV + 16 \int_{\partial N} k_1 k_2 k_3 dA + 8 \int_{\partial N} (k_1 K_{23} + k_2 K_{13} + k_3 K_{12})dA, \quad (4.3)$$

where the $k_i = \text{II}(e_i, e_i)$ are the principal curvatures of ∂N (here e_1, e_2, e_3 is an orthonormal basis of $T\partial N$ diagonalizing the second fundamental form) and the $K_{ij} = \text{Rm}(e_i, e_j, e_i, e_j)$ are sectional curvatures of N .

In a first step, we prove Theorem 1.2 under an extra assumption which ensures in particular that the cubic boundary term has the good sign.

Proposition 4.3 (Convexity implies Riemann energy condition)

Every sequence of 4-dimensional gradient shrinkers (M_i, g_i, f_i) (with normalization and basepoint as usual) with entropy bounded below, $\mu(g_i) \geq \underline{\mu}$, Euler characteristic bounded above, $\chi(M_i) \leq \bar{\chi}$, and convex potential at large distances,

$$\text{Hess}_{g_i} f_i(x) \geq 0 \quad \text{if } d(x, p_i) \geq r_0, \quad (4.4)$$

satisfies the energy condition (1.4).

Proof. Let us introduce some notation first. We suppress the index i and write $F(x) = e^{-f(x)}$ and define the level and superlevel sets

$$\Sigma_u = \{x \in M \mid F(x) = u\}, \quad M_u = \{x \in M \mid F(x) \geq u\}. \quad (4.5)$$

Note that $M_0 = M$ and $M_{u_2} \subset M_{u_1}$ if $u_2 \geq u_1$.

By the traced soliton equation (2.2) and assumption (4.4), we have $R \leq \frac{n}{2}$ at large distances. Using this, the auxiliary equation (2.4), Lemma 2.1, and the bounds $\underline{\mu} \leq -C_1(g) \leq \bar{\mu}$, we see that f does not have critical points at large distances. In fact, there is a constant $u_0 = u_0(r_0, \underline{\mu}) > 0$ such that $|\nabla f| \geq 1$ and $\nabla^2 f \geq 0$ if $F(x) \leq u_0$. Moreover, for $0 < u \leq u_0$ the Σ_u are smooth compact hypersurfaces, they are all diffeomorphic and we have $\partial M_u = \Sigma_u$ and $\chi(M_u) = \chi(M)$.

Define a cutoff function $\vartheta(x) := \min\{u_0, F(x)\}$, then

$$\int_M |\text{Rm}|^2 \vartheta dV = \int_M |\text{Rm}|^2 \int_0^{u_0} 1_{\{u \leq F\}} du dV = \int_0^{u_0} \int_{M_u} |\text{Rm}|^2 dV du. \quad (4.6)$$

Now, we can apply (4.3) for $N = M_u$. Note that $\chi(M_u) \leq \bar{\chi}$, and that the scalar curvature term and the cubic boundary term are nonnegative. Indeed,

$$\text{II} = -\nabla_{\perp}^2 F / |\nabla F| = \frac{1}{|\nabla f|} (\nabla^2 f - \nabla f \otimes \nabla f)_{\perp} = \frac{1}{|\nabla f|} \nabla_{\perp}^2 f \geq 0, \quad (4.7)$$

where \perp denotes the restriction of the Hessian to $T\Sigma_u$. Thus, we obtain

$$\begin{aligned} \int_{M_u} |\text{Rm}|^2 dV &\leq 32\pi^2 \bar{\chi} + 4 \int_{M_u} |\text{Rc}|^2 dV \\ &\quad - 8 \int_{\Sigma_u} (k_1 K_{23} + k_2 K_{13} + k_3 K_{12}) dA \end{aligned} \quad (4.8)$$

and undoing (4.6), using $\vartheta \leq e^{-f}$, $|k_i| \leq |\mathbb{I}|$ and $|K_{ij}| \leq |\text{Rm}|$, this implies

$$\int_M |\text{Rm}|^2 \vartheta dV \leq 32\pi^2 \bar{\chi} u_0 + 4 \int_M |\text{Rc}|^2 e^{-f} dV + 24 \int_0^{u_0} \int_{\Sigma_u} |\mathbb{I}| |\text{Rm}| dA du. \quad (4.9)$$

The Ricci term can be estimated as in (4.1). For the last term we use the coarea formula (observe the cancelation):

$$\begin{aligned} \int_0^{u_0} \int_{\Sigma_u} |\mathbb{I}| |\text{Rm}| dA du &\leq \int_{M \setminus M_{u_0}} \frac{|\nabla^2 f|}{|\nabla f|} |\text{Rm}| |\nabla f| \vartheta dV \\ &\leq \frac{1}{48} \int_M |\text{Rm}|^2 \vartheta dV + 12 \int_M |\nabla^2 f|^2 e^{-f} dV. \end{aligned} \quad (4.10)$$

The first term can be absorbed, the second one can be dealt with as in (4.1),

$$\begin{aligned} \int_M |\nabla^2 f|^2 e^{-f} dV &= \int_M \langle \nabla^2 f, \frac{1}{2}g - \text{Rc} \rangle e^{-f} dV \\ &= \frac{1}{2} \int_M \Delta f e^{-f} dV + \int_M \langle \nabla f, \text{div}(e^{-f} \text{Rc}) \rangle dV \\ &= \frac{1}{2} \int_M \left(\frac{n}{2} - R \right) e^{-f} dV, \end{aligned} \quad (4.11)$$

where we used the traced soliton equation and $\text{div}(e^{-f} \text{Rc}) = 0$ in the last step. Putting everything together, we obtain a uniform bound for $\int_M |\text{Rm}|^2 \vartheta dV$, and (1.4) follows. \square

Let us now replace the (unnatural) assumption (4.4) by the weaker assumption (1.5). Let $u_0 = u_0(r_0, \underline{\mu}) > 0$ such that $|\nabla f| \geq c$ if $F(x) \leq u_0$ and $\vartheta(x) := \min\{u_0, F(x)\}$ a cutoff function as before. The proof is essentially identical, except that in addition we have to estimate (the negative part of) the cubic boundary term in the Gauss-Bonnet formula. By the coarea formula

$$\left| \int_0^{u_0} \int_{\Sigma_u} \det \mathbb{I} dA du \right| \leq \int_{M \setminus M_{u_0}} \frac{|\nabla_{\perp}^2 f|^3}{|\nabla f|^2} e^{-f} dV \leq \frac{1}{c^2} \int_M |\text{Rc} - \frac{1}{2}g|^3 e^{-f} dV. \quad (4.12)$$

Note that the only difficult term is $\int_M |\text{Rc}|^3 e^{-f} dV$, since all other terms can be uniformly bounded using Lemma 4.1. Fortunately, we can bound this weighted L^3 -norm of Ricci by uniformly controlled terms and a weighted L^2 Riemann term that can be absorbed in the Gauss-Bonnet argument. Exploiting the algebraic structure of the equations for gradient shrinkers and the full strength of Lemma 4.1, we obtain the following key estimate.

Lemma 4.4 (Weighted L^3 estimate for Ricci)

For $\varepsilon > 0$ and $\underline{\mu} > -\infty$ there exist constants $C(\varepsilon) = C(\varepsilon, \underline{\mu}, n) < \infty$ such that for every gradient shrinker (M^n, g, f) with our usual normalization and $\mu(g) \geq \underline{\mu}$ we have the estimate

$$\int_M |\text{Rc}|^3 e^{-f} dV \leq \varepsilon \int_M |\text{Rm}|^2 e^{-f} dV + C(\varepsilon). \quad (4.13)$$

Proof. Analogous to (2.3), we have

$$\nabla_k R_{ij} - \nabla_i R_{kj} = -\nabla_k \nabla_i \nabla_j f + \nabla_i \nabla_k \nabla_j f = R_{ikj\ell} \nabla_\ell f \quad (4.14)$$

and as a direct consequence $\text{div}(e^{-f} \text{Rm}) = 0$. Moreover, analogous to (2.6), the shrinker version of the evolution equation for the Ricci tensor is

$$R_{ij} + \langle \nabla f, \nabla R_{ij} \rangle = \Delta R_{ij} + 2R_{ikj\ell} R_{k\ell}. \quad (4.15)$$

Now, for a cutoff function η_r as in the proof of Lemma 4.1, we compute

$$\begin{aligned} \int_M \eta_r |\text{Rc}|^3 e^{-f} dV &= \int_M \eta_r |\text{Rc}| \langle \tfrac{1}{2}g - \nabla^2 f, \text{Rc} \rangle e^{-f} dV \\ &= \int_M \left(\tfrac{1}{2} \eta_r |\text{Rc}| R + |\text{Rc}| \text{Rc}(\nabla f, \nabla \eta_r) + \eta_r \text{Rc}(\nabla f, \nabla |\text{Rc}|) \right) e^{-f} dV \\ &\leq \int_M \left(\tfrac{1}{2} \eta_r |\text{Rc}| R + |\nabla \eta_r| |\text{Rc}|^2 |\nabla f| \right) e^{-f} dV \\ &\quad + \delta \int_M \eta_r |\nabla \text{Rc}|^2 e^{-\frac{3}{2}f} dV + \frac{1}{4\delta} \int_M \eta_r |\text{Rc}|^2 |\nabla f|^2 e^{-\frac{1}{2}f} dV \\ &\leq \delta \int_M \eta_r |\nabla \text{Rc}|^2 e^{-\frac{3}{2}f} dV + C(\delta), \end{aligned}$$

for $\delta > 0$ to be chosen later. Here, we used Young's inequality, Kato's inequality, the growth estimates from Section 2 and Lemma 4.1 (note that $|\nabla f|^2 e^{-f/2} \leq C e^{-f/4}$ etc.). Note that the constant $C(\delta)$ does not depend on the scaling factor r of the cutoff function η_r , so by sending $r \rightarrow \infty$, we obtain

$$\int_M |\text{Rc}|^3 e^{-f} dV \leq \delta \int_M |\nabla \text{Rc}|^2 e^{-\frac{3}{2}f} dV + C(\delta). \quad (4.16)$$

Next, we estimate the weighted L^2 -norm of ∇Rc with a partial integration, equation (4.15), and Young's inequality,

$$\begin{aligned} \int_M \eta_r^2 |\nabla \text{Rc}|^2 e^{-\frac{3}{2}f} dV &= - \int_M \eta_r^2 (\Delta R_{ij} - \tfrac{3}{2} \langle \nabla f, \nabla R_{ij} \rangle) R_{ij} e^{-\frac{3}{2}f} dV \\ &\quad - \int_M 2\eta_r \langle \nabla \eta_r, \nabla R_{ij} \rangle R_{ij} e^{-\frac{3}{2}f} dV \\ &\leq - \int_M \eta_r^2 (R_{ij} - 2R_{ikj\ell} R_{k\ell}) R_{ij} e^{-\frac{3}{2}f} dV \\ &\quad + \int_M \left(\tfrac{1}{2} \eta_r^2 |\nabla \text{Rc}|^2 + \tfrac{1}{4} \eta_r^2 |\text{Rc}|^2 |\nabla f|^2 + 4|\nabla \eta_r|^2 |\text{Rc}|^2 \right) e^{-\frac{3}{2}f} dV. \end{aligned}$$

By absorption, the growth estimates from Section 2, Lemma 4.1 and the soliton equation, we obtain

$$\begin{aligned} \int_M \eta_r^2 |\nabla \text{Rc}|^2 e^{-\frac{3}{2}f} dV &\leq C - 2 \int_M \eta_r^2 (R_{ij} - 2R_{ikj\ell}R_{k\ell}) R_{ij} e^{-\frac{3}{2}f} dV \\ &= C - 4 \int_M \eta_r^2 R_{ikj\ell} R_{ij} \nabla_k \nabla_\ell f e^{-\frac{3}{2}f} dV. \end{aligned}$$

Finally, with another partial integration, $\text{div}(e^{-f} \text{Rm}) = 0$, and with

$$2R_{ikj\ell} \nabla_k R_{ij} \nabla_\ell f = R_{ikj\ell} \nabla_\ell f (\nabla_k R_{ij} - \nabla_i R_{kj}) = |R_{ikj\ell} \nabla_\ell f|^2, \quad (4.17)$$

which follows from (4.14) and which is the identity that makes the proof work, we get

$$\begin{aligned} \int_M \eta_r^2 |\nabla \text{Rc}|^2 e^{-\frac{3}{2}f} dV &\leq C + 2 \int_M \eta_r^2 |R_{ikj\ell} \nabla_\ell f|^2 e^{-\frac{3}{2}f} dV \\ &\quad + \int_M (8\eta_r \nabla_k \eta_r R_{ikj\ell} R_{ij} \nabla_\ell f - 2\eta_r^2 R_{ikj\ell} R_{ij} \nabla_k f \nabla_\ell f) e^{-\frac{3}{2}f} dV \\ &\leq C + C \int_M \eta_r^2 |\text{Rm}|^2 e^{-f} dV. \end{aligned} \quad (4.18)$$

In the last step, we used again Young's inequality, the growth estimates from Section 2, Lemma 4.1 and $|\nabla f|^2 e^{-\frac{3}{2}f} \leq C e^{-f}$. The claim now follows by sending $r \rightarrow \infty$, plugging into (4.16), and choosing $\delta > 0$ such that $C\delta \leq \varepsilon$ for the constant C in (4.18). \square

Now Theorem 1.2 is an immediate consequence.

Proof of Theorem 1.2. Picking $\varepsilon = \varepsilon(c, r_0, \mu) > 0$ so small that $\varepsilon e^{-f} \leq \vartheta c^2/100$ and applying Lemma 4.4, the theorem follows as explained in the discussion after Proposition 4.3. \square

A Proofs of the lemmas from Section 2

Proof of Lemma 2.1. From (2.4) and (2.5), we obtain

$$0 \leq |\nabla f|^2 \leq f + C_1, \quad (\text{A.1})$$

i.e. $|\nabla \sqrt{f + C_1}| \leq \frac{1}{2}$ whenever $f + C_1 > 0$. Hence $\sqrt{f + C_1}$ is $\frac{1}{2}$ -Lipschitz and thus

$$\sqrt{f(x) + C_1} \leq \frac{1}{2}(d(x, y) + 2\sqrt{f(y) + C_1}), \quad (\text{A.2})$$

for all $x, y \in M$, which will give the upper bound in (2.7). The idea to prove the lower bound is the same as in the theorem of Myers (which would give a diameter

bound if the shrinker potential was constant). Consider a minimizing geodesic $\gamma(s)$, $0 \leq s \leq s_0 := d(x, y)$, joining $x = \gamma(0)$ with $y = \gamma(s_0)$. Assume $s_0 > 2$ and let

$$\phi(s) = \begin{cases} s, & s \in [0, 1] \\ 1, & s \in [1, s_0 - 1] \\ s_0 - s, & s \in [s_0 - 1, s_0]. \end{cases}$$

By the second variation formula for the energy of γ ,

$$\int_0^{s_0} \phi^2 \operatorname{Rc}(\gamma', \gamma') ds \leq (n-1) \int_0^{s_0} \phi'^2 ds = 2n - 2,$$

where $\gamma'(s) = \frac{\partial}{\partial s} \gamma(s)$. Note that by the soliton equation (2.1)

$$\operatorname{Rc}(\gamma', \gamma') = \frac{1}{2} - \nabla_{\gamma'} \nabla_{\gamma'} f,$$

which implies

$$\begin{aligned} \frac{d(x, y)}{2} + \frac{4}{3} - 2n &\leq \int_0^{s_0} \phi^2 \nabla_{\gamma'} \nabla_{\gamma'} f ds \\ &= -2 \int_0^1 \phi \nabla_{\gamma'} f ds + 2 \int_{s_0-1}^{s_0} \phi \nabla_{\gamma'} f ds \\ &\leq \sup_{s \in [0, 1]} |\nabla_{\gamma'} f| + \sup_{s \in [s_0-1, s_0]} |\nabla_{\gamma'} f| \\ &\leq \sqrt{f(x) + C_1} + \frac{1}{2} + \sqrt{f(y) + C_1} + \frac{1}{2}, \end{aligned} \tag{A.3}$$

where we used (A.1) and the fact that $\sqrt{f + C_1}$ is $\frac{1}{2}$ -Lipschitz in the last step. By (A.3), every minimizing sequence is bounded and f attains its infimum at a point p . Since $\Delta f(p) \geq 0$, (2.2) and (2.5) imply

$$0 \leq R(p) \leq \frac{n}{2}. \tag{A.4}$$

Using this and $\nabla f(p) = 0$, equation (2.4) implies

$$0 \leq f(p) + C_1 \leq \frac{n}{2}. \tag{A.5}$$

Now the quadratic growth estimate (2.7) follows from (A.2), (A.3) and (A.5) by setting $y = p$. Finally, if $d(x, p) > 5n + \sqrt{2n}$, then

$$f(x) + C_1 \geq \frac{1}{4}(d(x, p) - 5n)^2 > \frac{n}{2} \geq f(p) + C_1,$$

which implies the last statement of the lemma. \square

Proof of Lemma 2.2. Let $\varrho(x) = 2\sqrt{f(x) + C_1}$. This grows linearly, since (2.7) implies

$$d(x, p) - 5n \leq \varrho(x) \leq d(x, p) + 5n. \tag{A.6}$$

Define ϱ -discs by $D(r) := \{x \in M \mid \varrho(x) < r\}$, let $V(r)$ be their volume and

$$S(r) := \int_{D(r)} R dV \quad (\text{A.7})$$

their total scalar curvature. Since $\int_{D(r)} \Delta f dV = \int_{\partial D(r)} |\nabla f| dA \geq 0$, integrating (2.2) gives

$$S(r) \leq \frac{n}{2} V(r), \quad (\text{A.8})$$

i.e. the average scalar curvature is bounded by $\frac{n}{2}$. Moreover, (2.2) and (2.4) imply

$$(r^{-n} V(r))' = 4r^{-(n+2)} S'(r) - 2r^{-(n+1)} S(r),$$

which yields the following estimate by integration

$$V(r) \leq \frac{V(r_0)}{r_0^n} r^n + \frac{4}{r^2} S(r) \quad (\text{A.9})$$

for $r \geq r_0 := \sqrt{2n+4}$, see [26, Eq. (3)] or [10, Eq. (3.6)] for details. Hence, if $r \geq \sqrt{4n}$, we get by absorption

$$V(r) \leq \frac{2V(r_0)}{r_0^n} r^n.$$

Thus, for every $r \geq 5n$ we obtain

$$\text{Vol } B_r(p) \leq V(r+5n) \leq V(2r) \leq \frac{2^{n+1}}{r_0^n} V(r_0) r^n \leq \frac{2^{n+1}}{r_0^n} \text{Vol } B_{r_0+5n}(p) r^n.$$

This proves the lemma up to the statement that C_2 depends only on the dimension and that (2.9) also holds for balls with $r < 5n$. To get this, note that $|\nabla f(x)| \leq \frac{1}{2} r_0 + 5n =: a$ for $d(x, p) \leq r_0 + 5n =: R_0$. Now, using the fact that the Bakry-Emery Ricci tensor $\text{Rc}_f = \text{Rc} + \text{Hess } f$ of the manifold with density $(M, g, e^{-(f+C_1)} dV)$ is nonnegative by the soliton equation, we obtain, see [40, Thm. 1.2a],

$$\frac{\int_{B_R(p)} e^{-(f+C_1)} dV}{\int_{B_\varepsilon(p)} e^{-(f+C_1)} dV} \leq e^{aR} \frac{R^n}{\varepsilon^n}. \quad (\text{A.10})$$

for $0 < \varepsilon < R \leq R_0$. Since $|f + C_1| \leq a^2$ on $B_{R_0}(p)$, this implies

$$\text{Vol } B_R(p) \leq e^{2a^2+aR} \frac{R^n}{\varepsilon^n} \text{Vol } B_\varepsilon(p) \quad (\text{A.11})$$

and by sending ε to zero the claim follows. \square

Proof of Lemma 2.3. Suppose towards a contradiction that there exist a sequence of gradient shrinkers (M_i, g_i, f_i) with $\mu(g_i) \geq \underline{\mu}$ and balls $B_{\delta_i}(x_i) \subset B_r(p_i)$ with $\delta_i^{-n} \text{Vol } B_{\delta_i}(x_i) \rightarrow 0$. We will not directly use $B_{\delta_i}(x_i)$ but consider the sequence of unit balls $B_1(x_i) \subset B_{r+1}(p_i)$ instead, which allows to work with the shrinker entropy

as defined above rather than with a version that explicitly involves a scaling or time parameter τ as it is necessary for the argument in [25]. Set $a := \frac{1}{2}(r+1+5n)$, then $|\nabla f_i| \leq a$, $|f_i + C_1(g_i)| \leq a^2$ and $R_{g_i} \leq a^2$ on $B_{r+1}(p_i)$. The volume comparison theorem for the Bakry-Emery Ricci tensor implies, as in (A.11),

$$\text{Vol } B_1(x_i) \leq e^{2a^2+a} \delta_i^{-n} \text{Vol } B_{\delta_i}(x_i) \rightarrow 0 \quad (\text{A.12})$$

for $i \rightarrow \infty$, as well as

$$\text{Vol } B_1(x_i) \leq 2^n e^{2a^2+a} \text{Vol } B_{1/2}(x_i), \quad \forall i \in \mathbb{N}. \quad (\text{A.13})$$

Define the test functions $\tilde{u}_i = c_i^{-1/2} \eta_i$ with $\eta_i(x) = \eta(d(x, x_i))$ for a cutoff function η as in Section 2 and with $\int_{M_i} \tilde{u}_i^2 dV = (4\pi)^{n/2}$, i.e.

$$c_i = (4\pi)^{-n/2} \int_{M_i} \eta_i^2 dV \leq (4\pi)^{-n/2} \text{Vol } B_1(x_i) \rightarrow 0.$$

Let C be an upper bound for $4\eta'^2 - \eta^2 \log \eta^2$. Using (A.13) and $c_i^{-1} \text{Vol } B_{1/2}(x_i) \leq c_i^{-1} \int_{M_i} \eta_i^2 = (4\pi)^{n/2}$, we obtain

$$c_i^{-1} \int_{M_i} \left(4|\nabla \eta_i|^2 - \eta_i^2 \log \eta_i^2 \right) dV \leq c_i^{-1} \text{Vol } B_1(x_i) C \leq (4\pi)^{n/2} 2^n e^{2a^2+a} C.$$

Hence

$$\begin{aligned} \mathcal{W}(g_i, \tilde{u}_i) &= (4\pi)^{-n/2} c_i^{-1} \int_{M_i} \left(4|\nabla \eta_i|^2 - \eta_i^2 \log \eta_i^2 \right) dV \\ &\quad + (4\pi)^{-n/2} \int_{M_i} (R - n + \log c_i) \tilde{u}_i^2 dV \\ &\leq 2^n e^{2a^2+a} C + a^2 - n + \log c_i, \end{aligned}$$

which tends to $-\infty$ as c_i tends to zero, contradicting the lower entropy bound $\mathcal{W}(g_i, \tilde{u}_i) \geq \mu(g_i) \geq \underline{\mu} > -\infty$. \square

References

- [1] U. Abresch and D. Gromoll. On complete manifolds with nonnegative Ricci curvature. *J. Amer. Math. Soc.*, 3(2):355–374, 1990.
- [2] M. Anderson. Ricci curvature bounds and Einstein metrics on compact manifolds. *J. Amer. Math. Soc.*, 2(3):455–490, 1989.
- [3] M. Anderson and J. Cheeger. Diffeomorphism finiteness for manifolds with Ricci curvature and $L^{n/2}$ -norm of curvature bounded. *Geom. Funct. Anal.*, 1(3):231–252, 1991.
- [4] L. Bessières, G. Besson, M. Boileau, S. Maillot, and J. Porti. Geometrisation of 3-manifolds. *EMS Tracts in Mathematics*, 13, Zürich, 2010.
- [5] S. Bando, A. Kasue, and H. Nakajima. On a construction of coordinates at infinity on manifolds with fast curvature decay and maximal volume growth. *Invent. Math.*, 97(2):313–349, 1989.

- [6] H.-D. Cao. Recent progress on Ricci solitons. In *Recent Advances in Geometric Analysis*, volume 11 of *Advanced Lectures in Mathematics (ALM)*. International Press, 2009.
- [7] B. Chow and D. Knopf. The Ricci flow: an introduction. *Mathematical Surveys and Monographs*, 110, AMS, Providence, 2004.
- [8] H.-D. Cao and N. Sesum. A compactness result for Kähler Ricci solitons. *Adv. Math.*, 211(2):794–818, 2007.
- [9] H.-D. Cao and X.-P. Zhu. Hamilton-Perelman’s proof of the Poincaré conjecture and the geometrization conjecture. *ArXiv:math/0612069v1*, 2006.
- [10] H.-D. Cao and D. Zhou. On complete gradient shrinking Ricci solitons. *ArXiv:0903.3932v2*, 2009.
- [11] J. Carrillo and L. Ni. Sharp logarithmic Sobolev inequalities on gradient solitons and applications. *Comm. Anal. Geom.*, 17(4):721–753, 2009.
- [12] J. Cheeger. Finiteness theorems for Riemannian manifolds. *Amer. J. Math.*, 92:61–74, 1970.
- [13] J. Cheeger. Degeneration of Einstein metrics and metrics with special holonomy. *Surveys Diff. Geom.*, VIII:29–73, 2003.
- [14] J. Cheeger, M. Gromov, and M. Taylor. Finite propagation speed, kernel estimates for functions of the Laplace operator, and the geometry of complete Riemannian manifolds. *J. Diff. Geom.*, 17(1):15–53, 1982.
- [15] X. Chen and B. Wang. Space of Ricci flows (I). *ArXiv:0902.1545v1*, 2009.
- [16] C. Croke. Some isoperimetric inequalities and eigenvalue estimates. *Ann. Sci. École Norm. Sup. (4)*, 13(4):419–435, 1980.
- [17] T. Eguchi and A. Hanson. Gravitational instantons. *Gen. Relativity Gravitation*, 11(5):315–320, 1979.
- [18] M. Feldman, T. Ilmanen, and D. Knopf. Rotationally symmetric shrinking and expanding gradient Kähler-Ricci solitons. *J. Diff. Geom.*, 65(2):169–209, 2003.
- [19] P. Gilkey. *Invariance theory, the heat equation, and the Atiyah-Singer index theorem*. Math. Lect. Series. Publish or Perish, 1984.
- [20] R. Greene and H. Wu. Lipschitz convergence of Riemannian manifolds. *Pacific J. Math.*, 131(1):119–141, 1988.
- [21] M. Gromov. *Metric structures for Riemannian and non-Riemannian spaces*, volume 152 of *Progress in Mathematics*. Birkhäuser Boston, 1999.
- [22] R. Hamilton. Three-manifolds with positive Ricci curvature. *J. Diff. Geom.* 17, no. 2, 255–306, 1982.
- [23] R. Hamilton. A compactness property for solutions of the Ricci flow. *Amer. J. Math.*, 117:545–572, 1995.
- [24] R. Hamilton. The formation of singularities in the Ricci flow. *Surveys in differential geometry*, Vol. II, 7–136, Int. Press, Cambridge, MA, 1995.
- [25] B. Kleiner and J. Lott. Notes on Perelman’s papers. *Geom. Topol.*, 12(5):2587–2855, 2008.
- [26] O. Munteanu. The volume growth of complete gradient shrinking Ricci solitons. *ArXiv:0904.0798v2*, 2009.
- [27] O. Munteanu and N. Sesum. On gradient Ricci solitons. *ArXiv:0910.1105v1*, 2009.
- [28] J. Morgan and G. Tian. Ricci flow and the Poincaré conjecture. *Clay Mathematics Monographs*, 3, AMS, Cambridge, MA, 2007.
- [29] H. Nakajima. Hausdorff convergence of Einstein 4-manifolds. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 35(2):411–424, 1988.
- [30] G. Perelman. The entropy formula for the Ricci flow and its geometric applications. *ArXiv:math/0211159v1*, 2002.
- [31] L. Sibner. The isolated point singularity problem for the coupled Yang-Mills equations in higher dimensions. *Math. Ann.*, 271(1):125–131, 1985.

- [32] J. Song and B. Weinkove. Contracting exceptional divisors by the Kähler-Ricci flow. *ArXiv:1003.0718v1*, 2010.
- [33] G. Tian. On Calabi’s conjecture for complex surfaces with positive first Chern class. *Invent. Math.*, 101(1):101–172, 1990.
- [34] G. Tian and J. Viaclovsky. Bach-flat asymptotically locally Euclidean metrics. *Invent. Math.*, 160(2):357–415, 2005.
- [35] G. Tian and J. Viaclovsky. Moduli spaces of critical Riemannian metrics in dimension four. *Adv. Math.*, 196(2):346–372, 2005.
- [36] G. Tian and J. Viaclovsky. Volume growth, curvature decay, and critical metrics. *Comment. Math. Helv.*, 83(4):889–911, 2008.
- [37] K. Uhlenbeck. Removable singularities in Yang-Mills fields. *Comm. Math. Phys.*, 83(1):11–29, 1982.
- [38] C. Villani. *Optimal transport. Old and new*, volume 338 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, 2009.
- [39] B. Weber. Convergence of compact Ricci Solitons. *Int. Math. Res. Not.*, 2010.
- [40] G. Wei and W. Wylie. Comparison geometry for the Bakry-Emery Ricci tensor. *J. Diff. Geom.*, 83(2):337–405, 2009.
- [41] X. Zhang. Compactness theorems for gradient Ricci solitons. *J. Geom. Phys.*, 56(12):2481–2499, 2006.
- [42] Z.H. Zhang. On the completeness of gradient Ricci solitons. *Proc. Amer. Math. Soc.*, 137(8):2755–2759, 2009.

Robert Haslhofer
 ETH ZÜRICH, 8092 ZÜRICH, SWITZERLAND

Reto Müller
 SCUOLA NORMALE SUPERIORE DI PISA, 56126 PISA, ITALY