



The connected prescription for form factors in twistor space

Brandhuber, A; Hughes, E; Panerai, R; Spence, B; Travaglini, G

© The Author(s) 2016

This article is distributed under the terms of the Creative Commons Attribution License (CC-BY 4.0), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

For additional information about this publication click this link.

<http://qmro.qmul.ac.uk/xmlui/handle/123456789/18787>

Information about this research object was correct at the time of download; we occasionally make corrections to records, please therefore check the published record when citing. For more information contact scholarlycommunications@qmul.ac.uk

The connected prescription for form factors in twistor space

A. Brandhuber, E. Hughes, R. Panerai, B. Spence and G. Travaglini

*Centre for Research in String Theory, School of Physics and Astronomy,
Queen Mary University of London,
Mile End Road, London E1 4NS U.K.*

E-mail: a.brandhuber@qmul.ac.uk, e.f.hughes@qmul.ac.uk,
r.panerai@qmul.ac.uk, w.j.spence@qmul.ac.uk, g.travaglini@qmul.ac.uk

ABSTRACT: We propose a connected prescription formula in twistor space for all tree-level form factors of the stress tensor multiplet operator in $\mathcal{N} = 4$ super Yang-Mills, which is a generalisation of the expression of Roiban, Spradlin and Volovich for superamplitudes. By introducing link variables, we show that our formula is identical to the recently proposed four-dimensional scattering equations for form factors. Similarly to the case of amplitudes, the link representation of form factors is shown to be directly related to BCFW recursion relations, and is considerably more tractable than the scattering equations. We also discuss how our results are related to a recent Grassmannian formulation of form factors, and comment on a possible derivation of our formula from ambitwistor strings.

KEYWORDS: Scattering Amplitudes, Supersymmetric gauge theory

ARXIV EPRINT: [1608.03277](https://arxiv.org/abs/1608.03277)

Contents

1	Introduction	1
2	The connected prescription formula	3
3	A link representation for form factors	6
4	Examples	7
4.1	The maximally non-MHV form factor	7
4.2	The MHV form factors	8
4.3	The $1^-2^-3^-4^+$ form factor and connection to BCFW diagrams	8
5	Connections to the Grassmannian	10
6	Form factors from ambitwistor strings	12

1 Introduction

The study of the S -matrix of $\mathcal{N} = 4$ supersymmetric Yang-Mills (SYM) has inspired remarkable progress and novel ideas, even at tree level. A first crucial result was the integral representation of this quantity, found by Roiban, Spradlin and Volovich (RSV) in [1] as an integral over the moduli space of degree $k - 1$ curves in super twistor space (for N^{k-2} MHV amplitudes), following Witten’s groundbreaking insights [2]. An important feature of the RSV formula, presented below in (2.3), is that the integral is in fact localised on a discrete set of solutions of certain polynomial equations. However, despite being conceptually beautiful, the RSV representation proved hard to work with because of the difficulty in determining these solutions. In this respect, the BCFW representation [3, 4] emerged as a much more tractable and generalisable approach to compute amplitudes, also applicable in different theories, including gravity [5–7].

Important progress was made later in [8], which accomplished two goals: firstly, it showed that by rewriting the RSV formula using the link variables introduced in [9] — which have the neat property of linearising momentum conservation — one can overcome the roadblocks due to the complexity of the algebraic equations arising in [1]; and furthermore, it proved that a certain precisely formulated change of integration contour in the RSV formula, rewritten using link variables, expresses the amplitudes as a sum of residues that are identical to BCFW diagrams (with appropriate shifts). This is an intriguing result, as it relates two a priori very different formulations of gauge theory amplitudes.

In this paper we wish to extend these observations to form factors, i.e. partially off-shell quantities of the form $\langle 1 \cdots n | \mathcal{O}(0) | 0 \rangle$, obtained by applying a local gauge-invariant

operator \mathcal{O} to the vacuum and then projecting the result onto an n -particle state of on-shell particles with momenta p_i obeying $q := p_1 + \dots + p_n$. As we shall see this task turns out to be surprisingly simple, suggesting also potential new directions to explore for correlation functions.

An additional, more recent motivation for our work stems from the CHY scattering equations [10, 11], which describe scattering amplitudes at tree level in a variety of theories with and without supersymmetry, and in different numbers of dimensions. Specialising to four dimensions, a new remarkable closed formula for the S -matrix of Yang-Mills theories with different amounts of supersymmetry was derived in [12] starting from ambitwistor strings. Taking gluon scattering as an example, these four-dimensional scattering equations treat positive and negative helicity gluons in a different, complementary way, similarly to the link representations of [8, 9]. It is then natural to ask how different representations of the same S -matrix of gauge theory can be related.

This question was answered in [13], which wrote down a map between the polynomial and rational form of the scattering equations, appearing in the RSV formula and in [12], respectively. A first observation we will make is that the connection is (and, in fact, was) immediate once one makes use of the link representation of the RSV formula discussed in [8]. We will then move on to discuss how to extend the RSV formula to form factors. Our starting point will be an interesting formula written down in [14] which conjectures an extension of the four-dimensional scattering equations for Yang-Mills theory to form factors of the local operator $\text{Tr } F_{\text{SD}}^2$. These form factors are of phenomenological importance, given their connection to Higgs + multi-gluon scattering amplitudes [15, 16], where the Higgs is represented by the insertion of the operator.¹ In that case, very few modifications to the formula for amplitudes are needed — specifically, two auxiliary gluons of positive helicity x and y are added.² Importantly, the amplitudes generated by this formula depend only on $p_x + p_y$ (or $p_x + p_y$ and the supermomentum $q_x + q_y$ in the supersymmetric version we introduce in section 2) rather than on the two momenta separately. We note another important feature of this formula: it contains certain Parke-Taylor like denominators of the form (ab) , with $\sigma_{a,b}$ parameterising punctures on the Riemann sphere,³ which only involve adjacent *physical* particles, i.e. they do not include x and y .

After establishing in section 2 a quick path to relate the RSV formula of [1] and the four-dimensional scattering equations of [12], our next goal is to write down a formula (2.2), analogous to the RSV result, describing supersymmetric form factors of the chiral stress tensor multiplet operator in twistor space.⁴ We will then show how this proposal is equivalent to a simple supersymmetric extension of the scattering equations formula for form factors presented in [14]. In section 3 we show that our formula can naturally be expressed in terms of link variables, in the same vein as the RSV formula. This link variable formula-

¹See also [17–21] for some recent related work also in the maximally supersymmetric theory.

²The choice of positive helicity is such that all-plus and single-minus gluon form factors of $\text{Tr } F_{\text{SD}}^2$ are now non-vanishing.

³The precise meaning of this notation will be explained in the next section.

⁴We also note the works [22–25] on representing and calculating form factors in twistor space, which would be interesting to relate to the present work.

tion turns out to be very advantageous from the point of view of simplifying calculations, as we demonstrate in several examples in section 4. Importantly, we confirm an important feature of the link variable representation of the RSV formula found in [8], namely that a simple deformation of the integration contour in the link variable space and the global residue theorem lead to an alternative representation of the amplitudes which coincides with the BCFW recursion relations for form factors [26].

Another important strand of research on amplitudes is that of the Grassmannian (see for example [27, 28]). These novel representations of amplitudes are derived from on-shell diagrams where familiar concepts such as locality and unitarity are only emergent, rather than manifest at each step of the calculation. In an interesting paper [29], a Grassmannian-based formula was conjectured which describes form factors of the stress tensor multiplet operator.⁵ A relevant feature of this conjecture is the appearance of certain Parke-Taylor denominators where all particles appear on the same footing, including the auxiliary particles x and y used to describe the form factor insertion, unlike the formula of [14]. Using a Veronese map [13, 31], we will see in section 5 how one can relate the Grassmannian formulae of [29] to the four-dimensional scattering equations for form factors of [14], and hence to our twistor-space and link-representation based formulae for form factors of section 2, at least in certain cases. Finally we conclude in section 6 with some observations: we comment on a possible derivation of form factors from ambitwistor strings, and then discuss possible extensions of the scattering equation approach beyond form factors.

2 The connected prescription formula

We begin by describing the main ingredients of the connected formula for form factors.

1. The first ingredient is a set of supertwistor variables \mathcal{Z}_a , $a = 1, \dots, n$ describing the n particles, with $\mathcal{Z} = (\lambda_\alpha, \mu^{\dot{\alpha}}, \eta^A)$. As in [14, 29], we describe the form factor insertion through two extra particles x and y . The momentum and supermomentum carried by the form factor will then be $q := \lambda_x \tilde{\lambda}_x + \lambda_y \tilde{\lambda}_y$ and $\gamma := \lambda_x \eta_x + \lambda_y \eta_y$, respectively. In twistor space, this amounts to introducing two extra super-twistors \mathcal{Z}_x and \mathcal{Z}_y .
2. As in [1], we introduce a degree $k - 1$ map from \mathbb{CP}^1 to $\mathbb{C}^{4|4}$, where $k - 2$ is the MHV degree of the superamplitude.⁶ This polynomial has the form

$$\mathcal{P}(\sigma, \{\mathcal{A}\}) := \sum_{d=0}^{k-1} \mathcal{A}_d \sigma^d, \tag{2.1}$$

where the supertwistors \mathcal{A}_d are the supermoduli of the curve.

We propose that all form factors of the supersymmetric stress tensor multiplet operator in $\mathcal{N} = 4$ SYM are described in twistor space by the following simple generalisation of the

⁵The same approach has been later extended in [30] to take into account other operator insertions.

⁶For pure gluon amplitudes, k is the number of negative helicity gluons.

RSV formula of [1]:

$$\begin{aligned} \mathcal{F}(\mathcal{Z}) &= \langle \mathcal{Z}_x \mathcal{I} \mathcal{Z}_y \rangle^2 \int \frac{d^{4k|4k} \mathcal{A} d^{n+2} \sigma d^{n+2} \xi}{\text{vol GL}(2)} \prod_{a=1}^n \frac{\delta^{(4|4)}(\mathcal{Z}_a - \xi_a \mathcal{P}(\sigma_a, \{\mathcal{A}\}))}{\xi_a (\sigma_a - \sigma_{a+1})} \\ &\times \frac{\prod_{a=x,y} \delta^{(4|4)}(\mathcal{Z}_a - \xi_a \mathcal{P}(\sigma_a, \{\mathcal{A}\}))}{\xi_x \xi_y (\sigma_x - \sigma_y)^2}, \end{aligned} \quad (2.2)$$

where \mathcal{I} is the infinity twistor, so that $\langle \mathcal{Z}_x \mathcal{I} \mathcal{Z}_y \rangle = \langle xy \rangle$, and $\sigma_{n+1} := \sigma_1$.

It is instructive to compare this formula to the corresponding one for amplitudes, which is

$$\mathcal{A}(\mathcal{Z}) = \int \frac{d^{4k|4k} \mathcal{A} d^n \sigma d^n \xi}{\text{vol GL}(2)} \frac{\prod_{a=1}^n \delta^{(4|4)}(\mathcal{Z}_a - \xi_a \mathcal{P}(\sigma_a, \{\mathcal{A}\}))}{\prod_{a=1}^n \xi_a (\sigma_a - \sigma_{a+1})}. \quad (2.3)$$

The only modifications needed to describe form factors are: the presence of the multiplicative factor $\langle \mathcal{Z}_x \mathcal{I} \mathcal{Z}_y \rangle^2$; the presence of the two extra twistors \mathcal{Z}_x and \mathcal{Z}_y ; and the inclusion of the corresponding integration variables ξ_x, ξ_y and σ_x, σ_y , with an integrand containing $1/(\xi_x \xi_y (\sigma_x - \sigma_y)^2)$. Note that we do not involve the coordinates for particles x and y in the string of Parke-Taylor type denominators, similarly to [14] (but at variance with e.g. (3.27) of [29], which includes terms of the type $(nx)(xy)(y1)$ in the denominator).

We now show how from (2.2) we can deduce the scattering equation representation of [14] for form factors (or, more precisely, its generalisation describing supersymmetric form factors of the stress tensor multiplet operator). The proof parallels closely that of [8].

1. To begin with, we divide the particles into two sets containing k and $n - k + 2$ particles, which we label with indices J and i , respectively, with the auxiliary particles belonging to the second set. We will denote by \mathfrak{m} the first set of k particles, and by \mathfrak{p} that of the remaining $n - k$ (physical) particles, and also define $\bar{\mathfrak{p}} = \mathfrak{p} \cup \{x, y\}$. A particularly convenient choice when working with, say, component gluon amplitudes is then to assign gluons of negative (positive) helicity to the first (second) group, with the fictitious particles x and y being included in the second set. This parallels the assignments made in [14] for the non-supersymmetric scattering equations for form factors, where these two particles are treated as gluons of positive helicity.

2. Next, one Fourier transforms all the twistor variables of the i -particles to dual twistor variables: $\mathcal{Z}_i \rightarrow \mathcal{W}_i$. Calling the resulting expression $\mathcal{F}(\mathcal{W}_i, \mathcal{Z}_J)$ we have

$$\begin{aligned} \mathcal{F}(\mathcal{W}_i, \mathcal{Z}_J) &= \int \frac{d^{4k|4k} \mathcal{A} d^{n+2} \sigma d^{n+2} \xi}{\text{vol GL}(2)} \frac{\prod_{J \in \mathfrak{m}} \delta^{(4|4)}(\mathcal{Z}_J - \xi_J \mathcal{P}(\sigma_J, \{\mathcal{A}\}))}{\xi_x \xi_y (\sigma_x - \sigma_y)^2 \prod_{a=1}^n \xi_a (\sigma_a - \sigma_{a+1})} \\ &\times \left\langle \frac{\partial}{\partial \mathcal{W}_x} \mathcal{I} \frac{\partial}{\partial \mathcal{W}_y} \right\rangle^2 \prod_{i \in \bar{\mathfrak{p}}} \exp(i \xi_i \mathcal{W}_i \cdot \mathcal{P}(\sigma_i, \{\mathcal{A}\})). \end{aligned} \quad (2.4)$$

3. The key observation of [8] is that this procedure has the advantage that there are now as many δ -functions as moduli, and the integration over the \mathcal{A} can be performed explicitly, with the net effect of localising the polynomial $\mathcal{P}(\sigma, \{\mathcal{A}\})$ onto

$$\mathcal{P}(\sigma) = \sum_{J \in \mathfrak{m}} \frac{\mathcal{Z}_J}{\xi_J} \prod_{K \neq J} \frac{\sigma_K - \sigma}{\sigma_K - \sigma_J}. \quad (2.5)$$

One is then left with

$$\begin{aligned} \mathcal{F}(\mathcal{W}_i, \mathcal{Z}_J) &= \int \frac{1}{\text{vol GL}(2)} \frac{d\sigma_x d\xi_x d\sigma_y d\xi_y}{\xi_x \xi_y (\sigma_x - \sigma_y)^2} \prod_{a=1}^n \frac{d\sigma_a d\xi_a}{\xi_a (\sigma_a - \sigma_{a+1})} \\ &\times \left\langle \frac{\partial}{\partial \mathcal{W}_x} \mathcal{I} \frac{\partial}{\partial \mathcal{W}_y} \right\rangle^2 \exp \left(i \sum_{i \in \bar{\mathfrak{p}}, J \in \mathfrak{m}} \mathcal{W}_i \cdot \mathcal{Z}_J \frac{\xi_i}{\xi_J} \prod_{K \neq J} \frac{\sigma_K - \sigma_i}{\sigma_K - \sigma_J} \right). \end{aligned} \quad (2.6)$$

4. The change of variables $(\xi_i, \xi_J) \rightarrow (t_i, t_J)$ with [8]

$$t_i := \xi_i \prod_K (\sigma_K - \sigma_i), \quad t_J^{-1} := \xi_J \prod_{K \neq J} (\sigma_K - \sigma_J), \quad (2.7)$$

simplifies (2.6) to

$$\begin{aligned} \mathcal{F}(\mathcal{W}_i, \mathcal{Z}_J) &= \int \frac{1}{\text{vol GL}(2)} \frac{d\sigma_x dt_x d\sigma_y dt_y}{t_x t_y (\sigma_x - \sigma_y)^2} \prod_{a=1}^n \frac{d\sigma_a dt_a}{t_a (\sigma_a - \sigma_{a+1})} \\ &\times \left\langle \frac{\partial}{\partial \mathcal{W}_x} \mathcal{I} \frac{\partial}{\partial \mathcal{W}_y} \right\rangle^2 \exp \left(i \sum_{i \in \bar{\mathfrak{p}}, J \in \mathfrak{m}} \mathcal{W}_i \cdot \mathcal{Z}_J \frac{t_i t_J}{\sigma_J - \sigma_i} \right). \end{aligned} \quad (2.8)$$

We now introduce spinor coordinates $\sigma_\alpha = t^{-1}(1, \sigma)$, with $(ab) := \epsilon_{\alpha\beta} \sigma_a^\alpha \sigma_b^\beta$. Then $d^2\sigma = t^{-3} dt d\sigma$, and

$$\prod_{a=1}^n \frac{dt_a d\sigma_a}{t_a (\sigma_a - \sigma_{a+1})} = \prod_{a=1}^n \frac{d^2\sigma_a}{(aa+1)}. \quad (2.9)$$

We then arrive at the very simple result

$$\mathcal{F}(\mathcal{W}_i, \mathcal{Z}_J) = \int \frac{1}{\text{vol GL}(2)} \frac{d^2\sigma_x d^2\sigma_y}{(xy)^2} \prod_{a=1}^n \frac{d^2\sigma_a}{(aa+1)} \left\langle \frac{\partial}{\partial \mathcal{W}_x} \mathcal{I} \frac{\partial}{\partial \mathcal{W}_y} \right\rangle^2 \exp \left(i \sum_{i \in \bar{\mathfrak{p}}, J \in \mathfrak{m}} \frac{\mathcal{W}_i \cdot \mathcal{Z}_J}{(iJ)} \right). \quad (2.10)$$

5. It is now easy to go back to spinor variables by performing a Fourier transform. The result is⁷

$$\begin{aligned} \mathcal{F}(\{\lambda, \tilde{\lambda}\}) &= \langle xy \rangle^2 \int \frac{1}{\text{vol GL}(2)} \frac{d^2\sigma_x d^2\sigma_y}{(xy)^2} \prod_{a=1}^n \frac{d^2\sigma_a}{(aa+1)} \\ &\times \prod_{i \in \bar{\mathfrak{p}}} \delta^{(2)}(\lambda_i - \lambda(\sigma_i)) \prod_{J \in \mathfrak{m}} \delta^{(2|4)}(\tilde{\lambda}_J - \tilde{\lambda}(\sigma_J), \eta_J - \eta(\sigma_J)), \end{aligned} \quad (2.11)$$

where we have defined the functions

$$\lambda(\sigma) := \sum_{J \in \mathfrak{m}} \frac{\lambda_J}{(\sigma \sigma_J)}, \quad \tilde{\lambda}(\sigma) := \sum_{i \in \bar{\mathfrak{p}}} \frac{\tilde{\lambda}_i}{(\sigma_i \sigma)}, \quad \eta(\sigma) := \sum_{i \in \bar{\mathfrak{p}}} \frac{\eta_i}{(\sigma_i \sigma)}. \quad (2.12)$$

Eq. (2.11) is nothing but the supersymmetric form of the scattering equation for form factors presented in [14]. By performing in reverse the same steps of this proof, one can of course derive the connected prescription for form factors (2.2) from the scattering equations.⁸

⁷We recall that $\mathcal{Z} = (\lambda, \mu, \eta)$ and $\mathcal{W} = (\tilde{\mu}, \tilde{\lambda}, \tilde{\eta})$, with $\mathcal{Z} \cdot \mathcal{W} := \lambda^\alpha \tilde{\mu}_\alpha + \mu_{\dot{\alpha}} \tilde{\lambda}^{\dot{\alpha}} + \tilde{\eta}_A \eta^A$.

⁸We comment that the four-dimensional version of the amplitude scattering equations was noted in [8].

3 A link representation for form factors

In [8], Spradlin and Volovich presented an interesting formula for the S -matrix of $\mathcal{N} = 4$ SYM using the link variables introduced in [9], and we now give the corresponding formula for the form factors of the stress tensor multiplet operator.

The link representation is obtained by introducing auxiliary variables

$$c_{iJ} := \frac{1}{(iJ)}, \quad (3.1)$$

where we note that the first and second index run over the sets $\bar{\mathfrak{p}}$ and \mathfrak{m} , respectively. This identification is achieved by introducing $1 = \int dc_{iJ} \delta(c_{iJ} - 1/(iJ))$. Doing so, we can recast (2.11) as⁹

$$\mathcal{F}(\{\lambda, \tilde{\lambda}\}) = \langle xy \rangle^2 \int \prod_{i \in \bar{\mathfrak{p}}, J \in \mathfrak{m}} dc_{iJ} U(c_{iJ}) \prod_{i \in \bar{\mathfrak{p}}} \delta^{(2)}(\lambda_i - c_{iJ} \lambda_J) \prod_{J \in \mathfrak{m}} \delta^{(2|4)}(\tilde{\lambda}_J + c_{iJ} \tilde{\lambda}_i, \eta_J + c_{iJ} \eta_i), \quad (3.2)$$

where

$$U(c_{iJ}) := \int \frac{1}{\text{vol GL}(2)} \frac{d^2 \sigma_x d^2 \sigma_y}{(xy)^2} \prod_{a=1}^n \frac{d^2 \sigma_a}{(aa+1)} \prod_{i \in \bar{\mathfrak{p}}, J \in \mathfrak{m}} \delta\left(c_{iJ} - \frac{1}{(iJ)}\right). \quad (3.3)$$

There are several reasons why it is interesting to study the link representation form (3.2). Firstly, it has the advantage of linearising momentum conservation in terms of the c_{iJ} variables. Secondly, the quantity $U(c_{iJ})$ defined in (3.3) appears to be much more easily computed for any k — this is a considerable advantage in comparison to the scattering equations. Finally, it was shown in [8] that by using the (global) residue theorem, one can arrive at an alternative representation of the amplitudes which precisely matches BCFW diagrams, thus establishing a direct connection between the twistor-string representation of amplitudes and on-shell recursion relations. We will see that the same is also true for our representation of form factors, as we will explain in section 4.

In performing explicit calculations, a natural way to fix the $\text{GL}(2)$ gauge freedom is to fix the four variables corresponding to the two auxiliary legs x and y ,

$$\sigma_x = (1, 0), \quad \sigma_y = (0, 1), \quad (xy) = 1. \quad (3.4)$$

We can then change variables from the spinors σ_a^α to the brackets (xa) and (ya) so that

$$U(c_{iJ}) = \int \prod_{a=1}^n \frac{d(xa) d(ya)}{(aa+1)} \prod_{i \in \bar{\mathfrak{p}}, J \in \mathfrak{m}} \delta\left(c_{iJ} - \frac{1}{(iJ)}\right). \quad (3.5)$$

All the other brackets can be obtained from those used as integration variables using the Schouten identity,

$$(xy)(ab) = (xa)(yb) - (ya)(xb). \quad (3.6)$$

⁹In the delta functions appearing in (3.2) and (3.7) we are implicitly summing over repeated indices.

In (3.5) we have $2n$ integration variables and $k(n+2-k)$ delta functions, which means that $U(c_{iJ})$ contains $(k-2)(n-k)$ delta functions after integration. In (3.2), four of the Grassmann-even delta functions enforce momentum conservation, leaving $2n$ delta functions and $k(n+2-k)$ variables c_{iJ} to integrate over. This leaves $(k-2)(n-k)$ integration variables, which we denote by τ_k . Thus (3.2) can be written as

$$\mathcal{F}(\{\lambda, \tilde{\lambda}\}) = J \langle xy \rangle^2 \delta^{(4)}\left(q - \sum_{a=1}^n p_a\right) \int d^{(k-2)(n-k)} \tau U(c_{iJ}) \prod_{J \in \mathfrak{m}} \delta^{(4)}(\eta_J + c_{iJ} \eta_i), \quad (3.7)$$

for some $c_{iJ}(\tau)$ linear in τ , and an appropriate Jacobian J .

4 Examples

In this section we work out explicitly the form of $U(c_{iJ})$ defined in (3.3) for various form factors. We will always use the gauge fixing (3.4) so that $U(c_{iJ})$ is computed using (3.5).

4.1 The maximally non-MHV form factor

The simplest form factor to discuss turns out to be the maximally non-MHV form factor, where $\bar{\mathfrak{p}} = \{x, y\}$ and $\mathfrak{m} = \{1, \dots, n\}$. This corresponds to the case where for the component operator $\mathcal{L}_{\text{on-shell}}$ the on-shell state contains n gluons all with negative helicity. In this case we find

$$U^{\text{N}^{\text{maxMHV}}} = \prod_{J=1}^n \frac{1}{c_{xy;JJ+1}}, \quad (4.1)$$

where we have defined

$$c_{ab;cd} := c_{ac} c_{bd} - c_{ad} c_{bc}. \quad (4.2)$$

The expression for the n -point maximally non-MHV form factor is

$$\begin{aligned} \mathcal{F}^{\text{N}^{\text{maxMHV}}} &= \langle xy \rangle^2 \int \prod_{J=1}^n dc_{xJ} dc_{yJ} \delta^{(2)}\left(\lambda_x - \sum_{J=1}^n c_{xJ} \lambda_J\right) \delta^{(2)}\left(\lambda_y - \sum_{J=1}^n c_{yJ} \lambda_J\right) \\ &\times \prod_{J=1}^n \delta^{(2|4)}(\tilde{\lambda}_J + c_{xJ} \tilde{\lambda}_x + c_{yJ} \tilde{\lambda}_y, \eta_J + c_{xJ} \eta_x + c_{yJ} \eta_y) \times U^{\text{N}^{\text{maxMHV}}}, \end{aligned} \quad (4.3)$$

with $U^{\text{N}^{\text{maxMHV}}}$ given in (4.1). We can first extract momentum conservation,

$$\delta^{(2)}\left(\lambda_x - \sum_{J=1}^n c_{xJ} \lambda_J\right) \delta^{(2)}\left(\lambda_y - \sum_{J=1}^n c_{yJ} \lambda_J\right) = [xy]^2 \delta^{(4)}\left(q - \sum_{J=1}^n p_J\right), \quad (4.4)$$

where we used $p_x + p_y = q$. It is then immediate to solve for the variables c_{xJ} and c_{yJ} . The result is

$$c_{xJ} = \frac{[Jy]}{[yx]}, \quad c_{yJ} = \frac{[Jx]}{[xy]}. \quad (4.5)$$

The Jacobian from the delta functions for the $\tilde{\lambda}$ variables in (4.3) contributes a factor of $[xy]^{-n}$, while a short calculation shows that

$$c_{xy;JJ+1} = \frac{[JJ+1]}{[xy]}. \quad (4.6)$$

Combining the Jacobian with the expression for (4.1) evaluated on (4.6) leads to the known supersymmetric expression from [32],

$$\mathcal{F}^{\text{N}^{\text{max}}\text{MHV}} = \frac{q^4}{[1\ 2] \cdots [n\ 1]} \prod_{J=1}^n \delta^{(4)} \left(\eta_J + \frac{[J\ y]}{[y\ x]} \eta_x + \frac{[J\ x]}{[x\ y]} \eta_y \right). \quad (4.7)$$

Note that the fermionic delta function in (4.7) imposes supermomentum conservation $\sum_{J=1}^n \lambda_J \eta_J = \lambda_x \eta_x + \lambda_y \eta_y$ (and we have dropped the obvious momentum conservation delta function). In particular the form factor of $\text{Tr}(F_{\text{SD}}^2)$ can be obtained by setting $\eta_x = \eta_y = 0$ [32]. For a state consisting only of gluons with negative helicity one obtains immediately the known result $q^4 / ([1\ 2] \cdots [n\ 1])$.

It is interesting to contrast the remarkable simplicity of this derivation from link variables with the original calculation presented in [32] of this component form factor, which required a more significant amount of work.

4.2 The MHV form factors

The next simplest case is that of an MHV form factor. In this case the two sets are $\bar{\mathbf{p}} = \{1, \dots, \widehat{J}_1, \dots, \widehat{J}_2, \dots, n, x, y\}$ and $\mathbf{m} = \{J_1, J_2\}$, where hatted entries are to be omitted from the set. This corresponds to the case where, for the component operator $\mathcal{L}_{\text{on-shell}}$ and a purely gluonic on-shell state, there are exactly two gluons with negative and $n - 2$ with positive helicity.

The U function in (3.5) is given by

$$U^{\text{MHV}} = \frac{1}{(c_{xy; J_1 J_2})^2 c_{J_1-1\ J_2} c_{J_1+1\ J_2} c_{J_1\ J_2-1} c_{J_1\ J_2+1}} \prod_{a \neq J_1-1, J_1, J_2-1, J_2} \frac{1}{c_{a\ a+1; J_1 J_2}}. \quad (4.8)$$

Performing the integration over the link variables is again straightforward, and one arrives at the MHV super form factor of the chiral part of the stress tensor multiplet

$$\mathcal{F}^{\text{MHV}} = \frac{1}{\langle 1\ 2 \rangle \cdots \langle n\ 1 \rangle} \delta^{(8)} \left(\sum_{a=1}^n \lambda_a \eta_a + \lambda_x \eta_x + \lambda_y \eta_y \right), \quad (4.9)$$

which agrees with the known result [32] if we identify $\gamma_+ = \lambda_x \eta_x + \lambda_y \eta_y$.

4.3 The $1^-2^-3^-4^+$ form factor and connection to BCFW diagrams

In this case we have $\mathbf{m} = \{1, 2, 3\}$ and $\bar{\mathbf{p}} = \{4, x, y\}$, and (3.5) reads

$$U^{1^-2^-3^-4^+} = \int \prod_{a=1}^4 \frac{d(x\ a)\ d(y\ a)}{(a\ a+1)} \prod_{J=1}^3 \delta \left(c_{xJ} - \frac{1}{(x\ J)} \right) \delta \left(c_{yJ} - \frac{1}{(y\ J)} \right) \delta \left(c_{4J} - \frac{1}{(4\ J)} \right). \quad (4.10)$$

With nine delta functions and eight integrations, there is one delta function remaining after all integrations are carried out. The integrations over $(x\ J)$ and $(y\ J)$ are straightforward, and one can then choose to solve the two delta functions involving (41) and (42), producing

a Jacobian, and insert this solution into the remaining delta function for (4.3). Collecting all terms from this process, one finds that that

$$U^{1-2-3-4+} = \frac{c_{x2} c_{y2}}{c_{42} c_{xy;21} c_{xy;23}} \delta(S_{123;4xy}), \quad (4.11)$$

where, following the notation introduced in [8], we define

$$S_{ijk;lmn} := c_{mi} c_{mj} c_{lk} c_{nk} c_{ln;ij} - c_{ni} c_{nj} c_{lk} c_{mk} c_{lm;ij} - c_{li} c_{lj} c_{mk} c_{nk} c_{mn;ij}. \quad (4.12)$$

We comment that in this case $S_{ijk,LMN} = (\prod_{i,J} C_{iJ}) \det([1/C_{iJ}])$ where $[1/C_{iJ}]$ is the matrix with elements $1/C_{iJ}$. As in (3.7), the form factor can be obtained by integrating out the remaining delta function. However, there is a more efficient way to derive the final result which avoids solving the constraint of $\delta(S_{123;4xy})$ altogether.

In general, the complex delta function has the property

$$\int dz_1 \dots dz_m g(z) \delta(f_1(z)) \dots \delta(f_m(z)) = \sum_{z_0 \in f^{-1}(0)} \text{Res } \omega|_{z_0}, \quad (4.13)$$

where

$$\omega := \frac{g(z) dz_1 \wedge \dots \wedge dz_m}{f_1(z) \dots f_m(z)}. \quad (4.14)$$

In our case, this means that the integral in (3.7) can be written as a sum of residues of

$$\omega_U = \frac{c_{x2} c_{y2}}{c_{42} c_{xy;21} c_{xy;23}} \frac{d\tau}{S_{123;4xy}}, \quad (4.15)$$

evaluated on the zeros of the quartic polynomial $S_{123;4xy}(\tau)$. However, since ω_U can be straightforwardly extended to a meromorphic form on \mathbb{CP}^1 , we can use the global residue theorem to compute the result in terms of the other poles of ω_U , which correspond to the simple zeros of $c_{42}(\tau)$, $c_{xy;21}(\tau)$ and $c_{xy;23}(\tau)$. Focusing on gluon scattering, the corresponding residues are

$$\begin{aligned} \mathcal{F}_{42} &= -\frac{\langle 13 \rangle q^4}{s_{134} \langle 14 \rangle \langle 34 \rangle \langle 3|q|2 \rangle \langle 1|q|2 \rangle}, \\ \mathcal{F}_{xy;21} &= -\frac{\langle 3|q|4 \rangle^3}{s_{124} [12] [14] \langle 3|q|2 \rangle}, \\ \mathcal{F}_{xy;23} &= -\frac{\langle 1|q|4 \rangle^3}{s_{324} [32] [34] \langle 1|q|2 \rangle}, \end{aligned} \quad (4.16)$$

and the complete result is obtained by adding the three terms,

$$\mathcal{F}^{1-2-3-4+} = \mathcal{F}_{42} + \mathcal{F}_{xy;21} + \mathcal{F}_{xy;23}. \quad (4.17)$$

It is notable that each term in (4.16) depends on p_x and p_y only through the combination $p_x + p_y = q$. Moreover, each term is a rational function of external kinematics. Interestingly, these two properties do not hold for the four terms arising from the solutions of the scattering equation $S_{123;4xy} = 0$, and are only recovered in the sum over the four solutions.

Perhaps more remarkably, each term in (4.16) corresponds to a BCFW diagram for a [1 2] shift, analogously to the amplitude case, as discussed in [8] (see also [33]). Specifically, we have found that the sum in (4.17) corresponds, term by term, to the sum

$$(4.18)$$

given by the BCFW expansion of the form factor.

5 Connections to the Grassmannian

In [29] it was conjectured that the n -point N^{k-2} MHV chiral stress-tensor super form factor in $\mathcal{N} = 4$ SYM can be obtained as an integral over the Grassmannian¹⁰ $G(k, n + 2)$ of the form

$$\langle n+1 \ n+2 \rangle^2 \int \frac{d^{k(n+2)} C \ d^{2k} \rho}{\text{vol GL}(k)} \sum_{\text{ins}} \frac{\Omega_{n,k}(C)}{(1 \cdots k) \cdots (n+2 \cdots k-1)} \times \delta^{2k}(C \cdot \tilde{\lambda}) \delta^{2(n+2)}(\rho \odot C - \lambda) \delta^{4k}(C \cdot \eta), \tag{5.1}$$

where in this section the auxiliary particles x and y are denoted as $n + 1$ and $n + 2$, respectively. C_{Ia} is a $k \times (n + 2)$ matrix, describing a k -plane in \mathbb{C}^{n+2} , ρ_J^α is a $k \times 2$ matrix, \cdot indicates contraction of the a indices, \odot denotes contraction of the J indices,¹¹ $(a \cdots b)$ is a $k \times k$ minor of C , and the sum runs over allowed insertions of $\{n + 1, n + 2\}$ into $\{1, \dots, n\}$. The integrand $\Omega_{n,k}(C)$ appearing in (5.1) is defined to be

$$\Omega_{n,k}(C) = \frac{Y}{1 - Y}, \quad Y = \frac{(n + 2 - k \cdots n \ n + 1)(n + 2 \ 1 \cdots k - 1)}{(n + 2 - k \cdots n \ n + 2)(n + 1 \ 1 \cdots k - 1)}. \tag{5.2}$$

We are interested in connecting this conjecture with the result (2.11) expressing the form factor as a sum over solutions to the rational scattering equations. An obvious approach is suggested by comparison to [31], in which Grassmannian amplitude formulae are mapped to CHY-type formulae courtesy of the Veronese map. The Veronese map is an embedding of $G(2, n + 2)$ in $G(k, n + 2)$ defined by taking

$$C_{Ja} = \xi_a \sigma_a^{J-1}. \tag{5.3}$$

¹⁰ $G(k, n)$ is defined to be the set of k -dimensional linear subspaces of \mathbb{C}^n .

¹¹For clarity, $a = 1, \dots, n + 2$ and $J \in \mathfrak{m}$. In this section we will also set $\mathfrak{m} = \{1, \dots, k\}$ and $\mathfrak{p} = \{k + 1, \dots, n\}$ for convenience.

We can perform a partial integration of (5.1), reducing it to an integral over $G(2, n + 2)$ in this embedding:¹²

$$\begin{aligned} \langle n+1 \ n+2 \rangle^2 \int \frac{d^{n+2}\sigma \ d^{n+2}\xi \ d^{2k}\rho}{\text{vol GL}(2)} \sum_{\text{ins}} \frac{\Omega_{n,k}^V(\sigma_a, \xi_a)}{\prod_{a=1}^{n+2} \xi_a(\sigma_a - \sigma_{a+1})} \\ \times \prod_{a=1}^{n+2} \delta^{(2)}\left(\lambda_a - \xi_a \sum_{J \in \mathfrak{m}} \rho_J \sigma_a^{J-1}\right) \prod_{J \in \mathfrak{m}} \delta^{(2|4)}\left(\sum_{a=1}^{n+2} \xi_a \sigma_a^{J-1} \{\tilde{\lambda}_a | \eta_a\}\right). \end{aligned} \quad (5.4)$$

The δ -functions enforce the polynomial scattering equations in the language of [13]. For practical purposes it is more convenient to gauge fix the $\text{GL}(k)$ symmetry before applying the Veronese map, enforcing the rational scattering equations of [12]. We also apply the change of variables (2.7) to yield

$$\begin{aligned} \langle n+1 \ n+2 \rangle^2 \int \frac{d^{2(n+2)}\sigma}{\text{vol GL}(2)} \sum_{\text{ins}} \frac{\Omega_{n,k}^V(\sigma_a, t_a)}{(\sigma_1 \sigma_2) \cdots (\sigma_{n+2} \sigma_1)} \\ \times \prod_{i \in \mathfrak{p}} \delta^{(2)}(\lambda_i - \lambda(\sigma_i)) \prod_{J \in \mathfrak{m}} \delta^{(2|4)}(\tilde{\lambda}_J - \tilde{\lambda}(\sigma_J), \eta_J - \eta(\sigma_J)), \end{aligned} \quad (5.5)$$

where the functions defining the scattering equations are given by (2.12).

Under the Veronese map, Y then becomes

$$Y^V(\sigma_a, \xi_a) = \prod_{j=n+2-k}^n \frac{\sigma_j - \sigma_{n+1}}{\sigma_j - \sigma_{n+2}} \prod_{i=1}^{k-1} \frac{\sigma_{n+2} - \sigma_i}{\sigma_{n+1} - \sigma_i}, \quad (5.6)$$

after using the Vandermonde determinant formula. Note immediately that this is independent of the ξ_a , thus the transition to the rational scattering equation version is simply the identity map.

In relation to (5.1), with $Y_k = Y^V(\sigma_a, \xi_a)$ and $C_n = (\sigma_1 - \sigma_2)(\sigma_1 - \sigma_2) \cdots (\sigma_n - \sigma_1)$ it may be of interest to note the following recursive expression

$$\frac{1}{C_{n+2}} \frac{Y_k}{1 - Y_k} = \frac{1}{(\sigma_{n+1} - \sigma_{n+2})^2 C_n} \frac{N_k}{D_k}, \quad (5.7)$$

with

$$N_k = (\sigma_n - \sigma_1) \prod_{i=1}^{k-2} (\sigma_{n-i} - \sigma_{n+1})(\sigma_{n+2} - \sigma_{n+3+i}), \quad (5.8)$$

and

$$D_k = N_k + (\sigma_n - \sigma_{n+2})(\sigma_{n+1} - \sigma_1) D_{k-1}(\hat{\sigma}_n, \hat{\sigma}_1), \quad (5.9)$$

with $D_{k-1}(\hat{\sigma}_n, \hat{\sigma}_1)$ meaning D_{k-1} with variables $\sigma_1, \sigma_2, \dots, \sigma_{n+2}$ but omitting σ_1 and σ_n . We have checked (5.7) algebraically with `Mathematica` up to $k = 10$ and numerically for various higher values.

¹²Naively one might worry that there are four fewer integration variables than δ -functions. However the leftover constraints combine to form the δ -function of momentum conservation in the final answer, as required.

In terms of the homogeneous coordinates, (5.6) is

$$Y^V(\sigma_a) = \prod_{j=n+2-k}^n \frac{(j \ n+1)}{(j \ n+2)} \prod_{i=1}^{k-1} \frac{(n+2 \ i)}{(n+1 \ i)}. \quad (5.10)$$

The authors of [14] conjectured a simpler formula for the chiral stress tensor super form factor, namely (2.11). In the cases $k = 2$ and $k = n$ a short calculation shows agreement with the formula (5.10) obtained from the Grassmannian. Indeed, these cases correspond to the MHV and maximally non-MHV form factor, where the sum in (5.1) consists of a single term. More generally, one must sum over terms arising from several top-cell forms constructed via on-shell diagrams. These correspond to particular cyclic shifts of the insertion point of the additional legs representing the form factor.

The first non-trivial case in which we wish to show agreement between (5.5) and (2.11) is $n = 4, k = 3$, which corresponds to the helicity assignment $1^- 2^- 3^- 4^+$ in our chosen convention. For this case it was shown in [29] that the appropriate insertions are $\{1, 2, 3, 4, 5, 6\}$ and $\{1, 2, 5, 6, 3, 4\}$. A little algebra suffices to prove that

$$\begin{aligned} & \frac{Y_1}{1 - Y_1} \frac{1}{(1 \ 2)(2 \ 3)(3 \ 4)(4 \ 5)(5 \ 6)(6 \ 1)} + \frac{Y_2}{1 - Y_2} \frac{1}{(1 \ 2)(2 \ 5)(5 \ 6)(6 \ 3)(3 \ 4)(4 \ 1)} \\ &= \frac{1}{(1 \ 2)(2 \ 3)(3 \ 4)(4 \ 1)(5 \ 6)^2}, \end{aligned} \quad (5.11)$$

where

$$Y_1 = \frac{(3 \ 5)(4 \ 5)(6 \ 1)(6 \ 2)}{(3 \ 6)(4 \ 6)(5 \ 1)(5 \ 2)}, \quad Y_2 = \frac{(1 \ 5)(2 \ 5)(6 \ 3)(6 \ 4)}{(1 \ 6)(2 \ 6)(5 \ 3)(5 \ 4)}. \quad (5.12)$$

Note that in (5.11) we have obtained the expected integrand, where the auxiliary particles associated to the form factor now only appear in the factor $(5 \ 6)^2$.

The next non-trivial case is $n = 5, k = 3$. In this case, we have checked numerically that no combination of insertions reproduces the formula (2.11). This is not so surprising, since in this case different residues are required from each top-cell diagram, whereas the Veronese map treats terms democratically. It would be interesting to determine whether there is an improved choice of top-cells compatible with a Veronese reduction. We leave this question to future work.

6 Form factors from ambitwistor strings

The result (2.11) bears a close resemblance to the formula¹³

$$\begin{aligned} \mathcal{A}(\{\lambda, \tilde{\lambda}\}) &= \int \frac{1}{\text{vol GL}(2)} \prod_{a=1}^n \frac{d^2 \sigma_a}{(a \ a+1)} \\ &\times \prod_{i \in \bar{\mathfrak{p}}} \delta^{(2)}(\lambda_i - \lambda(\sigma_i)) \prod_{J \in \mathfrak{m}} \delta^{(2|4)}(\tilde{\lambda}_J - \tilde{\lambda}(\sigma_J), \eta_J - \eta(\sigma_J)), \end{aligned} \quad (6.1)$$

¹³Our superamplitudes have η^0 for positive helicity and η^4 for negative helicity gluons, which is the opposite of the convention employed in [12].

first derived in [12] from an ambitwistor-string model, describing the tree-level n -particle scattering in four-dimensional $\mathcal{N}=4$ SYM.

In this construction the Parke-Taylor denominator of the measure emerges from a current algebra on the worldsheet, similarly to the standard heterotic string construction. Each vertex operator is dressed with a current J^a built from N free complex fermions ψ^i and $SU(N)$ generators T^a . More explicitly, we define

$$J^a(\sigma) = \frac{i}{2} T_{ij}^a : \psi^i(\sigma) \bar{\psi}^j(\sigma) :, \tag{6.2}$$

where i, j are fundamental representation indices and a is an adjoint representation index. Recall that the only non-vanishing Wick contraction between complex fermions takes the form

$$\langle \psi^i(\sigma_1) \bar{\psi}^j(\sigma_2) \rangle = \frac{\delta^{ij}}{\sigma_1 - \sigma_2}, \tag{6.3}$$

so we may immediately evaluate the correlator of n currents to be

$$\langle J^{a_1} \dots J^{a_n} \rangle = \frac{\text{Tr}(T^{a_1} \dots T^{a_n})}{(\sigma_1 - \sigma_2) \dots (\sigma_n - \sigma_1)} + \text{perms} + \dots, \tag{6.4}$$

where we have ignored multiple trace terms. Keeping only the first term corresponds to computing a certain colour-ordered amplitude.

We may construct the measure of formula (2.11) from ambitwistor strings in a similar way, at least up to an overall factor. We must include two additional vertex operators, corresponding to the punctures σ_{n+1} and σ_{n+2} on the Riemann sphere. These are dressed with additional currents defined as above. However, in order to obtain the chiral stress tensor super form factor, we now do not require the single trace term. Rather we extract from Wick's theorem the double trace term displayed below,

$$\langle J^{a_1} \dots J^{a_{n+2}} \rangle = \dots + \frac{\text{Tr}(T^{a_1} \dots T^{a_n})}{(\sigma_1 - \sigma_2) \dots (\sigma_n - \sigma_1) (\sigma_{n+1} - \sigma_{n+2})^2} \cdot \text{Tr}(T^{a_{n+1}} T^{a_{n+2}}) + \dots, \tag{6.5}$$

providing the appropriate denominator and colour factor for the on-shell state. It would be very interesting to have a complete derivation of (2.11) from ambitwistor strings, also explaining the $\langle xy \rangle^2$ prefactor.

Of course, the current algebra in the four-dimensional ambitwistor string construction is identical to that in the ten-dimensional formula of [34] which reproduces standard CHY formulae. We might thus recast the formula (2.11) as a sum over solutions to the standard scattering equations [10]. To do this would require an appropriate prescription for the polarisation vector associated with the off-shell insertion.

Given that form factors emerge so naturally from an ambitwistor string construction, it is tempting to speculate that appropriate current algebra modifications might allow the construction of still more general objects, namely correlation functions.

An obvious generalisation of the approach followed for form factors would be to include additional auxiliary particles to represent further operator insertions. The simplest example would be that of a two-point correlator of $\mathcal{O} = \text{Tr} F_{\text{SD}}^2$ with the vacuum as the external

state. In order to contract the two operators, we choose the two pairs of auxiliary particles to have opposite helicity, (x^+, y^+) and (u^-, v^-) . The corresponding quantity is

$$\langle xy \rangle^2 [uv]^2 \int \frac{1}{\text{vol GL}(2)} \frac{d^2\sigma_x d^2\sigma_y d^2\sigma_u d^2\sigma_v}{(xy)^2 (uv)^2} \prod_{i=x,y} \delta^{(2)}(\lambda_i - \lambda(\sigma_i)) \prod_{J=u,v} \delta^{(2)}(\tilde{\lambda}_J - \tilde{\lambda}(\sigma_J)), \tag{6.6}$$

with

$$\lambda(\sigma) := \sum_{J=u,v} \frac{\lambda_J}{(\sigma \sigma_J)}, \quad \tilde{\lambda}(\sigma) := \sum_{i=x,y} \frac{\tilde{\lambda}_i}{(\sigma_i \sigma)}. \tag{6.7}$$

An explicit calculation shows that (6.6) is equal to

$$q^4 \delta^{(4)}(p_x + p_y + p_u + p_v). \tag{6.8}$$

with $q = p_x + p_y$. This is not quite the result one expects to find for $\langle \mathcal{O}(q) \bar{\mathcal{O}}(q') \rangle$ from Equation (26) of [35], namely $\langle \mathcal{O}(q) \bar{\mathcal{O}}(q') \rangle \sim \delta^{(4)}(q + q') q^4 \log(q^2) + \text{analytic terms}$. Note in particular that the $\log q^2$ term is absent. In order to be able to derive such terms one may need to understand scattering equations for off-shell quantities at loop level, along the lines of [36–38]. It would also be very interesting to find concrete vertex operators in the ambitwistor string construction of (2.11) that correspond to the operator insertions on the field theory side, possibly making contact with the recent ideas in [22–25]. We hope to report on some of these ideas in a future publication.

Acknowledgments

We would like to thank Paul Heslop and Brenda Penante for very interesting discussions. This work was supported by the Science and Technology Facilities Council (STFC) Consolidated Grant ST/L000415/1 “String theory, gauge theory & duality”. The work of EH was supported by an STFC quota studentship.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] R. Roiban, M. Spradlin and A. Volovich, *On the tree level S matrix of Yang-Mills theory*, *Phys. Rev. D* **70** (2004) 026009 [[hep-th/0403190](#)] [[INSPIRE](#)].
- [2] E. Witten, *Perturbative gauge theory as a string theory in twistor space*, *Commun. Math. Phys.* **252** (2004) 189 [[hep-th/0312171](#)] [[INSPIRE](#)].
- [3] R. Britto, F. Cachazo and B. Feng, *New recursion relations for tree amplitudes of gluons*, *Nucl. Phys. B* **715** (2005) 499 [[hep-th/0412308](#)] [[INSPIRE](#)].
- [4] R. Britto, F. Cachazo, B. Feng and E. Witten, *Direct proof of tree-level recursion relation in Yang-Mills theory*, *Phys. Rev. Lett.* **94** (2005) 181602 [[hep-th/0501052](#)] [[INSPIRE](#)].

- [5] J. Bedford, A. Brandhuber, B.J. Spence and G. Travaglini, *A Recursion relation for gravity amplitudes*, *Nucl. Phys. B* **721** (2005) 98 [[hep-th/0502146](#)] [[INSPIRE](#)].
- [6] F. Cachazo and P. Svrček, *Tree level recursion relations in general relativity*, [hep-th/0502160](#) [[INSPIRE](#)].
- [7] N. Arkani-Hamed and J. Kaplan, *On Tree Amplitudes in Gauge Theory and Gravity*, *JHEP* **04** (2008) 076 [[arXiv:0801.2385](#)] [[INSPIRE](#)].
- [8] M. Spradlin and A. Volovich, *From Twistor String Theory To Recursion Relations*, *Phys. Rev. D* **80** (2009) 085022 [[arXiv:0909.0229](#)] [[INSPIRE](#)].
- [9] N. Arkani-Hamed, F. Cachazo, C. Cheung and J. Kaplan, *The S-matrix in Twistor Space*, *JHEP* **03** (2010) 110 [[arXiv:0903.2110](#)] [[INSPIRE](#)].
- [10] F. Cachazo, S. He and E.Y. Yuan, *Scattering of Massless Particles in Arbitrary Dimensions*, *Phys. Rev. Lett.* **113** (2014) 171601 [[arXiv:1307.2199](#)] [[INSPIRE](#)].
- [11] F. Cachazo, S. He and E.Y. Yuan, *Scattering of Massless Particles: Scalars, Gluons and Gravitons*, *JHEP* **07** (2014) 033 [[arXiv:1309.0885](#)] [[INSPIRE](#)].
- [12] Y. Geyer, A.E. Lipstein and L.J. Mason, *Ambitwistor Strings in Four Dimensions*, *Phys. Rev. Lett.* **113** (2014) 081602 [[arXiv:1404.6219](#)] [[INSPIRE](#)].
- [13] S. He, Z. Liu and J.-B. Wu, *Scattering Equations, Twistor-string Formulas and Double-soft Limits in Four Dimensions*, *JHEP* **07** (2016) 060 [[arXiv:1604.02834](#)] [[INSPIRE](#)].
- [14] S. He and Y. Zhang, *Connected formulas for amplitudes in standard model*, [arXiv:1607.02843](#) [[INSPIRE](#)].
- [15] F. Wilczek, *Decays of Heavy Vector Mesons Into Higgs Particles*, *Phys. Rev. Lett.* **39** (1977) 1304 [[INSPIRE](#)].
- [16] M.A. Shifman, A.I. Vainshtein, M.B. Voloshin and V.I. Zakharov, *Low-Energy Theorems for Higgs Boson Couplings to Photons*, *Sov. J. Nucl. Phys.* **30** (1979) 711 [[INSPIRE](#)].
- [17] L.J. Dixon, E.W.N. Glover and V.V. Khoze, *MHV rules for Higgs plus multi-gluon amplitudes*, *JHEP* **12** (2004) 015 [[hep-th/0411092](#)] [[INSPIRE](#)].
- [18] D. Neill, *Two-Loop Matching onto Dimension Eight Operators in the Higgs-Gluon Sector*, [arXiv:0908.1573](#) [[INSPIRE](#)].
- [19] A. Brandhuber, G. Travaglini and G. Yang, *Analytic two-loop form factors in $N = 4$ SYM*, *JHEP* **05** (2012) 082 [[arXiv:1201.4170](#)] [[INSPIRE](#)].
- [20] S. Dawson, I.M. Lewis and M. Zeng, *Effective field theory for Higgs boson plus jet production*, *Phys. Rev. D* **90** (2014) 093007 [[arXiv:1409.6299](#)] [[INSPIRE](#)].
- [21] A. Brandhuber, M. Kostacinska, B. Penante, G. Travaglini and D. Young, *The $SU(2|3)$ dynamic two-loop form factors*, *JHEP* **08** (2016) 134 [[arXiv:1606.08682](#)] [[INSPIRE](#)].
- [22] L. Koster, V. Mitev, M. Staudacher and M. Wilhelm, *Composite Operators in the Twistor Formulation of $N = 4$ Supersymmetric Yang-Mills Theory*, *Phys. Rev. Lett.* **117** (2016) 011601 [[arXiv:1603.04471](#)] [[INSPIRE](#)].
- [23] D. Chicherin and E. Sokatchev, *Demystifying the twistor construction of composite operators in $N = 4$ super-Yang-Mills theory*, [arXiv:1603.08478](#) [[INSPIRE](#)].
- [24] L. Koster, V. Mitev, M. Staudacher and M. Wilhelm, *All tree-level MHV form factors in $N = 4$ SYM from twistor space*, *JHEP* **06** (2016) 162 [[arXiv:1604.00012](#)] [[INSPIRE](#)].

- [25] D. Chicherin and E. Sokatchev, *Composite operators and form factors in $N = 4$ SYM*, [arXiv:1605.01386](#) [[INSPIRE](#)].
- [26] A. Brandhuber, B. Spence, G. Travaglini and G. Yang, *Form Factors in $N = 4$ Super Yang-Mills and Periodic Wilson Loops*, *JHEP* **01** (2011) 134 [[arXiv:1011.1899](#)] [[INSPIRE](#)].
- [27] N. Arkani-Hamed, F. Cachazo, C. Cheung and J. Kaplan, *A Duality For The S Matrix*, *JHEP* **03** (2010) 020 [[arXiv:0907.5418](#)] [[INSPIRE](#)].
- [28] N. Arkani-Hamed, J.L. Bourjaily, F. Cachazo, A.B. Goncharov, A. Postnikov and J. Trnka, *Scattering Amplitudes and the Positive Grassmannian*, Cambridge University Press, Cambridge U.K. (2012) [[arXiv:1212.5605](#)] [[INSPIRE](#)].
- [29] R. Frassek, D. Meidinger, D. Nandan and M. Wilhelm, *On-shell diagrams, Grassmannians and integrability for form factors*, *JHEP* **01** (2016) 182 [[arXiv:1506.08192](#)] [[INSPIRE](#)].
- [30] L.V. Bork and A.I. Onishchenko, *Wilson lines, Grassmannians and Gauge Invariant Off-shell Amplitudes in $N = 4$ SYM*, [arXiv:1607.02320](#) [[INSPIRE](#)].
- [31] N. Arkani-Hamed, J. Bourjaily, F. Cachazo and J. Trnka, *Unification of Residues and Grassmannian Dualities*, *JHEP* **01** (2011) 049 [[arXiv:0912.4912](#)] [[INSPIRE](#)].
- [32] A. Brandhuber, O. Gürdogan, R. Mooney, G. Travaglini and G. Yang, *Harmony of Super Form Factors*, *JHEP* **10** (2011) 046 [[arXiv:1107.5067](#)] [[INSPIRE](#)].
- [33] L.V. Bork and A.I. Onishchenko, *Grassmannians and form factors with $q^2 = 0$ in $N = 4$ SYM theory*, [arXiv:1607.00503](#) [[INSPIRE](#)].
- [34] L. Mason and D. Skinner, *Ambitwistor strings and the scattering equations*, *JHEP* **07** (2014) 048 [[arXiv:1311.2564](#)] [[INSPIRE](#)].
- [35] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from noncritical string theory*, *Phys. Lett. B* **428** (1998) 105 [[hep-th/9802109](#)] [[INSPIRE](#)].
- [36] Y. Geyer, L. Mason, R. Monteiro and P. Tourkine, *Loop Integrands for Scattering Amplitudes from the Riemann Sphere*, *Phys. Rev. Lett.* **115** (2015) 121603 [[arXiv:1507.00321](#)] [[INSPIRE](#)].
- [37] Y. Geyer, L. Mason, R. Monteiro and P. Tourkine, *One-loop amplitudes on the Riemann sphere*, *JHEP* **03** (2016) 114 [[arXiv:1511.06315](#)] [[INSPIRE](#)].
- [38] Y. Geyer, L. Mason, R. Monteiro and P. Tourkine, *Two-Loop Scattering Amplitudes from the Riemann Sphere*, [arXiv:1607.08887](#) [[INSPIRE](#)].