



## **Reduced Primary Cilia Length and Altered Arl13b Expression Are Associated with Deregulated Chondrocyte Hedgehog Signalling in Alkaptonuria.**

Thorpe, SD; Gambassi, S; Thompson, CL; Chandrakumar, C; Santucci, A; Knight, MM

© The Author(s) 2017

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

For additional information about this publication click this link.

<http://qmro.qmul.ac.uk/xmlui/handle/123456789/19413>

Information about this research object was correct at the time of download; we occasionally make corrections to records, please therefore check the published record when citing. For more information contact [scholarlycommunications@qmul.ac.uk](mailto:scholarlycommunications@qmul.ac.uk)

## ORIGINAL RESEARCH ARTICLE

Reduced primary cilia length and altered Arl13b expression are associated with deregulated chondrocyte hedgehog signalling in alkaptonuria<sup>††</sup>

Stephen D Thorpe<sup>1</sup>, Silvia Gambassi<sup>2</sup>, Clare L Thompson<sup>1</sup>, Charmilie Chandrakumar<sup>1</sup>, Annalisa Santucci<sup>2†</sup>, Martin M Knight<sup>1\*</sup>

<sup>1</sup>Institute of Bioengineering, School of Engineering and Materials Science, Queen Mary University of London, London, UK

<sup>2</sup>Dipartimento di Biotecnologie, Chimica e Farmacia, Università degli Studi di Siena, Siena, Italy

<sup>†</sup>Correspondence to: Annalisa Santucci

Address: Università degli Studi di Siena, Dipartimento di Biotecnologie, Chimica e Farmacia, Via Aldo Moro 2, 53100 Siena, Italy

E-mail: annalisa.santucci@unisi.it

Tel: +390577-234958

Fax: +390577-234954

\*Correspondence to: Martin M Knight

Address: School of Engineering and Materials Science, Queen Mary University of London, Mile End Road, London E1 4NS, United Kingdom

Email: m.m.knight@qmul.ac.uk

Tel: +44 (0)20 7882 8868

Running head: Alkaptonuria alters primary cilia

Keywords:

- Chondrocyte
- Actin
- Primary cilium
- Hedgehog signalling
- Alkaptonuria

Contract grant sponsor: Medical Research Council; Contract grant number: MR/L002876/1

Contract grant sponsor: Royal College of Surgeons of England; Intercalated Bachelor of Science Degree in Surgery Award 2016

Contract grant sponsor: Fondazione Telethon; Contract grant number: GGP10058

Contract grant sponsor: Associazione Italiana Malati di Alcaptonuria (AimAKU); Contract grant number: ORPHA263402

The authors have no conflicts of interest to declare.

<sup>††</sup>This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/jcp.25839]

**Additional Supporting Information may be found in the online version of this article.**

**Received 25 November 2016; Revised 11 January 2017; Accepted 2 February 2017**

**Journal of Cellular Physiology**

**This article is protected by copyright. All rights reserved**

**DOI 10.1002/jcp.25839**

## Abstract

Alkaptonuria (AKU) is a rare inherited disease resulting from a deficiency of the enzyme homogentisate 1,2-dioxygenase which leads to the accumulation of homogentisic acid (HGA). AKU is characterised by severe cartilage degeneration, similar to that observed in osteoarthritis. Previous studies suggest that AKU is associated with alterations in cytoskeletal organisation which could modulate primary cilia structure/function. This study investigated whether AKU is associated with changes in chondrocyte primary cilia and associated Hedgehog signalling which mediates cartilage degradation in osteoarthritis. Human articular chondrocytes were obtained from healthy and AKU donors. Additionally, healthy chondrocytes were treated with HGA to replicate AKU pathology (+HGA). Diseased cells exhibited shorter cilia with length reductions of 36% and 16% in AKU and +HGA chondrocytes respectively when compared to healthy controls. Both AKU and +HGA chondrocytes demonstrated disruption of the usual cilia length regulation by actin contractility. Furthermore, the proportion of cilia with axoneme breaks and bulbous tips was increased in AKU chondrocytes consistent with defective regulation of ciliary trafficking. Distribution of the hedgehog-related protein Arl13b along the ciliary axoneme was altered such that its localisation was increased at the distal tip in AKU and +HGA chondrocytes. These changes in cilia structure/trafficking in AKU and +HGA chondrocytes were associated with a complete inability to activate hedgehog signalling in response to exogenous ligand. Thus, we suggest that altered responsiveness to Hedgehog, as a consequence of cilia dysfunction, may be a contributing factor in the development of arthropathy highlighting the cilium as a novel target in AKU. This article is protected by copyright. All rights reserved

## Introduction

Alkaptonuria (AKU) is an ultra-rare metabolic disease in which a deficiency of the enzyme homogentisate 1,2-dioxygenase leads to the accumulation of homogentisic acid (HGA) (La Du et al., 1958); for review see Gallagher et al. (2016). The oxidised HGA derivative benzoquinone acetic acid (BQA) forms a melanin-like pigmentation known as 'ochronosis', causing dramatic tissue degeneration. The primary clinical manifestation of AKU is a severe form of early onset arthropathy, characterised by articular cartilage degeneration (Helliwell et al., 2008; Mannoni et al., 2004). Within the joint, articular cartilage covers the bone surfaces where it has a protective, low friction, load distributing function (Mow et al., 1999). This cartilage tissue matrix is largely comprised of a dense collagen network in which the highly-hydrated glycosaminoglycan, aggrecan, is immobilised. Chondrocytes reside within this matrix and are responsible for maintaining tissue homeostasis in response to mechanical cues. Disruption of this process results in the degenerative disease osteoarthritis (OA) which is characterised by progressive failure of the articular cartilage accompanied by changes in the synovium and subchondral bone that result in loss of mobility and significant pain (Goldring and Goldring, 2007). Despite being the world's most common form of arthropathy, there is currently no treatment to prevent OA.

Many similarities between the pathogenesis of AKU and OA are apparent. A significant inflammatory contribution is present in OA (Goldring and Otero, 2011), while it has also been shown that inflammatory cytokine expression is altered in AKU chondrocytes resulting in severe inflammation (Braconi et al., 2012; Mannoni et al., 2004). Moreover, significant protein oxidation and aggregation is observed in these cells which, in addition to promoting the production of ochronotic pigments, results in the altered expression of proteins involved in cell defence, protein folding and cell organisation (Braconi et al., 2011). In both OA and AKU tissue, chondrocytes undergo a particular form of cell death termed chondroptosis (Millucci et al., 2015; Perez et al., 2005). Chondrocytes first appear enlarged and hypertrophic with a prominent Golgi apparatus, characterised by swollen cisternae. This organelle is structurally and functionally connected with the cytoskeleton, as microtubules and actin filaments are fundamental for the maintenance of its structural integrity (Thyberg and Moskalewski, 1999; Valderrama et al., 1998). It is well established that the cytoskeleton is disrupted in OA chondrocytes (Capin-Gutierrez et al., 2004; Haudenschild et al., 2011; Holloway et al., 2004). Similarly, Geminiani *et al.* (2016) recently reported that both actin and microtubule cytoskeletal elements are disordered in AKU chondrocytes. In addition to alterations in cell shape, actin stress fibre formation is increased in AKU chondrocytes accompanied by the formation of actin bundles in the nuclear periphery (Geminiani et al., 2016). The disruption of actin organisation and dynamics may have consequences for the maintenance of chondrocyte phenotype (Brown and Benya, 1988), mechanotransduction (Ohashi et al., 2006; Wright et al., 1997) and may also influence chondrocyte function through the primary cilium.

The primary cilium is a singular, immotile organelle present in the majority of cells during interphase. It consists of a microtubule-based axoneme covered by a specialized plasma membrane; for review see Satir and Christensen (2008). This tubulin-based structure is involved in many biological processes, including differentiation and vertebrate development (Huangfu et al., 2003; Tummala et al., 2010), cell cycle control (Tucker et al., 1979), cancer signalling (Reilova-Velez and Seiler, 1984; Wong et al., 2009), sensory function and migration (Schneider et al., 2005), and mechano-signalling (Khayyeri et al., 2015; Praetorius and Spring, 2001; Wann et al., 2012). The cilium is assembled and maintained by a process called intraflagellar transport (IFT), a microtubule-based motility present in the axoneme. IFT transports axonemal precursors and signalling proteins along the length of the cilium towards the distal tip and returns them to the basal body (Qin et al., 2004). This mechanism of transport is essential not only for maintenance of cilia structure but for the regulation of cilia-mediated signalling pathways such as Hedgehog (Huangfu et al., 2003), Wnt (Ross et al., 2005) and TGF (Clement et al., 2013). Alterations in IFT lead to disruption of cilia structure and function and are responsible for a group of related disorders termed ciliopathies (Waters and Beales, 2011). The regulation of ciliogenesis is dependent on a number of actin-regulatory molecules (Kim et al., 2010). Moreover, the regulation of primary cilia length has been linked to cell shape and the level of actin contractility such that greater actin tension is associated with cilia shortening (McMurray et al., 2013; Pitaval et al., 2010). Cilia length modifications have been reported in several pathological conditions including OA where an increase in axoneme length is observed (McGlashan et al., 2008).

Hedgehog signalling is regulated through the primary cilium (Corbit et al., 2005; Huangfu and Anderson, 2005; Rohatgi et al., 2007). Binding of hedgehog ligands to the receptor Patched (Ptch1) triggers the IFT-dependent accumulation of the transmembrane protein Smoothed (Smo) within the cilium (Corbit et al., 2005). This process is mediated by several proteins, among which is the GTPase Arl13b (Caspary et al., 2007; Larkins et al., 2011). Smo ciliary distribution regulates the activity of transcription factors belonging to the Gli family (Gli1, Gli2, Gli3), the effectors of this pathway (Buttitta et al., 2003). Non-canonical hedgehog signalling can also regulate actin cytoskeleton organisation via the RhoA pathway, independent of transcription (Polizio et al., 2011; Renault et al., 2010; Sasaki et al., 2010). Recent studies have demonstrated that chondrocyte hedgehog signalling can be influenced by alterations in cilia length (Thompson et al., 2014; Thompson et al., 2016) as the size of this signalling compartment is expected to influence the rate at which proteins are delivered to or removed from the cilia tip (Engel et al., 2009; Ludington et al., 2013). Given its association with cilia length, actin cytoskeletal organisation may be implicated in aberrant hedgehog signalling. Several studies have now identified activated hedgehog signalling as an important pathological factor in OA (Lin et al., 2009; Wei et al., 2012). Activation of this pathway in osteoarthritic cartilage promotes chondrocyte hypertrophy and matrix catabolism *in vivo* (Lin et al., 2009; Wei et al., 2012). However, the contribution of hedgehog signalling to arthropathy in AKU has not previously been reported.

This study aims to investigate the effects of HGA accumulation in AKU on primary cilia structure and hedgehog signalling. In addition to chondrocytes isolated from AKU donor cartilage, we treated healthy donor chondrocytes with HGA (+HGA) to replicate the disease state. Given the paucity of AKU disease tissue, this model of AKU with HGA addition has been implemented *in vivo* through direct injection into rabbit knee joints and chick embryos (Moran and Yunis, 1962), and more widely *in vitro* in cells from tissues including cartilage and the nervous system (Bernardini et al., 2015; Braconi et al., 2015; Mistry et al., 2015). These studies are the first to demonstrate that primary cilia length and prevalence are reduced in AKU chondrocytes, and chondrocytes treated with HGA, relative to healthy control cells. Consistent with previous reports we show that actin organisation is disrupted in AKU and HGA treated chondrocytes and report that actin-dependent cilia length regulation is disrupted. This is accompanied by altered cilia structure and trafficking such that there is an increase in the proportion of cilia exhibiting axoneme breaks and bulbous tips in AKU cells. Super resolution microscopy reveals an increase in the volume of cilioplasm at the ciliary tip which is associated with accumulation of the hedgehog-related protein Arl13b within the tip region. Furthermore, we identify that AKU and HGA treated cells exhibit a complete inability to activate hedgehog signalling in response to exogenous ligands which we propose may be the result of the observed dysfunction of cilia structure and trafficking.

## Materials and Methods

### Isolation and culture of human articular chondrocytes

Human articular chondrocytes were obtained from knee cartilage of healthy donors and those with alkaptonuria (AKU) with full local ethics approval (CEL AOUS 21.07.2010). Chondrocytes were isolated by enzymatic digestion as previously described (Braconi et al., 2012) and cultured in Dulbecco's Modified Eagle Medium (DMEM; Life Technologies, Paisley, UK) supplemented with 10% foetal bovine serum (FBS) and penicillin (100 U/ml) – streptomycin (100 µg/ml; all Sigma-Aldrich, Dorset, UK). To induce an AKU-like state, healthy chondrocytes were also treated with 66 µM HGA (Sigma-Aldrich) for 7 days (+HGA). Culture media was exchanged every 2-3 days and cultures were maintained at 37°C and 5% CO<sub>2</sub>. Chondrocytes were seeded at approx. 10×10<sup>3</sup> cells/cm<sup>2</sup> onto serum coated glass coverslips in 24-well plates for immunofluorescence experiments, and in 6-well plates for gene expression studies. To disrupt actin stress fibre contractility, chondrocytes were treated with the Rho-associated protein kinase (ROCK) inhibitor Y27632 (Y27, 10 µM; Sigma-Aldrich) in the absence of FBS for 24 hrs and fixed.

### Immunofluorescent staining

Confluent chondrocytes were serum starved (0% FBS) for 24 hours prior to fixation in 4% paraformaldehyde for 10 min. Cells were permeabilised with 0.5% Triton X-100 in phosphate buffered saline (PBS) for 5 min and blocked with 5% goat serum in 0.1% bovine serum albumin-PBS (0.1% BSA-PBS; all Sigma-Aldrich) for 1 hour. Cells were incubated with primary antibodies in 0.1% BSA-PBS at 4°C overnight. Mouse monoclonal

anti-acetylated  $\alpha$ -tubulin (clone 611B-1, 1:2000; Sigma Aldrich, Cat# T7451, RRID: AB\_609894) and rabbit polyclonal anti-Arl13b (1:2000; Proteintech, Manchester, UK, Cat# 17711-1-AP, RRID: AB\_2060867) were used for the detection of the ciliary axoneme. Rabbit polyclonal anti-pericentrin (1:500; Abcam, Cambridge, UK, Cat# ab4448, RRID: AB\_304461) was used alongside with anti-acetylated  $\alpha$ -tubulin for the detection of centriole-derived basal bodies. Following repeated washing in 0.1% BSA-PBS, cells were incubated with appropriate Alexa Fluor-conjugated secondary antibodies (1:1000; Life Technologies) for 1 hour at room temperature. F-actin was detected using CruzFluor 488-conjugated phalloidin (1:1000; Santa Cruz Biotechnology, Heidelberg, Germany, Cat# sc-363791, RRID: AB\_2631056) in 1% BSA-PBS and nuclei were detected with 1  $\mu$ g/ml 4',6-diamidino-2-phenylindole (DAPI; Sigma-Aldrich) in PBS. Following washes in PBS, coverslips were mounted with Fluoromount-G (Cambridge Bioscience, Cambridge, UK).

### Confocal and super-resolution microscopy and image analysis

For primary cilia length measurement, samples were imaged using a Leica TCS SP2 confocal microscope with a  $\times 63$  1.4 NA objective (Leica Microsystems, Milton Keynes, UK). Confocal z-stacks were generated containing the entire cell depth with a voxel size of  $116 \times 116 \times 500$  nm in x-y-z. The maximum intensity projection in the z-axis was used for the measurement of cilia length and assessment of morphology using ImageJ software (NIH, Bethesda, MD, USA, RRID: SCR\_003070). For primary cilia prevalence, samples were imaged using a Leica DMI4000B epifluorescent microscope with a  $\times 63$  1.25 NA objective (Leica Microsystems). The proportion of ciliated cells was determined across multiple fields of view for each condition. Structural abnormalities were manually assessed on acetylated  $\alpha$ -tubulin stained cilia as a proportion of total cilia for each condition. Cilia tips were defined as the region distal to pericentrin staining of the basal body region (Fig. 3A). Bulbous tipped cilia were defined as having a tip diameter  $>1.5 \times$  diameter at axoneme midpoint. Fragmented cilia were defined as cilia with clear breaks in acetylated  $\alpha$ -tubulin staining along the axoneme. Examples are presented in Fig. 3A.

Arl13b and acetylated  $\alpha$ -tubulin stained cilia were imaged using super-resolution structured illumination microscopy (SR-SIM). Samples were imaged on a Zeiss 710 ELYRA PS.1 microscope with a  $\times 63$  1.4 NA objective (Carl Zeiss, Cambridge, UK). SR-SIM processing and channel alignment were performed to obtain a voxel size of  $25 \times 25 \times 100$  nm in x-y-z. Maximum intensity projections in the z-axis were used to generate intensity profiles perpendicular to the axoneme close to the base and tip of individual primary cilia as shown in Fig. 3D. Cilia diameter was assessed from Arl13b and acetylated  $\alpha$ -tubulin signals as full width at half maximum of the intensity profile; representatives of which are shown in Fig. 3E. Mean and integrated intensity of each marker was assessed for each profile at both the base and tip of the cilium.

### Evaluation of hedgehog pathway gene expression in response to ligand stimulation

The expression of the hedgehog target proteins GLI1 and PTCH1 is increased upon pathway activation (Buttitta et al., 2003; Dai et al., 1999). Gene expression was evaluated in untreated chondrocytes from the

three groups (Healthy, AKU and +HGA) and chondrocytes from each group treated for 24 hrs with 1 µg/ml recombinant Indian hedgehog (Ihh; R&D Systems, Abingdon, UK) to measure pathway activity. Total RNA was extracted from isolated human chondrocytes cultured in single wells using an RNeasyKit and converted to cDNA using the QuantiTect Reverse Transcription Kit (both Qiagen, Manchester, UK) according to the manufacturer's instructions. Both RNA and cDNA were quantified using the Nanodrop ND-1000 spectrophotometer (LabTech, East Sussex, UK). Quantitative real-time PCR reactions were performed in 10 µl volumes containing 1 µl cDNA (diluted 1:2), 5 µl KAPA SYBR<sup>®</sup> FAST Universal 2× qPCR Master Mix containing SYBR-green dye and ROX reference dye (KAPA Biosystems, London, UK), and 1 µl optimised primer pairs (Table 1). An annealing temperature of 60°C was used for all PCR reactions and fluorescence data was collected using the MX3000P QPCR instrument (Agilent Technologies, Cheshire, UK). Samples were run in triplicate to minimise pipetting errors. Data was analysed using the relative standard curve method and target genes were normalised to 18s RNA.

### Statistical analysis

Statistical analyses were performed using Minitab 17 software (Minitab, Coventry, UK, RRID: SCR\_014483). When data sets adhered to a normal distribution, two sample t-test, or a general linear model for analysis of variance with Fisher tests for multiple comparisons were used. For non-parametric data sets, Mann-Whitney tests were used to compare conditions. Data quoted in the text are presented as mean±s.e.m. Details of specific statistical tests and *n* values can be found in the figure legends.

## Results

### F-actin cytoskeleton is altered in AKU chondrocytes

Chondrocytes isolated from healthy donors exhibited some faint stress fibre formation with minimal formation of f-actin aggregates (Fig. 1A). In contrast, AKU and HGA treated (+HGA) chondrocytes exhibited an increased intensity of f-actin labelling with more stress fibres present and increased incidence of punctate f-actin aggregates, suggesting a more contractile phenotype (Fig. 1A).

### Chondrocyte primary cilia length and prevalence is reduced in AKU

Immunofluorescent labelling of primary cilia (Fig. 1B) revealed that the proportion of cells exhibiting a cilium was dramatically reduced from 90.4±1.5% in healthy control cells to 55.2±2.9% in AKU chondrocytes (Fig. 1C). Moreover, cilia length was significantly reduced in these cells relative to the healthy control (Fig. 1D). Cilia prevalence and length were significantly reduced in +HGA chondrocytes, but to a lesser degree than observed in AKU cells (Fig. 1C,D).

### Actin-dependent regulation of cilia length is disrupted in AKU chondrocytes

It is well established that actin contractility influences primary cilia length (Kim et al., 2010; McMurray et al., 2013; Pitaval et al., 2010). Due to the observed alterations in actin cytoskeleton (Fig. 1), we investigated



the effects of ROCK inhibition on cilia length and prevalence in these cells (Fig. 2A). Chondrocytes were treated with the ROCK inhibitor Y27632 (Y27) to disrupt actin stress fibre contractility. ROCK inhibition resulted in a reduction in stress fibre formation in all cultures (data not shown). No significant effects on primary cilia prevalence were observed in control, AKU or +HGA chondrocytes (Fig. 2B). Consistent with previous reports, primary cilia length was found to be significantly increased in healthy control chondrocytes by 8.5% from  $2.55 \pm 0.07$  to  $2.76 \pm 0.07$   $\mu\text{m}$  (Fig. 2C). However, ROCK inhibition did not affect cilia length in AKU chondrocytes while in +HGA chondrocytes a significant reduction in length was observed (Fig. 2C). These data suggest that the regulation of cilia length through alterations in actin cytoskeletal tension and actin dynamics is disrupted in AKU and reversed in +HGA chondrocytes.

### Primary cilia structure and protein localisation is altered in AKU chondrocytes

Primary cilia structure was found to be altered in AKU and +HGA chondrocytes compared to healthy control cells. A significant increase in the proportion of cilia exhibiting bulbous tips (Fig. 3A & B) and axoneme breaks (Fig. 3A & C) was observed in AKU suggesting that ciliary trafficking and maintenance are disrupted. +HGA chondrocytes did not exhibit an increase morphological alterations compared to healthy controls. Consistent with these findings we report that the ciliary localisation of the hedgehog related protein ARL13B is altered such that a significant increase in protein localisation at the ciliary tip was observed in both AKU and +HGA chondrocytes, with +HGA resulting in the greatest increase (Fig. 3D-F and Supplementary Fig. 1).

Closer examination of the structure of normal (not bulbous or fragmented) primary cilia using super-resolution structured illumination microscopy (SR-SIM) revealed that microtubule diameter was increased at the ciliary tip in AKU and +HGA chondrocytes (Fig. 3G). Similarly, examination of ciliary diameter using the membrane protein Arl13b confirmed that total ciliary tip diameter was also increased in these cells (Fig. 3H). Consequently, we report that the amount of cilioplasm within the tip region (calculated as the difference in Arl13b and Ac- $\alpha$ -Tub diameters) is increased (Fig. 3I). This trend is consistent with the increased presence of bulbous tips in AKU chondrocytes (Fig. 3B). These data further support the hypothesis that ciliary trafficking is altered in AKU and that aspects of this ciliary phenotype can be replicated by treatment with HGA.

### AKU chondrocytes are unresponsive to exogenous hedgehog ligands

Arl13b is a key regulator of the ciliary trafficking of hedgehog proteins (Caspary et al., 2007; Larkins et al., 2011; Mariani et al., 2016). Moreover we have recently demonstrated that alterations in cilia length and Arl13b cilia localisation are associated with altered hedgehog signalling in chondrocytes (Thompson et al., 2016). Hedgehog signalling was therefore examined in AKU and +HGA chondrocytes under basal conditions and in response to recombinant Indian hedgehog (Ihh) using real time PCR. Under basal conditions, the mRNA levels of GLI1 were significantly decreased in AKU but not +HGA chondrocytes relative to healthy

control cells (Fig. 4A) while PTCH1 expression was not altered (Fig. 4B). In healthy control cells, expression of GLI1 and PTCH1 were significantly increased in response to Ihh treatment for 24hrs relative to untreated cells by 1.77- and 1.54-fold respectively (Fig. 4). By contrast the expression of GLI1 and PTCH1 were not significantly altered in response to Ihh in AKU or +HGA chondrocytes, demonstrating that these cells are unresponsive to ligand treatment.

## Discussion

These studies are the first to show that primary cilia structure and function are disrupted in the ultra-rare metabolic disorder AKU, and that aspects of this cilia phenotype can be replicated by treatment with HGA. Subsequent studies by the authors have investigated the effect of different Hedgehog pathway antagonists on cilia structure and hedgehog signalling in HGA treated chondrocytes. In the present study, we report variations in primary cilia prevalence and length for AKU chondrocytes which may be associated with alterations in the organisation of actin (Fig. 1) and other cytoskeletal components (Geminiani et al., 2016). The filamentous microtubule structure is transformed in AKU chondrocytes, forming aggregates at the cell periphery (Geminiani et al., 2016). Furthermore, the vimentin network appears granular in these cells resulting in alterations in cell shape and cell shrinkage. Previous studies have also shown that AKU induces 4-HNE expression and lipid peroxidation which alters microtubule function (Neely et al., 2005; Neely et al., 1999). Interestingly in this study, we demonstrate that the normal regulation of cilia length through the contractility of the actin cytoskeleton is disrupted in AKU cells.

It is worth noting that the cilium itself can regulate cytoskeletal organisation. Hypomorphic mutation of IFT88 results in a reduction in cortical actin organisation and cell stiffness in articular chondrocytes (Wang et al., 2016). Additionally, mutation of several components of the BBSome (a complex of Bardet-Biedl syndrome (BBS) proteins involved in cilia formation) influence actin polymerisation and have been shown to produce dramatic alterations in stress fibre organisation that impact on cell migration, adhesion and division in other cell types (Hernandez-Hernandez et al., 2013). Thus, it is also possible that changes in actin cytoskeletal organisation in AKU chondrocytes are downstream of alterations in primary cilia structure and trafficking.

In addition to ciliary shortening, other structural changes to the axoneme were observed. In healthy cells, primary cilia diameter, as measured from microtubule staining, typically decreases towards the ciliary tip. Electron micrographs have revealed that this is due to the conversion of microtubule doublets to singlets in the distal portion of the axoneme (Silverman and Leroux, 2009; Tanuma and Ohata, 1978; Wen et al., 1982). Indeed, it is suggested that these singlet microtubules themselves have dedicated motors for transporting proteins along this distal region of the axoneme (Silverman and Leroux, 2009). In AKU and +HGA chondrocytes, ciliary diameter did not decrease to the same extent as healthy controls, perhaps

suggesting more doublet microtubules extend to the tip in these shorter cilia. Furthermore, thickening of the cilium was observed in AKU cells, as measured from the ciliary membrane marker ARL13B and an increased proportion of cilia exhibited swollen, or bulbous tips (Fig. 3). Together these findings are consistent with disrupted ciliary trafficking in AKU resulting in reduced cilia length and accumulation of ARL13B at the cilia tip.

Due to the shared nature of the IFT system, alterations in ciliary trafficking in AKU would consequently influence the import/export of ciliary proteins and therefore impact upon the dynamics of cilia signalling. This may provide an explanation for the abolition of hedgehog signalling observed in AKU and +HGA chondrocytes. However, the relationship between cilia length and Hedgehog signalling is complex. The current literature suggests that cilia – in a state of disassembly due to reduced trafficking – will have reduced Hedgehog signalling associated with shorter cilia. However, cells with a stable short cilium that have achieved their 'set length' will have an increased rate of IFT delivery and hence exhibit increased hedgehog signalling. This relationship is demonstrated by our previous work showing that loading-induced disassembly of primary cilia results in hedgehog inhibition (Thompson et al., 2014), while in separate studies, cells induced to have longer cilia also have reduced hedgehog signalling (Cruz et al., 2010; Mahjoub and Stearns, 2012; Thompson et al., 2016). Furthermore, complete cilia loss results in over activation of the pathway (Huangfu and Anderson, 2005; Huangfu et al., 2003). Thus, our data presented here suggests that reduced cilia length in AKU and +HGA chondrocytes is most likely the direct result of disrupted ciliary trafficking which also suppresses ligand-induced hedgehog signalling.

An increase in the proportion of cilia that exhibited fragmentation within the distal region of the axoneme was observed in AKU chondrocytes (Fig. 3C). It has been postulated that this fragmentation may represent ciliary ectosomes containing molecules for long distance intercellular signalling (Hogan et al., 2009; Nager et al., 2016). Alternatively, this vesicle shedding may function as a means to remove excess or unwanted ciliary proteins as a mechanism of disposal and thus modulate cilia-dependent signalling pathways (for review see Wood and Rosenbaum (2015)). Indeed, the latter is consistent with both a reduction in hedgehog signalling and the reduced cilia length observed in this study and therefore may represent an alternative hypothesis for the observations reported in this study.

Upon activation of the hedgehog signalling pathway, Gli proteins are trafficked to the ciliary tip where a complex of proteins, including the kinesin-4 protein Kif7, is localised and regulates Gli activity (He et al., 2014). Mislocalisation of these tip proteins results in pathway disruption. One possibility therefore is that within the shorter cilia of AKU cells, the formation of this tip complex is disrupted resulting in defective hedgehog signalling. The effect of disease-dependent disruption of cilia on the hedgehog related protein Arl13b was also analysed and cilia localisation found to be altered. This observation may have important consequences for downstream signalling as Arl13b is required for the ciliary traffic of Ptch1, Smo, Gli2, Gli3, and other Shh-associated proteins in response to ligand stimulation (Larkins et al., 2011). Loss of this

protein can also influence basal hedgehog signalling by promoting the constitutive production of Gli activator proteins, so that an increased ciliary concentration of Arl13b, as observed for AKU cells, might be expected to result in altered levels of Gli1 (Casparly et al., 2007). Of note, Arl13b has also been implicated in cilia length control itself (Casparly et al., 2007; Lu et al., 2015); overexpression of Arl13b results in abnormally long cilia, and increased levels of Arl13b induce protrusion of the ciliary membrane which is rapidly followed by extension of axonemal microtubules (Lu et al., 2015). In the absence of Arl13b, cilia are short and exhibit structural defects in the axonemal microtubules (Casparly et al., 2007), suggesting a complex inter-relationship.

Throughout this study, in addition to AKU cells from patient donors, we have also used a well-established *in vitro* model recapitulating aspects of AKU through supplementation with HGA (+HGA) (Braconi et al., 2015) to investigate the effects of HGA accumulation on primary cilia. While results for +HGA chondrocytes were broadly similar to those observed for AKU chondrocytes, a reduced ciliary response was apparent in the +HGA model despite comparable inhibition of Hedgehog signalling (Figs. 1-4). This difference is likely the result of the extended exposure of AKU cells to increased levels of HGA, whereas in our model, treatment was only conducted for 7 days which is sufficient to reduce proliferation and proteoglycan production, and induce significant deposition of ochronotic pigment (Tinti et al., 2011). However, while deposition of serum amyloid A (SAA), a severe complication of chronic inflammatory conditions such as rheumatoid arthritis and AKU, is present after HGA treatment, this is to a far lesser extent than in AKU cells (Geminiani et al., 2016). This, combined with our data indicating significant differences between AKU and +HGA chondrocytes, demonstrates that while HGA treatment can replicate many facets of AKU pathology, it is an incomplete disease model.

Several studies have now shown that very minor changes in cilia structure and trafficking can have dramatic effects upon functionality (Cruz et al., 2010; Thompson et al., 2014; Thompson et al., 2016; Tran et al., 2008). Thus, while the cilia disruption in +HGA cells does not influence cilia structure to the same extent as in AKU cells, it is sufficient to considerably disrupt function. Indeed, despite a much smaller reduction in length relative to the healthy control for +HGA when compared to AKU (a cilia length reduction of 16% vs. 36% respectively, Fig. 1D), greater accumulation of Arl13b at the ciliary tip was in fact observed in +HGA chondrocytes (Fig. 3F).

It is widely accepted that the disruption of hedgehog signalling in articular cartilage leads to the development of OA (Li et al., 2015b; Lin et al., 2009; Wei et al., 2012). Moreover, the level of activation and aberrant expression of Ihh protein correlates with disease severity (Lin et al., 2009; Shuang et al., 2015). Targeting hedgehog signalling in disease models has shown some efficacy in the prevention of articular cartilage degeneration (Li et al., 2015a; Lin et al., 2009), osteophyte formation (Ruiz-Heiland et al., 2012) and synovitis and synovial hyperplasia (Zhu et al., 2015). Given the dependence of hedgehog signalling upon primary cilia integrity and the disruptions to cilia structure reported in this study, the concept of

'ciliotherapy' (targeting disease pathways through the cilium) is one that could be readily applied to the treatment of cartilage degeneration not only in OA, but in the arthropathy associated with AKU.

### **Acknowledgements**

The authors thank Prof. P. Mariani and Dr. D. Gambera for providing cartilage samples, and Miss R. Patel for technical assistance. This work was supported by the Medical Research Council (MR/L002876/1), the Royal College of Surgeons of England, Fondazione Telethon Italy (GGP10058), and Associazione Italiana Malati di Alcaptonuria (AimAKU, ORPHA263402).

Accepted Article

## Literature cited

- Bernardini G, Laschi M, Geminiani M, Braconi D, Vannuccini E, Lupetti P, Manetti F, Millucci L, Santucci A. 2015. Homogentisate 1,2 dioxygenase is expressed in brain: implications in alkaptonuria. *J Inherit Metab Dis* 38(5):807-814.
- Braconi D, Bernardini G, Bianchini C, Laschi M, Millucci L, Amato L, Tinti L, Serchi T, Chellini F, Spreafico A, Santucci A. 2012. Biochemical and proteomic characterization of alkaptonuric chondrocytes. *J Cell Physiol* 227(9):3333-3343.
- Braconi D, Bianchini C, Bernardini G, Laschi M, Millucci L, Spreafico A, Santucci A. 2011. Redox-proteomics of the effects of homogentisic acid in an in vitro human serum model of alkaptonuric ochronosis. *J Inherit Metab Dis* 34(6):1163-1176.
- Braconi D, Millucci L, Bernardini G, Santucci A. 2015. Oxidative stress and mechanisms of ochronosis in alkaptonuria. *Free Radic Biol Med* 88(Pt A):70-80.
- Brown PD, Benya PD. 1988. Alterations in chondrocyte cytoskeletal architecture during phenotypic modulation by retinoic acid and dihydrocytochalasin B-induced reexpression. *J Cell Biol* 106(1):171-179.
- Buttitta L, Mo R, Hui CC, Fan CM. 2003. Interplays of Gli2 and Gli3 and their requirement in mediating Shh-dependent sclerotome induction. *Development* 130(25):6233-6243.
- Capin-Gutierrez N, Talamas-Rohana P, Gonzalez-Robles A, Lavalle-Montalvo C, Kouri JB. 2004. Cytoskeleton disruption in chondrocytes from a rat osteoarthrotic (OA) -induced model: its potential role in OA pathogenesis. *Histol Histopathol* 19(4):1125-1132.
- Caspary T, Larkins CE, Anderson KV. 2007. The graded response to Sonic Hedgehog depends on cilia architecture. *Dev Cell* 12(5):767-778.
- Clement CA, Ajbro KD, Koefoed K, Vestergaard ML, Veland IR, Henriques de Jesus MP, Pedersen LB, Benmerah A, Andersen CY, Larsen LA, Christensen ST. 2013. TGF-beta signaling is associated with endocytosis at the pocket region of the primary cilium. *Cell Rep* 3(6):1806-1814.
- Corbit KC, Aanstad P, Singla V, Norman AR, Stainier DY, Reiter JF. 2005. Vertebrate Smoothed functions at the primary cilium. *Nature* 437(7061):1018-1021.
- Cruz C, Ribes V, Kutejova E, Cayuso J, Lawson V, Norris D, Stevens J, Davey M, Blight K, Bangs F, Mynett A, Hirst E, Chung R, Balaskas N, Brody SL, Marti E, Briscoe J. 2010. Foxj1 regulates floor plate cilia architecture and modifies the response of cells to sonic hedgehog signalling. *Development* 137(24):4271-4282.
- Dai P, Akimaru H, Tanaka Y, Maekawa T, Nakafuku M, Ishii S. 1999. Sonic Hedgehog-induced Activation of the Gli1 Promoter Is Mediated by GLI3. *J Biol Chem* 274(12):8143-8152.
- Engel BD, Ludington WB, Marshall WF. 2009. Intraflagellar transport particle size scales inversely with flagellar length: revisiting the balance-point length control model. *J Cell Biol* 187(1):81-89.

Gallagher JA, Dillon JP, Sireau N, Timmis O, Ranganath LR. 2016. Alkaptonuria: An example of a "fundamental disease"--A rare disease with important lessons for more common disorders. *Semin Cell Dev Biol* 52:53-57.

Geminiani M, Gambassi S, Millucci L, Lupetti P, Collodel G, Mazzi L, Frediani B, Braconi D, Mazzocchi B, Laschi M, Bernardini G, Santucci A. 2016. Cytoskeleton Aberrations in Alkaptonuric Chondrocytes. *J Cell Physiol* In press.

Goldring MB, Goldring SR. 2007. Osteoarthritis. *J Cell Physiol* 213(3):626-634.

Goldring MB, Otero M. 2011. Inflammation in osteoarthritis. *Current opinion in rheumatology* 23(5):471-478.

Haudenschild DR, Chen J, Pang N, Steklov N, Grogan SP, Lotz MK, D'Lima DD. 2011. Vimentin contributes to changes in chondrocyte stiffness in osteoarthritis. *J Orthop Res* 29(1):20-25.

He M, Subramanian R, Bangs F, Omelchenko T, Liem KF, Jr., Kapoor TM, Anderson KV. 2014. The kinesin-4 protein Kif7 regulates mammalian Hedgehog signalling by organizing the cilium tip compartment. *Nat Cell Biol* 16(7):663-672.

Helliwell TR, Gallagher JA, Ranganath L. 2008. Alkaptonuria--a review of surgical and autopsy pathology. *Histopathology* 53(5):503-512.

Hernandez-Hernandez V, Pravin Kumar P, Diaz-Font A, May-Simera H, Jenkins D, Knight M, Beales PL. 2013. Bardet-Biedl syndrome proteins control the cilia length through regulation of actin polymerization. *Hum Mol Genet* 22(19):3858-3868.

Hogan MC, Manganelli L, Woollard JR, Masyuk AI, Masyuk TV, Tammachote R, Huang BQ, Leontovich AA, Beito TG, Madden BJ, Charlesworth MC, Torres VE, LaRusso NF, Harris PC, Ward CJ. 2009. Characterization of PKD protein-positive exosome-like vesicles. *J Am Soc Nephrol* 20(2):278-288.

Holloway I, Kayser M, Lee DA, Bader DL, Bentley G, Knight MM. 2004. Increased presence of cells with multiple elongated processes in osteoarthritic femoral head cartilage. *Osteoarthritis and Cartilage* 12(1):17-24.

Huangfu D, Anderson KV. 2005. Cilia and Hedgehog responsiveness in the mouse. *Proc Natl Acad Sci U S A* 102(32):11325-11330.

Huangfu D, Liu A, Rakeman AS, Murcia NS, Niswander L, Anderson KV. 2003. Hedgehog signalling in the mouse requires intraflagellar transport proteins. *Nature* 426(6962):83-87.

Khayyeri H, Barreto S, Lacroix D. 2015. Primary cilia mechanics affects cell mechanosensation: A computational study. *J Theor Biol* 379:38-46.

Kim J, Lee JE, Heynen-Genel S, Suyama E, Ono K, Lee K, Ideker T, Aza-Blanc P, Gleeson JG. 2010. Functional genomic screen for modulators of ciliogenesis and cilium length. *Nature* 464(7291):1048-1051.

La Du BN, Zannoni VG, Laster L, Seegmiller JE. 1958. The nature of the defect in tyrosine metabolism in alcaptonuria. *J Biol Chem* 230(1):251-260.

- Larkins CE, Aviles GD, East MP, Kahn RA, Caspary T. 2011. Arl13b regulates ciliogenesis and the dynamic localization of Shh signaling proteins. *Mol Biol Cell* 22(23):4694-4703.
- Li R, Cai L, Ding J, Hu CM, Wu TN, Hu XY. 2015a. Inhibition of hedgehog signal pathway by cyclopamine attenuates inflammation and articular cartilage damage in rats with adjuvant-induced arthritis. *J Pharm Pharmacol* 67(7):963-971.
- Li R, Cai L, Hu CM, Wu TN, Li J. 2015b. Expression of hedgehog signal pathway in articular cartilage is associated with the severity of cartilage damage in rats with adjuvant-induced arthritis. *Journal of inflammation* 12:24.
- Lin AC, Seeto BL, Bartoszko JM, Khoury MA, Whetstone H, Ho L, Hsu C, Ali SA, Alman BA. 2009. Modulating hedgehog signaling can attenuate the severity of osteoarthritis. *Nat Med* 15(12):1421-1425.
- Lu H, Toh MT, Narasimhan V, Thamilselvam SK, Choksi SP, Roy S. 2015. A function for the Joubert syndrome protein Arl13b in ciliary membrane extension and ciliary length regulation. *Dev Biol* 397(2):225-236.
- Ludington WB, Wemmer KA, Lechtreck KF, Witman GB, Marshall WF. 2013. Avalanche-like behavior in ciliary import. *Proc Natl Acad Sci U S A* 110(10):3925-3930.
- Mahjoub MR, Stearns T. 2012. Supernumerary centrosomes nucleate extra cilia and compromise primary cilium signaling. *Curr Biol* 22(17):1628-1634.
- Mannoni A, Selvi E, Lorenzini S, Giorgi M, Airo P, Cammelli D, Andreotti L, Marcolongo R, Porfirio B. 2004. Alkaptonuria, ochronosis, and ochronotic arthropathy. *Seminars in arthritis and rheumatism* 33(4):239-248.
- Mariani LE, Bijlsma MF, Ivanova AI, Suci SK, Kahn RA, Caspary T. 2016. Arl13b regulates Shh signaling from both inside and outside the cilium. *Molecular biology of the cell*.
- McGlashan SR, Cluett EC, Jensen CG, Poole CA. 2008. Primary cilia in osteoarthritic chondrocytes: from chondrons to clusters. *Dev Dyn* 237(8):2013-2020.
- McMurray RJ, Wann AK, Thompson CL, Connelly JT, Knight MM. 2013. Surface topography regulates wnt signaling through control of primary cilia structure in mesenchymal stem cells. *Sci Rep* 3:3545.
- Millucci L, Giorgetti G, Viti C, Ghezzi L, Gambassi S, Braconi D, Marzocchi B, Paffetti A, Lupetti P, Bernardini G, Orlandini M, Santucci A. 2015. Chondroptosis in alkaptonuric cartilage. *J Cell Physiol* 230(5):1148-1157.
- Mistry JB, Jackson DJ, Bukhari M, Taylor AM. 2015. A role for interleukins in ochronosis in a chondrocyte in vitro model of alkaptonuria. *Clin Rheumatol*.
- Moran TJ, Yunis EJ. 1962. Studies on Ochronosis: 2. Effects of Injection of Homogentisic Acid and Ochronotic Pigment in Experimental Animals. *The American Journal of Pathology* 40(3):359-369.
- Mow VC, Wang CC, Hung CT. 1999. The extracellular matrix, interstitial fluid and ions as a mechanical signal transducer in articular cartilage. *Osteoarthritis Cartilage* 7(1):41-58.
- Nager AR, Goldstein JS, Herranz-Perez V, Portran D, Ye F, Garcia-Verdugo JM, Nachury MV. 2016. An Actin Network Dispatches Ciliary GPCRs into Extracellular Vesicles to Modulate Signaling. *Cell*.



- Neely MD, Boutte A, Milatovic D, Montine TJ. 2005. Mechanisms of 4-hydroxynonenal-induced neuronal microtubule dysfunction. *Brain Res* 1037(1-2):90-98.
- Neely MD, Sidell KR, Graham DG, Montine TJ. 1999. The Lipid Peroxidation Product 4-Hydroxynonenal Inhibits Neurite Outgrowth, Disrupts Neuronal Microtubules, and Modifies Cellular Tubulin. *Journal of Neurochemistry* 72(6):2323-2333.
- Ohashi T, Hagiwara M, Bader DL, Knight MM. 2006. Intracellular mechanics and mechanotransduction associated with chondrocyte deformation during pipette aspiration. *Biorheology* 43(3-4):201-214.
- Perez HE, Luna MJ, Rojas ML, Kouri JB. 2005. Chondroptosis: an immunohistochemical study of apoptosis and Golgi complex in chondrocytes from human osteoarthritic cartilage. *Apoptosis : an international journal on programmed cell death* 10(5):1105-1110.
- Pitaval A, Tseng Q, Bornens M, They M. 2010. Cell shape and contractility regulate ciliogenesis in cell cycle-arrested cells. *J Cell Biol* 191(2):303-312.
- Polizio AH, Chinchilla P, Chen X, Kim S, Manning DR, Riobo NA. 2011. Heterotrimeric Gi proteins link Hedgehog signaling to activation of Rho small GTPases to promote fibroblast migration. *J Biol Chem* 286(22):19589-19596.
- Praetorius HA, Spring KR. 2001. Bending the MDCK cell primary cilium increases intracellular calcium. *J Membr Biol* 184(1):71-79.
- Qin H, Diener DR, Geimer S, Cole DG, Rosenbaum JL. 2004. Intraflagellar transport (IFT) cargo: IFT transports flagellar precursors to the tip and turnover products to the cell body. *J Cell Biol* 164(2):255-266.
- Reilova-Velez J, Seiler MW. 1984. Abnormal cilia in a breast carcinoma. An ultrastructural study. *Archives of pathology & laboratory medicine* 108(10):795-797.
- Renault MA, Roncalli J, Tongers J, Thorne T, Klyachko E, Misener S, Volpert OV, Mehta S, Burg A, Luedemann C, Qin G, Kishore R, Losordo DW. 2010. Sonic hedgehog induces angiogenesis via Rho kinase-dependent signaling in endothelial cells. *J Mol Cell Cardiol* 49(3):490-498.
- Rohatgi R, Milenkovic L, Scott MP. 2007. Patched1 regulates hedgehog signaling at the primary cilium. *Science* 317(5836):372-376.
- Ross AJ, May-Simera H, Eichers ER, Kai M, Hill J, Jagger DJ, Leitch CC, Chapple JP, Munro PM, Fisher S, Tan PL, Phillips HM, Leroux MR, Henderson DJ, Murdoch JN, Copp AJ, Eliot MM, Lupski JR, Kemp DT, Dollfus H, Tada M, Katsanis N, Forge A, Beales PL. 2005. Disruption of Bardet-Biedl syndrome ciliary proteins perturbs planar cell polarity in vertebrates. *Nat Genet* 37(10):1135-1140.
- Ruiz-Heiland G, Horn A, Zerr P, Hofstetter W, Baum W, Stock M, Distler JH, Nimmerjahn F, Schett G, Zwerina J. 2012. Blockade of the hedgehog pathway inhibits osteophyte formation in arthritis. *Ann Rheum Dis* 71(3):400-407.
- Sasaki N, Kurisu J, Kengaku M. 2010. Sonic hedgehog signaling regulates actin cytoskeleton via Tiam1-Rac1 cascade during spine formation. *Mol Cell Neurosci* 45(4):335-344.

- Satir P, Christensen ST. 2008. Structure and function of mammalian cilia. *Histochem Cell Biol* 129(6):687-693.
- Schneider L, Clement CA, Teilmann SC, Pazour GJ, Hoffmann EK, Satir P, Christensen ST. 2005. PDGFR $\alpha$  signaling is regulated through the primary cilium in fibroblasts. *Curr Biol* 15(20):1861-1866.
- Shuang F, Zhou Y, Hou SX, Zhu JL, Liu Y, Zhang CL, Tang JG. 2015. Indian Hedgehog signaling pathway members are associated with magnetic resonance imaging manifestations and pathological scores in lumbar facet joint osteoarthritis. *Sci Rep* 5:10290.
- Silverman MA, Leroux MR. 2009. Intraflagellar transport and the generation of dynamic, structurally and functionally diverse cilia. *Trends Cell Biol* 19(7):306-316.
- Tanuma Y, Ohata M. 1978. Transmission electron microscope observation of epithelial cells with single cilia in intrahepatic biliary ductules of bats. *Archivum histologicum Japonicum = Nihon soshikigaku kiroku* 41(4):367-376.
- Thompson CL, Chapple JP, Knight MM. 2014. Primary cilia disassembly down-regulates mechanosensitive hedgehog signalling: a feedback mechanism controlling ADAMTS-5 expression in chondrocytes. *Osteoarthritis Cartilage* 22(3):490-498.
- Thompson CL, Wiles A, Poole CA, Knight MM. 2016. Lithium chloride modulates chondrocyte primary cilia and inhibits Hedgehog signaling. *FASEB J* 30(2):716-726.
- Thyberg J, Moskalewski S. 1999. Role of microtubules in the organization of the Golgi complex. *Exp Cell Res* 246(2):263-279.
- Tinti L, Taylor AM, Santucci A, Wlodarski B, Wilson PJ, Jarvis JC, Fraser WD, Davidson JS, Ranganath LR, Gallagher JA. 2011. Development of an in vitro model to investigate joint ochronosis in alkaptonuria. *Rheumatology (Oxford)* 50(2):271-277.
- Tran PV, Haycraft CJ, Besschetnova TY, Turbe-Doan A, Stottmann RW, Herron BJ, Chesebro AL, Qiu H, Scherz PJ, Shah JV, Yoder BK, Beier DR. 2008. THM1 negatively modulates mouse sonic hedgehog signal transduction and affects retrograde intraflagellar transport in cilia. *Nat Genet* 40(4):403-410.
- Tucker RW, Pardee AB, Fujiwara K. 1979. Centriole ciliation is related to quiescence and DNA synthesis in 3T3 cells. *Cell* 17(3):527-535.
- Tummala P, Arnsdorf EJ, Jacobs CR. 2010. The Role of Primary Cilia in Mesenchymal Stem Cell Differentiation: A Pivotal Switch in Guiding Lineage Commitment. *Cell Mol Bioeng* 3(3):207-212.
- Valderrama F, Babia T, Ayala I, Kok JW, Renau-Piqueras J, Egea G. 1998. Actin microfilaments are essential for the cytological positioning and morphology of the Golgi complex. *Eur J Cell Biol* 76(1):9-17.
- Wang Z, Wann AK, Thompson CL, Hassen A, Wang W, Knight MM. 2016. IFT88 influences chondrocyte actin organization and biomechanics. *Osteoarthritis Cartilage* 24(3):544-554.

- Wann AK, Zuo N, Haycraft CJ, Jensen CG, Poole CA, McGlashan SR, Knight MM. 2012. Primary cilia mediate mechanotransduction through control of ATP-induced Ca<sup>2+</sup> signaling in compressed chondrocytes. *FASEB J* 26(4):1663-1671.
- Waters AM, Beales PL. 2011. Ciliopathies: an expanding disease spectrum. *Pediatr Nephrol* 26(7):1039-1056.
- Wei F, Zhou J, Wei X, Zhang J, Fleming BC, Terek R, Pei M, Chen Q, Liu T, Wei L. 2012. Activation of Indian hedgehog promotes chondrocyte hypertrophy and upregulation of MMP-13 in human osteoarthritic cartilage. *Osteoarthritis Cartilage* 20(7):755-763.
- Wen GY, Soifer D, Wisniewski HM. 1982. The doublet microtubules of rods of the rabbit retina. *Anatomy and embryology* 165(3):315-328.
- Wong SY, Seol AD, So PL, Ermilov AN, Bichakjian CK, Epstein EH, Jr., Dlugosz AA, Reiter JF. 2009. Primary cilia can both mediate and suppress Hedgehog pathway-dependent tumorigenesis. *Nat Med* 15(9):1055-1061.
- Wood CR, Rosenbaum JL. 2015. Ciliary ectosomes: transmissions from the cell's antenna. *Trends Cell Biol* 25(5):276-285.
- Wright MO, Nishida K, Bavington C, Godolphin JL, Dunne E, Walmsley S, Jobanputra P, Nuki G, Salter DM. 1997. Hyperpolarisation of cultured human chondrocytes following cyclical pressure-induced strain: evidence of a role for alpha 5 beta 1 integrin as a chondrocyte mechanoreceptor. *J Orthop Res* 15(5):742-747.
- Zhu SL, Huang JL, Peng WX, Wu DC, Luo MQ, Li QX, Li ZX, Feng XX, Liu F, Wang MX, Chen WQ, Olsen N, Zheng SG. 2015. Inhibition of smoothed decreases proliferation of synoviocytes in rheumatoid arthritis. *Cellular & molecular immunology*.

## Figure Legends

Fig. 1. Alkaptonuria (AKU) alters articular chondrocyte F-actin cytoskeleton, and primary cilia prevalence and length. A: Representative F-actin cytoskeletal organisation in Healthy, AKU and homogentisic acid treated (+HGA) chondrocytes, stained for F-actin (green) and cell nuclei (blue). Scale bar 20  $\mu\text{m}$ . B: Representative confocal maximum intensity z-projections of primary cilia from Healthy, AKU and +HGA chondrocytes immunofluorescently stained for acetylated  $\alpha$ -tubulin (Ac  $\alpha$ -Tub; red) and pericentrin (green). Scale bar 20  $\mu\text{m}$ , inset scale bar 2  $\mu\text{m}$ . C: Primary cilia prevalence in populations of Healthy, AKU and +HGA chondrocytes. Mean $\pm$ s.e.m,  $n\geq 23$  random fields of view, Mann-Whitney tests: \*\* $P<0.01$ , \*\*\* $P<0.001$ . D: Primary cilia length in Healthy, AKU and +HGA chondrocytes; boxes represent median and interquartile range, with whiskers extending to 1.5 $\times$  interquartile range or the max/min data points,  $n\geq 316$  cilia, General Linear Model with Fisher pairwise comparisons: \*\*\* $P<0.001$ .

**Fig. 2. ROCK modulation of cilia length is abrogated in alkaptonuria (AKU) and homogentisic acid treated (+HGA) chondrocytes.** A: Representative confocal maximum intensity z-projections of primary cilia from Healthy, AKU and +HGA chondrocytes, with (Y27) or without (Ctrl) addition of ROCK inhibitor Y27632, immunofluorescently stained for acetylated  $\alpha$ -tubulin (Ac  $\alpha$ -Tub; green). Scale bar 20  $\mu\text{m}$ , inset scale bar 5  $\mu\text{m}$ . B: Primary cilia prevalence in populations of Healthy, AKU and +HGA chondrocytes treated with Y27632 (Y27) normalised to control prevalence values;  $n\geq 120$  cells. ROCK inhibition did not significantly reduce cilia prevalence. C: Primary cilia length in Healthy, AKU and +HGA chondrocytes treated with or without ROCK inhibitor (Y27) normalised to control cilia length for each cell type. Mean $\pm$ s.e.m.,  $n\geq 106$  cilia, Two-sample T-test: \* $P<0.05$ , <sup>ns</sup> $P>0.05$ .


**Fig. 3. Cilia structure and Arl13b protein distribution are altered in Alkaptonuria (AKU) consistent with disrupted ciliary trafficking.** A: Representative confocal maximum intensity z-projections of primary cilium demonstrating typical structural alterations with normal, bulbous tipped and fragmented cilia. Chondrocytes were immunofluorescently stained for acetylated  $\alpha$ -tubulin (Ac  $\alpha$ -Tub; green) and pericentrin (red). Arrowhead: primary cilium tip. Scale bar 2  $\mu\text{m}$ . B-C: Proportions of Healthy, AKU and homogentisic acid treated (+HGA) chondrocytes presenting bulbous tipped (B) or fragmented (C) primary cilia. Fisher's exact test: \*\*\* $P<0.001$ . D: Representative super resolution structured illumination microscopy (SR-SIM) maximum intensity z-projections of representative Healthy, AKU and +HGA chondrocyte primary cilia stained for Ac  $\alpha$ -Tub (green) and ARL13B (red). Arrowhead: primary cilium tip. Scale bar 2  $\mu\text{m}$ . E: Pixel intensity profiles at base and tip of primary cilium as indicated by white lines across cilia in D. F: ARL13B integrated intensity at base and tip of Healthy, AKU and +HGA chondrocyte primary cilia. G-I: Diameter of ciliary components Ac  $\alpha$ -Tub (G), ARL13B (H) and the difference in ARL13B and Ac  $\alpha$ -Tub diameters (I) assessed as full width at half maximum on transverse profiles across the base and tip of primary cilia as in E. Mean $\pm$ s.e.m.,  $n\geq 11$ , Mann-Whitney tests: \* $P<0.05$ , \*\* $P<0.01$ ; # $P<0.05$ , ## $P<0.01$  and ### $P<0.001$  vs. Base.

**Fig. 4. Response to Indian Hedgehog ligand (Ihh) addition is abrogated in Alkaptonuria (AKU).** GLI1 (A) and PTCH1 (B) gene expression in Healthy, AKU and homogentisic acid treated (+HGA) chondrocytes relative to endogenous control gene GAPDH and normalised to Healthy Ctrl. Mean $\pm$ s.e.m.,  $n\geq 3$ , General Linear Model with Fisher pairwise comparisons: \* $P<0.05$ , <sup>ns</sup> $P>0.05$ , # $P<0.05$ , ## $P<0.01$  and ### $P<0.001$  vs. Healthy; \*\* $P<0.01$  vs. AKU.


Table 1. Primer sequences used in qPCR.

Gene	Sequence	
18S RNA	F-	CGGCTACCACATCCAAGGAA
	R-	AGCTGGAATTACCGCGGC
PTCH1	F-	GGGTGGCACAGTCAAGAACAG
	R-	TACCCCTTGAAGTGCTCGTACA
GLI1	F-	GCGTTGTAGAGAGGTAACCC
	R-	TGATGAAAGCTACGAGGGAG

Accepted Article




**Figure 1**



**Figure 2**





**Figure 3**




Figure 4