



# THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **Contribution of walking to school to individual and population moderate-vigorous intensity physical activity**

**Citation for published version:**

Martin, A, Boyle, J, Corlett, F, Kelly, P & Reilly, J 2016, 'Contribution of walking to school to individual and population moderate-vigorous intensity physical activity: Systematic review and meta-analysis' *Pediatric Exercise Science*, pp. 353 -363.

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

*Pediatric Exercise Science*

**Publisher Rights Statement:**

© HK, as accepted for publication

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



**Contribution of walking to school to individual and population moderate-vigorous intensity physical activity: systematic review and meta-analysis**

<sup>1</sup>Anne Martin, <sup>2</sup>Jessica Boyle, <sup>2</sup>Fenella Corlett, <sup>1</sup>Paul Kelly, <sup>2</sup>John J Reilly\*

<sup>1</sup>University of Edinburgh, Physical Activity for Health Research Centre, St Leonard's Land, Edinburgh, EH8 8AQ, Scotland; <sup>2</sup>University of Strathclyde, Physical Activity for Health Group, George Street, Glasgow G1 1XQ, Scotland.

**\*Author for correspondence:** John J Reilly; [john.j.reilly@strath.ac.uk](mailto:john.j.reilly@strath.ac.uk)

**Keywords:** active commuting; walking; physical activity; accelerometer; child; adolescent; youth; systematic review

**Running Head:** MVPA during walking to school

### Abstract

1 **Purpose.** This study estimated the contribution of walking to/from school to  
2 objectively measured daily moderate-vigorous intensity physical activity (MVPA) in  
3 individuals and populations. **Methods.** MEDLINE, PsycINFO and SPORTDiscus  
4 were systematically searched up to February 2015. Two reviewers independently  
5 screened titles/abstracts/full-text articles, and assessed study quality. **Results.** Of  
6 2430 records, 116 were eligible for full-text screening. Twelve studies met the  
7 inclusion criteria of reporting objectively obtained measures of MVPA (total and  
8 while walking to/from school) in children and adolescents. The weighted mean  
9 MVPA accumulated in walking to and from school was 17 minutes per day in primary  
10 school pupils (9 samples, n=3422) and 13 minutes per day in high school pupils (4  
11 samples, n=2600). Pooled analysis suggested that walking to and from school  
12 contributed 23% and 36% of MVPA on schooldays in primary school age children  
13 and high school pupils, respectively. All included studies were of high  
14 methodological quality. **Conclusions.** Walking to and from school makes a  
15 meaningful contribution to individual schoolday MVPA for active commuters in  
16 western countries. Since schooldays represent only around half of all days, and  
17 prevalence of walking to school is low in many countries, the contribution of walking  
18 to school to population MVPA is probably low.

19

20

21

## Introduction

22 Recent reviews have suggested that active commuting (walking, cycling) to school  
23 has a number of health and non-health benefits, including potentially reduced  
24 adiposity, and environmental gains arising from reduced car use (15, 26,40,42).

25 Active commuting to school is an important element of physical activity and health  
26 policy in many parts of the world. Active commuting to school is widely considered  
27 as an important contributor to the achievement of daily moderate-vigorous intensity  
28 physical activity (MVPA) recommendations (5,50), but there has been a marked  
29 decline in the prevalence of active commuting to school internationally (11,12,49).

30

31 A great deal of research and policy effort has focused on interventions to increase the  
32 prevalence of active commuting to school, with the implicit assumption that such  
33 interventions will produce a meaningful increase in population MVPA. Assessing the  
34 extent to which future research and policy should focus on active commuting to  
35 school depends on an improved –more quantitative– understanding of the contribution  
36 which it actually makes to MVPA. The MVPA accumulated during active commuting  
37 is usually seen by researchers and policymakers in terms of the individual child who  
38 is commuting actively, and with a focus on schooldays. A population perspective of  
39 the contribution of active commuting to school to MVPA would place less emphasis  
40 on individual active commuters and schooldays, by considering both the fact that not  
41 all days are schooldays (18), and the prevalence of active commuting, which is very  
42 low in many countries (49).

43

44 Recent reviews and original studies on the topic of active commuting to school have  
45 asked research questions about secular trends in, and prevalence of, active commuting

46 (11,12,31,41,42,49) ; the main determinants and correlates of active commuting  
47 (12,27,47); the efficacy of interventions to promote active commuting (29), and the  
48 health effects of active commuting (8,15,24,26, 34,40,41). These reviews have  
49 generally made the implicit or explicit assumption that active commuting to school  
50 makes a meaningful contribution to individual and/or population MVPA. To date, no  
51 systematic review has asked a research question about the amount of MVPA which is  
52 actually being accumulated by children and adolescents in active commuting to  
53 school, and the extent to which MVPA accumulated during the active commute  
54 contributes to population MVPA. The primary aims of the present study were  
55 therefore to systematically review and critically appraise the evidence on the amount  
56 of MVPA being accumulated while walking to/ from school, and to examine the  
57 contribution of MVPA while walking to school to overall MVPA on schooldays in  
58 those individuals. The secondary aim was to consider the contribution of walking to  
59 and from school to population MVPA, by allowing for days on which children and  
60 adolescents do not attend school, and the prevalence of active commuting in the  
61 population (18).

62

63

## Methods

### 64 Literature searching and study inclusion criteria

65 The literature search was conducted in February 2015 using the three most relevant  
66 electronic databases: MEDLINE; PsycINFO and SPORTDiscus. The search strategy  
67 was based on the following components: population (children and adolescents);  
68 exposure (active commuting to and from school via walking); and outcome  
69 (objectively measured MVPA). While we originally considered the inclusion of data  
70 from children and adolescents who cycled to/from school the focus of the present

71 review was walking to/from school, because in all eligible studies the prevalence of  
72 cycling to school was negligible and it was possible to extract data only from those  
73 who walked to/from school. In any case, accelerometry as used in most previous  
74 studies is unsuitable for measurement of MVPA during cycling (46). Studies  
75 published from 2004 were eligible for inclusion so that any evidence would be  
76 generalizable given recent rapid secular trends in active commuting (49). The search  
77 strategy in MEDLINE is given in **Table 1**, and was adapted as required for the other  
78 two databases. Full literature search details are available from the corresponding  
79 author on request. Reference lists of eligible studies were also examined for  
80 potentially eligible studies.

81

82 To be eligible for inclusion in the review, papers had to: report information on school-  
83 age children and adolescents (4 – 19 years of age); use objective methods for  
84 measuring MVPA (heart rate monitoring; accelerometry; combined heart rate  
85 monitoring-accelerometry; direct observation); report MVPA while walking to/from  
86 school, with any accelerometry cut-point or other objective method, be original  
87 research, published in a peer reviewed journal; be observational in design, though  
88 intervention studies were considered for inclusion if pre-intervention data and/or  
89 control group data were given separately; be published in the English language. There  
90 is no ideal (or even consensus) definition of active commuting to and from school,  
91 and so studies were not excluded on the basis of how they defined or operationalised  
92 active commuting. In most studies active commuting was operationalised as the time  
93 periods before and after school (typically the 1 hour before school and 1 hour after  
94 school), and so these will include some MVPA spent in domains other than active  
95 commuting. In all eligible studies the data extracted for the present review was

96 considered to represent the estimated MVPA accumulated during the walk to and  
97 from school (and the MVPA on schooldays) among those who regularly walked  
98 to/from school. There is also no certainty (or even consensus) over which  
99 accelerometer data reduction decisions are ideal for minimising biases in MVPA  
100 estimates (9,13,17,21,37), and so studies were not excluded on the basis of the data  
101 reduction decisions they made.

102

103 Two authors independently considered the titles/abstracts of all papers identified by  
104 the search, referring to a third author for discussion and mediation where required.

105 Two authors also examined the papers identified for full-text screening, and referred  
106 to a third author where necessary for discussion/mediation.

107

#### 108 **Data extraction**

109 Three authors examined every eligible study and used a standard data extraction form  
110 in order to populate the evidence tables. The extracted items were: first author,  
111 publication year, country, objective measurement type, cut point for MVPA, sample  
112 size, mean age, summary MVPA data (minutes/day) walking to and from school and  
113 daily MVPA during schooldays for those who walked to and from school. Eligible  
114 studies included only participants who walked to school regularly and/or provided  
115 data for such individuals-for the present study data were extracted only from children  
116 and adolescent study participants who regularly walked to/from school.

117

#### 118 **Data analysis and synthesis**

119 The eligible studies fell logically into two categories: studies of primary school pupils  
120 (elementary and middle school); studies of high school pupils, and so data were  
121 synthesised for these two age groups separately.

122

123 **Contribution of active commuting to *individual* MVPA for those who walk to**  
124 **school**

125 In some studies the MVPA content of commuting time was expressed as a percentage,  
126 and so absolute MVPA (minutes) was recalculated based on data on the percentage of  
127 time spent in MVPA and commuting time provided by each eligible study. For each  
128 individual study the proportional contribution of walking to and from school to total  
129 daily MVPA was calculated. An overall pooled estimate was calculated for primary  
130 school pupils and high school pupils by averaging the proportional contributions from  
131 each study. A weighting factor based on study sample size was used to weight  
132 proportional contributions in the pooled estimate.

133

134 **Contribution of walking to/from school to *population* MVPA, allowing for non-**  
135 **school days and prevalence of active commuting**

136

137 Since children and adolescents who walk to/from school can only do so on school  
138 days, and since not all children and adolescents walk to/from school, the contribution  
139 which walking to school makes to the overall population MVPA cannot be  
140 determined by considering active commuters and schooldays alone (18). In order to  
141 estimate the contribution of walking to/from school to *population* total MVPA, data  
142 on the proportion of days per year when children and adolescents attend school were



143 used, along with data on the population prevalence of walking to school. Data on the  
144 number of schooldays attended per year vary both within-nations and between-  
145 nations. For the economically developed nations from which eligible studies were  
146 found in the present review, around half of all days per year are school days (33). To  
147 estimate the contribution of walking to/from school MVPA to total population  
148 MVPA, the schoolday commuting data can therefore be reduced by around half for  
149 those in the population who walk to/from school (18).

150

151 The contribution which walking to and from school makes to population MVPA will  
152 also depend on the population prevalence of active commuting-for children who do  
153 not walk to/from school the contribution which this behavior makes is negligible. The  
154 impact of the prevalence of walking to/from school on population MVPA was  
155 illustrated with two examples, taken from nations with studies eligible in the review  
156 and of interest because of the contrast they provide in the prevalence of walking  
157 to/from school: Scotland, where current prevalence of regular walking to school is  
158 around 50% (36); the USA, where prevalence of walking to school in children is  
159 <15% (11).

160

### 161 **Assessment of quality of the eligible studies**

162 Studies identified as eligible were assessed independently for quality by three authors,  
163 resolving any disagreements by discussion. The Tooth et al (48) tool for assessing the  
164 quality of observational studies was considered initially-it consists of over 30 items,  
165 and some items of particular importance to the quality of accelerometry studies are  
166 not included. The Tooth et al tool has been used previously, with substantial  
167 modification, in recent systematic reviews of physical activity studies with an 11-item

168 (19), or 8-item (45) checklist. In the present study the Tooth et al tool (48) was  
169 modified for use as a 15-item checklist, scored out of 6, as shown in **Table 2**. Each  
170 eligible study therefore received a score out of 6, with higher scores reflecting higher  
171 study quality.

172

173

## Results

### 174 Study selection and characteristics of eligible studies

175 The study flow diagram is provided in **Figure 1**. Of 2430 records identified in the  
176 initial review of the three databases, 116 were identified for full text screening. Of  
177 these, 12 records were eligible for inclusion, reporting on 13 samples. Reasons for  
178 exclusion are reported in Figure 1.

179 All studies used the ActiGraph, though with a variety of different models as well as  
180 different approaches to data collection and reduction.

181 Nine samples involved primary school pupils, with a total sample size of 3422  
182 children, in Denmark (1 study), England (4 studies), Scotland (1 study), and USA (3  
183 studies). Study characteristics are summarized in **Table 3**. Four eligible samples  
184 involved high school pupils (**Table 4**), with a total sample size of 2600 adolescents, in  
185 three nations: Denmark (1 study); England (1 study); USA (2 studies).

186

### 187 Results on walking to and from school in primary school pupils

188 The mean daily MVPA accumulated during the walk to and from school in these  
189 studies ranged from a low of 4 minutes/day in one study to 24 minutes/day in another

190 (Table 3). The weighted mean MVPA across the nine studies was 17 minutes per  
191 school day.

192 Figure 2A displays the proportional contribution of walking to and from school to  
193 total daily MVPA for each study. For those children who walked regularly to/from  
194 school, pooled analysis showed that the commute represented 23% of daily MVPA on  
195 schooldays.

196

### 197 Results on walking to and from school in high school pupils

198 The mean daily MVPA accumulated while walking to and from school ranged from a  
199 low of 9 minutes/day in one study to a high of 18 minutes/day in another (Table 4).

200 Weighted mean MVPA in walking to and from school across the four studies was 13  
201 minutes per day.

202 Proportion of walking to and from school to total daily MVPA for individual studies  
203 is summarized in Figure 2B. For those adolescents who walked to and from school  
204 regularly, pooled analysis showed that the contribution of the commute represented  
205 36% of total daily MVPA on schooldays.

206

### 207 Study quality assessment

208 On quality assessment (Tables 2 and 3), all eligible studies scored at least 5/6.

209

210

211

## Discussion

212 Main findings and implications

213 The present study suggests that walking to and from school contributes about one  
214 quarter of *individual* total daily MVPA *on schooldays* for active commuters to  
215 primary school (contributing up to around a third of the recommended MVPA of 60  
216 minutes/day on schooldays), and around one-third of total *school day* MVPA for  
217 active commuters to high school (contributing up to around a fifth of the  
218 recommended MVPA of 60 minutes/day on schooldays) in western countries. These  
219 findings illustrate the importance of active commuting to MVPA, for those  
220 individuals who commute actively, on schooldays.

221

222 For an understanding of the importance of active commuting to *population* MVPA the  
223 number of school days actually attended per year matters (18), as does active  
224 commuting prevalence. As an example, the only eligible study from Scotland (29)  
225 reported that primary school age children who walked to and from school  
226 accumulated around 16 minutes MVPA per school day while doing so, the equivalent  
227 of around 8 minutes MVPA per day when averaged over a whole year for individuals  
228 who commute actively. Since the population prevalence of regular walking to school  
229 in Scottish primary school children is currently around 50% (36), this means that the 8  
230 minutes/day MVPA contribution to overall population MVPA (ie when those who do  
231 not walk to and from school are included) is reduced further.

232

233 In the USA, with a prevalence of walking to and from school of around 13% in 5-11  
234 year olds (11), and mean MVPA during walking to school of around 4-14 minutes per  
235 school day (Tables 3 and 4), the current contribution of walking to and from school to  
236 *population* MVPA will be very low. In accelerometry studies of nationally  
237 representative samples of US children, mean daily MVPA estimates vary from a low

238 of 75-95 minutes (3, 50), to a high of around 180 minutes (32). If these estimates are  
239 accurate, the present study suggests that nearly all population MVPA must be  
240 accumulated in domains other than active commuting to school (at home; in active  
241 and outdoor play; in school based physical activity-recess and physical education; in  
242 organized sport).

243

244 Walking to/from school may be associated with higher overall physical activity and  
245 may provide health and non-health benefits (15,40,43), but the present study suggests  
246 that it makes only a small contribution to *population* MVPA , probably a combination  
247 of low prevalence of active commuting to school, limited MVPA during the commute,  
248 short commuting distances (18, 22, 27,35,42), and the fact that so many days are not  
249 schooldays (12,18). If walking to school is going to make a much greater contribution  
250 to population MVPA in future, the prevalence, duration, and MVPA content of  
251 walking to school must all be increased substantially. A discussion of policy and  
252 strategy options and arguments for improving surveillance of active commuting to  
253 school, and for increasing the prevalence and MVPA content of active commuting to  
254 school, would go beyond the scope of the present study, but these issues are dealt with  
255 elsewhere (31,42,49, 51,52). Researchers and policymakers should also consider  
256 whether focusing on domains of physical activity other than active commuting to  
257 school might be more effective in the promotion of population MVPA (18).

258

259

## 260 **Comparisons with other studies**

261 Since previous systematic reviews on active commuting to school have asked research  
262 questions distinct from the present study, there are no directly comparable reviews.

263 Janssen (18) recently examined the relative public health gain in Canada, of targeting  
264 different physical activity domains (active commuting vs. physical education, active  
265 play, school recess, and organised sport). He concluded that successful promotion of  
266 active commuting to school might make only a relatively small contribution, in part  
267 because school days represent only around half of all days, and in part because walks  
268 to school were typically short.

269

#### 270 **Review and evidence strengths and weaknesses**

271 The present study represents a high-level of evidence. The study had an *a priori*  
272 protocol and followed PRISMA guidelines in conduct and reporting (30). The  
273 evidence considered by the present review had a number of strengths too. In  
274 particular, eligible studies were all rated as being of high or very high quality. The  
275 studies included were in some cases based in large, nationally representative, surveys  
276 or cohorts, a strength in terms of generalizability.

277

278 The present study also had a number of weaknesses. First, studies eligible for  
279 inclusion had to be published in peer reviewed journals in English language, and this  
280 may have excluded relevant evidence. Literature searching was restricted by starting  
281 the search for papers published from 2004: this may be seen as a weakness, but was  
282 intended as a strength, to focus the review on more recent, and more generalizable  
283 evidence given rapid secular declines in active commuting to school. The present  
284 study used a 15-item quality assessment measure, but collapsing this to 6 items for  
285 scoring purposes might have reduced the ability to discriminate between studies on  
286 the grounds of quality.

287

288 Various limitations probably led to overestimates of the estimated MVPA content of  
289 the walk to/from school. In most eligible studies the walk to and from school was  
290 operationalized as specified periods before and after school (typically in the hour  
291 before school and the hour after school), so MVPA accumulated will be greater than  
292 the MVPA during the walk *per se*, by including some MVPA in domains other than  
293 active commuting (e.g. play, sport). Walking to and from school might provide  
294 opportunities for active play which would not be available when commuting  
295 passively, though these opportunities may not always be realised (44). Removal of the  
296 accelerometer by study participants before the end of the day may have biased eligible  
297 studies towards an overestimate of the contribution of the commute to schoolday  
298 MVPA in some cases. The use of low accelerometer cutpoints may also have inflated  
299 the absolute amount of MVPA during the commute.

300

301 The present study did not consider light intensity physical activity during walking to  
302 and from school, but there is an emerging body of evidence that light intensity  
303 physical activity may have a number of health benefits for children and adolescents  
304 (4,14,23,25). It is unlikely that all walking to school is MVPA, and indeed several  
305 studies of the energy cost of walking in children and adults give mean values for  
306 walking of less than three times resting energy expenditure, and so categorise walking  
307 as a light intensity activity (1,2).

308

309 One major gap in the eligible evidence reviewed by the present study was the absence  
310 of data from low-middle income and non-western countries. The decision to restrict  
311 the search to studies in the English language may have contributed to this. Developing  
312 countries around the world are undergoing a 'physical activity transition' (20,31) and

313 recent international surveillance of active commuting to school (49) has suggested  
314 that the secular decline in active commuting to school seen in high-income countries  
315 may also be occurring in low-middle income countries. In many countries a minority  
316 of children will be commuting actively to school, and prevalence of active commuting  
317 will be declining (15,41,49). In a recent study of children in rural South Africa,  
318 walking long distances to school was the norm, but the speed of walking was low and  
319 so the MVPA accumulated during the walk to/from school was limited (10). It is not  
320 clear whether these findings from South Africa apply to other low and middle-income  
321 countries.

322

### 323 **Conclusions**

324 The present study suggests that walking to and from school may make a meaningful  
325 contribution to individual schoolday MVPA in western countries for those individuals  
326 who commute actively. If walking to school is going to make a more substantial  
327 contribution to *population* MVPA, then the prevalence will have to be increased  
328 markedly. The extent to which walking to school is contributing to individual or  
329 population MVPA among children and adolescents in low and middle-income  
330 countries is less clear.



## References Cited

1. Adolph AL, Puyau MR, Vohra FA, Nicklas TA, Zakeri IF, Butte NF. Validation of uniaxial and triaxial accelerometers for the assessment of physical activity in preschool children. *J Phys Act & Health* 2012; 9: 944-953.
2. Ainsworth BE, Haskell WL, Whitt MC, Irwin ML, Swartz AM, Strath SJ et al. Compendium of physical activities and update of activity codes and MET intensities. *Med Sci Sports Exerc* 2000; 32:s498-s504.
3. Belcher BR, Berrigan D, Dodd KW, Emken BA, Chan CP, Spruit-Metz D. Physical activity in US youth: effect of race/ethnicity, age, gender, and weight status. *Med Sci Sports Exerc* 2010; 42: 2211-2221.
4. Carson V, Ridgers, ND, Howard, BJ, Winkler EAH, Healy GN, Owen N, Salmon J. (2013). Light-intensity physical activity and cardiometabolic biomarkers in US adolescents. *PLoS ONE* 2013; 8(8).
5. Chillon P, Ortega FB, Ruiz JR, Veidebaum T, Oja L, Maestu J, Sjostrom, M. Active commuting to school in children and adolescents: An opportunity to increase physical activity and fitness. *Scand J Publ Health* 2010; 38, 873–879.
6. Cooper AR, Jago R., Southward E, Page AS. Active travel and physical activity across the school transition. *Med Sci Sports Exerc* 2012; 44: 1890–1897.

7. Cooper AR, Page AS, Wheeler BW., Griew P, Davis L , Hillsdon M, Jago R. Mapping the walk to school using accelerometry combined with a global positioning system. *Am J Prev Med* 2010; 38: 178–183.
8. Cooper AR, Andersen LB, Wedderkopp N., Page AS, Froberg K. Physical activity levels of children who walk, cycle, or are driven to school. *Am J Prev Med.* 2005; 29: 179–184.
9. Corder, K., Brage, S., Ramachandran, A., Snehalatha, C., Wareham, N. & Ekelund, U. Comparison of two Actigraph models for assessing free-living physical activity in Indian adolescents, *J Sports Sci* 2007; 25: 1607-1611.
10. Craig EM, Bland RM, Reilly JJ. Objectively measured physical activity levels of children and adolescents in rural South Africa: high volume of physical activity at low intensity. *Applied Physiol Nutr Metab* 2013; 38:81-4.
11. Dentre KN, Beals K, Crouter SE, Eisenmann JC, Mackenzie TL, Pate RR, Saelens BE, Sisson SB, Spruit-Metz D, Sothorn M, Katzmarzyk PT. Results for the US 2014 Report Card on Physical Activity for Children and Youth. *J Phys Act & Health* 2014 11 (suppl 1): s105-112.
12. Dessing D, de Vries SI, Graham J, Pierik FH. Active transport between home and school assessed with GPS: cross-sectional study amongst Dutch elementary school children. *BMC Public Health* 2014; 14: 227.

13. Dorsey K, Herrin J, Krumholz ZH, Irwin M. The utility of shorter epochs in direct motion monitoring. *Res Q Exerc Sport* 2009; 80: 460-468.
14. Dowd KP, Harrington DM, Hannigan A, Donnelly AE. Light intensity physical activity is associated with adiposity in adolescent females. *Med Sci Sports Exerc* 2014; 46: 2295-2300.
15. Faulkner GE, Buliung RN, Flora PK, Fusco C. Active school transport, physical activity levels and body weight of children and youth: a systematic review. *Prev Med* 2009; 48:3-8.
16. Freedson PS, Sirard J, Debold EP. Calibration of the CSA Inc accelerometer. *Med Sci Sports Exerc* 1997; 29: 45.
17. Hislop J, Bulley C, Mercer T, Reilly JJ. Comparison of accelerometer cutpoints for physical activity and sedentary behavior in pre-school children: a validation study. *Pediatr Exerc Sci* 2012; 24: 563-576.
18. Janssen I. Active play: an important physical activity strategy in the fight against childhood obesity. *Can J Publ Health* 2014; 105: e22-27.
19. Jones RA, Hinkley T, Okely AD, Salmon J. Tracking physical activity and sedentary behavior in childhood: a systematic review. *Am J Prev Med* 2013; 44:651-658.

20. Katzmarzyk PT. Increasing global research capacity in physical activity for children and youth. *Res Exerc Epidemiol* 2014; 16: 71-75.
21. Kim Y, Beets MW, Pate RR, Blair SN. The effect of reintegrating Actigraph accelerometer cutpoints in pre-school children: comparison using different epoch lengths. *J Sci Med Sport* 2013; 16: 129-134.
22. Klinker CD, Schipperijn J, Christian H, Kerr J, Ersbøll AK, Troelsen, J. Using accelerometers and global positioning system devices to assess gender and age differences in children's school, transport, leisure and home based physical activity. *Int J Behav Nutr Phys Act.* 2014; 11:8.
23. Kwon S, Janz KF, Burns TL & Levy SM . Association between light-intensity physical activity and adiposity in childhood. *Pediatr Exerc Sci*, 2011; 23:218-229.
24. Larouche R., Faulkner GEJ, Fortier M., Tremblay M. Active transportation and adolescents' health. *Am J Prev Med* 2014; 46: 507–515.
25. Leary SD, Ness AR, Smith GD, Mattocks C, Deere K, Blair SN, Riddoch C. Physical activity and blood pressure in childhood. *Hypertension* 2008; 31: 92-98.

26. Lee MC, Orenstein MR, Richardson MJ. Systematic review of active commuting to school and childrens physical activity and weight. *J Phys Act Health* 2008; 5:930-49.
27. Lee C, Li L. Demographic, physical activity, and route characteristics related to school transportation: an exploratory study. *Am J Health Promot* 2014; 28: 77–88.
28. Mendoza JA, Watson K, Nguyen N, Cerin E, Baranowski T, Nicklas TA. Active commuting to school and association with physical activity and adiposity among US youth. *J Phys Act Health*. 2011; 8: 488-495.
29. McMinn D, Rowe D. A, Murtagh S, Nelson NM. The effect of a school-based active commuting intervention on children’s commuting physical activity and daily physical activity. *Prev Med* 2012; 54: 316–318.
30. Moher D, Liberati A, Tetzlaff J, Altman DG, the PRISMA Group (2009). Preferred reporting items for systematic reviews and meta-analyses: the PRISMA Statement. *Br Med J* 2009; 339:b2535.
31. Muthuri SK, Wachira LJM, LeBlanc AG, Francis CE, Sampson M, Onywera VO, Tremblay MS. Temporal trends and correlates of physical activity, sedentary behaviour, and physical fitness among school-aged children and youth in sub-Saharan Africa: a systematic review. *Int J Env Res Pub Health* 2014; 11: 3327-3359.

32. Nader PR, Bradley RG, Houts RM, McRitchie SL, O'Brien M. Moderate-vigorous physical activity from ages 9-15 years. *JAMA* 2008; 300: 295-305.
33. OECD Education at a glance 2013. OECD Indicators, OECD Publishing 2013.
34. Owen CG, Nightingale CM, Rudnicka AR, van Sluijs EMF. Travel to school and physical activity levels in 9–10 year-old UK children of different ethnic origin. *PLoS ONE*. 2012: e30932.
35. Panter J, Jones A, Van Sluijs EMF, Griffin S. Associations between active commuting and physical activity. *Pediatr Exerc Sci* 2011; 23, 72-86.
36. Reilly JJ, Dick S, McNeill G, Tremblay MS. Results from the Scottish Report Card on Physical Activity for Children and Youth. *J Phys Act & Health* 2014; 11 (suppl 1), s93-s97.
37. Reilly JJ, V Penpraze, J Hislop, G Davies, S Grant, JY Paton. Objective measurement of physical activity and sedentary behaviour: review with new data *Arch Dis Child* 2008; 93:614-619.
38. Saksvig BI, Catellier DJ, Pfeiffer K, Schmitz KH, Conway T, Going S, Ward D, Strikmiller P. & Treuth MS. Travel by walking before and after school and physical activity among adolescent girls. *Arch Pediatr Adolesc Med* 2007; 161: 153-158.

39. Saksvig BI, Webber LS, Elder JP, Ward D, Evenson KR, Dowda M, Chae SE, Treuth, MS. A cross-sectional and longitudinal study of travel by walking before and after school among eighth-grade girls. *J Adolesc Health*. 2012; 51: 608–614.
40. Schoeppe S, Duncan MJ, Badland H, Oliver M, Curtis C. Associations of children's independent mobility and active travel with physical activity, sedentary behaviour and weight status: a systematic review. *J Sci Med Sport* 2013; 16:312-9.
41. Sirard JR, Riner WF, McIver KL, Pate RR. Physical activity and active commuting to elementary school, *Med Sci Sports Exerc*. 2005; 37: 2062–2069.
42. Sirard J, Slater ME. Walking and bicycling to school: a review. *Am J Lifestyle Med* 2008; 2:372-96.
43. Stanley RM, Maher C, Dollman J. Modelling the contribution of walking between home and school to daily physical activity in primary age children. *BMC Publ Health* 2015; 15: 445.
44. Southward EF, Page AS, Wheeler BW, Cooper AR. Contribution of the school journey to daily physical activity in children aged 11-12 years. *Am J Prev Med* 2012; 43: 201-204.

45. Tanaka C, Reilly JJ, Huang W. Longitudinal changes in objectively measured sedentary behaviour and their relationship with adiposity in children and adolescents: systematic review and evidence appraisal. *Obes Rev* 2014; 15: 791–803.
46. Tarp J, Anderson LB, Østergaard L. Quantification of Underestimation of Physical Activity During Cycling to School When Using Accelerometry. *J Phys Act Health*, 2015, 12, 701 -707.
47. Timperio A, Ball K, Salmon J, Roberts R, Giles-Corti B., Simmons D, Baur LA, Crawford D. Personal, family, social, and environmental correlates of active commuting to school. *Am J Prev Med* 2006; 30:45-51.
48. Tooth I, Ware R., Bain C., Purdie D M, Dobson, A. Quality of reporting of observational longitudinal research. *Am J Epidemiol* 2005; 161:240-248.
49. Tremblay MS, Gray CE, Akinroye K, Harrington DM, Katzmarzyk PT, Lambert EV et al .Physical activity of children: a global matrix of grades comparing 15 countries. *J Phys Act & Health* 2014; 11 (suppl 1): s113-s125.
50. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the US measured by accelerometer. *Med Sci Sports Exerc* 2008; 40: 181-188



51. Tudor-Locke C, Ainsworth BE, Popkin BM, 2001. Active commuting to school: an overlooked source of childrens' physical activity ? *Sport Med* 2001; 31:309-13.
52. Voss E, Waters M, Fraser A, McKay H. School travel by public transport: rethinking active transport. *Prev Med Rep* 2015; 2:65-70.

## Table 1 Search Strategy in MEDLINE

---

exp child/

exp adolescent/

child\*.tw.

adolesc\*.tw.

(boy\* or girl\*).tw.

teen\*.tw.

youth\*.tw.

(pupil\* or student\* or schoolchild\*).tw.

(young adj2 (person\* or people)).tw.

p?diatr\*.tw.

school\*.tw.

1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11

exp Walking/ or exp Bicycling/

\*Travel/

(active adj2 (commut\* or transport\* or travel\* or lifestyle\* or life-style\* or living)).tw.

(walk\* or cycl\*).tw.

13 or 14 or 15 or 16

exp Motor Activity/

exp Exercise/

\*physical endurance/ or exp physical fitness/

\*Sports/

21 or 18 or 19 or 20

(physical\* adj2 activ\*).tw.

exercis\*.tw.

"physical fitness".tw.

"physical endurance".tw.

(physical activity adj2 (level\* or intensit\* or energy expenditure)).tw.

"MVPA".tw.

moderate-to-vigorous.tw.

"moderate to vigorous".tw.

22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30

objectiv\* measur\*.tw.

exp actigraphy/ or \*monitoring, ambulatory/

exp Accelerometry/

32 or 33 or 34

("GPS" or global positioning system or "GIS" or global information system).tw.

acceleromet\*.tw.

(activpal or activgraph or activity monitor\*).tw.

heart rate monitor\*.tw.

35 or 36 or 37 or 38 or 39

12 and 17 and 31 and 40

limit 41 to english language

limit 42 to yr="2004 -Current"

---

**TABLE 2 Study Quality Assessment Criteria**

<b><u>Criterion</u></b>	<b><u>Definition</u></b>	<b><u>Mark Allocation</u></b>
Sample recruitment	Sample: How were they recruited e.g. poster Time: When was the study conducted Place: Where did the recruitment take place	1 point for listing 3 criteria
Sample description (n, age, gender)	Number of participants recruited Mean age of participants % Gender male and female	1 point for listing all 3 criteria
Attrition	Number of participants recruited and the number actually measured	1 point for listing both criteria
Data collection and reduction	Type of device; epoch; no of days of active commuting specified as minimum; duration of monitoring time; monitor placement; data reduction decisions	1 point for listing 3 criteria
MVPA definition given	MVPA defined and accelerometry cut-off or other method given	1 point for listing both criteria
Results	Adequate description of numbers actually analysed, with summary MVPA data	1 point for listing both criteria

**TABLE 3 Contribution of Walking to and from School to Daily MVPA in Primary School Studies**

<b>Study, Year, Setting</b>	<b>Accelerometer, MVPA cut-off Point</b>	<b>Sample Size; Mean age (SD)</b>	<b>Total mean schoolday MVPA [minutes/day]</b>	<b>Mean MVPA Walking to/from school [minutes/day]</b>	<b>Quality Rating</b>
McMinn et al 2012, Scotland (29)	ActiGraph GT1M, Freedson cutpoint <sup>d</sup>	166; 8.6y (0.5)	90 (SD 27)	16 (SD 8)	5
Owen et al 2012, England (34)	ActiGraph GT1M, $\geq 2000$ cpm	1393; 9.9y (0.4)	74 (95% CI 71-76)	22 (95% CI 21-23)	6
Panter et al 2011, England (35)	ActiGraph GT1M, $\geq 2000$ cpm	723; 10.2y (0.3)	74 (SD 23)	15 (SD 7)	6
Lee and Li 2014, USA (27)	ActiGraph GT1M and GT3X, Freedson cutpoint <sup>d</sup>	109; 9.5y(not given)	63 (SD 11)	7 (SD 10)	5
Cooper et al 2005, Denmark (8)	ActiGraph 7164, cut-point unclear	328; 11y (0.4)	193 (SD 59)	7 (SD not given)	5
Cooper et al 2012 <sup>a</sup> , England (6)	ActiGraph GT1M, $\geq 2295$ cpm	500; 11y (0.4)	62 (SD 22)	14 (SD not given)	6
Cooper et al 2010, England (7)	ActiGraph GT1M, $\geq 3200$ cpm	70; 11y (0.3)	43 (SD 18)	11 <sup>b</sup> (SD 5)	6

Sirard et al 2005, USA (41)	ActiGraph, model not given, $\geq 1017$ cmp	21; 10y(0.6)	102 (SD not given)	24 (SD not given)	6
Saksvig et al 2007 <sup>c</sup> , USA (38)	ActiGraph 7164, $\geq 1500$ counts per 30 seconds	112; 6 <sup>th</sup> grade	29 (SD 2)	11 (SD 1)	6

<sup>a</sup> Primary school age sample from Cooper et al 2012 (6). <sup>b</sup>Paper provided MVPA during route to school only, so has been doubled. <sup>c</sup> Study of girls only. <sup>d</sup>

Freedson MVPA cutpoint (16) equivalent to 906cpm and 1018cpm for 9 and 10y olds respectively, using the following equation

$$\text{METS} = 2.757 + (0.0015 \times \text{counts/min}) - (0.08957 \times \text{age (yr)}) - (0.000038 \times \text{counts/min} \times \text{age (yr)})$$

In all cases daily MVPA data refer to schooldays only among children who walked to school regularly. MVPA: moderate-to-vigorous physical activity, cpm: counts per minute



**TABLE 4 Contribution of Walking to and From School to Daily MVPA in High School Studies**


<b>Study, Year, Setting</b>	<b>Method and Cut Point</b>	<b>Sample Size; Mean (SD) age</b>	<b>Mean total schoolday MVPA [minutes/day]</b>	<b>Mean MVPA Walking to/from school [minutes/day]</b>	<b>Quality Rating</b>
Mendoza et al 2011, USA (28)	ActiGraph 7164, Freedson cutpoint <sup>a</sup>	789; 14.4y (SE 0.1)	30 (SE 2)	9 (SE 1)	6
Klinker et al 2014, Denmark (22)	ActiGraph GT3X, $\geq 2296$ cpm	367; 13.2y (0.2)	Median 64 (IQR 42-97)	Median 10 (IQR 5-16)	5
Cooper et al 2012 <sup>b</sup> , England (6)	ActiGraph GT1M, $> 2295$ cpm	500; 12y (0.4)	63 (SD 23)	18 (SD not given)	6
Saksvig et al 2012, USA (39)	ActiGraph 7164, $\geq 3000$ cpm	944; 14y(SD not given)	26 (SD 2)	14 (SD 1)	6



<sup>a</sup>Freedson MVPA cut-point (16) equivalent to 1546cpm in 14y olds , using the following equation

$$\text{METs} = 2.757 + (0.0015 \times \text{counts/min}) - (0.08957 \times \text{age (yr)}) - (0.000038 \times \text{counts/min} \times \text{age (yr)})$$

<sup>b</sup>Secondary school data from Cooper et al 2012 (6).

Daily MVPA estimates are schoolday MVPA in study participants who walked to school regularly. MVPA: moderate-to-vigorous physical activity, cpm: counts per minutes



**A****B**

**Figure 2** Proportional contribution (mean, SD) of daily walking to and from school to total moderate-to-vigorous physical activity (MVPA) on schooldays. A: Primary school pupils, B: High school pupils.