



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Cloning of the African indigenous cattle breed Kenyan Boran

Citation for published version:

Yu, M, Muteti, C, Ogugo, M, Ritchie, WA, Raper, J & Kemp, S 2016, 'Cloning of the African indigenous cattle breed Kenyan Boran' *Animal Genetics*, vol. 47, no. 4, pp. 510-511. DOI: 10.1111/age.12441

Digital Object Identifier (DOI):

[10.1111/age.12441](https://doi.org/10.1111/age.12441)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Animal Genetics

Publisher Rights Statement:

© 2016 The Authors. *Animal Genetics* published by John Wiley & Sons Ltd on behalf of Stichting International Foundation for Animal Genetics.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Cloning of the African indigenous cattle breed Kenyan Boran

Mingyan Yu*, Charity Muteti*, Moses Ogugo*, William A Ritchie†, Jayne Raper‡ and Stephen Kemp*

*International Livestock Research Institute; The Joint Centre for Tropical Livestock Genetics and Health of the University of Edinburgh, International Livestock Research Institute and Scotland's Rural College, Nairobi, P.O. Box 30709-00100, Kenya; †Roslin Embryology, (21) EH331QB, Scotland, UK; ‡Department of Biological Sciences, Hunter College, City University of New York, New York, NY 10065, USA

Accepted for publication 11 March 2016

Description: Kenyan Boran, an indigenous East African zebu (*Bos indicus*) breed, is kept mostly for beef production in semiarid areas of Kenya.¹ The breed is well adapted to high ambient temperature, poor quality feed and high disease challenges compared to European exotic *Bos taurus* breeds.² However, they are susceptible to African tsetse fly-transmitted trypanosomiasis (ATT) caused by parasites (*Trypanosoma* spp.), which are also the cause of human African trypanosomiasis or sleeping sickness. The ATT is a major constraint to livestock production in sub-Saharan Africa.³

It is known that serum from baboons kills both animal- and human-infective African trypanosomes through serum trypanosome lytic factors (TLFs).⁴ The generation of trypanosomiasis-resistant transgenic cattle carrying baboon-derived TLFs may have the potential to improve livestock productivity in Kenya and Africa.⁴ As a precursor to such a study, we cloned a Kenyan Boran bull by somatic cell nuclear transfer (SCNT) using primary embryonic fibroblasts. This successful cloning represents an important first step towards the establishment of genetically modified Kenyan Boran through SCNT with genome-modified fibroblasts.

Methods: Boran embryonic fibroblasts were isolated from a 3-month-old male Kenyan Boran foetus and cultured with DMEM+10% FBS. Karyotyping was performed to confirm that the cell line had no major chromosome anomaly.

Nuclei from an early passage (P6) of the fibroblasts were used for SCNT into oocytes aspirated from the ovaries of *B. indicus* cattle after slaughter. SCNT was performed as described,⁵ and the reconstructed blastocysts were transferred to suitable surrogate mothers. Around 273 days post-embryo transfer, a cloned calf was born by caesarean section operation after dexamethasone induction. Ten microsatellite markers (Table S1) were used to verify the parentage of the cloned calf and its offspring.

Results: In total, three cloned embryos (60%) aborted by the second trimester, and two cloned calves (40%) were born (Table S2). Both of the born calves showed a low blood glucose level (<30 mg/dl) and difficulty in standing alone with tendon laxity at birth. The first cloned calf died apparently due to low blood glucose. The second cloned calf was supplemented with 2.5 g glucose through the jugular vein within 8 h of birth followed by abdominal injection of glucose solution (5%) when the glucose level fell below 50 mg/dl.⁶ The blood glucose level and the tendon laxity normalised after 1 week, and the calf remained healthy afterwards. Two calves were born after introducing the cloned bull to a female Boran herd (Fig. 1). Genotyping confirmed that both of the calves were offspring of the cloned bull.

Comments: With the boom of genome editing tools, for example transcription activator-like effector nucleases and clustered regularly interspaced short palindromic repeats/Cas9, there are unprecedented opportunities for improving livestock genetics efficiently through the introduction of superior traits between breeds by precise genome modification. The successful cloning of a Kenyan Boran bull has opened the possibility of making genetically modified Kenyan Boran with foreign genes or desired traits through genome editing at the fibroblast level followed by SCNT.

Conflict of interest: None of the authors have any conflict of interest to declare.

References

- 1 Rischkowsky B. & Pilling D.(Eds). (2007) *The State of the World's Animal Genetic Resources for Food and Agriculture*. FAO, Rome.



Figure 1 Cloned Boran bull and its offspring.

2 Brief Note

- 2 Rewe T.O. *et al.* (2006) *Anim Sci J* **77**, 163–77.
- 3 Williams D.J.L. *et al.* (1991) *Parasite Immunology* **13**, 171–85.
- 4 Thomson R. *et al.* (2009) *PNAS* **106**, 19509–14.
- 5 Ross P.J. & Cibelli B. (2010) *Methods in Molecular Biology*, **636**, 155–77.
- 6 Meirelles F.V. *et al.* (2010) *Reprod Fertil and Dev* **22**, 88–97.

Correspondence: M. Yu (m.yu@cgiar.org)

Supporting information

Additional Supporting Information may be found online in the supporting information tab for this article:

Table S1 Microsatellite markers used for parentage identification of the cloned calf and its offspring

Table S2 Summary of somatic cell nuclear transfer with Boran embryonic fibroblasts
