



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Pneumonia hospitalisations in Scotland following the introduction of pneumococcal conjugate vaccination in young children

Citation for published version:

Nair, H, Watts, AT, Williams, L, Omer, SB, Simpson, C, Willocks, L, Cameron, JC & Campbell, H 2016, 'Pneumonia hospitalisations in Scotland following the introduction of pneumococcal conjugate vaccination in young children' BMC Infectious Diseases. DOI: 10.1186/s12879-016-1693-x

Digital Object Identifier (DOI):

[10.1186/s12879-016-1693-x](https://doi.org/10.1186/s12879-016-1693-x)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Published In:

BMC Infectious Diseases

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



BMC Infectious Diseases

Pneumonia hospitalisations in Scotland following the introduction of pneumococcal conjugate vaccination in young children

--Manuscript Draft--

Manuscript Number:	INFD-D-15-00955R2
Full Title:	Pneumonia hospitalisations in Scotland following the introduction of pneumococcal conjugate vaccination in young children
Article Type:	Research article
Section/Category:	Vaccines
Funding Information:	
Abstract:	<p>Background Scotland introduced PCV7 and PCV13 immunisation in young children in 2006 and 2010 respectively. One recent study from the United States reported a decrease in hospitalisation rates for all-cause pneumonia most notably in adults older than 75 years of age following PCV7 introduction in the US child population. We aimed to examine the effect of PCV7 and PCV13 on hospitalisation rates for all-cause pneumonia across all age groups in Scotland.</p> <p>Methods We linked hospital records and death certification datasets for the entire Scottish population for the period 2000 to 2012. We included all cases where the primary / secondary diagnosis was pneumonia. Differences in hospital admission rates for pneumonia by age group were calculated using the difference in average annual rates for each period.</p> <p>Results We estimated that all-cause pneumonia hospitalisation rates in children <2 years decreased by about 30% in the post-PCV-13 period compared with the pre-PCV period. However, in adults aged 75-84 years and ≥85 years, all-cause pneumonia hospitalisation rates increased by 63% and 46 % respectively in the post-PCV 13 period compared to the pre-PCV period. This resulted in an additional 7000 hospitalisations across all age groups in Scotland in 2012 about half of which were in adults >75 years. At the same time, the median length of hospital stay decreased by a third in children <2 years and by about 20% in adults >75 years in the post-PCV13 period compared to the pre-PCV period. Additionally, there was an 11% reduction in deaths due to all-cause pneumonia, and 30% reduction in pneumococcal hospitalisations across all age groups in the post-PCV13 period compared with pre-PCV period.</p> <p>Interpretation The modest and sustained decline in the rates of hospitalisation for all-cause pneumonia in children and the reduction in proportion of pneumonia hospitalisations in children coded as pneumococcal disease in the post-PCV period should alleviate concerns that pneumococcal serotype replacement may have resulted in an increased pneumonia burden in this age group. The indirect impact of child PCV immunisation in those not vaccinated (in terms of reduction in all-cause pneumonia hospitalisations in the elderly) has not been seen in Scotland. Our results are likely to be confounded by changes in clinical coding and healthcare practices over the same period and illustrate that health care planners cannot, with confidence, predict indirect PCV vaccine impacts on hospitalisations. IPD surveillance across all age groups is needed to assess the indirect effects of PCV in the community.</p>
Corresponding Author:	Harish Nair, MBBS, PhD University of Edinburgh UNITED KINGDOM
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	University of Edinburgh
Corresponding Author's Secondary Institution:	

First Author:	Harish Nair, MBBS, PhD
First Author Secondary Information:	
Order of Authors:	Harish Nair, MBBS, PhD
	Arun Thor Watts, BMedSci
	Linda J Williams, PhD
	Saad B Omer, PhD
	Colin R Simpson, PhD
	Lorna J Willocks, MD
	J Claire Cameron, FFPH
	Harry Campbell, MD
Order of Authors Secondary Information:	
Response to Reviewers:	<p>Dear Editor, Thank you for considering our manuscript for publication and offering us an opportunity to revise the manuscript based on the comments from peer-reviewers and the editorial comments. We have now revised the manuscript and provide herewith the version in track changes as well as a clean version. We provide our point by point response to reviewer's comments below:</p> <p>Reviewer 1</p> <p>1.Abstract- Final sentence – might want to revise according to revision in last paragraph of discussion. Response- We thank the reviewer for their comments. We have now made appropriate changes in the manuscript to address their concerns. Changes in manuscript – Page 3, line 33-35</p> <p>2.Pg 4 Introduction (not sure which line numbers are correct!) – try line 20 or 28 – delete in 2+1 dose as has been added a few lines further down. Response- We thank the reviewer for their comments. We have now made appropriate changes in the manuscript to address their concerns. Changes in manuscript – Page 3, line 20</p> <p>3.Statistical analysis page 5 – this revision is not clear. "Differences in hospital admission rates by age group were calculated using the difference in average annual rates for each period., and the expected difference calculated to illustrate where there were significant differences from expected rates." If I understand rightly, perhaps put something like: 'average annual rates for each period, and observed vs expected were compared' Response- We thank the reviewer for their comments. We have now made appropriate changes in the manuscript to address their concerns. Changes in manuscript – Page 5, line 35-37</p> <p>4.Pg 10 line 27 - correct close bracket. The failure to note any positive indirect effect of infant PCV vaccination on (pneumococcal) pneumonia in non-vaccinated individuals observed through measurement..... Response- We thank the reviewer for their comments. We have now made appropriate changes in the manuscript to address their concerns. Changes in manuscript – Page 10, line 18</p> <p>5.Figure 1 – y axis label has been changed to Mean per 100K (mean of what?). The label however has not been changed in Figure 2. Response- We thank the reviewer for their comments. We have now made appropriate changes in the manuscript to address their concerns. Changes in manuscript – Figure 1 and Figure 2</p> <p>6.Number 3 – response – might be worth commenting to this effect in the manuscript. Response- We thank the reviewer for their comments. We however feel this is not really relevant here. Changes in manuscript – No changes required</p>

7.Number 4 –“ The percentage of cases with pneumonia as the secondary cause of hospitalisation has remained almost static at 2.5% in the 2000-2005 period, 2.4% in the 2007-2009 period and 2.5% in the 2010-2012 period.” Also worth mentioning in discussion perhaps.

Response- We thank the reviewer for their comments. We have now made appropriate changes in the manuscript to address their concerns.

Changes in manuscript – Page 8, line 42-44

Reviewer 2

8.Page 4 line 20: please replace "in 2+1 dose" with "using a 2+1 dose schedule".

Response- We thank the reviewer for their comments. This directly contradicts the advice by reviewer 1 (comment #2). We can only implement ONE of these changes. We have gone with the advice by reviewer 1, as it makes more sense to talk about PCV7 and PCV13 dosing schedules together. .

Changes in manuscript – Page 3, line 20

9.remove the newly introduced changes about 2+1 schedule

Response- As above

Changes in manuscript – Page 3, line 20

Editorial revisions requested

10.Conclusions- Could you please add a section "Conclusions" after section "Discussion". This should state clearly the main conclusions of the research article and give a clear explanation of their importance and relevance. You can use the last paragraph(s) of your Discussion if relevant.

Response- We have now made appropriate changes in the manuscript to meet your requirements

11.Declarations- Please ensure that your manuscript contains a Declarations section, with all of the following subsections:

Response- We have now made appropriate changes in the manuscript to meet your requirements

12.Please ensure all Figure legends are placed at the end of the text (after the References). Figures should just be the images themselves.

Response- We have now made appropriate changes in the manuscript to meet your requirements

Best wishes,
Dr Harish Nair

Dear Editor,

Thank you for considering our manuscript for publication and offering us an opportunity to revise the manuscript based on the comments from peer-reviewers and the editorial comments. We have now revised the manuscript and provide herewith the version in track changes as well as a clean version.

We provide our point by point response to reviewer's comments below:

Reviewer 1

1. Abstract- Final sentence – might want to revise according to revision in last paragraph of discussion.
Response- We thank the reviewer for their comments. We have now made appropriate changes in the manuscript to address their concerns.
Changes in manuscript – Page 3, line 33-35
2. Pg 4 Introduction (not sure which line numbers are correct!) – try line 20 or 28 – delete in 2+1 dose as has been added a few lines further down.
Response- We thank the reviewer for their comments. We have now made appropriate changes in the manuscript to address their concerns.
Changes in manuscript – Page 3, line 20
3. Statistical analysis page 5 – this revision is not clear. “Differences in hospital admission rates by age group were calculated using the difference in average annual rates for each period., and the expected difference calculated to illustrate where there were significant differences from expected rates.” If I understand rightly, perhaps put something like: ‘average annual rates for each period, and observed vs expected were compared’
Response- We thank the reviewer for their comments. We have now made appropriate changes in the manuscript to address their concerns.
Changes in manuscript – Page 5, line 35-37
4. Pg 10 line 27 - correct close bracket. The failure to note any positive indirect effect of infant PCV vaccination on (pneumococcal) pneumonia in non-vaccinated individuals observed through measurement.....
Response- We thank the reviewer for their comments. We have now made appropriate changes in the manuscript to address their concerns.
Changes in manuscript – Page 10, line 18
5. Figure 1 – y axis label has been changed to Mean per 100K (mean of what?). The label however has not been changed in Figure 2.
Response- We thank the reviewer for their comments. We have now made appropriate changes in the manuscript to address their concerns.
Changes in manuscript – Figure 1 and Figure 2
6. Number 3 – response – might be worth commenting to this effect in the manuscript.
Response- We thank the reviewer for their comments. We however feel this is not really relevant here.
Changes in manuscript – No changes required

7. Number 4 –“ The percentage of cases with pneumonia as the secondary cause of hospitalisation has remained almost static at 2.5% in the 2000-2005 period, 2.4% in the 2007-2009 period and 2.5% in the 2010-2012 period.” Also worth mentioning in discussion perhaps.

Response- We thank the reviewer for their comments. We have now made appropriate changes in the manuscript to address their concerns.

Changes in manuscript – Page 8, line 42-44

Reviewer 2

8. Page 4 line 20: please replace "in 2+1 dose" with "using a 2+1 dose schedule".

Response- We thank the reviewer for their comments. This directly contradicts the advice by reviewer 1 (comment #2). We can only implement ONE of these changes. We have gone with the advice by reviewer 1, as it makes more sense to talk about PCV7 and PCV13 dosing schedules together. .

Changes in manuscript – Page 3, line 20

9. remove the newly introduced changes about 2+1 schedule

Response- As above

Changes in manuscript – Page 3, line 20

Editorial revisions requested

10. Conclusions- Could you please add a section "Conclusions" after section "Discussion". This should state clearly the main conclusions of the research article and give a clear explanation of their importance and relevance. You can use the last paragraph(s) of your Discussion if relevant.

Response- We have now made appropriate changes in the manuscript to meet your requirements

11. Declarations- Please ensure that your manuscript contains a Declarations section, with all of the following subsections:

Response- We have now made appropriate changes in the manuscript to meet your requirements

12. Please ensure all Figure legends are placed at the end of the text (after the References). Figures should just be the images themselves.

Response- We have now made appropriate changes in the manuscript to meet your requirements

Best wishes,
Dr Harish Nair

[Click here to view linked References](#)

Pneumonia hospitalisations in Scotland following the introduction of pneumococcal conjugate vaccination in young children

Harish Nair PhD^{1,7}, Arun Thor Watts BMedSci¹, Linda J Williams PhD², Saad B Omer PhD³, Colin R Simpson PhD⁴, Lorna J Willocks MD⁵, J Claire Cameron FFPH⁶, Harry Campbell MD^{1,2}

¹Centre for Global Health Research, Usher Institute of Population Health Sciences and Informatics, Edinburgh Medical School, University of Edinburgh

²Centre for Population Health Sciences, Usher Institute of Population Health Sciences and Informatics, Edinburgh Medical School, University of Edinburgh

³Hubert Department of Global Health, Rollins School of Public Health, Emory University

⁴Centre for Medical Informatics, Usher Institute of Population Health Sciences and Informatics, Edinburgh Medical School, University of Edinburgh

⁵Directorate for Public Health and Health Policy, NHS Lothian, Edinburgh

⁶Health Protection Scotland, NHS National Services Scotland, Glasgow

⁷Public Health Foundation of India, New Delhi, India

Corresponding author:

Dr. Harish Nair,
Reader in Global Health Epidemiology,
Centre for Global Health Research,
Usher Institute of Population Health Sciences and Informatics,
The University of Edinburgh,
Medical School, Teviot Place,
Edinburgh EH8 9AG
United Kingdom
Ph: +44 (0) 131 6506963
E-mail: Harish.Nair@ed.ac.uk

Author emails:

Harish Nair: Harish.Nair@ed.ac.uk

- 1 Arun Thor Watts: arun_watts@hotmail.co.uk
- 2 Linda J Williams: Linda.Williams@ed.ac.uk
- 3 Saad B Omer: somer@emory.edu
- 4 Colin R Simpson: c.simpson@ed.ac.uk
- 5 Lorna J Willocks: Lorna.Willocks@nhslothian.scot.nhs.uk
- 6 J Claire Cameron: clairecameron1@nhs.net
- 7 Harry Campbell: harry.campbell@ed.ac.uk

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Abstract

2 Background

3 Scotland introduced PCV7 and PCV13 immunisation in young children in 2006 and 2010 respectively.
4 One recent study from the United States reported a decrease in hospitalisation rates for all-cause
5 pneumonia most notably in adults older than 75 years of age following PCV7 introduction in the US
6 child population. We aimed to examine the effect of PCV7 and PCV13 on hospitalisation rates for all-
7 cause pneumonia across all age groups in Scotland.

8 Methods

9 We linked hospital records and death certification datasets for the entire Scottish population for the
10 period 2000 to 2012. We included all cases where the primary / secondary diagnosis was
11 pneumonia. Differences in hospital admission rates for pneumonia by age group were calculated
12 using the difference in average annual rates for each period.

13 Results

14 We estimated that all-cause pneumonia hospitalisation rates in children <2 years decreased by
15 about 30% in the post-PCV-13 period compared with the pre-PCV period. However, in adults aged
16 75-84 years and ≥85 years, all-cause pneumonia hospitalisation rates increased by 63% and 46 %
17 respectively in the post-PCV 13 period compared to the pre-PCV period. This resulted in an
18 additional 7000 hospitalisations across all age groups in Scotland in 2012 about half of which were in
19 adults >75 years. At the same time, the median length of hospital stay decreased by a third in
20 children <2 years and by about 20% in adults >75 years in the post-PCV13 period compared to the
21 pre-PCV period. Additionally, there was an 11% reduction in deaths due to all-cause pneumonia, and
22 30% reduction in pneumococcal hospitalisations across all age groups in the post-PCV13 period
23 compared with pre-PCV period.

24 Interpretation

25 The modest and sustained decline in the rates of hospitalisation for all-cause pneumonia in children
26 and the reduction in proportion of pneumonia hospitalisations in children coded as pneumococcal
27 disease in the post-PCV period should alleviate concerns that pneumococcal serotype replacement
28 may have resulted in an increased pneumonia burden in this age group. The indirect impact of child
29 PCV immunisation in those not vaccinated (in terms of reduction in all-cause pneumonia
30 hospitalisations in the elderly) has not been seen in Scotland. Our results are likely to be confounded
31 by changes in clinical coding and healthcare practices over the same period and illustrate that health
32 care planners cannot, with confidence, predict indirect PCV vaccine impacts on hospitalisations. IPD
33 surveillance across all age groups is needed to assess the indirect effects of PCV in the community.

34 Keywords:

35 pneumococcal conjugate vaccine; indirect effects; all-cause pneumonia hospitalisations;
36 pneumococcal hospitalisations; pneumonia mortality

1 Introduction

2 Globally, lower respiratory tract infections (primarily pneumonia) were the second leading cause of
3 morbidity and fourth leading cause of mortality across all age groups in 2013 [1]. The burden of
4 pneumonia is disproportionately higher in young children and the elderly (above 65 years of age) [2,
5 3]. In general, mortality due to pneumonia across all ages has declined substantially (22 (95% CI 16
6 to 30) percent) between 1990 and 2013 [1]. Prior to the introduction of the pneumococcal conjugate
7 vaccine (PCV) in the US, hospitalisation rates for pneumonia in the elderly (65 to 84 years) increased
8 by about 20% in the 15 year period between 1988-1990 to 2000-2002 [3]. Similar increases (over a
9 similar though abbreviated period) have been reported in England, Denmark, and the Netherlands
10 [4-6]. The increase in this age group has been attributed to an ageing population, increase in the
11 prevalence of comorbidities, and changes in coding and hospital admission practices [4].

12 The prevalence of *Streptococcus pneumoniae* in adults with community-acquired pneumonia (CAP)
13 has declined sharply since the advent of antibiotics. Data from Medicare (for year 2009) indicate that
14 about 7.6% of CAP hospitalisations in the elderly (aged 65 years and above) in the United States can
15 be attributed to *Streptococcus pneumoniae* [7]. Therefore, pneumococcal polysaccharide vaccine
16 (PPV) was introduced in Scotland in the winter of 2003 to persons aged 65 years and above, and high
17 risk individuals (i.e. those with co-morbidities). Following the introduction of PCV7 into the national
18 child immunisation schedules in the US and England and Wales, there has been a decrease in the
19 overall incidence of invasive pneumococcal disease (IPD) across all age groups [8, 9]. PCV7 was
20 introduced into the Scottish child immunisation schedule in September 2006 at the age of 2 and 4
21 months with a booster dose at around 13 months. Severe pneumococcal disease (using IPD as a
22 surrogate) was increasing in Scotland in the period prior to introduction of PCV7 (from 559 cases in
23 2000 to 753 cases in 2006) [10]. After the introduction of PCV7, and PCV13 later in 2010 (both using
24 2+1 dose schedule), there has been a decrease in the number of IPD cases in children younger than 5
25 years (from 753 cases in 2006 to 482 cases in 2012). It was expected that with the increased uptake
26 of PCV, there would be an increase in the circulating non-vaccine serotypes (NVT) [11, 12]. In the US,
27 after the introduction of PCV7, there was an increase in disease (across all age groups especially in
28 young children and elderly) caused by NVT, most notably serotype 19 A [13]. In addition, in 2013,
29 Griffin and colleagues demonstrated a decrease in hospitalisation rates for all-cause pneumonia
30 (henceforth referred to as pneumonia) not only in children but also across all age groups, most
31 notably in adults older than 75 years of age following introduction PCV7 in the US child population
32 [14]. We aimed to examine the effect of PCV7 and PCV13 on hospitalisation rates for pneumonia
33 across all age groups in Scotland and examine the generalizability of Griffin and colleagues' findings.

34 Methods

35 Data Sources

36 Scotland has an excellent health service data system with an ability to perform high quality linkage
37 (through a unique personal identifier attached to health records – the Community Health Index [CHI]
38 number) of individual hospitalisation, laboratory and death certification data which permits
39 individual patient data analysis and follow-up. National Services Scotland is responsible for
40 maintaining these data and ensuring high data quality, consistency and coverage and appropriate
41 levels of data governance (<http://www.isdscotland.org/>). The Scottish Morbidity Record (SMR01)
42 comprises episode level data on hospital inpatient and day case discharges from acute specialties

1 from hospitals in Scotland. The dataset contains patient identifiers such as the CHI number,
2 demographic details such as postcode together with health record data (episode management
3 details and general clinical information). These data have been available electronically (for research
4 purposes) since 1981. Usually, when inpatients and day cases are transferred to another specialty
5 within the same hospital or are transferred to another hospital, a new record on SMR01 is
6 generated. However, in this analysis, we used continuous inpatient stay (CIS) data by linking SMR01
7 episodes for each patient. This created “linked” individual patient histories (which includes transfers
8 between hospitals, specialties, and consultants). This integrated measure documents the length of
9 continuous hospital care episodes. Data from SMR01 on hospital admissions for pneumonia,
10 hospitalisation, discharge diagnosis of pneumococcal pneumonia, readmissions within 14 days, sex,
11 age, health board area, length of stay; and mortality data from General Register Office of Scotland
12 (GRO99B) were collected.

13 Definition of hospitalisation for pneumonia

14 Hospitalisation for pneumonia was defined in accordance with ICD 10 procedures. CIS data were
15 selected for inclusion in the study if the patient had a primary diagnosis of pneumonia OR a primary
16 diagnosis of sepsis, meningitis or empyema, with a diagnosis of pneumonia in any subsequent
17 diagnostic position. A list of ICD 10 codes used to qualify diagnosis for all-cause pneumonia,
18 meningitis, empyema, septicemia and pneumococcal pneumonia are detailed in Supplementary
19 table 1. We considered hospital readmissions within 14 days of the primary admission as part of the
20 same event[15].

21 Statistical analysis

22 SMR01 data from 2000 through 2012 (epidemiological year July-June) were used as numerator data
23 in estimating annual hospitalisation rates for pneumonia. We constructed a cohort of patients by
24 linking hospital records and death certification datasets. We used the following age groups for
25 analysis: children aged below two years, 2 to 4 years, 5 to 17 years, adults aged 18 to 39 years, 40 to
26 64 years, 65 to 74 years, 75 to 84 years, and 85 years or older. We stratified the analysis into three
27 different time periods: 2000 through 2005 were considered as the pre-PCV period; 2006 was treated
28 as transition year, as this was the year that PCV was introduced nationally and therefore excluded
29 from the analysis; 2007 through 2009 as the post-PCV7 period; and 2010 through 2012 as the post-
30 PCV13 period (as PCV 13 was introduced nationally in April 2010). We used population by Scottish
31 NHS Health Board of residence provided by ISD (based on official national census data and
32 estimates) as population denominator data in the calculation of annual hospital admission rates (by
33 age group) and reported these per 100,000 persons per year. Differences in hospital admission rates
34 by age group were calculated using the average annual rates for each period, and the observed and
35 expected rates were compared. We used these data to derive the percentage difference in relative
36 rates. Where no data were available for a particular health board (age/sex/year combination), most
37 frequently seen in the case of the smallest Health Boards, we assumed that there were no cases of
38 pneumonia, rather than regarding these as missing data. Where the Health Board was given as
39 “Other”, we considered that these could be due to two reasons: data were missing but the patient
40 was resident in Scotland; or patient was a resident of another country but was treated in Scotland.
41 We considered both scenarios in the analysis.

42 All data analyses were conducted using SPSS V19.0 (IBM, NY, USA) and SAS V9.2 (SAS institute Inc.,
43 NC, USA).

Results

From 2000 to 2012, there were 153,636 new hospitalisation episodes for pneumonia across Scotland. In addition to these hospitalisations, there were 4551 readmissions within 14 days of discharge. Pneumonia was the first listed diagnosis for 97.6% of these hospitalisations. Sepsis was the first listed diagnosis for 1.9%, and meningitis and empyema the first listed diagnosis each for 0.3% and 0.2% respectively. Prior to the introduction of PCV, we observed a gradual increase in the rates of pneumonia hospitalisation in children aged below 5 years (Figure 1). A similar pattern was seen in adults aged over 75 years (Figure 2). Across the twelve year period (2000-2012), the annual rates of pneumonia hospitalisations were highest in adults aged above 65 years and children aged below two years (Table 1). In the pre-PCV period, the rates for pneumonia hospitalisation in adults aged 75 to 84 years and 85 years and above, were approximately 3 and 6.5 fold higher respectively than those in children aged less than two years. In the post-PCV 7 period, this difference increased to about 5 and 9 fold respectively; and further increased to about 7 and 13 fold respectively in the post-PCV 13 period. This pattern was sustained even when data from “other” Health Boards were excluded (Supplementary figure 1 and 2).

The pneumonia hospitalisation rates in children aged less than 2 years decreased by about 30% in the post-PCV 13 period compared to the pre-PCV period (Figure 3). Pneumonia hospitalisations in this age group declined by about 19 percent in the post-PCV7 period (compared to pre-PCV) and by about 14 percent in the post-PCV13 period (compared to post-PCV7). However, the rates remained largely unchanged in the age groups 2-39 years. We saw that in the older age groups, the pneumonia hospitalisation rates increased steadily in the post-PCV period (greater increase post-PCV13 compared to post-PCV7). In adults aged 75 -84 years and 85 years and above, the pneumonia hospitalisation rates increased by 63 and 46 percent respectively in the post-PCV 13 period compared to the pre-PCV period (Table 1, Supplementary table 2). This translates to an absolute increase of about 7000 hospitalisations across all age groups in Scotland in 2012. Half of this increase was observed in age groups older than 74 years (increase of 2200 and 1300 hospitalisations in the age group 75-84 years and 85 years and above respectively (Table 2).

We observed that during 2000-2012, the mean and median length of stay (LOS) increased with age (Supplementary figure 3 and supplementary table 3). The pattern of longer stays (defined as more than 30 days) is the familiar “hook” pattern with the lowest number being in the 2-4 year age group and then increasing with age (Supplementary table 4). This pattern is sustained even if a lower threshold (of stays more than 14 days) is used to define longer hospitalisation. In the post-PCV 13 period, median length of hospital stay decreased by a third in children aged below 2 years and by 22 and 18 percent in the age groups 75-84 years and 85 years and above respectively compared with the pre-PCV period (Table 1). We also observed that although the distribution of length of stay in hospital (across all age groups) remained the same in the pre-PCV and post-PCV13 period, there was a significantly higher proportion of patients with stays longer than 14 days in the former (24.8% vs 20.9%, $p < 0.001$) (supplementary table 5). When the analysis was stratified by age, the pattern of significantly higher proportion of patients with stays longer than 14 days was seen only in adults aged over 40 years (Supplementary table 5).

The proportion of pneumonia hospitalisations coded as pneumococcal disease decreased substantially across all age groups in the post-PCV13 period (Table 1). The most dramatic reduction

1 was seen in children aged below 5 years (about a two-thirds reduction) while in the elderly aged 85
2 years only a one-third reduction was demonstrated. At a population level, we observed 29 (95% CI
3 22 to 35) percent reduction in pneumococcal admissions across all age groups in Scotland in the
4 4 post-PCV13 period compared to pre-PCV period. However, this varied by age group -while there was
5 5 a 70% reduction in pneumococcal hospitalisations in children aged below 5 years in the post-PCV13
6 6 period compared to pre-PCV period, older age groups experienced a more modest reduction (21
7 7 and 25 percent in those 65-74 years and 75 years and above respectively) (Supplementary table 6).

8 We observed a substantial reduction in the proportion of deaths due to pneumonia (as a primary
9 cause) across all age groups in the post-PCV13 period when compared to pre-PCV period. In elderly
10 patients aged 75-84 and 85 years and above there was a decrease of about 53% and 40%
11 respectively (Table 1). This pattern was sustained when deaths from pneumonia were defined more
12 broadly and included all deaths in which pneumonia was mentioned on the death certificate but was
13 not recorded as the underlying cause of death (Supplementary table 7). At a population level, there
14 was an overall 11% reduction in deaths in post-PCV13 period compared to pre-PCV period, with a 19
15 and 12 percent reduction in those aged 65-74 years and 75 years and above respectively
16 (supplementary table 8). We also observed a 7 percent reduction in pneumococcal deaths across all
17 age groups with an 18 and 12 percent reduction in the 65-74 years and 75 years and above age
18 groups respectively (Supplementary table 9).

19 Discussion

20 This is the first national study to report the impact of a child PCV immunisation programme on
21 pneumonia hospitalisations across all age groups. Our results are based on a Scottish national
22 dataset for a 12 year period (2000-2012). We observed modest and sustained declines in the
23 childhood hospitalisation rates for pneumonia in the six years following the introduction of PCV 7
24 and 13. This should alleviate concerns regarding pneumococcal serotype replacement, as we have
25 found no evidence of increased pneumonia burden in children following the introduction of PCV. We
26 also observed a 33% reduction in the median length of stay in hospital for pneumonia in children
27 aged below 2 years. However, we noted a dramatic and progressive increase in pneumonia
28 hospitalisations in adults, most notably in those aged above 65 years. This translates to an annual
29 increase of about 7000 hospital admissions for pneumonia across Scotland. This increase in hospital
30 admissions is accompanied by contrasting decreases in the length of stay in hospital, the proportion
31 of hospital admissions coded as pneumococcal pneumonia and pneumonia mortality in adults aged
32 over 75 years.

33 We found a gradual increase in the annual pneumonia hospitalisation rates for children aged below
34 5 years in the pre-PCV period (Figure 1). These results are consistent with those reported in previous
35 analyses of pneumonia trends in Scotland and England prior to the introduction of PCV [16, 17].
36 Koshy and colleagues concluded that this was likely due to a true increase in disease incidence
37 (reflected in hospitalisation rates), as it was consistent with similar reports elsewhere.

38 We reported a 30% decline in pneumonia hospitalisations in children aged below 2 years following
39 the introduction of first PCV 7 and later PCV 13, which is reassuringly consistent with results of
40 clinical trials investigating the effectiveness of PCV against radiographically confirmed WHO Primary
41 Endpoint Pneumonia (bacterial) pneumonia [18, 19]. A previous study from England reported 19 to
42 22 percent decline in hospitalisation rates for bacterial pneumonia in children aged below 15 years

1 in the two years post-PCV7 (2007 and 2008) [17]. Our results for the six years post-PCV in children
2 aged below 18 years are consistent with those reported by Griffin and colleagues for US populations
3 in the first decade after introduction of PCV 7 [14] (Panel 1).

4 However, in older adults, our results contrast markedly from findings reported by Griffin and
5 colleagues [14]. We estimated that in Scotland, the hospitalisation rate for all-cause pneumonia in
6 adults aged 65-74 years in the post-PCV13 period was 640.4 per 100,000 persons per year compared
7 to 376.6 per 100,000 persons per year in the pre-PCV period. Similar increases were seen in the age
8 groups 75 years and above. By contrast, Griffin and colleagues reported a slight decrease in the rate
9 of hospitalisation in the post-PCV7 period (1208 compared to 1293 per 100,000 persons per year) in
10 the 65-74 year age group. More substantial decreases were reported in the older age groups (75
11 years and above). In Scotland, this translates to an additional 7000 hospital admissions for
12 pneumonia in post-PCV period, compared to 112,000 fewer hospitalisations in U.S adults. Griffin and
13 colleagues attributed these declines to the indirect “herd immunity” effect of PCV on pneumococcal
14 pneumonia. We did however observe a substantial decline in the proportion of pneumonia
15 hospitalisations coded as pneumococcal disease across all age groups in the post-PCV period.
16 However, it is very likely that we may have underestimated the impact of PCV on pneumococcal
17 pneumonia. This is because hospital discharge data are based on clinical diagnoses recorded in
18 patients’ medical records and accurate diagnosis of pneumococcal pneumonia is difficult without
19 isolation of *S. pneumoniae* [20]. Notably, since 2010, we observed an increase in both the
20 hospitalisations for pneumococcal pneumonia (numerator) and all-cause pneumonia (denominator)
21 in adults older than 65 years even though there was a decrease in the proportion of pneumonia
22 cases coded as pneumococcal pneumonia (Supplementary table 11). Extrapolating from the
23 pneumonia hospitalisation rates in the pre-PCV period, we calculated the (expected) number of
24 pneumonia hospitalisations assuming absence of PCV immunisation in the period 2007-12. Unlike in
25 younger children, the expected number of pneumonia hospitalisations during this period would have
26 been lower in older children (aged above 5 years) and adults. Therefore, it appears that the impact
27 of child PCV immunization in Scotland on pneumonia hospitalisations has been limited to children
28 aged younger than 5 years (Table 2).

29 There are several reasons why our results are not consistent with those observed by Griffin and
30 colleagues. Firstly, this could be attributed to differences in coding practices in both countries. The
31 US Nationwide Inpatient Sample (NIS) database uses up to 15 diagnostic positions, whereas ISD only
32 uses up to 6 diagnostic positions. Therefore for a first diagnosis of meningitis, septicaemia or
33 empyema, pneumonia could have been a secondary diagnosis in one of 14 diagnostic positions in
34 the US study, but a secondary diagnosis in only one of 5 diagnostic positions in the present study.
35 This could mean that fewer cases of septicaemia, meningitis and empyema were included as
36 pneumonia cases in Scotland. It has been reported that during 2003-09 in the US there was an
37 observed decrease in the coding for a primary diagnosis of pneumonia, which was partially
38 compensated for by increases in coding for primary diagnoses of other diseases e.g. sepsis and
39 respiratory failure [22]. Therefore, the decline in hospitalisations for pneumonia reported in the US
40 may be partially due to an artefact of changes in diagnostic coding. In Scotland, the percentage of
41 hospitalisations with pneumonia as a secondary diagnosis has remained almost static at 2.5% in the
42 2000-2005 period, 2.4% in the 2007-2009 period and 2.5% in the 2010-2012 period. Although the
43 study by Griffin and colleagues would have included cases with a primary diagnosis of sepsis as
44 pneumonia hospitalisation, they would not have included diagnoses that were coded as respiratory

1 failure [14]. Indeed, Lindenauer and colleagues have reported that when combined with diagnoses
2 of sepsis and respiratory failure, much more modest declines were seen in the hospitalisation rates
3 for pneumonia [22]. Secondly, the US study was based on a 20% sample of US hospitals, whereas our
4 study was based on inpatient data from the whole of Scotland. Thirdly, the increasing trend of adult
5 pneumonia hospitalisations in Scotland may be an artefact of changing healthcare practices. This
6 could have overcompensated for any declines in adult all-cause pneumonia hospitalisations that may
7 have been seen as a result of the known herd effect of PCV [13]. Indeed, there was a 10 percent
8 increase in all-cause hospitalisations (defined using CIS) during our study period (2000-2012) (data
9 source: Information Services Division, National Services Scotland in response to data request IR2015-
10 01696). Fourthly, the increase in pneumonia hospitalisations could be attributed to improvement in
11 coding practices (even though ICD-10 codes have been in use in Scotland since 1996). Coding is also
12 reported to have improved in England[23] and may have contributed to the apparent increases in
13 pneumonia hospitalisations reported there [17]. In England and Wales, in the period 2000-10, there
14 was an improvement in case ascertainment for invasive bacterial infections which is corroborated by
15 a concomitant increase in cases of other invasive bacterial infections reported to Public Health
16 England. Miller and colleagues corrected for these changes by retrospectively inflating the case
17 count to the projected 2009-10 level of ascertainment [9]. They found that without adjusting for
18 these changes there was little reported change in IPD incidence (across all age groups except
19 children younger than 5 years) after PCV introduction although PCV has been demonstrated to have
20 a substantial impact on IPD incidence both in children and adults [24].

21 The National Audit Office (NAO) has reported an increasing trend of emergency hospital admissions
22 in Scotland and the rest of the UK [25]. The increase was thought to be partially due to an ageing
23 population and due to a tendency to treat patients as day cases or give patients earlier discharge to
24 reduce bed pressure, resulting in more readmissions. This is consistent with our observation of a
25 trend towards decreasing length of hospital stay for patients with pneumonia. In England there was
26 a 40% increase in pneumonia hospitalisations for children aged less than 15 years between 1999
27 through 2010 [26]. Moreover, another study from England has reported a 34% increase in hospital
28 admissions for pneumonia before the introduction of PCV, mainly for pneumonias coded with an
29 unspecified diagnosis [4]. Subsequent to changes to the general medical services contracts in 2004,
30 General Practitioners in the UK (including Scotland) were no longer required to provide out of hours
31 services. In England and Wales, this has resulted in a 32% increase in A&E attendances between
32 2003-04 and 2013-14, translating into an 8% growth in the likelihood of being admitted to hospital
33 after attending A&E during this period [27]. Indeed, patients may be more likely to be sent to
34 hospital by an unfamiliar doctor [4]. This is likely to be true for Scotland as well. With increased
35 emergency department visits and referrals there would be a greater diagnosis of pneumonias that
36 would have otherwise gone undiagnosed. We also observed an overall increase in the percentage of
37 hospital readmissions for pneumonia from 2.0% in 2000 to 3.6% in 2012. We therefore considered
38 episodes generated from readmissions within 14 days to be part of the same event and
39 consequently excluded them from our primary analysis. However, when disaggregated by age
40 groups, we observed that the proportion of early readmissions (within 14 days of discharge)
41 decreased in those younger than 40 years (the most substantial decrease of about 74 percent being
42 in children aged below 2 years) (Supplementary table 9). The US study also found a trend of
43 decreasing hospital stays before and after the introduction of PCV and found decreases in
44 pneumonia hospitalisation rates, even without adjusting for readmissions [14].

1 The difference between our findings and those reported by Griffin cannot be attributed to
2 differences in pneumococcal polysaccharide vaccine (PPV) coverage in the two populations as they
3 are likely to be similar for Scotland and U.S. Limited PPV coverage data for Scotland are available –
4 the initial vaccination uptake was estimated to be 65% [28]. Estimates for England indicate a
5 coverage of about 69 percent [29]. Similarly, PPV was introduced in the US in 1997 for all individuals
6 aged 65 years and above. Although recent coverage data are not available, the data for 2005
7 estimate PPV coverage to be about 64 percent [30].

8 Conclusions

9 Our findings support the evidence on the impact of PCV on pneumonia hospitalisations in children,
10 indicating that vaccine impact in this national programme is broadly consistent with predictions from
11 vaccine trials. It is reassuring that reductions in hospitalization have persisted over 6 years with no
12 evidence of negative indirect effects (due to serotype replacement) eroding this impact over this
13 time period. This reduced proportion of pneumonia hospitalisations in children coded as
14 pneumococcal disease post PCV13 introduction gives further reassurance that there is not a
15 substantial serotype replacement effect over the study period. The failure to note any positive
16 indirect effect of infant PCV vaccination on (pneumococcal) pneumonia in non-vaccinated individuals
17 observed through measurement of hospitalisations among older individuals is likely due to
18 contemporaneous changes in clinical practice and coding as discussed above. This suggests that
19 indirect effects of PCV on observed hospitalisations may be context specific and demonstrates that
20 health care planners and health economists cannot, with confidence, predict positive indirect PCV
21 vaccine impacts on hospitalisations in their estimates of vaccine impact on health service costs. IPD
22 surveillance in all age groups is needed to assess the indirect effects of PCV in the community.

23 Abbreviations

24 CAP – community-acquired pneumonia

25 CHI – Community Health Index

26 CI – confidence interval

27 CIS – continuous inpatient stay

28 ICD – International classification of diseases

29 IPD – invasive pneumococcal disease

30 ISD – Information Services Division

31 LOS – length of stay

32 NAO – National Audit Office

33 NHS – National Health Service

34 NIS – Nationwide Inpatient Sample

35 NVT – non-vaccine serotypes

- 1 PCV – pneumococcal conjugate vaccine
2
3
4 2 PPV – pneumococcal polysaccharide vaccine
5
6
7 3 SHLMPRL – Scottish Haemophilus, Legionella, Meningococcus and Pneumococcus Reference
8 Laboratory
9
10 4 UK – United Kingdom
11
12 5 US – United States
13

14 7 **Declarations**

15 8 **Ethics approval and consent to participate**

16 9 This research is in compliance with the Helsinki Declaration
17 (http://www.wma.net/en/30publications/10policies/b3/index.html). A waiver for ethical approval
18 10 was obtained from the Caldicott Guardian as the data on human subjects used in this analysis was
19 11 not having any personal identifiers and therefore deemed to be “low risk”.
20 12

21 13 **Consent to publish**

22 14 The results have been reviewed internally by Health Protection Scotland and have been approved
23 15 for publication.

24 16 **Availability of data and materials**

25 17 The primary data used in this analysis can be requested from the ISD, National Services Scotland.
26

27 18 **Author’s Contributions**

28 19 HN and HC conceptualised the study. ATW conducted literature review. LJ Williams and ATW
29 20 conducted the analysis. All authors contributed to data interpretation. HN prepared the initial draft
30 21 of the manuscript. All contributed to report writing and critically reviewed the manuscript. All
31 22 authors read and approved the final draft of the manuscript.

32 23 **Acknowledgement**

33 24 We would like to acknowledge the contribution of Dr. Rachael Wood, ISD, National Services Scotland
34 25 for their helpful advice and facilitating data access.
35

36 26 **Competing interest**

37 27 All authors declare that they have no conflicts of interest.
38

39 28 **Funding**

40 29
41 30 None
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

1. GBD 2013 Mortality and Causes of Death Collaborators: **Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013.** *Lancet* 2015, **385**(9963):117-171.
2. Rudan I, O'Brien KL, Nair H, Liu L, Theodoratou E, Qazi S, Luksic I, Fischer Walker CL, Black RE, Campbell H: **Epidemiology and etiology of childhood pneumonia in 2010: estimates of incidence, severe morbidity, mortality, underlying risk factors and causative pathogens for 192 countries.** *J Glob Health* 2013, **3**(1):10401.
3. Fry AM, Shay DK, Holman RC, Curns AT, Anderson LJ: **Trends in hospitalizations for pneumonia among persons aged 65 years or older in the United States, 1988-2002.** *JAMA* 2005, **294**(21):2712-2719.
4. Trotter CL, Stuart JM, George R, Miller E: **Increasing hospital admissions for pneumonia, England.** *Emerg Infect Dis* 2008, **14**(5):727-733.
5. Thomsen RW, Riis A, Norgaard M, Jacobsen J, Christensen S, McDonald CJ, Sorensen HT: **Rising incidence and persistently high mortality of hospitalized pneumonia: a 10-year population-based study in Denmark.** *J Intern Med* 2006, **259**(4):410-417.
6. Oosterheert JJ, Bonten MJ, Hak E, Lammers JW, Schneider MM, Hoepelman IM: **[The increase in pneumonia-related morbidity and mortality among adults in the Netherlands and possible explanations for it].** *Ned Tijdschr Geneesk* 2004, **148**(36):1765-1769.
7. Bartlett JG: **Diagnostic tests for agents of community-acquired pneumonia.** *Clin Infect Dis* 2011, **52** Suppl 4:S296-304.
8. Lexau CA, Lynfield R, Danila R, Pilishvili T, Facklam R, Farley MM, Harrison LH, Schaffner W, Reingold A, Bennett NM *et al*: **Changing epidemiology of invasive pneumococcal disease among older adults in the era of pediatric pneumococcal conjugate vaccine.** *JAMA* 2005, **294**(16):2043-2051.
9. Miller E, Andrews NJ, Waight PA, Slack MP, George RC: **Herd immunity and serotype replacement 4 years after seven-valent pneumococcal conjugate vaccination in England and Wales: an observational cohort study.** *Lancet Infect Dis* 2011, **11**(10):760-768.
10. **Respiratory Infections: Pneumococcal Disease**
[\[http://www.hps.scot.nhs.uk/resp/pneumococcaldisease.aspx#\]](http://www.hps.scot.nhs.uk/resp/pneumococcaldisease.aspx#)
11. Pelton SI, Huot H, Finkelstein JA, Bishop CJ, Hsu KK, Kellenberg J, Huang SS, Goldstein R, Hanage WP: **Emergence of 19A as virulent and multidrug resistant Pneumococcus in Massachusetts following universal immunization of infants with pneumococcal conjugate vaccine.** *Pediatr Infect Dis J* 2007, **26**(6):468-472.
12. Moore MR, Gertz RE, Jr., Woodbury RL, Barkocy-Gallagher GA, Schaffner W, Lexau C, Gershman K, Reingold A, Farley M, Harrison LH *et al*: **Population snapshot of emergent Streptococcus pneumoniae serotype 19A in the United States, 2005.** *J Infect Dis* 2008, **197**(7):1016-1027.
13. Pilishvili T, Lexau C, Farley MM, Hadler J, Harrison LH, Bennett NM, Reingold A, Thomas A, Schaffner W, Craig AS *et al*: **Sustained reductions in invasive pneumococcal disease in the era of conjugate vaccine.** *J Infect Dis* 2010, **201**(1):32-41.
14. Griffin MR, Zhu Y, Moore MR, Whitney CG, Grijalva CG: **U.S. hospitalizations for pneumonia after a decade of pneumococcal vaccination.** *N Engl J Med* 2013, **369**(2):155-163.
15. Flasche S, Slack M, Miller E: **Long term trends introduce a potential bias when evaluating the impact of the pneumococcal conjugate vaccination programme in England and Wales.** *Euro Surveill* 2011, **16**(20):19868.
16. Roxburgh CS, Youngson GG, Townend JA, Turner SW: **Trends in pneumonia and empyema in Scottish children in the past 25 years.** *Arch Dis Child* 2008, **93**(4):316-318.
17. Koshy E, Murray J, Bottle A, Sharland M, Saxena S: **Impact of the seven-valent pneumococcal conjugate vaccination (PCV7) programme on childhood hospital admissions**

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- for bacterial pneumonia and empyema in England: national time-trends study, 1997-2008.** *Thorax* 2010, **65**(9):770-774.
18. Hansen J, Black S, Shinefield H, Cherian T, Benson J, Fireman B, Lewis E, Ray P, Lee J: **Effectiveness of heptavalent pneumococcal conjugate vaccine in children younger than 5 years of age for prevention of pneumonia: updated analysis using World Health Organization standardized interpretation of chest radiographs.** *Pediatr Infect Dis J* 2006, **25**(9):779-781.
19. Cutts FT, Zaman SMA, Enwere G, Jaffar S, Levine OS, Okoko JB, Oluwalana C, Vaughan A, Obaro SK, Leach A *et al*: **Efficacy of nine-valent pneumococcal conjugate vaccine against pneumonia and invasive pneumococcal disease in The Gambia: randomised, double-blind, placebo-controlled trial.**[Erratum appears in *Lancet*. 2005 Jul 2-8;366(9479):28]. *Lancet* 2005, **365**(9465):1139-1146.
20. Obaro SK, Madhi SA: **Bacterial pneumonia vaccines and childhood pneumonia: are we winning, refining, or redefining?** *Lancet Infect Dis* 2006, **6**(3):150-161.
21. **Scottish Haemophilus, Legionella, Meningococcus & Pneumococcus Reference Laboratory** [[http://library.nhsggc.org.uk/mediaAssets/SMRL/SHLMPRL_UM_2014_v2.1_User_Manual\[1\].pdf](http://library.nhsggc.org.uk/mediaAssets/SMRL/SHLMPRL_UM_2014_v2.1_User_Manual[1].pdf)]
22. Lindenauer PK, Lagu T, Shieh MS, Pekow PS, Rothberg MB: **Association of diagnostic coding with trends in hospitalizations and mortality of patients with pneumonia, 2003-2009.** *JAMA* 2012, **307**(13):1405-1413.
23. Bottle A, Aylin P, Majeed A: **Identifying patients at high risk of emergency hospital admissions: a logistic regression analysis.** *J R Soc Med* 2006, **99**(8):406-414.
24. Moore MR, Link-Gelles R, Schaffner W, Lynfield R, Lexau C, Bennett NM, Petit S, Zansky SM, Harrison LH, Reingold A *et al*: **Effect of use of 13-valent pneumococcal conjugate vaccine in children on invasive pneumococcal disease in children and adults in the USA: analysis of multisite, population-based surveillance.** *Lancet Infect Dis* 2015, **15**(3):301-309.
25. Comptroller and Auditor General: **Emergency admissions to hospital: managing the demand.** In. London: National Audit Office; 2013: 53.
26. Gill PJ, Goldacre MJ, Mant D, Heneghan C, Thomson A, Seagroatt V, Harnden A: **Increase in emergency admissions to hospital for children aged under 15 in England, 1999-2010: national database analysis.** *Arch Dis Child* 2013, **98**(5):328-334.
27. Blunt I, Edwards N, Merry L: **What's behind the A&E crisis?** In. London: Nuffield Trust; 2015.
28. Mooney JD, Weir A, McMenamin J, Ritchie LD, Macfarlane TV, Simpson CR, Ahmed S, Robertson C, Clarke SC: **The impact and effectiveness of pneumococcal vaccination in Scotland for those aged 65 and over during winter 2003/2004.** *BMC Infect Dis* 2008, **8**:53.
29. Public Health England: **Pneumococcal Polysaccharide Vaccine (PPV) coverage report, England, April 2013 to March 2014.** In: *Health Protection Report: weekly report*. vol. 8; 2014.
30. Centers for Disease Control and Prevention: **Pneumococcal Disease.** In: *Epidemiology and Prevention of Vaccine-Preventable Diseases*. Edited by Hamborsky J, Kroger A, Wolfe S, 13th edn. Washington D.C: Public Health Foundation; 2015.

Figure title and legends

Figure 1: Pneumonia hospitalisations in Scotland from July 2000 to June 2012 in children. Panel A shows annual rates of hospitalisation for pneumonia from 2000 through 2012. PCV 7 was introduced in 2006 and PCV 13 in 2010. Panel B shows annual rates of hospitalisation for 2000 to 2005, 2007 to 2009 and 2010 to 2012 for all children (including those coded as “other” health boards)

Figure 2: Pneumonia hospitalisations in Scotland from July 2000 to June 2012 in adults. Panel A shows annual rates of hospitalization for pneumonia from 2000 through 2012. PCV 7 was introduced in 2006 and PCV 13 in 2010. Panel B shows annual rates of hospitalization for 2000 to 2005, 2007 to 2009 and 2010 to 2012 for all adults (including those coded as “other” health boards)

Figure 3: Average Annual Rates of Scottish hospitalisations for Pneumonia before and after the introduction of PCV7 and PCV13

Tables

Table 1: Rates of hospitalisation, length of hospital stay, and rates of in-hospital death related to pneumonia in Scotland 5 years before introduction of PCV 7 and three years after introduction of PCV 13

Age (years)	<2	2-4	5-17	18-39	40-64	65-74	75-84	85+
Pneumonia Hospitalisation rate /100K* (95% CI)								
2000-2005	293 (280-306)	181 (173-190)	43 (41-45)	53 (51-54)	112 (110-114)	373 (365-380)	875 (861-889)	1909 (1872-1946)
2010-2012	206 (191-221)	177 (166-188)	45 (42-48)	62 (59-64)	182 (179-186)	634 (621-647)	1426 (1402-1451)	2785 (2728-2842)
Lengths of hospital stay (days). Median (IQR)								
2000-2005	3 (1-5)	2 (1-4)	3 (1-5)	4 (2-6)	5 (3-10)	7 (4-14)	9 (4-19)	11 (5-25)
2010-2012	2 (1-5)	2 (1-4)	2 (1-5)	3 (1-6)	5 (2-9)	6 (3-12)	7 (4-15)	9 (4-19)
Lengths of hospital stay (days) incl. readmissions within 14 days. Median, IQR								
2000-2005	3 (1-5)	2 (1-4)	3 (1-5)	4 (2-7)	5 (3-11)	7 (4-15)	9 (4-20)	12 (4-25)
2010-2012	2 (1-5)	2 (1-4)	2 (1-5)	3 (1-7)	5 (2-9)	6 (3-12)	8 (4-16)	9 (4-19)
Reasons for admission coded as pneumococcal disease. %, 95% CI								
2000-2005	9.3 (8.1-10.7)	3.3 (2.5-4.2)	4.1 (3.4-5.1)	6.8 (6.1-7.5)	6.1 (5.7-6.6)	3.8 (3.4-4.1)	2.3 (2.0-2.5)	1.1 (1.0-1.4)
2010-2012	3.3 (2.3, 4.9)	1.1 (0.6-1.9)	2.5 (1.7-3.6)	3.5 (2.9-4.3)	2.9 (2.6-3.3)	1.7 (1.4-2.0)	1.0 (0.8-1.2)	0.7 (0.5-0.9)

* The hospitalisation rates have been calculated after excluding data where patient's health board was coded as "other"

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Age (years)	<2	2-4	5-17	18-39	40-64	65-74	75-84	85+
Deaths from pneumonia (primary cause, all patients)[†] %, 95% CI								
2000-2005	0	0.06 (0.008, 0.4)	0.2 (0.1, 0.6)	0.7 (0.5, 1.0)	3.5 (3.2, 3.8)	6.9 (6.5, 7.4)	13.1 (12.5, 13.6)	25.8 (24.9, 26.6)
2010-2012	0	0	0.3 (0.1, 0.9)	0.3 (0.1, 0.6)	1.4 (1.2, 1.7)	3.4 (3.0, 3.7)	6.2 (5.8, 6.6)	15.7 (15.0, 16.5)
Deaths from pneumonia as proportion of all deaths (primary cause)[‡] %, 95% CI								
2000-2005	0	2.4 (0.3, 15.1)	3.1 (1.3, 7.4)	5.2 (3.7, 7.2)	7.4 (6.7, 8.2)	8.7 (8.1, 9.3)	14.1 (13.5, 14.7)	26.3 (25.4, 27.2)
2010-2012	0	0	11.1 (3.6, 29.3)	4.3 (2.2, 8.4)	6.0 (5.1, 7.1)	8.2 (7.4, 9.2)	12.0 (11.2, 12.8)	24.0 (22.9, 25.1)
Deaths from pneumonia (any cause, all patients)[§] %, 95% CI								
2000-2005	0.8 (0.5, 1.3)	0.6 (0.3, 1.0)	2.3 (1.8, 3.0)	3.7 (3.2, 4.3)	18.8 (18.1, 19.5)	36.9 (35.9, 37.8)	52.8 (52.0, 53.6)	66.1 (56.1, 67.0)
2010-2012	0.7 (0.3, 1.6)	0.2 (0.05, 0.8)	2.1 (1.4, 3.2)	2.4 (1.9, 3.1)	8.5 (8.0, 9.1)	14.9 (14.2, 15.6)	21.6 (20.9, 22.3)	36.1 (35.1, 37.1)

[†] Deaths with pneumonia as a primary cause as a proportion of the whole population
[‡] Deaths with pneumonia as a primary cause as a proportion of the deaths reported in this population
[§] Deaths with pneumonia as any cause as a proportion of the whole population

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65


Table 2: Difference in Pneumonia Hospitalisation Rates in Scotland Comparing pre-PCV7 and post-PCV13 Periods. With Absolute Change in Hospitalisations by 2012

Age group	Pneumonia Hospitalisations in 2012	Scottish Population 2012	Difference in hospitalisation rates per 100,000 population, PrePCV7 vs LatePCV7/PCV13 (n, 95% CI)	Difference in Hospitalisations, PrePCV7 to Late PCV7/PCV13 (% , 95% CI) ¹	Expected number of hospitalisations from 2000-2006 distribution ²	Estimated Absolute Change in Number of Hospitalisations, 2012 (observed – expected), n, 95% CI
<2	240	119,275	-87.0 (-107.0, -67.0)	-29.7 (-36.5, -22.8)	349.8	-109.8 (-125.7, -94.0)
2-4	283	176,596	-4.8 (-19.0, 9.0)	-2.8 (-10.5, 5.1)	320.9	-37.9 (-52.7, -23.0)
5-17	262	742,593	2.1 (-1.0, 5.4)	4.9 (-2.3, 12.6)	317.8	-55.8 (-69.9, -42.5)
18-39	783	1,506,987	9.1 (6.4, 11.9)	17.4 (12.2, 22.7)	789.7	-6.7 (-29.3, 15.9)
40-64	3521	1,842,398	70.3 (66.2, 74.4)	62.9 (59.2, 66.6)	2058.0	1463.0 (1424.4, 1500.0)
65-74	3467	507,265	262.0 (247.2, 276.7)	70.4 (66.4, 74.3)	1888.5	1578.5 (1541.4, 1615.5)
75-84	4888	309,244	551.6 (523.4, 579.7)	63.0 (59.8, 66.3)	2705.6	2182.4 (2138.8, 2226.0)
85+	3372	109,242	876.4 (808.7, 944.1)	45.9 (42.4, 49.5)	2085.4	1286.6 (1246.2, 1327.0)
All Ages	16816	5,313,600	110.3 (107.2, 113.3)	59.6 (57.9, 61.2)	9835.5	6980.5 (6895.5, 7060.2)


¹ As percentage of 2000-2006 (PrePCV7) data

² Derived from hospitalisation rate from PrePCV7 data applied to the 2012 population figures


15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Panel A




Panel B



Panel A




Panel B





Click here to access/download
Supplementary Material
Supplementary material-030516.docx





[Click here to access/download](#)

Supplementary Material

[PCV in Scotland- revised-300616-track changes.docx](#)

