



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Equipoise and the technology curve

Citation for published version:

Simpson, H, Murray, I & Duckworth, AD 2016, 'Equipoise and the technology curve: Relevance in the design of Surgical trials' Bone & Joint Research, vol. 5, no. 10, pp. 520-522. DOI: 10.1302/2046-3758.510.2000655

Digital Object Identifier (DOI):

[10.1302/2046-3758.510.2000655](https://doi.org/10.1302/2046-3758.510.2000655)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Bone & Joint Research

Publisher Rights Statement:

©2016 Simpson, Murray and duckworth. this is an open-access article distributed under the terms of the creative commons Attributions licence (cc-bY-Nc), which permits unrestricted use, distribution, and reproduction in any medium, but not for commercial gain, provided the original author and source are credited.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Equipoise and the technology curve

RELEVANCE IN THE DESIGN OF SURGICAL TRIALS

Keywords: Clinical trials, Equipoise, Technology curve, Methodology

**A. H. R. W. Simpson,
I. R. Murray,
A. D. Duckworth**

*Royal Infirmary of
Edinburgh, Edinburgh,
United Kingdom*

■ A. H. R. W. Simpson, MA(Cantab), DM (Oxon), FRCS(England & Edinburgh), George Harrison Law Professor of Orthopaedic Surgery, Royal Infirmary of Edinburgh, Editor-in-Chief, Bone & Joint Research, 22 Buckingham Street, London WC2N 6ET, UK.

■ I. R. Murray, BSc(Hons), MBChB, MSc, FRCSEd(Tr&Orth), PhD, ECAT Clinical Lecturer and Honorary Specialty Registrar (StR) Trauma and Orthopaedics, Department of Orthopaedics and Trauma

■ A. D. Duckworth, BSc(Hons), MBChB, MSc, FRCSEd(Tr&Orth), PhD, Professor of Orthopaedics and Trauma, Department of Orthopaedics and Trauma, University of Edinburgh, Royal Infirmary of Edinburgh, 51 Little France Crescent, Edinburgh, EH16 4SU, UK.

Correspondence should be sent to A. H. R. W. Simpson; email: e.vodden@boneandjoint.org.uk

doi:10.1302/2046-3758.510.2000655

Bone Joint Res 2016;5:520–522.

Elimination of bias is a core element of high quality clinical trials.¹ In addition to selection, detection, attrition, reporting and spectrum bias,² a further type of bias has been reported, known as ‘design bias’.³ This occurs before the trial is begun and is inconsistent with the principle of ‘equipoise’ and has relevance for surgical trials in a different manner to the bias described in drug trials. Equipoise is defined as the ‘even balance of weight or other forces’, or alternatively as an ‘equilibrium’. Equipoise is an important concept in clinical trials, not only from an ethical standpoint but also with regards to feasibility and recruitment on the part of both the clinicians involved and the patients who may be recruited. ‘Clinical equipoise’ exists “if there is genuine uncertainty within the expert medical community – not necessarily on the part of the individual investigator – about the preferred treatment”.^{4,5}

A clinician or researcher enrolling and/or consenting patients into a prospective randomised clinical trial must believe that the available evidence does not indicate that the new treatment or intervention being studied is either superior or inferior to the existing standard treatment.⁶ However, an individual’s equipoise can be altered by personal experience, anecdotal evidence or even by single case studies. Thus the term clinical equipoise is used to reflect the views of the wider medical community, rather than the individual clinician. In order for patients to agree to participate in a clinical trial, they need to be well informed and understand there is equipoise, otherwise this will inevitably result in inadequate recruitment.

The technology curve

When a new treatment is introduced whether it is a new drug or implant, the point of uncertainty within the medical community is

not constant and fluctuates with time from the point of introduction. This can be compared with the introduction to other new technologies, such as 3D printing. For these innovations, the uptake of these new devices has been described as the technology life cycle or ‘hype’ curve, which has five phases (Fig. 1).

The technology curve can be adapted to the uptake of medical interventions and follows the levels of evidence of research available, as well as the evolution of data for a given subject area. The first phase is the ‘technology trigger’ when a new intervention appears on the market and there is initial proof of concept data, commonly from the medical company or innovators bringing the product to market. The initial sharp rise in uptake of an intervention is caused by early adopters, followed by a growing number of clinicians using the technology based on positive data from tightly controlled case series, and explanatory/efficacy randomised trials that use strict indications for the intervention, and which are often carried out by experts in the given area. This represents the second phase known as the ‘peak of inflated expectations’.

There is an inevitable decline in use due to a combination of the expansion of indications for employing the intervention in both case series and trials, leading to an inevitable rise in associated complications and poorer outcomes reported in the literature. This results in phase 3 - the ‘trough of disillusionment’. These data are eventually drawn together in the form of systematic reviews and meta-analyses, and lead to the ‘slope of enlightenment’ (phase 4), where the indications and limitations of the intervention become clear. Phase 5 represents the ‘plateau of productivity’ when a ‘steady state’ is reached, the indications for the intervention




Fig. 1

Technology curve applied to the expectations of emerging treatments in orthopaedic surgery. The levels of evidence of research often available at each phase are outlined.




Fig. 2

Graph showing the response of differently damped systems to 'response of the system'. The solid line represents an undamped system, the dotted line a moderately damped system and the dashed line, a heavily damped system. In clinical practice, it has been suggested that the uptake of agents such as platelet rich plasma follow the solid line, whereas others such as metal-on-metal hip arthroplasties follow the dotted line, and vitamin C uptake the dashed line.

are clear, and the techniques and technology involved has been evolved and adapted as necessary. At this stage, clinical equipoise is less likely to change during the recruitment phase, and is therefore less likely to confound the results of trials comparing the new intervention with the current benchmark. A diverse spectrum of products, ranging from bone morphogenetic proteins⁷ and pulsed electromagnetic fields⁸ to meniscal transplantation⁹ are thought to be following this pattern.

When a system is perturbed by some agent (or data) it will either return to the established 'steady state' or settle at a new steady state. The path this dynamic system follows depends on how heavily 'damped' the system is (Fig. 2). In relation to medical interventions, the damping of the system (and thus the path the system follows) will depend on a variety of factors including how 'conservative' any group of clinicians and patients are, the marketing of the

product to both clinicians and patients, the simplicity of implementing the treatment, and the perceived and reported complications with which it is associated.

Emerging and established technologies and treatments in orthopaedic surgery can be considered to exist at different points along this curve. For example, emerging animal and exploratory clinical studies have raised expectations that mesenchymal stem cell-based therapies may be used to regenerate bone and cartilage, although such therapies are not yet part of mainstream treatment.¹⁰⁻¹³ An increasing number of clinical trials evaluating the use of platelet rich plasma (PRP) across a wide range of applications have not supported promising initial *in vitro* and early clinical data.¹⁴⁻¹⁶ While a number of clinicians have become disillusioned with PRP, it is possible that further analysis of emerging and published literature may reveal particular indications in which such therapies are effective. These new therapies are likely to follow the 'technology/hype' curve, whereas others have followed a more damped pattern such as metal-on-metal joint arthroplasties,^{17,18} which has been cited as an example of Scott's parabola.¹⁹ In some cases, such as the introduction on vitamin C for preventing scurvy,²⁰ the curve is even more damped (Fig. 2).

Implications for clinical trials

The technology curve for a novel product reflects the change in perceived benefit of the clinical community concerning that new treatment, i.e., it reflects the clinical equipoise of the community. During the recruitment phase of a clinical trial it is desirable that clinical equipoise remains constant. Strict inclusion and exclusion criteria are used to achieve this in 'explanatory trials', which aim to determine the efficacy of an intervention in 'ideal

conditions²¹ and are, by definition, tightly controlled clinical trials. In contrast, pragmatic trials aim to test the effectiveness between an established intervention (the current benchmark) against the new intervention in a setting most representative of day-to-day clinical practice.²¹ This often results in broader inclusion/exclusion criteria, making the trial more prone to changes in clinical equipoise.

However, even in explanatory trials, a change in equipoise may occur, for example when new interventions are being initially tested and an unexpected adverse event rate becomes apparent on interim monitoring and data analysis.²² If a 'steady state' has been reached prior to carrying out a large multi-centre pragmatic trial,²³ this will not only give equipoise, but will also inform the investigators which groups of patients should be included, thus establishing robust inclusion and exclusion criteria. If an established plateau 'steady state' has not been reached at the time of commencing a pragmatic trial, there is a risk that the intervention being assessed will not be evaluated in the clinical circumstances most relevant to routine contemporary practice.

Therefore, to take into account the effect of changing equipoise during a clinical study, it would be beneficial to monitor the uptake of the technology in a control group that is not in the trial.

This will allow changes that result from the trial to be distinguished from changes that result from the technology life cycle. In addition, appreciation of this curve allows us to introduce treatments in a manner that more rapidly reaches 'plateau'. This may be by ensuring that treatments are optimised in a pre-clinical phase, or through a more concerted effort to avoid 'widening' applications of treatments without robust rationale or preliminary data.

References

- Mundi R, Chaudhry H, Mundi S, Godin K, Bhandari M. Design and execution of clinical trials in orthopaedic surgery. *Bone Joint Res* 2014;3:161-168.
- Kleinlugtenbelt YV, Hoekstra M, Ham SJ, et al. Spectrum bias, a common unrecognised issue in orthopaedic agreement studies: do CT scans really influence the agreement on treatment plans in fractures of the distal radius? *Bone Joint Res* 2015;4:190-194.
- Fries JF, Krishnan E. Equipoise, design bias, and randomized controlled trials: the elusive ethics of new drug development. *Arthritis Res Ther* 2004;6:R250-R255.
- Freedman B. Equipoise and the ethics of clinical research. *N Engl J Med* 1987;16:317:141-145.
- Perry DC, Griffin XL, Parsons N, Costa ML. Designing clinical trials in trauma surgery: overcoming research barriers. *Bone Joint Res* 2014;3:123-129.
- Katz JN, Losina E, Lohmander LS. OARSI Clinical Trials Recommendations: design and conduct of clinical trials of surgical interventions for osteoarthritis. *Osteoarthritis Cartilage* 2015;23:798-802.
- Bodalía PN, Balaji V, Kaila R, Wilson L. Effectiveness and safety of recombinant human bone morphogenetic protein-2 for adults with lumbar spine pseudarthrosis following spinal fusion surgery: A systematic review. *Bone Joint Res* 2016;5:145-152.
- van der Jagt OP, van der Linden JC, Waarsing JH, Verhaar JA, Weinans H. Electromagnetic fields do not affect bone micro-architecture in osteoporotic rats. *Bone Joint Res* 2014;3:230-235.
- Smith NA, Achten J, Parsons N, et al. Meniscal Transplantation and its Effect on Osteoarthritis Risk: an abridged protocol for the MeTEOR study: a comprehensive cohort study incorporating a pilot randomised controlled trial. *Bone Joint Res* 2015;4:93-98.
- Murray IR, Corselli M, Petrigliano FA, Soo C, Péault B. Recent insights into the identity of mesenchymal stem cells: implications for orthopaedic applications. *Bone Joint Res* 2014;3:291-298.
- Singh A, Goel SC, Gupta KK, et al. The role of stem cells in osteoarthritis: an experimental study in rabbits. *Bone Joint Res* 2014;3:32-37.
- Hogendoorn S, Duijnisveld BJ, van Duinen SG, et al. Local injection of autologous bone marrow cells to regenerate muscle in patients with traumatic brachial plexus injury: a pilot study. *Bone Joint Res* 2014;3:38-47.
- Ismail HD, Phedy P, Kholinne E, et al. Mesenchymal stem cell implantation in atrophic nonunion of the long bones: A translational study. *Bone Joint Res* 2016;5:287-293.
- Murray IR, LaPrade RF. Platelet-rich plasma: renewed scientific understanding must guide appropriate use. *Bone Joint Res* 2016;5:92-94.
- Murray IR, LaPrade RF, Musahl V, et al. Biologic Treatments for Sports Injuries II Think Tank-Current Concepts, Future Research, and Barriers to Advancement, Part 2: rotator Cuff. *Orthop J Sports Med* 2016;4:eCollection 2016.
- LaPrade RF, Geeslin AG, Murray IR, et al. Biologic Treatments for Sports Injuries II Think Tank-Current Concepts, Future Research, and Barriers to Advancement, Part 1: Biologics Overview, Ligament Injury, Tendinopathy. *Am J Sports Med* 2016 March 29. (Epub ahead of print)
- Takamura KM, Maher P, Nath T, Su EP. Survivorship of standard versus modified posterior surgical approaches in metal-on-metal hip resurfacing. *Bone Joint Res* 2014;3:150-154.
- Berber R, Skinner J, Board T, et al; ISCCoMH. International metal-on-metal multidisciplinary teams: do we manage patients with metal-on-metal hip arthroplasty in the same way? An analysis from the International Specialist Centre Collaboration on MOM Hips (ISCCoMH). *Bone Joint Res* 2016;98-B:179-186.
- Hamilton DF, Howie C, Gaston P, Simpson AH. Scott's parabola and the rise and fall of metal-on-metal hip replacements. *BMJ* 2012;345:e8306.
- Levine M. New concepts in the biology and biochemistry of ascorbic acid. *N Engl J Med* 1986;3:314:892-902.
- Sedgwick P. Explanatory trials versus pragmatic trials. *BMJ* 2014;349:g6694.
- Hamid N, Ashraf N, Bosse MJ, et al. Radiation therapy for heterotopic ossification prophylaxis acutely after elbow trauma: a prospective randomized study. *J Bone Joint Surg [Am]* 2010;92-A:2032-2038.
- Carr AJ, Rees JL, Ramsay CR, et al. Protocol for the United Kingdom Rotator Cuff Study (UKUFF): a randomised controlled trial of open and arthroscopic rotator cuff repair. *Bone Joint Res* 2014;3:155-160.

Funding Statement

- None declared

ICMJE conflict of interest

- None declared

©2016 Simpson, Murray and Duckworth. This is an open-access article distributed under the terms of the Creative Commons Attribution licence (CC-BY-NC), which permits unrestricted use, distribution, and reproduction in any medium, but not for commercial gain, provided the original author and source are credited.