

BMJ Open Vaginal birth after caesarean section: why is uptake so low? Insights from a meta-ethnographic synthesis of women's accounts of their birth choices

Mairead Black,¹ Vikki A Entwistle,² Siladitya Bhattacharya,² Katie Gillies²

To cite: Black M, Entwistle VA, Bhattacharya S, et al. Vaginal birth after caesarean section: why is uptake so low? Insights from a meta-ethnographic synthesis of women's accounts of their birth choices. *BMJ Open* 2016;**6**:e008881. doi:10.1136/bmjopen-2015-008881

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2015-008881>).

Received 24 May 2015

Revised 1 October 2015

Accepted 12 October 2015



CrossMark

¹Division of Applied Health Sciences, University of Aberdeen, Aberdeen Maternity Hospital, Aberdeen, UK

²Division of Applied Health Sciences, University of Aberdeen, Aberdeen, UK

Correspondence to

Dr Mairead Black;
Mairead.black@abdn.ac.uk

ABSTRACT

Objective: To identify what women report influences their preferred mode of birth after caesarean section.

Design: Systematic review of qualitative literature using meta-ethnography.

Data sources: Medline, EMBASE, ASSIA, CINAHL and PsycINFO (1996 until April 2013; updated September 2015). Hand-searched journals, reference lists and abstract authors.

Study selection: Primary qualitative studies reporting women's accounts of what influenced their preferred mode of birth after caesarean section.

Data extraction and synthesis: Primary data (quotations from study participants) and authors' interpretations of these were extracted, compared and contrasted between studies, and grouped into themes to support the development of a 'line of argument' synthesis.

Results: 20 papers reporting the views of 507 women from four countries were included. Distinctive clusters of influences were identified for each of three groups of women. Women who confidently sought vaginal birth after a caesarean section were typically driven by a long-standing anticipation of vaginal birth. Women who sought a repeat caesarean section were strongly influenced by distressing previous birth experiences, and at times, by encouragement from social contacts. Women who were more open to information and professional guidance had fewer strong preconceptions and concerns, and viewed a range of considerations as potentially important.

Conclusions: Women's attitudes towards birth after caesarean section appear to be shaped by distinct clusters of influences, suggesting that opportunities exist for clinicians to stratify and personalise decision support by addressing relevant ideas, concerns and experiences from the first caesarean section birth onwards.

INTRODUCTION

Caesarean section (CS) births are described as being at epidemic levels across middle-income and high-income countries.^{1 2}

Strengths and limitations of this study

- Meta-ethnographic methods ensured sensitivity to contextual factors surrounding the influences reported by women planning birth after caesarean section.
- The contextual factors that were taken into consideration included the circumstances under which women were recruited and interviewed, and the timing of the interventions or exposures that influenced their views.
- The iterative process of reciprocal translation of study findings facilitated a higher level of understanding than previous mixed-method review methodology has allowed.
- The focus on women's perspectives is consistent with woman-centred approaches to care, but this review did not consider the views of health professionals and family.
- The identification of clustering of influences was robust to 'testing back the fit' which confirmed that primary authors' interpretations supported the synthesis 'line of argument'.

One in three babies in the USA are born by CS.¹ South American rates of CS exceed 50% in many areas, with over 70% of births in private healthcare settings being by CS.^{3 4} Concern to reduce overall rates of CS is in tension with efforts to promote patient choice, as women themselves often request this mode of birth.⁵

The greatest contribution to current high rates of CS comes from repeat CS procedures.⁶ Worldwide rates of vaginal birth after CS (VBAC) have dropped dramatically in recent years. Between 1999 and 2002, US VBAC attempts fell from 48.3% in 2000 to 30.7% in 2002, with 73.4% of VBAC attempts being successful.⁷ The UK saw actual VBAC rates fall from 45.9% in 1988 to 36% between 2004 and 2011.^{8 9} Health service support for VBAC diminished after retrospective data published in 1996 favoured the

maternal safety profile of repeat CS.¹⁰ Although more evidence for the relative safety of VBAC has emerged in recent years,¹¹ and efforts have been made to increase VBAC attempts, rates have never fully recovered.^{12 13}

Enthusiasm to reduce rates of CS stems from policy concerns about the relatively high financial costs and the greater maternal morbidity and mortality of CS when compared with vaginal birth.¹⁴ It can also be linked to broader concerns about unnecessary medical intervention (too much medicine).¹⁵ However, the costs and harms that are evident when CS is considered at a population level are much less apparent at the level of individual women. Absolute rates of serious morbidity from CS are low,^{2 16} and there is little evidence that women themselves regret CS when they have requested this mode of birth.¹⁷ At the same time, potential benefits of CS can often be identified for (and by) individual women.¹⁸ Population data suggest that an increase in rates of CS does not contribute to parallel improvements in neonatal outcomes.¹⁹

Broad policy consensus in high-income countries supports offering women who become pregnant after CS a choice between repeat CS and attempting VBAC, unless clinical circumstances or available services preclude this (eg, when a high risk of CS scar rupture contraindicates VBAC).^{12 16 20} UK guidance outlines which risks (including probabilities) should be discussed by women and health professionals before agreeing on the planned mode of birth by 36 weeks gestation.²⁰ Although probabilistic information about the physical health outcomes of VBAC and repeat CS might seem to support VBAC, the introduction of decision support interventions in the latter part of pregnancy after CS has made little difference to women's choices.^{21 22} There are several plausible explanations for this, including the likelihood that decision-making is influenced by a much broader range of cultural values and social and emotional considerations than are addressed through existing decision support. It is known, for example, that some women have a strong desire to experience vaginal birth,^{23 24} and that some fear dissatisfaction if they choose VBAC but their attempt fails.²⁵⁻²⁷ However, the insights that have emerged from studies, to date, have been somewhat fragmented. A more comprehensive and nuanced understanding of the complex range of influences on women's decisions is needed to support informed ethical judgements about efforts either to reduce rates of CS or to support women's decision-making. Development of public health policy and clinical practice would benefit from as robust as possible an understanding of the diverse perspectives that women bring to decisions about mode of birth following a previous caesarean, as would debate about what range of options, information, advice and decision support could be appropriately provided by health services. To address this need, we aimed to identify, contextualise and synthesise an understanding of the reasons why women prefer VBAC or elective repeat CS (ERCS).

METHODS

A systematic literature search and meta-ethnography was conducted. The seven steps of meta-ethnography described by Noblit and Hare, as listed in [box 1](#), were followed to synthesise the available primary research studies.²⁸

A systematic search was conducted using Medline, EMBASE, ASSIA, CINAHL and PsycINFO in April 2013 (updated in September 2015) using multiple subject headings and free text key words relating to modes of birth and exploration of women's preferences or choices (the full search strategy for Medline is provided as online supplementary appendix 1, and further search strategies are available from the authors on request). Inclusion and exclusion criteria are outlined in [table 1](#).

Titles, abstracts and, where necessary, full papers were screened for potential eligibility. Inclusion and exclusion criteria were applied to full papers. Authors were contacted when only abstracts were published and studies appeared relevant. Three journals containing the greatest number of relevant studies in the 2013 search (*British Journal of Midwifery*, *International Journal of Nursing Practice*, and *BJOG: an International Journal of Obstetrics and Gynaecology*) were hand searched to identify any further relevant papers. High-quality translation of two abstracts and one full article was obtained. Quality assessment was performed using the Critical Appraisal Skills Programme checklist for qualitative studies²⁹ to prompt reflection on study quality, but studies were not excluded on the basis of quality if they contained some qualitative data of value to our research question.

The key characteristics of included studies were extracted and summarised (see [table 2](#)). The studies were initially read individually, in chronological order, and relevant points from the primary data (first-order constructs) and the study authors' descriptions and interpretations (second-order constructs) were extracted. First-order constructs were obtained from quotations from women reported in the 'results' section of each study, while second-order constructs (primary authors' account and interpretation of their findings) were obtained from 'results' and 'discussion' sections. All first and second-order constructs were tabulated in the form of primary quotes, or exact author interpretations, to support the identification of key themes.

Box 1 Meta-ethnography steps as described by Noblit and Hare²⁸

1. Identify the research question
2. Identify relevant studies
3. Read the studies
4. Identify themes
5. Translate the findings of each study into those of the others
6. Synthesise the findings
7. Express the synthesis

Table 1 Inclusion and exclusion criteria

Inclusion criteria	
Study population	Comprised or included an identifiable subgroup of women who have had at least one previous caesarean section
Study design	Primary research that included and clearly reported a qualitative element
Study findings	<ul style="list-style-type: none"> ▶ Included accounts of influences on preferred mode of birth after a previous caesarean section, from the women's perspectives ▶ Primary data provided relevant to the research question and target population of this synthesis
Language	Any; no language restrictions applied
Exclusion criteria	
Date of publication	Studies published before 1996.

Searching was conducted by one author (MB), with input from an information specialist. Screening and identification of studies, followed by coding of constructs were conducted by two authors (one clinical (MB), one non-clinical (KG)) independently, with regular meetings to establish agreement. During these meetings, provisional third-order constructs (our interpretation of both primary authors' interpretations and primary data) and key themes were identified. The third and fourth authors (VAE and SB) were involved in further development of these themes, having each reviewed a different sample of included studies.

The key interpretive aspect, step five of Noblit and Hare's approach, involved one author comparing and contrasting the constructs and themes that featured in the different studies in an iterative manner. The findings of each study were interpreted in light of each of the other relevant studies in turn. This allowed for detailed consideration of how study design and context could have shaped study findings (eg, which women were included and when they were interviewed in relation to their original CS and/or subsequent birth). During this process, third-order constructs were confirmed, and a line-of-argument synthesis developed. All four authors contributed to the development of the line of argument.

The potential for the clinical background of two authors (MB and SB) in particular to influence the findings was recognised from the outset. All team members' interpretations and preconceptions were continually challenged and used in a constructive manner during discussions throughout the synthesis process to ensure that all reported perspectives were fairly considered, and that the line of argument developed was robust.

Following the updated search in September 2015, additional eligible papers were identified. Relevant findings were used to test the fit of the line of argument. This involved identification of first and second-order constructs (primary data and authors' interpretations,

respectively) in the additional papers, and analysing these for relevant themes of influence on birth preferences after CS. These themes were compared and contrasted with the content of the line of argument to assess the extent to which they appeared to 'fit' together or 'conflict' with one another.

RESULTS

The search results are outlined in [figure 1](#). Of 2391 citations obtained in the original search, 1174 duplicates were excluded. Screening of 1217 titles and/or abstracts resulted in a further 1092 exclusions for lack of relevance; 71 full papers and two sets of conference proceedings were obtained, and attempts made to contact four authors, of which two were unsuccessful. A total of 57 titles lacked relevant primary data or were published before 1996 and were excluded. Twenty papers reporting from 15 primary studies were included following resolution of disagreement over eligibility of two papers.

The focus and key study characteristics for the 20 included papers are outlined in [table 2](#).

The identified studies were conducted in four countries (UK, USA, China and Australia) and each included between 4 and 170 women, with findings from 507 women in total reported across the papers. Six papers reported on women who planned VBAC, four reported on women who planned ERCS, nine reported on both, and one reported on women who planned ERCS but would have desired VBAC in other circumstances.

Quality assessment of the papers is presented in online supplementary appendix 2. All papers had a clear statement of study aim which deemed qualitative methods to be appropriate. Common quality concerns included lack of information on: justification for the theoretical approach; lack of information about women who declined to take part; the interview guide used; and data saturation. Only one paper included a discussion of the potential for the researcher's role to influence the study's findings, although two further papers described involvement of a multidisciplinary team to perform the data analysis, mitigating the risk of dominance of a single interpretive perspective.

Our initial grouping of first and second-order constructs resulted in 40 subthemes. These were then categorised into six key themes which characterised the main kinds of consideration and features of decision-making processes that appeared to influence preferences for mode of birth. These themes were: long-standing anticipation of vaginal birth; responses to previous birth experiences (positive and/or negative); encouragement or dissuasion from influential people for either birth mode; fear or reassurance from risk-related information on VBAC; perceived net benefit or harm of birth options; and extent and nature of involvement in decision-making. As the labels suggest, several of these themes accommodate a spectrum of views or experiences.

Table 2 Characteristics of included studies

ID number	Author	Year	Country	Study aim	Data collection method	Planned birth method at time of study	Participants (n)	Timing of interview
M1	Ridley ³⁰	2002	USA	Discover what influences women in the decision to deliver via VBAC	Interview (FTF)	VBAC	5	Postnatal (2–4/12)
M2	York ³¹	2005	UK	Describe childbirth expectations, influences and knowledge in women who had experienced emergency CS and planned subsequent CS	Interview (FTF)	CS	10	Antenatal (third trimester)
M3	Liu ²³	2006	China	Investigate the decision factors involved and experience of women who had successful VBAC	Interview (FTF), researcher diary, field notes	VBAC	10	Postnatal (1–2/7)
M4*	Fenwick ¹⁸	2006	Australia	Describe childbirth expectations, influences and knowledge in women who had experienced emergency CS and planned subsequent CS	Interview (T), field notes	CS	49	Pre-pregnancy, antenatal and postnatal (no limits)
M5	Emmett ³²	2006	UK	Explore women's experience of decision-making regarding mode of delivery after having a previous CS	Interview (FTF)	VBAC and CS	21	Postnatal (2–8/12)
M6	Cheung ³³	2006	China	Understand Chinese women's perceptions and interpretations of their own CS decision-making, and to investigate how their negotiation with healthcare professionals may be improved	Interview (FTF), field notes	CS	52	Postnatal (1/52 or 8/12)
M7	Meddings ³⁴	2006	UK	Examine the lived experience of women who elected to attempt a vaginal birth following a previous CS delivery	Interview (FTF) *2	VBAC	8	Antenatal (>34/40) and postnatal (~6/52)
M8	Moffat ³⁵	2007	UK	Prospectively explore women's decision-making regarding mode of delivery after a previous CS	Consultation observation, patient diaries, interview (FTF)	VBAC and CS	26	Antenatal (from 20/40) and postnatal (6/52)
M9*	Fenwick ³⁶	2007	Australia	Explore childbirth expectations and knowledge of women who had experienced a CS and would prefer a vaginal birth in a subsequent pregnancy	Interview (T)	VBAC	35	Pre-pregnancy, Antenatal and Postnatal (no limits)
M10	Farnworth ³⁷	2007	UK	Identify and describe factors which influence women making a choice regarding mode of delivery after a previous CS delivery in a UK setting, and to identify the role of the obstetrician in this process	Interview (FTF)	VBAC and CS	10	Antenatal (36/40)
M11	Cox ³⁸	2007	UK	Explore issues around the choices between VBAC and elective CS based on the nature and extent of the information women actually received when making a decision between elective CS and VBAC, the sources of that	Interview (type not clear)	VBAC and CS	7	Postnatal (timing not clear)

Continued

Table 2 Continued

ID number	Author	Year	Country	Study aim	Data collection method	Planned birth method at time of study	Participants (n)	Timing of interview
M12	Farnworth ³⁹	2008	UK	information, and its importance in terms of the influence it had on their decision Examine the impact of a decision support intervention designed for women choosing mode of delivery after one previous CS	Interview (FTF)	VBAC and CS	18	Antenatal (37/40)
M13†	McGrath ⁴⁰	2009 (a)	Australia	Explore, from the mother's perspective, the decision-making experience with regard to subsequent birth choice for women who had delivered previously by CS	Interview (FTF)	CS	16	Postnatal (6/52)
M14†	McGrath ⁴¹	2009 (b)	Australia	Describe the perspective of mothers who underwent elective CS on risks associated with the delivery modes of VBAC and elective CS, and their experience discussing such risks with their health professionals	Interview (FTF)	CS	16	Postnatal (6/52)
M15	Goodall ⁴²	2009	UK	Explore women's perceptions of the role of health professionals in their decision regarding mode of delivery, following previous delivery by CS	Interview (FTF)	VBAC and CS	8	Antenatal (20–40/40)
M16	Frost ⁴³	2009	UK	Obtain the views of women on their experiences of decision-making about the method of delivery following a previous CS, and the role of decision aids in this process	Interview (FTF)	VBAC and CS	30	Antenatal (37/40), postnatal (6–8/52)
M17†	Phillips ²⁴	2009	Australia	Explore, from a phenomenological perspective, the reasons motivating women to try for or achieve VBAC	Interview (FTF)	VBAC	4	Postnatal (6/52)
M18†	McGrath ⁴⁴	2010 (a)	Australia	Explore, from the mothers' perspective, the process of decision-making about mode of delivery for a subsequent birth after a previous CS	Interview (FTF)	VBAC	4	Postnatal (6/52)
M19	David ⁴⁵ Originates from same study as	2010	Australia	Provide maternity healthcare providers with an increased understanding of, and insight into, the different information needs of this specific group of maternity care consumers.	Telephone log and field notes	VBAC	170	Antenatal (various gestations)
M20†	McGrath ⁴⁶	2010 (b)	Australia	To focus on findings which recorded the frustration of women who valued a vaginal delivery but who delivered by CS	Interview (FTF)	CS	8	Postnatal (6/52)

*Originates from same study (M4 and M9).

†Originates from same study (M13, M14, M17, M18 and M20).

CS, caesarean section; FTF, face-to-face; M, manuscript; T, telephone; VBAC, vaginal birth after CS.




Figure 1 Flow diagram of search results' caesarean section. CS, caesarean section.

Key themes

The six key themes identified as shaping birth preferences after CS are illustrated with example data in [table 3](#). Primary study participant quotes illustrating first-order constructs are displayed in bold text, and primary author interpretations illustrating second-order constructs are presented in italics.

Patterns of influence: a line of argument

We noted that some kinds of views and experiences (specific instances of the six key themes) tended to cluster together in support of the main birth preferences. These clusterings are discussed in the context of the line of argument we developed using the process of meta-ethnography to synthesise knowledge of influences on women's birth preferences after CS.

Women approaching a birth after a CS generally have either a clear preference for VBAC or ERCS, or a relatively open mind to either option. Although some studies by design included women from only one or two of these categories, looking across the studies, we were able to develop a line of argument to explain how their findings were related. In summary, the line of argument is that three distinctive clusters of influences support the three attitudinal positions that women adopt towards mode of birth after CS.

The three positions and the distinctive influences on these are summarised in [figure 2](#) and described below. We note that the influences could be operative from different times, and that some were significant before and around the first CS.

Preferences for vaginal birth

Preferences for vaginal birth could be shaped by influences acting over a period of time, which for some

women reached several years, and for many was linked to key events or periods of their lives. With respect to women's long-standing anticipation of vaginal birth, some women had a personal ambition to achieve vaginal birth that predated their first pregnancy and drove them to pursue VBAC (M17 and M3 (subject ID numbers)). This could act synergistically with negative responses to a previous birth experience. For example, unpleasant memories of the initial CS experience, particularly where women had felt a loss of control over that birth, led some women to view VBAC as a potentially life-enriching experience that met their ambitions and avoided further negative emotions (M1, M19, M9, M3, M8 and M14). This impression was often enhanced by interpregnancy social interaction with influential others, including women who provided encouragement by sharing accounts of their own positive VBAC experiences (M19). For some, the probability of successful VBAC was pivotal (M1 and M3).

Future considerations could also play an important role in the shaping of preferences for VBAC, as women considered implications beyond the birth itself when evaluating their expected net gain from VBAC. Several women believed that VBAC offered physiological benefits to physical and emotional health of themselves and their offspring, with particular emphasis on the facilitation of bonding and breastfeeding (M17 and M3). This was a particularly dominant issue among women who experienced breastfeeding difficulties after a previous planned CS, especially in those who had successfully breast fed their babies born vaginally in prior pregnancies (M3). The social benefits of being able to return to usual family roles and resume driving as soon as possible in the postnatal period were also cited as reasons for preferring to avoid CS particularly within UK study settings (M7, M8 and M9).

Further, influential people included health professionals who provided support, advice or encouragement in favour of VBAC. Women's perception of the extent to which they themselves should make the decision regarding planned mode of birth was important. Although some women, particularly in the UK and Australia, were confident about their right to decide how to plan the birth (M18, M7, M17 and M1), others judged any personal reasons they had in favour of ERCS to be unimportant or unjustified when considered in light of medical advice in favour of VBAC (M8).

Preferences for ERCS

Response to the previous birth experience was the central theme among women who demonstrated a clear preference to have an ERCS. A previous emergency CS in labour appeared to lead many women to believe their bodies were incapable of vaginal birth (M8, M10 and M13). Some women sought an ERCS to actively avoid any possibility of a repeat emergency CS (M8, M10 and M13), while others feared the possibility of a recurrence of the factors which led to the previous CS. Others

Table 3 Key themes of influence on birth preferences after CS, with corresponding example data

Theme	Exemplary quote
Long-standing anticipation of vaginal birth	'Right from the start I wanted a natural delivery. All the women in my family just gave birth naturally and so I was very disappointed when it didn't work out that way for the first baby' (M17) <i>'Despite their CS they still considered women's bodies were 'designed' to give birth vaginally' (M9)'Some of the study cases believed, due to their own notions, that there was only one way to feel like a real mother, ie. experiencing vaginal birth and the delivery pain in person. This was why they chose VBAC' (M3)</i>
Responses to previous birth experience (positive and/or negative)	'If my body can't do it [vaginal birth], why put myself and bub [baby] through all the stress and heartache' (M13) <i>'Many of these women also expressed that the CS experience had made them feel powerless and helpless; 'taking away total control'(M9) 'In the end we said, look, we're going to go with what we know. What we did first time worked out okay' (M13)</i>
Encouragement or dissuasion from influential people for either birth mode	'they [doctors] said you can try normally, but they didn't seem very positive that it would work and I think they preferred me to have a caesarean'. (M11) <i>'Horror stories' and the knowledge and/or personal experience of friends also worked to reinforce their emerging view that CS was the safest birthing option' (M4) <i>'..other sources of information were noted as mothers groups and/or playgroups.[where] .. sharing of knowledge 'inspired' them' to pursue VBAC (M19)</i></i>
Fear or reassurance from risk-related information on VBAC	'I like to gather as much information as I can about things and then make my own decisions from that' (M17) <i>'A persistent theme appeared to be the lack of both local written information and professional opinion...this led the women to base their knowledge on a mixture of media, professional and personal sources' (M2) <i>'Some women described feeling very sure about their preferred mode of delivery from the beginning of pregnancy and those women generally needed little in the way of decisional support' (M8). 'Information and support gave women confidence in their decision, and ultimately, the power to own and justify the decision that they had made' (M12) 'Oh yeah, the riskiest approach was to try a vaginal delivery. Yeah, no I wouldn't even have attempted it. And everything I read backed that up, yes.' (M14) 'supposed to have all that stuff squeezed out and that's not done in a CS but it's probably less risky for the baby' (M4) 'About the biggest thing for me was the success rate.. . . There was more positive than negative.. . . 80% of the women who tried it were able to do it'. (M1) <i>'When deciding whether to accept the VBAC or not, in most cases patients would first evaluate the advantages and disadvantages which included the recovery time after delivery, time of hospitalisation, potential harms to the mother and baby.' (M3) <i>'women...considered CS a physical, emotional and lifestyle disruption that was risky and had the potential to cause harm to both mother and baby; separated them from their baby; and interrupted the postnatal period' (M9)</i></i></i></i>
Extent and nature of involvement in decision-making	'I was basically told they would prefer for me to try vaginal delivery but I could have a section if I really wanted' (M8) 'I feel every time I go and see the doctor or the midwife they keep talking about elective Caesareans...they keep finding reasons why I'll probably need an elective Caesarean so yeah it feels like choice is lot more limited this time' (M15) <i>'The important point is that the mothers who tried for a VBAC were clear and focused in their determination to own the decision-making process' (M1)</i>

Primary study participant quotes are displayed in bold text and primary author interpretations are presented in italics. CS, caesarean section; VBAC, vaginal birth after CS.

opted for ERCS on the grounds that it was a familiar and positive birth experience (M19, M5 and M6).

The previous birth and its outcome could also shape women's perceptions of the safety of VBAC (as outlined, it could lead to an assessment of net harm from planning VBAC), moderate the influence of social contacts (favouring those who encouraged ERCS and/or discouraged from planning VBAC) and limit the degree to

which they felt they had a choice to make in the subsequent pregnancy (role in decision-making).

Safety concerns were described as particularly influential among some women in Australia who wished to avoid VBAC due to fear of the uterine scar 'splitting', or 'rupturing' during labour. This feeling dominated their preference for ERCS despite awareness of neonatal breathing problems being more common following this

Sought Vaginal birth	Sought repeat CS	Open minded
<ul style="list-style-type: none"> • Longstanding anticipation of vaginal birth • Responses to previous birth experiences (negative) • Perceived net benefit or harm of birth options (postnatal issues predominating) • Encouragement and dissuasion from influential people [for VBAC and CS respectively] • Extent and nature of involvement in decision-making 	<ul style="list-style-type: none"> • Responses to previous birth experiences (negative and positive) • Fear or reassurance from risk-related information on VBAC • Encouragement and dissuasion from influential people [for CS and VBAC respectively] • Extent and nature of involvement in decision-making • Perceived net benefit or harm of birth options 	<ul style="list-style-type: none"> • Encouragement and dissuasion from influential people for either birth mode • Fear or reassurance from risk-related information on VBAC • Perceived net benefit or harm of birth options (mix of health and social considerations predominating) • Extent and nature of involvement in decision-making

Figure 2 Summary attitudinal positions of women early in the pregnancy after CS and clusters of key influences acting on their eventual birth preferences. CS, caesarean section; VBAC, vaginal birth after caesarean.

mode of birth (M4). Some women with a strong preference for VBAC had been influenced, sometimes powerfully, by family, friends and health professionals who recommended ERCS as a safer and more predictable mode of birth than VBAC (M13 and M4).

Ownership of choice, or lack of the same, appeared crucial in determining whether or not some women opted for ERCS. Many women perceived that their health professionals would prefer this option, and as such, that VBAC was not available to them (M15). Others choosing ERCS felt happy to exercise their preference as they had been positively encouraged to opt for the mode of birth that felt right for them (M5).

Open-minded approach

Women who did not have a firm preference for either VBAC or ERCS appeared to be less strongly influenced by prior expectations about childbirth or by their previous birth experience than those who were more committed to one particular mode of birth. Influential others were apparently key to the decisions made in this context. These women valued and often actively sought the opinion of health professionals during their pregnancy, processed information on the options available and put considerable effort into weighing up the attributes of the birth options available to assess net benefit. An exception to this involved women who felt overwhelmed by the decision-making responsibility, and preferred to follow health professionals' advice (M19, M8 and M18). Obstetricians, and, at times, midwives, appeared to have particular influence over women who

were open to considering either mode of birth, even when women were not actively advised as to how to deliver, but perceived subtle signals that their health professional had a preference (M11). Some women said their choice should be based on information alone, rather than the input or opinions of others, recognising that other people are not necessarily impartial (M17).

Robustness of findings

On 'testing back the fit' of our line of argument, we found that the clusters of influence we identified were consistent with the findings of each of the individual included studies, but that none of these studies included a broad enough mix of participants to have enabled the development of this level of understanding in isolation.

Further 'testing' of the line of argument was made possible by the publication of the three new studies identified in the update of the search conducted in 2015 which are summarised in [table 4](#). Shorten *et al* analysed written text in which women explained their reasons for choosing either mode of birth after CS. They highlighted the significance of previous birth experience, safety concerns and speed of recovery along with health professionals' preferences in shaping eventual decisions. Although they did not describe a clear distinction between the attitudinal groups, their findings were broadly supportive of the conclusions of this synthesis, with no evidence of confliction or contradiction.⁴⁷ Kennedy *et al*⁴⁸ performed an institutional ethnography exploring the complexity of choice around elective CS. This included interviews with women within the English

Table 4 Studies identified in the updated search which were used to 'test the fit' of the line of argument

Author	Year	Country	Study aim	Data collection method	Planned birth method at time of study	Participants (n)	Timing of data collection
Shorten ⁴⁷	2014	Australia	explore values and expectations that guide women during decision-making about the next birth after caesarean, and identify factors that influence consistency between women's choices and actual birth experiences	Written surveys and narrative accounts	VBAC and CS	187	36–37 weeks' gestation and postnatal
Kennedy ⁴⁸	2013	UK	To explore the complexities of women's and clinicians' choices around elective caesarean delivery	Interview (FTF) and consultation observations.	CS and vaginal birth (sample not restricted to birth after CS)	27 women, of whom three had VBAC and 19 had no history of prior CS. Previous obstetric history of 5 participants who underwent CS was not clear	Not specified; appears to span antenatal and postnatal period
Tully ⁴⁹	2013	UK	To document the circumstances in which caesarean section was deemed to be appropriate in one UK hospital through the eyes of the women and their partners experiencing the operative delivery of their infant	Interview (FTF)	Not applicable (postnatal)	115 women	Postnatal hospital stay

CS, caesarean section; FTF, face-to-face; VBAC, vaginal birth after CS.

National Health Service provider settings. The authors identified that women planning birth after CS negotiated with clinicians to reach a 'comfortable compromise' which facilitated a plan for VBAC that included adequate assurance of early recourse to CS if labour progress was suboptimal. This supports our findings of the crucial role of health professionals in influencing VBAC decisions by providing support for this option. Further author interpretation echoed our emphasis on the importance of predicted VBAC success in influencing women to aim for this mode of birth. Finally, the authors highlighted the desire for information among some women, providing an exemplary quote which supported our impression that women with an open mind to mode of birth after CS place great emphasis on the content, and in this case, quality of information accessed:

When I was getting told about the 0.3% chance of a scar rupturing, you know, when I was asking people about how that statistic was arrived at no one could tell me, so I kept digging for more and more information, 'and there's just not enough research, there's not enough studies that have been done, the women aren't in the same circumstances, they're not all in even one country, it's international, it's in under-developed countries, so you're pulling together these statistics from a complete diverse set of sample set, and how can you make judgements on what an individual's circumstances are going to be based on that? There's just not enough there's not enough information out there to be able to say you're going to be one of those statistics. (P108; woman pondering VBAC decision)

Tully and Ball⁴⁹ presented findings of an interview study of 115 mothers recently delivered by CS over a 3-year period in England. Although minimal primary or secondary constructs related to birth after CS were presented, there was evidence that predicted VBAC success was important to women aiming for a vaginal birth, and that a negative previous birth experience drove women to seek control and predictability in the form of an ERCS. These observations are consistent with our findings, and no evidence of contradictory interpretations was identified.

DISCUSSION

Summary of main findings

This study sought to answer the research question 'What influences women's preferred mode of birth after previous caesarean section?' We have identified distinct clusters of influences that tend to underpin the three main positions that pregnant women adopt towards modes of birth. After an initial CS, women tend to approach childbirth with one of three broad attitudinal positions meaning that they: (1) seek vaginal birth (2) seek repeat caesarean or (3) are open minded to consideration of either mode of birth. These positions reflect thought processes which are likely to evolve from at least as early



as the primary CS, with some influential cultural norms in operation well before that time. A strong preference for VBAC appears to be driven by a belief that vaginal birth is 'normal' and has some intrinsic value. This belief is often accompanied by a keen desire to resume a normal life soon after vaginal birth. By contrast, a clear preference for ERCS from early in pregnancy can be driven by a previous negative experience of attempting but failing to achieve vaginal birth, and a positive emphasis on the predictability of ERCS. Finally, there are women who embark on their next pregnancy undecided about mode of birth. These women are more open to external influence: they appreciate the benefits of written information and personalised expert advice which they use to weigh up what they see as the advantages and disadvantages of their options. The recognition of these clusters of influences, according to attitude towards birth from early in the pregnancy after CS, is a novel finding made possible by looking across the range of relevant studies. Historical and contemporary studies, have highlighted influences on birth preferences after CS which resonate with those identified in this synthesis, but without identification of attitudinal groups or attention to the multiple influences and the ways these may vary over time.^{27 50 51} The importance of timing of influence has, however, been highlighted recently by prospective work which found that first-trimester preferences for either ERCS or VBAC persist by early in the third trimester in over 70% of women.⁵²

Benefits of a meta-ethnographic approach

Meta-ethnography enabled an interpretation of the available research that incorporated a sensitivity to the contextual factors surrounding the influences reported by specific groups of women planning birth after CS.⁵³ Contextual factors considered included key time points at which influences took hold, fundamental study characteristics (setting; eligibility criteria; recruitment processes; timing of interviews; healthcare systems) and factors unique to individual women. These contextual considerations limit the likelihood that findings would be generalised inappropriately. The iterative process of reciprocal translation used to build on emergent themes facilitated a higher level of understanding than previous mixed-method review methodology has allowed, particularly that of quantitative work, where presence or absence of potential influences has been the focus.²⁵ The clustering of influences identified within specific attitudinal groups provided clinically relevant insight into the nature of women's decision-making behaviour. In addition, the identification of clustering was considered robust in light of the 'testing back the fit', which confirmed that primary authors' interpretations supported specific attitudinal clusters.

Women's perspectives

The specific focus on *women's* perspectives on what influences birth preferences after CS complements the

current focus on joint healthcare decision-making in which informed patients contribute to decisions which reflect their beliefs and preferences.⁵⁴ This, therefore, provides insight which has maximal clinical application in settings where every effort should be made to ensure decisions about mode of birth after CS incorporate women's values and preferences. Given that health professionals have a variable level of input into shaping the eventual mode of birth, it is possible that consideration of health professionals' perspectives may have further developed our understanding of the decision-making process.⁵⁵ However, women's insights were considered central to achieving the goal of informing future efforts to optimise and support woman-centred planning of birth after CS.

Clinical and research implications

Reflection on current practice

The strength of evidence supporting the first CS birth experience as a key influence on future birth preferences demands immediate attention. Women should be effectively supported in dealing with the unexpected and potentially traumatic nature of a primary CS. Efforts to promptly address any inaccurate perceptions of their CS birth events, and to provide personally specific information about the risks and benefits of future birth options could be made following the first CS, and be reiterated early in the pregnancy after CS. The findings of this synthesis suggest that women's concerns about serious maternal or offspring health risks (beyond those of CS scar rupture) are not important influences on their birth choices after CS. This is of particular interest because information currently provided by health professionals for women planning birth after CS focuses largely on these risks and clinical health considerations.²⁰ Recognition of this mismatch between what women and health professionals prioritise should prompt health professionals to engage in discussion with women which allows identification of their main concerns and places sufficient emphasis on the psychological and social, as well as the physical health consequences of modes of birth after CS. The heterogeneity of influences on birth choices after CS demonstrated in this synthesis highlight why approaching all women planning birth after CS with, for example, the same decision support tool in the latter part of pregnancy, is unlikely to alter their prior attitudinal positions.

Implications for future research and practice

Recognition of the diverse range of influences on, and attitudes towards birth after CS enables us to understand why decision support interventions have had limited effects on ERCS so far,^{21 22} and opens up the possibility of a more targeted approach. We suggest that future interventions should aim to promote positive experiences of informed and shared decision-making, while minimising maternal and fetal morbidity, and avoiding

unnecessary healthcare costs. Insights from this synthesis suggest that future strategies should ensure early consideration of women's concerns and preferences, and their likelihood of achieving good physical birth outcomes. Women may be broadly categorised in early pregnancy after CS as being in favour of either VBAC or ERCS, or being open to either option. At the same time, their prognosis for successful VBAC may also be assessed based on factors such as their age, body mass index and indication for previous CS.^{8 56} To support high-quality decision-making and increase VBAC success rates, efforts could be made to ensure design of decision support which reflects women's prognosis for VBAC success and is sensitive to any early preferences regarding mode of birth after CS. The six main prognosis/preference categories are represented in [figure 3](#).

Decision support for women may be delivered via conversations with health professionals, advice and information, including decision aids.⁵⁷ Decision aids provide women with information about options relevant to their health status, while helping them to reflect and draw on their personal values. Previous research has demonstrated that use of some such tools in supporting birth choices after CS improved decision satisfaction but had minimal impact on VBAC rates.²² The lack of success in increasing VBAC rates may reflect that the tools that were tested were not tailored to women's early attitudes towards each birth mode, but instead delivered advice according to outcomes which women prioritised. Faced with a choice of surgery and less invasive options, decision aids have been shown to lead patients to choose conservative or less invasive treatments.⁵⁸

In the context of planning birth after CS, decision aids might usefully be stratified according to predicted VBAC success and also be responsive to individual women's early preferences and priorities of mode of birth. It is likely to be particularly important to engage women who are open minded (groups E and F on [figure 3](#)), and women with a VBAC prognosis which is at odds with their preferred mode of birth (groups B and C in [figure 3](#)) by the second trimester, in conversations with health professionals, to ensure sufficient time to explore their views and discuss and allow them to consider their options. In such situations, a 'consider a recommendation' approach may be warranted, explaining why either ERCS or VBAC is recommended, but leaving sufficient scope and ensuring sufficient support for

women to assess and discuss the recommendation before making their own mind up about it.⁵⁹ In those pursuing VBAC despite a poor prognosis for success, there could be a discussion about criteria for conversion to CS, and adequate counselling in preparation for the possible psychological impact of such an outcome. Those in whom VBAC prognosis is in keeping with their preferred mode of birth (groups A and D in [figure 3](#)) might need less in the way of information, conversation and recommendations from health professionals, but their needs for information and reassurance about their decisions should not be neglected: balanced written information regarding the risks and benefits of both birth options, and clarification/confirmation of ongoing preferences are still likely to be important. As events unfold during subsequent pregnancies, ongoing communication and decision support for all women would need to be tailored to accommodate new clinical information, concerns and preferences, but a broad pathway identified following the first CS would ensure timely and relevant intervention to address modifiable influences.

CONCLUSIONS

Forming a preference for repeat CS or VBAC is a dynamic process shaped by many influences which appear to cluster distinctively in the development of strongly held positions. Long-standing expectations of childbirth and perceptions of previous birth experiences appear particularly influential on VBAC and ERCS preferences, respectively. This suggests that early communication to discuss women's prospects for VBAC success and explore and discuss their attitudes towards future births may be valuable, and could perhaps start from as early as the first CS. This might help increase the proportion of women who approach birth after CS with an open mind, being receptive to written information, and the advice of health professionals. Our synthesis has highlighted why current care models involving provision of information in pregnancy after CS may not lead to the birth choices which could help reduce the unnecessary rate of CS. It suggests a need to address women's social and psychological concerns, and not just the currently recommended information, both to support women's autonomy in decision-making, and to address public health concerns about rising rates of clinically unnecessary CS.

Twitter Follow Mairead Black at @maireadblack and Katie Gillies at @GilliesKatie

Contributors MB and SB conceived the idea of the study. MB, SB and KG planned the study. MB and KG conducted the literature search and analysed all data. VAE and SB contributed to data analysis and interpretation. MB wrote all drafts of the manuscript, and is the guarantor. VAE, SB and KG contributed to all drafts of the manuscript.

Funding MB is a research training fellow funded by The Wellcome Trust.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

	Good VBAC prognosis	Poor VBAC prognosis
Prefers VBAC	A	B
Prefers ERCS	C	D
Open minded	E	F

Figure 3 Table represents how women may be categorised according to their preferred mode of birth in early pregnancy and their prognosis for VBAC success' VBAC, vaginal birth after caesarean; ERCS, elective repeat caesarean section.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- World Health Organisation. World health statistics. 2013. http://www.who.int/gho/publications/world_health_statistics/2013/en/
- Stavrou EP, Ford JB, Shand AW, *et al*. Epidemiology and trends for caesarean section births in New South Wales, Australia: a population-based study. *BMC Pregnancy Childbirth* 2011;11:8.
- Lumbiganon P, Laopaiboon M, Gülmezoglu AM, *et al*. World Health Organization Global Survey on Maternal and Perinatal Health Research Group. Method of delivery and pregnancy outcomes in Asia: the WHO global survey on maternal and perinatal health 2007–08. *Lancet* 2010;375:490–9.
- Almeida S, Bettiol H, Barbieri MA, *et al*. Significant differences in caesarean section rates between a private and a public hospital in Brazil. *Cad Saude Publica* 2008;24:2909–18.
- Minkoff H, Powderly KR, Chervenak F, *et al*. Ethical dimensions of elective primary caesarean delivery. *Obstet Gynecol* 2004;103:387–92.
- Timofeev J, Reddy UM, Huang CC, *et al*. Obstetric complications, neonatal morbidity, and indications for caesarean delivery by maternal age. *Obstet Gynecol* 2013;122:1184–95.
- Landon MB, Hauth JC, Leveno KJ, *et al*. Maternal and perinatal outcomes associated with a trial of labor after prior caesarean delivery. *N Engl J Med* 2004;351:2581–9.
- Knight HE, Gurol-Urganci I, van der Meulen JH, *et al*. Vaginal birth after caesarean section: a cohort study investigating factors associated with its uptake and success. *BJOG* 2014;121:183–92.
- Paterson CM, Saunders NJ. Mode of delivery after one caesarean section: audit of current practice in a health region. *BMJ* 1991;303:818–21.
- McMahon MJ, Luther ER, Bowes WA Jr, *et al*. Comparison of a trial of labor with an elective second caesarean section. *N Engl J Med* 1996;335:689–95.
- Dodd JM, Crowther CA, Huertas E, *et al*. Planned elective repeat caesarean section versus planned vaginal birth for women with a previous caesarean birth. *Cochrane Database Syst Rev* 2013;12:CD004224..
- American College of Obstetricians and Gynecologists. ACOG Practice bulletin no. 115: Vaginal birth after previous caesarean delivery. *Obstet Gynecol* 2010;116(2 Pt 1):450–63.
- Menacker F, Curtin SC. Trends in caesarean birth and vaginal birth after previous caesarean, 1991–99. *Natl Vital Stat Rep* 2001; 49:1–16.
- Main EK, Morton CH, Melsop K, *et al*. Creating a public agenda for maternity safety and quality in caesarean delivery. *Obstet Gynecol* 2012;120:1194–8.
- Macdonald H, Loder E. Too much medicine: the challenge of finding common ground. *BMJ* 2015;350:h1163.
- National Institute for Health and Clinical Excellence. Caesarean section; clinical guideline no. 132. 2011. <https://www.nice.org.uk/guidance/CG132/chapter/About-this-guideline>
- Dursun P, Yanik FB, Zeyneloglu HB, *et al*. Why women request caesarean section without medical indication? *J Matern Fetal Neonatal Med* 2011;24:1133–7.
- Fenwick J, Gamble J, Hauck Y. Reframing birth: a consequence of caesarean section. *J Adv Nurs* 2006;56:121–30.
- Guise JM, Eden K, Emeis C, *et al*. Vaginal birth after caesarean: New insights. evidence report/technology assessment no.191. 2010; AHRQ Publication No. 10-E003. <http://www.ncbi.nlm.nih.gov/pubmed/20629481>
- Royal College of Obstetricians and Gynaecologists. Birth after previous caesarean birth; green top guideline. 2007;45. <https://www.rcog.org.uk/en/guidelines-research-services/guidelines/gtg45/>
- Shorten A, Shorten B, Keogh J, *et al*. Making choices for childbirth: a randomized controlled trial of a decision-aid for informed birth after caesarean. *Birth* 2005;32:252–61.
- Montgomery AA, Emmett CL, Fahey T, *et al*. Two decision aids for mode of delivery among women with previous caesarean section: Randomised controlled trial. *BMJ* 2007;334:1305–9.
- Liu S, Li H, Lee S. The experiences of multipara who chose to undergo vaginal birth after caesarean [In Chinese]. *J Evid Based Nurs* 2006;2:241–9.
- Phillips E, McGrath P, Vaughan G. 'I wanted desperately to have a natural birth': mothers' insights on vaginal birth after caesarean (VBAC). *Contemp Nurse* 2009;34:77–84.
- Eden KB, Hashima JN, Osterweil P, *et al*. Childbirth preferences after caesarean birth: a review of the evidence. *Birth* 2004;31:49–60.
- Shorten A, Shorten B. The importance of mode of birth after previous caesarean: success, satisfaction, and postnatal health. *J Midwifery Womens Health* 2012;57:126–32.
- Flannagan C, Reid B. Repeat CS or VBAC? A systematic review of the factors influencing pregnant women's decision-making processes. *Evidence Based Midwifery* 2012;10:80–6.
- Noblit G, Hare R. *Meta-ethnography: synthesising qualitative studies*. Sage Publications, 1988.
- Collaboration for qualitative methodologies. Critical appraisal skills program. 1998. <http://www.casp-uk.net>
- Ridley RT, Davis PA, Bright JH, *et al*. What influences a woman to choose vaginal birth after caesarean? *J Obstet Gynecol Neonatal Nurs* 2002;31:665–72.
- York S, Briscoe L, Walkinshaw S, *et al*. Why women choose to have a repeat caesarean section. *Br J Midwifery* 2005;13:440–5.
- Emmett CL, Shaw AR, Montgomery AA, *et al*. Women's experience of decision making about mode of delivery after a previous caesarean section: the role of health professionals and information about health risks. *BJOG* 2006;113:1438–45.
- Cheung NF, Mander R, Cheng L, *et al*. Caesarean decision-making: negotiation between Chinese women and healthcare professionals. *Evid Based Midwifery* 2006;4:24–30.
- Meddings F, Phipps FM, Haith-Cooper M, *et al*. Vaginal birth after caesarean section (VBAC): exploring women's perceptions. *J Clin Nurs* 2007;16:160–7.
- Moffat MA, Bell JS, Porter MA, *et al*. Decision making about mode of delivery among pregnant women who have previously had a caesarean section: a qualitative study. *BJOG* 2007;114:86–93.
- Fenwick J, Gamble J, Hauck Y. Believing in birth—choosing VBAC: the childbirth expectations of a self-selected cohort of Australian women. *J Clin Nurs* 2007;16:1561–70.
- Farnworth A, Pearson PH. Choosing mode of delivery after previous caesarean birth. *Br J Midwifery* 2007;15:188.
- Cox B. Hot topic. Women's perceptions of their access to, and value of, information as part of their decision making on mode of birth following a previous caesarean section delivery. *Midwifery Digest* 2007;17:159–68.
- Farnworth A, Robson SC, Thomson RG, *et al*. Decision support for women choosing mode of delivery after a previous caesarean section: a developmental study. *Patient Educ Couns* 2008;71:116–24.
- McGrath P, Ray-Barruel G. The easy option? Australian findings on mothers' perception of elective caesarean as a birth choice after a prior caesarean section. *Int J Nurs Pract* 2009;15:271–9.
- McGrath P, Phillips E, Ray-Barruel G. Bioethics and birth: insights on risk decision-making for an elective caesarean after a prior caesarean delivery. *Monash Bioeth Rev* 2009;28:22.1–19.
- Goodall KE, McVittie C, Magill M. Birth choice following primary caesarean section: mothers' perceptions of the influence of health professionals on decision-making. *J Reprod Infant Psychol* 2009;27:4–14.
- Frost J, Shaw A, Montgomery A, *et al*. Women's views on the use of decision aids for decision making about the method of delivery following a previous caesarean section: qualitative interview study. *BJOG* 2009;116:896–905.
- McGrath P, Phillips E, Vaughan G. Vaginal birth after caesarean risk decision-making: Australian findings on the mothers' perspective. *Int J Nurs Pract* 2010;16:274–81.
- David S, Fenwick J, Bayes S, *et al*. A qualitative analysis of the content of telephone calls made by women to a dedicated 'next birth after caesarean' antenatal clinic. *Women Birth* 2010;23:166–71.
- McGrath P, Phillips E, Vaughan G. Speaking out! qualitative insights on the experience of mothers who wanted a vaginal birth after a birth by caesarean section. *Patient* 2010;3:25–32.
- Shorten A, Shorten B, Kennedy HP. Complexities of choice after prior caesarean: a narrative analysis. *Birth* 2014;41:178–84.
- Kennedy HP, Grant J, Walton C, *et al*. Elective caesarean delivery: a mixed method qualitative investigation. *Midwifery* 2013;29:e138–44.
- Tully K, Ball H. Misrecognition of need: women's experiences of and explanations of caesarean delivery. *Soc Sci Med* 2013;85:103–11.
- McClain CS. Patient decision making: the case of delivery method after a previous caesarean section. *Cult Med Psychiatry* 1987;11:495–508.
- Lundgren I, Begley C, Gross MM, *et al*. 'Groping through the fog': a metasynthesis of women's experiences on VBAC (vaginal birth after caesarean section). *BMC Pregnancy Childbirth* 2012;12:85.

52. Shorten A, Shorten B. Timing the provision of a pregnancy decision-aid: temporal patterns of preference for mode of birth during pregnancy. *Patient Educ Couns* 2014;97:108–13.
53. Barnett-Page E, Thomas J. Methods for the synthesis of qualitative research: a critical review. *BMC Med Res Methodol* 2009;9:59.
54. Dy SM, Purnell TS. Key concepts relevant to quality of complex and shared decision-making in health care: a literature review. *Soc Sci Med* 2012;74:582–7.
55. Kamal P, Dixon-Woods M, Kurinczuk JJ, *et al*. Factors influencing repeat caesarean section: Qualitative exploratory study of obstetricians' and midwives' accounts. *BJOG* 2005;112:1054–60.
56. Grobman WA, Lai Y, Landon MB, *et al*. Development of a nomogram for prediction of vaginal birth after cesarean delivery. *Obstet Gynecol* 2007;109:806–12.
57. Say R, Murtagh M, Thomson R. Patients' preference for involvement in medical decision making: a narrative review. *Patient Educ Couns* 2006;60:102–14.
58. Knops AM, Legemate DA, Goossens A, *et al*. Decision aids for patients facing a surgical treatment decision: a systematic review and meta-analysis. *Ann Surg* 2013;257:860–6.
59. Entwistle VA, Carter SM, Trevena L, *et al*. Communicating about screening. *BMJ* 2008;337:a1591.

BMJ Open

Vaginal birth after caesarean section: why is uptake so low? Insights from a meta-ethnographic synthesis of women's accounts of their birth choices

Mairead Black, Vikki A Entwistle, Siladitya Bhattacharya and Katie Gillies

BMJ Open 2016 6:

doi: 10.1136/bmjopen-2015-008881

Updated information and services can be found at:
<http://bmjopen.bmj.com/content/6/1/e008881>

These include:

Supplementary Material

Supplementary material can be found at:
<http://bmjopen.bmj.com/content/suppl/2016/01/08/bmjopen-2015-008881.DC1.html>

References

This article cites 53 articles, 4 of which you can access for free at:
<http://bmjopen.bmj.com/content/6/1/e008881#BIBL>

Open Access

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See:
<http://creativecommons.org/licenses/by/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Obgyn](#) (209)
[Qualitative research](#) (409)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>