

Research Paper

PAIN[®]

OPEN

Cross-centre replication of suppressed burrowing behaviour as an ethologically relevant pain outcome measure in the rat: a prospective multicentre study

Rachel Wodarski^{a,b}, Ada Delaney^c, Camilla Ultenius^d, Rosie Morland^a, Nick Andrews^e, Catherine Baastrup^f, Luke A. Bryden^g, Ombretta Caspani^h, Thomas Christophⁱ, Natalie J. Gardiner^j, Wenlong Huang^a, Jeffrey D. Kennedy^k, Suguru Koyama^l, Dominic Li^k, Marcin Ligocki^b, Annika Lindsten^m, Ian Machinⁿ, Anton Pekcec^g, Angela Robens^l, Sanziana M. Rotariuⁱ, Sabrina Voß^h, Marta Segerdahl^{d,m}, Carina Stenfors^d, Camilla I. Svensson^c, Rolf-Detlef Treede^h, Katsuhiko Uto^l, Kazumi Yamamoto^l, Kris Rutten^{h,*}, Andrew S.C. Rice^a

Abstract

Burrowing, an ethologically relevant rodent behaviour, has been proposed as a novel outcome measure to assess the global impact of pain in rats. In a prospective multicentre study using male rats (Wistar, Sprague-Dawley), replication of suppressed burrowing behaviour in the complete Freund adjuvant (CFA)-induced model of inflammatory pain (unilateral, 1 mg/mL in 100 µL) was evaluated in 11 studies across 8 centres. Following a standard protocol, data from participating centres were collected centrally and analysed with a restricted maximum likelihood-based mixed model for repeated measures. The total population (TP—all animals allocated to treatment; n = 249) and a selected population (SP—TP animals burrowing over 500 g at baseline; n = 200) were analysed separately, assessing the effect of excluding “poor” burrowers. Mean baseline burrowing across studies was 1113 g (95% confidence interval: 1041–1185 g) for TP and 1329 g (1271–1387 g) for SP. Burrowing was significantly suppressed in the majority of studies 24 hours (7 studies/population) and 48 hours (7 TP, 6 SP) after CFA injections. Across all centres, significantly suppressed burrowing peaked 24 hours after CFA injections, with a burrowing deficit of –374 g (–479 to –269 g) for TP and –498 g (–609 to –386 g) for SP. This unique multicentre approach first provided high-quality evidence evaluating suppressed burrowing as robust and reproducible, supporting its use as tool to infer the global effect of pain on rodents. Second, our approach provided important informative value for the use of multicentre studies in the future.

Keywords: Non-evoked pain, Validation, Reproducibility, Preclinical controlled trials

1. Introduction

Prospective multicentre controlled trials are an important clinical research tool, providing high-quality evidence to inform health care systems about the validity of new treatments and outcome measures. In contrast, preclinical studies are predominantly single-centre studies conducted using experimental protocols varying significantly across laboratories. This likely contributes to the well-recognized poor level of experimental reproducibility.⁵⁴

Implementing the concept, ethos, and design of multicentre clinical trials into preclinical studies could significantly increase reproducibility and translatability by standardising experimental design, monitoring data, and improving reporting standards.

The use of multicentre designs in preclinical studies is rare; however, a few pioneering studies in mice have demonstrated the effects of environmental conditions, strain, and study design on behavioural outcomes across centres.^{37,40,59} A multicentre approach

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

^a Pain Research Group, Department of Surgery and Cancer, Imperial College, London, United Kingdom, ^b Eli Lilly and Company, Erl Wood Manor, Windlesham, United Kingdom, ^c Department of Physiology and Pharmacology, Karolinska Institute, Stockholm, Sweden, ^d Neuroscience CNSP iMED, AstraZeneca R&D Södertälje, Södertälje, Sweden, ^e Department of Neurobiology, Boston Children’s Hospital, MA, USA, ^f Danish Pain Research Center, Aarhus University Hospital, Aarhus, Denmark, ^g CNS Disease Division Research Germany, Boehringer Ingelheim Pharma GmbH and Co KG, Biberach an der Riss, Germany, ^h Department of Neurophysiology, Centre for Biomedicine and Medical Technology Mannheim (CBTM), Heidelberg University, Mannheim, Germany, ⁱ Department of Pharmacology and Biomarker Development, Translational Science and Strategy, Grünenthal GmbH, Aachen, Germany, ^j Faculty of Life Sciences, University of Manchester, Manchester, United Kingdom, ^k Eli Lilly and Company, Indianapolis, IN, USA, ^l Laboratory for Pharmacology, Pharmaceutical Research Center, Asahi Kasei Pharma Corporation, Shizuoka, Japan, ^m H. Lundbeck A/S, Valby, Denmark, ⁿ Deal, Kent, United Kingdom. L. A. Bryden is now with the Department of Experimental Psychology, University of Oxford, Oxford, United Kingdom. W. Huang is now with the Institute of Medical Sciences, University of Aberdeen, United Kingdom. C. Stenfors is now with the R&D CNS Research, Orion Corporation, Orion Pharma, Espoo, Finland.

*Corresponding author. Address: Department of Pharmacology and Biomarker Development, Translational Science and Strategy, Grünenthal GmbH, 52078 Aachen, Germany. Tel.: +49-241-569-2077; fax: +49-241-569-2852. E-mail address: kris.rutten@grunenthal.com (K. Rutten).

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal’s Web site (www.painjournalonline.com).

PAIN 157 (2016) 2350–2365

© 2016 International Association for the Study of Pain. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially.

<http://dx.doi.org/10.1097/j.pain.0000000000000657>

for preclinical pain studies could equally provide robust validation and evidence for reproducibility of models, outcome measures, and pharmacological interventions. This could be achieved by systematically assessing replication and interlaboratory variability and identifying factors that may be associated with such variability.

A major shortcoming in chronic pain research has been the lack of newly developed analgesics, because translation from promising preclinical studies to successful clinic trials has been poor.^{49,65,75} Animal studies mostly rely on stimulus-evoked measures of sensory gain, whereas clinical trials use patient-reported outcomes focusing on spontaneous pain intensity, functioning, and quality of life. Although spontaneous pain and decreased quality of life are reported across chronic pain conditions,^{11,16,52,56} sensory gain is common only in a subset of clinical pain conditions caused by peripheral nerve trauma and inflammation but is rare in others, such as polyneuropathies caused by diabetes and HIV.^{39,51,52,56} This has brought into question the translational strength of relying on sensory gain as the sole pain outcome measure preclinically.^{3,48,57} We have previously proposed ethologically relevant rodent behaviours, such as predator avoidance behaviours, as novel nonevoked pain outcome measures to assess the global impact of pain on animals.^{10,23,25,60,72–74}

Among these ethologically relevant behaviours is burrowing. Burrowing, a social behaviour of rats that is important for building underground habitats and nests, is highly conserved in laboratory rats.⁵ Suppressed burrowing, quantified by a reduced amount of displaced substrate from an artificial burrow, is indicative of behavioural dysfunction.¹⁴ Pain-depressed behaviours, such as reduction in feeding, locomotion, or operant behaviours, have been proposed as pain-relevant measures that assess functioning instead of sensory changes.⁴⁵ Although burrowing is a relatively novel pain outcome measure, a few studies have reported decreased burrowing in rat models of inflammatory and neuropathic pain; importantly, burrowing deficits were reversed by known analgesics, suggesting a degree of predictive validity for this outcome measure.^{2,9,20,35,61,62}

We hypothesise that by using a prospective multicentre design, the reliability of novel outcome measures can be efficiently and rapidly evaluated. As part of the IMI European collaboration (<http://www.imieuropain.org>), we investigated rat burrowing behaviour after induction of complete Freund adjuvant (CFA)-associated inflammation across 8 centres to assess the reliability of suppressed burrowing as a pain-related outcome measure. We hope this will facilitate future routine use of multicentre studies to evaluate novel outcome measures and models.

2. Materials and methods

The purpose of this study was to identify whether the impact of CFA-induced inflammation on burrowing behaviour can be reproduced across multiple centres when following a basic protocol (Supplemental digital content 1, <http://links.lww.com/PAIN/A302>). For this, a draft protocol was written and then reviewed by the participating centres before it was finalised. Minor changes to the protocol were permitted to accommodate for local variations in laboratory practice and procedures (eg, different animal suppliers, variations in environmental conditions, substrate and equipment availability, and precise randomisation procedures).

2.1. Ethical statement

All animal experiments conformed to local Government and Institutional guidelines on the care and use of animals in research and the IASP guidelines for in vivo research.^{26,76} Guidance was given on Good Laboratory Practice standards,^{38,58} but exact

methods were not specified, and local variations were allowed and recorded (**Table 1**). Experiments are reported in accordance with the ARRIVE Guidelines³² (Supplemental digital content 6, <http://links.lww.com/PAIN/A307>).

2.2. Experimental animals and environmental conditions

Adult male rats were used for all experiments. Animals were group-housed in standard cages in temperature and humidity-controlled conditions (**Table 1**). Bedding and environmental enrichment was provided according to local convention, including nesting material, jolly balls, or cardboard/plastic tubes. Animals were provided with standard rat chow pellets and tap water ad libitum and were allowed to acclimatise to their housing environment for a minimum of 4 to 7 days before experiments started. Animals were monitored during regular husbandry activity to ensure their well-being.

2.3. Complete Freund's adjuvant—induced inflammation

Inflammation was induced under isoflurane/O₂ anaesthesia by unilateral intraplantar injection of CFA. Sham animals received a saline or incomplete CFA injection of the same volume. The CFA model was chosen because of the high likelihood of ongoing spontaneous pain, as reported in humans after an accidental CFA injection¹⁹ and in animals.^{46,47}

2.4. Burrowing

Burrowing experiments were performed as previously described.^{2,62} For this, a burrowing tube (320 × 100 mm; open end elevated by 60 mm), made of either steel (Boehringer Ingelheim) or plastic (all other groups), was filled with 2500 g substrate and placed in an empty cage (**Fig. 1**). No floor bedding was provided during the burrowing task, as this could hinder cleaning of the substrate. In some studies, however, the cage floor was covered with paper towels to create a more comfortable environment for the animals. If multiple animal cohorts were tested per day, test cages were emptied of displaced substrate, faecal boli and tissue paper, and were wiped clean or replaced with a clean testing cage before starting the next session. As free access to food and water could distract animals from burrowing, most studies opted not to provide food and water during testing. At studies performed at Eli Lilly (United Kingdom), water was accessible during testing, as local regulations only allowed a maximal period of 2 hours without water. Details of the experimental set-ups can be found in **Table 2**.

2.4.1. Training—social facilitation

For animals to learn the task, 2 to 3 training sessions were conducted over consecutive days (apart from weekends). After acclimatisation to the testing room and/or experimental set-up without a burrowing tube, a filled burrowing tube was placed in the test cage, and animals, in pairs, were allowed to burrow for 60 to 120 minutes. If a pair showed poor burrowing behaviour during the first training session, one of the animals was swapped with an animal from a known burrowing pair to facilitate burrowing in subsequent sessions. As no criterion was set a priori to define a poor burrowing pair, experimenters determined this on a case-by-case basis. At Boehringer Ingelheim, pairs burrowing less than 1500 g were classified as poor burrowers; because strong burrowers would displace nearly all of the 2500 g substrate, this high limit was set to ensure the behaviour was strongly expressed.

Table 1

Major domains of Good Laboratory Practice.

	Description of the procedure							
	Boehringer Ingelheim Pharma GmbH	Grünenthal GmbH	Heidelberg University	Asahi Kasei Pharma	University of Manchester	Eli Lilly and Company (United Kingdom)	Imperial College London	Eli Lilly and Company (USA)
Animals	Animal characteristics for each study can be found in Table 3 and in the results in "3.1 Study profile, animal characteristics, and experimental design." Details about species, strain, sex, weight, and supplier are given							
Housing environment/experimental lighting conditions	20-22°C; 45%-60% humidity; 12/12 h light/dark cycle/15 lux	19-23°C; 15%-50% humidity; 12/12 h light/dark cycle/dimmed lights	23-24°C; 47%-63% humidity; 13/11 h light/dark cycle/lights 0.8-7.6 lux	24-26°C; 49%-66% humidity; 12/12 h light/dark cycle/dimmed lights (<50 lux)	19-21°C; 50% humidity; 12/12 h light/dark cycle/lights 30-60 lux	20-22°C; 30%-70% humidity; 12/12 h light/dark cycle/bright lights (study 1 and 2); dimmed lights (study 3)	23°C; 35%-40% humidity; 12/12 h light/dark cycle/lights 30-60 lux	23°C; 35%-40% humidity; 12/12 h light/dark cycle/lights 30-60 lux
Sample size	Sample size estimation (SigmaPlot 11.0 [analysis of variance]) was calculated with pilot burrowing data at 3 d after induction of inflammation (n = 6 and n = 5 for control and complete Freund adjuvant groups, respectively). Parameters were difference in means = 951.79 g, SD = 549.29 g, α = 0.05, and power = 0.8. An adequately powered study required an n = 7 per group and n = 8 when testing 2 and 3 groups, respectively. N numbers for each study are reported in the results and in Figure 2							
Inclusion/exclusion criteria	In the basic protocol, animals that burrowed more or less than 2 SD of the total mean at baseline sessions should have been excluded before model induction. However, because of high variability, this never occurred. Alternative criteria were discussed including the exclusion of animals that burrowed less than 500g at baseline. Thus, at Eli Lilly (United Kingdom), 5 of 32 animals in study 1 and at Grünenthal 1 of 22 animals were excluded before model induction as they burrowed less than 500 g during baseline measurements. No data from these animals was included. At Boehringer Ingelheim, rats were excluded when burrowing on average <1000 g or had an SD >450 g after 3 individual training days; no animals had to be excluded according to this criterion. For all other studies investigators opted to not exclude animals before model induction but instead analyse burrowing behaviour with a total population and selected population approach as described below. Exclusions are reported in Figure 2							
Randomisation to groups: naive, sham, CFA or sham, CFA	Animals were randomly allocated to groups by using a random number generator in Excel. Baseline values (mean 3 baseline sessions) were then used to shift individual values, in a blinded fashion, from group to group until each group had a comparable baseline in terms of mean value and variation	Baseline values were ranked from low to high performers and pseudorandomised by allocating them to groups following a repeated sequence. Individual values were shifted from group to group until each group had a comparable baseline in terms of mean value and variation	Baseline values were used to rank animals from low to high performers and pseudorandomised by allocating them to groups in an alternating pattern	Last baseline session values were used to randomly allocate animals to groups using StatLight#11 (Yukms Co, Ltd)	Baseline values were used to rank animals from low to high performers and pseudorandomised by allocating to groups after a repeated sequence	Baseline values were used to rank animals from low to high performers before randomly allocating animals to groups by using a random sequence generator in Excel. Individual values were shifted blinded from group to group until each group had a comparable baseline in terms of mean value and variation	Baseline values were used to rank animals from low to high performers before randomly allocating animals to groups by using a random sequence generator in Excel. Individual values were shifted blinded from group to group until each group had a comparable baseline in terms of mean value and variation	Last baseline session values were used to randomly allocate animals to groups using a random sequence generator in Excel
Allocation concealment	No allocation concealment procedure was followed	No allocation concealment procedure was followed	No allocation concealment procedure was followed because only CFA animals were injected because of the control group being naive animals	Syringes for model induction were prepared by an independent investigator. The investigator performing model induction was only aware of animal ID	No allocation concealment procedure was followed as the investigator found that colour/viscosity of CFA solutions would have revealed treatment groups	Syringes for model induction were prepared by the investigator only aware of the letter coded group name. During model induction, the investigator was not aware of the treatment as animals and syringes were picked by an independent researcher outside the experimental room	Syringes for model induction were prepared by the investigator only aware of the letter coded group name. During model induction, the investigator was not aware of the treatment as animals and syringes were picked by an independent researcher outside the experimental room	Syringes for model induction were prepared by the investigator. An independent investigator performing model induction was only aware of animal ID

(continued on next page)

Table 1 (continued)

Description of the procedure		Boehringer Ingelheim Pharma GmbH	Grünenthal GmbH	Heidelberg University	Asahi Kasei Pharma	University of Manchester	Eli Lilly and Company (United Kingdom)	Imperial College London	Eli Lilly and Company (USA)
Animals excluded from analysis	For analysis, a total population and selected population approach was exerted. The total population contained all animals randomised to treatment groups (n = 249), whereas the selected population (n = 200) was defined as all rats in the total population that burrowed at least 500 g on average at baseline. Numbers of how many animals were excluded from the total population in each study are given in Figure 2								
Blinding	The investigator who measured burrowing performance was blinded to group allocation	The investigator who measured burrowing performance was blinded to group allocation	The investigator who measured burrowing performance was not blinded to group allocation as investigator performing model induction also performed behavioural analysis	The investigator who measured burrowing performance was blinded to group allocation. The code was broken only at the end of the study	The investigator who measured burrowing performance and assessment of burrowing behaviour were blinded to group allocation. The code was broken only at the end of the study with the help of allocation concealment spreadsheet and safely stored the noted piece of paper with group information	Investigators who performed measurement and assessment of burrowing behaviour were blinded to group allocation. The code was broken at the end of study with the help of allocation concealment spreadsheet and safely stored the envelope containing group information	Investigators who performed measurement and assessment of burrowing behaviour were blinded to group allocation. The code was broken only at the end of the study	Investigators who performed measurement and assessment of burrowing behaviour were blinded to group allocation. The code was broken only at the end of the study	The investigator who measured burrowing performance was blinded to group allocation. The code was broken only at the end of the study
Conflicts of interest	All potential conflicts of interest are stated in the Acknowledgements section including affiliations with pharmaceutical companies, consulting work, and information about study funding								
CFA, complete Freund adjuvant.									




Figure 1. Burrowing tube in the test cage (gravel 2-5 mm).

2.4.2. Baseline performance and testing

For baseline performance and testing, animals were acclimatised to the testing room/experimental set-up as described above. Importantly, animals were placed into the burrowing cage alone rather than in pairs, and allowed to burrow for 60 minutes. At the end of the session, the weight of the displaced substrate was measured. For this, each burrowing tube was weighed before and after burrowing to allow for accurate measurement. In each study, 1 to 3 baseline sessions were performed. Animals were randomised into experimental groups according to their baseline performance by evenly distributing “poor” and “good” burrowers between groups; it was ensured that each experimental group burrowed on average a similar mean amount to reduce the impact of this source of variability on following experiments (Table 1). Burrowing performance was measured on days 1, 2, 3, 7, and 10 after CFA/sham injections.

2.5. Study design

This prospective multicentre validation study was divided into 2 parts. The first part comprised the experimental work at participating laboratories following the basic protocol (Supplemental digital content 1, <http://links.lww.com/PAIN/A302>). All data were centrally collected at Grünenthal and then processed for statistical analysis at H. Lundbeck A/S, who did not participate in the experimental work or data collection.

All burrowing experiments were performed during the light phase. The primary outcome was the change from baseline in the amount burrowed, with a negative value representing a decrease in burrowing behaviour. For this, the mean of the last 2 baseline sessions was used as the baseline value. In cases in which only one session was performed, this was used as the baseline value. The secondary outcome was the effect of protocol variations on burrowing behaviour. Because the study was not designed to investigate the effect of protocol variations, no statistical hypothesis testing has been performed for these elements.

2.6. Statistical analysis

Details for sample size calculation are given in Table 1. The study was powered for individual self-contained experiments at each centre.

Table 2
Experimental set-ups.

	Substrate	Cage floor	Cleaning of the substrate
Boehringer Ingelheim Pharma GmbH	Quartz sand	Bare	Autoclaved at the beginning of the study; removal of wet clumps and faecal boli through sieving; changed over every 6 wk
Grünenthal GmbH	Gravel (2-5 mm)	Paper towel	Every 3 mo with tap water and soap
Heidelberg University	Gravel (2-3 mm)	Bare	Before study with tap water; faecal boli were removed after each session
Asahi Kasei Pharma	Gravel (4-8 mm)	Thick paper towel	Before study with tap water; faecal boli were removed at the end of study
University of Manchester	Gravel (5-8 mm)	Bare	After each session rinsed with tap water and faecal boli were removed; autoclaved at the beginning and end of the study
Eli Lilly and Company (United Kingdom)			
Study 1	Gravel (5-7 mm)	Thick paper towel	After study with tap water
Study 2	Gravel (5-7 mm)	Thick paper towel	After study with tap water
Study 3	Gravel (5-7 mm)	Thick paper towel	After study with tap water
Imperial College London			
Study 1	Gravel (2-5 mm)	Paper towel	After study with tap water
Study 2	Gravel (2-5 mm)	Paper towel	After study with tap water
Eli Lilly and company (USA)	Gravel (size unspecified)	Paper towel	After study rinsed with tap water and diluted bleach

For statistical analyses, we used an approach adapted from an intention-to-treat and per-protocol analysis.²² For this, we separately analysed 2 populations: the total population (TP), including all animals that were allocated to treatment groups, and the selected population (SP) population, including all animals that were allocated to treatment groups and burrowed over 500 g at baseline, a cutoff that has been previously described for burrowing.⁶²

Change from baseline was analysed using a restricted maximum likelihood (REML)-based mixed model for repeated measures (MMRM). Experimental group (naive, sham, CFA), time of assessment (days 1, 2, 3, 7, and 10), group-by-time interaction, and baseline burrowing-by-time interaction were used as fixed effects for analysis of individual studies. Laboratory ID (8 participating centres) was added as the fixed effect for combined analysis across all studies. An unstructured covariance design was used to model the within-animal errors (type 3 tests). A Kenward–Roger approximation was used to estimate denominator degrees of freedom. The analysis was based on the missing-at-random assumption and performed using all available observations. The mean differences between naive and CFA, and sham and CFA, were estimated based on the least squares means for the treatment-by-time interaction in the MMRM model. The estimates are presented with *P* values and 95% confidence intervals (CIs). A *P* value <0.05 was considered statistically significant. Because of the exploratory nature of the study, no adjustment for multiplicity was made. Within the text, burrowing data are presented as mean with 95% CI.

Strain (Sprague-Dawley, Wistar [Wistar and Wistar Hannover]), animal weight at the start of study (200–225 g, 225–250 g, >250 g), substrate size (2-5 mm, 4-8 mm, <1 mm [sand]), number of training sessions (2, 3), weight of substrate provided (2000 g, 2500 g), and sex of the experimenter (female, male, mixed) were investigated as protocol variations. As many of these factors were given over a range, with equal values for all animals within a study or laboratory, ranges have been redefined to fit a set of studies. This could result in overlapping values but always with clear cutoff values given by the original ranges. When applicable, the mean values of the minimum and maximum were used as numeric

values. No statistical analyses have been performed on explanatory factors, and data were presented as observational findings with descriptive statistics only.

For sample size recommendations, sample sizes for a range of mean differences and SD were calculated. Sample size calculations were based on a 2-sample *t* test informed by the descriptive statistics and MMRM analysis for both TP and SP populations 24 hours after CFA injections. Analyses were performed following the exact method under a fixed scenario adopting normal distribution, 2 sides, a nominal power of 0.8, $\alpha = 0.05$, and a null hypothesis assuming no difference between groups. For data manipulation and analysis, SAS software Version 9.4 (SAS Institute Inc, Cary, NC, USA) was used.

3. Results

3.1. Study profile, animal characteristics, and experimental design

Eleven studies, completed between June 2013 and September 2014, were performed at 8 different laboratories across 4 countries (Fig. 2). Of 255 recorded animals, 249 were randomly assigned to 3 treatment groups: naive (49), sham (96), and CFA-treated (104). No adverse events were reported across the studies, with animals tolerating sham injections, CFA injections, and behavioural testing well. Six animals that displayed no burrowing behaviour during baseline sessions were excluded before group assignment. The study protocol regarding animal exclusion was adjusted over the course of the project, and no further animals were excluded at this stage, as a TP and SP analysis approach was favoured instead. All 249 animals allocated to treatment groups were part of the TP analysis, whereas 200 animals were included in the SP analysis; the latter was defined as all rats in the TP population that burrowed at least 500 g on average at baseline (Fig. 2). Although all eleven studies included a CFA group, control groups varied. One study chose naive as the control group, 4 studies chose sham as the control group, and 6 studies chose naive and sham as control groups.




Figure 2. Study profile.

All 11 studies reported randomisation procedures for group allocation. In 4 studies, no allocation concealment was performed. In the remaining 7 studies, allocation concealment procedures were followed during model induction; however, because of the occasional presence of a slight yellowish colour and the higher viscosity of CFA suspensions, allocation concealment could be maintained only in 2 studies. Ten studies reported that outcome assessment was performed blinded. Oedema of the paw in some CFA animals potentially revealed the identity of the experimental group to the investigator in 7 of these studies.

In all studies, male rats from outbred albino strains with a starting weight range of 200 to 335 g were used (Table 3). All CFA-injected animals received 100 µg CFA (1 mg/mL in 100 µL, intraplantar), whereas sham animals received a saline or incomplete Freund adjuvant injection of the same volume (Table 3). Animals were tested at days 1, 2, 3, 7, and 10 after CFA/sham

injections with the exception of the study performed at Eli Lilly (USA), in which animals were only tested up to 7 days. Because of a weighing error, in study 3 at Eli Lilly (United Kingdom), only 2000 g substrate was provided.

Some variations in the features of experimental design were permitted across studies (Fig. 3). Time for animals to acclimatise to the test room ranged from 0 minutes to 60 minutes. Time for acclimatisation to the experimental set-up (empty test cage) again ranged from 0 minutes to 60 minutes. The study performed at the University of Heidelberg introduced a 3-hour habituation session for animals in their home cages in the experimental room before training started. On the first training day, another habituation session was performed in the morning, during which animals were allowed to acclimatise to the experimental room for 30 minutes and for a further 30 minutes in the empty test cage. An empty test tube was then added to

Table 3**Animal and model characteristics.**

	Animal characteristics				
	Strain	Sex	Weight at start of study, g	Supplier	Country of origin
Boehringer Ingelheim Pharma GmbH	Wistar Hannover	♂	200-220	Charles River	Germany
Grünenthal GmbH	Sprague-Dawley	♂	260-310	Janvier	France
Heidelberg University	Wistar Hannover	♂	250-290	Charles River	Germany
Asahi Kasei Pharma	Sprague-Dawley	♂	213-247	Charles River	Japan
University of Manchester	Sprague-Dawley	♂	243-335	Charles River	United Kingdom
Eli Lilly and Company (United Kingdom)					
Study 1	Sprague-Dawley	♂	200-250	Charles River	United Kingdom
Study 2	Wistar	♂	200-250	Charles River	United Kingdom
Study 3	Wistar	♂	200-250	Charles River	United Kingdom
Imperial College London					
Study 1	Wistar	♂	235-300	Charles River	United Kingdom
Study 2	Wistar	♂	235-300	Charles River	United Kingdom
Eli Lilly and Company (USA)	Sprague-Dawley	♂	260-323	Harlan	USA
	Model characteristics				
	CFA supplier	Form of CFA	Concentration, mg/mL	Volume, μ L	Vehicle used for CFA solution
Boehringer Ingelheim Pharma GmbH	Sigma (Germany)	Liquid	1	100	NA
Grünenthal GmbH	DIFCO Laboratories (United Kingdom)	Powder	1	100	Incomplete Freund adjuvant
Heidelberg University	Sigma (Germany)	Liquid	1	100	NA
Asahi Kasei Pharma	Sigma (Japan)	Liquid	1	100	NA
University of Manchester	Sigma (United Kingdom)	Liquid	1	100	NA
Eli Lilly and Company (United Kingdom)					
Study 1	Sigma (United Kingdom)	Liquid	1	100	NA
Study 2	Sigma (United Kingdom)	Liquid	1	100	NA
Study 3	Sigma (United Kingdom)	Liquid	1	100	NA
Imperial College London					
Study 1	Sigma (United Kingdom)	Liquid	1	100	NA
Study 2	Sigma (United Kingdom)	Liquid	1	100	NA
Eli Lilly and Company (Indianapolis, USA)	Sigma (St. Louis, USA)	Liquid	1	100	NA


CFA, complete Freund adjuvant; NA, not applicable because CFA solution was provided in liquid form by the supplier and was injected neat; when CFA was purchased from Sigma, saline was used for sham injections.

the cage, and animals spent another 60 minutes to acclimatise. The training session was then conducted in the afternoon. As animals were still in the behavioural room, they were allowed to acclimatise for only 30 minutes to the experimental set-up before an empty burrowing tube was placed in the test cage for 30 minutes. Subsequently, a filled burrowing tube was added, and animals were allowed to burrow for 60 minutes. All subsequent training sessions followed the standard protocol (**Fig. 3**). For all training, baseline, and testing sessions, animals were allowed to burrow for 60 minutes. The only exception was for training sessions performed at Boehringer Ingelheim, in which animals were allowed to burrow for 120 minutes. Across studies, 2 to 3 training sessions and 1 to 3 baseline sessions were performed (**Fig. 3**). Testing sessions followed the same protocol as outlined for baseline sessions.

3.2. Burrowing behaviour over time

Baseline burrowing behaviour was highly variable between individual animals and ranged from 0 to 2286 g at baseline. Mean baseline burrowing across studies was 1113 g (95%

confidence interval: 1041-1185 g) and 1329 g (1271-1387 g) in the TP and SP population, respectively (Supplemental digital content 2, <http://links.lww.com/PAIN/A303>). To reduce the impact of baseline variability between studies, mean change from baseline in the amount burrowed was chosen as the primary outcome, with a negative value representing a decrease in burrowing behaviour. A summary of the nonnormalised data can be found in the supplemental material (Supplemental digital content 2, <http://links.lww.com/PAIN/A303>). Statistical analysis of individual studies showed, for both TP and SP populations, that in 7 of 11 studies, burrowing behaviour was significantly suppressed 24 hours after CFA injections as compared with a control group (naive or sham) (**Tables 4 and 5**). Forty-eight hours after CFA injections, significantly suppressed burrowing behaviour was observed in 7 studies for the TP and in 6 studies for the SP population. Burrowing performance over time in naive and sham groups from individual studies was largely within the 95% CI of the overall mean (all animals combined of relevant group) (**Fig. 4A–D**). In contrast, burrowing behaviour in CFA groups was more variable (**Fig. 4E, F**). Burrowing behaviour was comparable between the TP and SP population, with a burrowing deficit of



* N/A not applicable as animals were already in the test room from the acclimatisation session in the morning
 † The first number represents time spent in the empty burrowing cage and the second number represents the time spent in the burrowing cage with an empty burrowing tube

Figure 3. Experimental designs across studies.

–374 g (–479 to –269 g) and –498 g (–609 to –386 g), at 24 hours after CFA injection, respectively.

Statistical analysis of the combined data showed suppressed burrowing peaked 24 hours after CFA injections and, although less pronounced, was present for up to 10 days, gradually regressing to baseline values (Fig. 5). Each of the fixed effects, namely, group, time of assessment, laboratory ID, group-by-time interaction, and baseline burrowing-by-time interactions, significantly contributed to heterogeneity (Table 6).

3.3. Effect of protocol variations on burrowing behaviour—observational data

The secondary outcome was the effect of local protocol variations on burrowing behaviour. Because of the nature of the study design, no statistical hypothesis testing was performed, and data are presented as observational findings only with descriptive statistics. Observations were made including all animals allocated to a treatment group (TP population), focusing on the 24-hour time point at which burrowing deficits were most pronounced. Data summarising burrowing behaviour at all time points by protocol variation can be found in the supplemental material (Supplemental digital content 3, <http://links.lww.com/PAIN/A304>). There seemed to be no strain difference between Wistar and Sprague-Dawley rats, but there was a tendency towards a more prominent CFA-associated burrowing deficit in animals of lower body weight. Burrowing deficits after CFA injection were also more noticeable when a substrate of smaller size was used (Fig. 6). An increased number of training sessions negatively affected burrowing behaviour in naive animals. In addition, when a larger amount of substrate was provided, the burrowing deficit was more pronounced. The sex of the experimenter also might have affected burrowing behaviour; increased burrowing behaviour was observed with a male experimenter in naive animals, whereas in contrast, a more pronounced burrowing deficit was

present in CFA animals with a male experimenter (Supplemental digital content 4, <http://links.lww.com/PAIN/A305>). It should be noted that because the study was not designed to identify variables that affect burrowing outcome, confounding factors were identified, and observational data should be interpreted cautiously. In particular, animals were unevenly distributed across groups. An n = 4 was recorded for the naive group that underwent 3 training sessions and for the naive group that was tested by a male experimenter. Furthermore, sand as the burrowing substrate and provision of 2000 g of the substrate was only reported by one centre in one study each.

4. Discussion

A prospective preclinical multicentre study across 8 laboratories assessed the reliability of CFA-associated suppressed burrowing in rats. Overall, reduced burrowing was partially replicated at 6 of the 8 participating centres with an element of variability between and within centres. The prospective multicentre approach was important in that it enabled the evaluation of variability of suppressed burrowing, and it could prove important for future studies aiming to identify the factors underlying such variability.

We showed prominent CFA-associated suppressed burrowing in 7 of 11 studies. Consistent with our results, CFA-associated suppression of behaviour previously has been shown in feeding behaviours,^{31,33,36} locomotion,^{21,30} and operant behaviours.^{18,69} Analysis of the combined data demonstrated that burrowing deficits peaked 24 hours after CFA injection, although with high variability between individual studies, but some studies show no suppression of burrowing. Notably, the original sample size calculation was based on a pilot study with a large effect size; however, across centres, burrowing deficits ranged from 1570 to 273 g. Therefore, some studies were underpowered using the originally estimated sample size. This could result in a reduced

Table 4

Time course of change from baseline in the amount burrowed of individual studies (total population).

	Group	N	24 h	48 h	72 h	7 d	10 d
		number					
Boehringer Ingelheim Pharma GmbH	Naive	9	-38 (-269 to 193), P = 0.0002	86 (-132 to 304), P = 0.006	-91 (-320 to 138), P = 0.2	85 (-91 to 261), P = 0.002	-16 (-179 to 147), P = 0.2
	Sham	10	-60 (-280 to 160), P = 0.0002	-39 (-245 to 167), P = 0.04	20 (-198 to 238), P = 0.046	37 (-130 to 204), P = 0.005	122 (-31 to 275), P = 0.02
	CFA	10	-740 (-960 to -520)	-369 (-575 to -163)	-308 (-526 to -90)	-331 (-498 to -164)	-151 (-306 to 4)
Grünenthal GmbH	Naive	7	49 (-163 to 261), P < 0.0001	227 (-18 to 472), P = 0.008	70 (-308 to 448), P = 0.02	354 (185 to 523), P = 0.08	112 (-126 to 314), P = 0.5
	Sham	7	84 (-126 to 294), P < 0.0001	56 (-187 to 299), P = 0.06	-27 (-403 to 349), P = 0.04	207 (40 to 374), P = 0.5	203 (-15 to 421), P = 1.0
	CFA	7	-898 (-1110 to -686)	-307 (-552 to -62)	-631 (-1009 to -253)	126 (-43 to 295)	205 (-15 to 425)
Heidelberg University	Naive	8	196 (-161 to 553), P = 0.3	172 (-163 to 507), P = 0.7	376 (70 to 682), P = 0.9	297, (30 to 564), P = 0.4	337 (-41 to 715), P = 0.7
	CFA	8	-98 (-455 to 259)	282 (-53 to 617)	338 (32 to 644)	461 (194 to 728)	246 (-132 to 624)
Asahi Kasei Pharma	Sham	12	256 (-77 to 589), P = 0.006	237 (-102 to 576), P = 0.03	235 (-106 to 576), P = 0.02	328 (-70 to 726), P = 0.4	112 (-260 to 484), P = 1.0
	CFA	12	-488 (-821 to -155)	-337 (-676 to 2)	-408 (-749 to -67)	58 (-328 to 444)	120 (-241 to 481)
University of Manchester	Naive	8	15 (-342 to 372), P = 0.03	207 (-212 to 626), P = 0.08	271 (-199 to 741), P = 0.2	510 (85 to 935), P = 0.2	282 (-226 to 790), P = 0.4
	Sham	8	470 (113 to 827), P = 0.06	728 (309 to 1147), P = 0.01	482 (12 to 952), P = 0.9	991 (566 to 1416), P = 0.4	650 (142 to 1158), P = 0.6
	CFA	8	392 (35 to 749)	244 (-175 to 663)	542 (72 to 1012)	200 (-225 to 625)	0 (-508 to 508)
Eli Lilly and Company (United Kingdom)							
Study 1	Naive	9	-305 (-813 to 203), P = 0.03	-386 (-627 to -145), P = 0.08	-2 (-431 to 427), P = 0.2	216 (-209 to 641), P = 0.2	-114 (-678 to 450), P = 0.4
	Sham	9	-400 (-700 to -100), P = 0.06	-203 (-452 to 46), P = 0.01	-331 (-760 to 98), P = 0.9	-520 (-945 to -95), P = 0.4	-247 (-811 to 317), P = 0.6
	CFA	9	-833 (-1133 to -533)	-707 (-948 to -466)	-380 (-809 to 49)	-236 (-661 to 189)	-446 (-1010 to 118)
Study 2	Sham	16	187 (7 to 367), P = 0.04	197 (-30 to 424), P = 0.1	171 (-82 to 424), P = 0.8	291 (70 to 512), P = 0.7	162 (-89 to 413), P = 0.7
	CFA	16	-99 (-279 to 81)	-56 (-283 to 171)	119 (-134 to 372)	351 (130 to 572)	241 (-10 to 492)
Study 3	Sham	16	261 (-72 to 594), P = 0.06	403 (125 to 681), P = 0.045	192 (-143 to 527), P = 0.8	340 (3 to 677), P = 0.4	230 (-89 to 549), P = 0.3
	CFA	16	-203 (-526 to 120)	-17 (-295 to 261)	139 (-196 to 474)	139 (-198 to 476)	-39 (-358 to 280)
Imperial College London							
Study 1	Naive	4	721 (235 to 1207), P = 0.002	221 (-312 to 754), P = 0.6	390 (-223 to 1003), P = 0.9	-80 (-782 to 622), P = 0.8	-267 (-1047 to 513), P = 0.6
	Sham	4	-178 (-664 to 308), P = 0.09	133 (-400 to 666), P = 0.8	288 (-325 to 901), P = 0.7	310 (-392 to 1012), P = 0.7	414 (-366 to 1194), P = 0.1
	CFA	4	-849 (-1335 to -363)	34 (-499 to 567)	438 (-177 to 1053)	72 (-630 to 774)	-560 (-1342 to 222)
Study 2	Naive	4	-670 (-1189 to -151), P = 0.9	-183 (-681 to 315), P = 0.07	16 (-619 to 651), P = 0.1	327 (-347 to 1001), P = 0.3	-522 (-1177 to 133), P = 0.9
	Sham	4	-86 (-607 to 435), P = 0.1	117 (-383 to 617), P = 0.02	-135 (-772 to 502), P = 0.2	119 (-557 to 795), P = 0.5	177 (-478 to 832), P = 0.1
	CFA	4	-736 (-1253 to -219)	-956 (-1452 to -460)	-847 (-1475 to -216)	-186 (-856 to 484)	-581 (-1232 to 70)
Eli Lilly and Company (USA)	Sham	10	48 (-164 to 260), P = 0.0001	46 (-109 to 201), P = 0.005	226 (8 to 444), P = 0.3	12 (-186 to 210), P = 0.4	—
	CFA	10	-742 (-954 to -530)	-335 (-490 to -180)	37 (-181 to 255)	-127 (-325 to 71)	—
% of naive vs CFA comparisons reporting significant suppression of burrowing			71	29	14	14	0
% of sham vs CFA comparisons reporting significant suppression of burrowing			60	80	30	10	10

Data shown as mean (95% confidence interval) and analysed using a restricted maximum likelihood (REML)-based mixed model for repeated measures (MMRM) approach with fixed effects (group [naive, sham, CFA]), time of assessment (days 1, 2, 3, 7, 100), group-by-time, and baseline burrowing-by-time interactions). P values given for naive and sham groups refer to comparison with the complete Freund adjuvant (CFA) group. Significance level has been set at $P < 0.05$, bold values indicate significant differences to the CFA treated groups. CFA, complete Freund adjuvant.

Table 5

Time course of change from baseline in the amount burrowed of individual studies (selected population).

	Group	N	24 h	48 h	72 h	7 d	10 d		
		number							
Boehringer Ingelheim Pharma GmbH	Naive	9	-38 (-269 to 193), P = 0.0002	86 (-132 to 304), P = 0.006	-91 (-320 to 138), P = 0.2	85 (-91 to 261), P = 0.002	-16 (-179 to 147), P = 0.2		
	Sham	10	-60 (-280 to 160), P = 0.0002	-39 (-245 to 167), P = 0.04	20 (-198 to 238), P = 0.046	37 (-130 to 204), P = 0.0005	122 (-31 to 275), P = 0.02		
	CFA	10	-740 (-960 to -520)	-369 (-575 to -163)	-308 (-526 to -90)	-331 (-498 to -164)	-151 (-306 to 4)		
Grünenthal GmbH	Naive	7	49 (-163 to 261), P < 0.0001	227 (-18 to 472), P = 0.008	70 (-308 to 448), P = 0.02	354 (185 to 523), P = 0.08	112 (-126 to 314), P = 0.5		
	Sham	7	84 (-126 to 294), P < 0.0001	56 (-187 to 299), P = 0.06	-27 (-403 to 349), P = 0.04	207 (40 to 374), P = 0.5	203 (-15 to 421), P = 1.0		
	CFA	7	-898 (-1110 to -686)	-307 (-552 to -62)	-631 (-1009 to -253)	126 (-43 to 295)	205 (-15 to 425)		
Heidelberg University	Naive	6	303 (-142 to 748), P = 0.2	265 (-162 to 692), P = 0.8	554 (197 to 911), P = 0.5	472 (225 to 719), P = 0.8	297 (-160 to 754), P = 0.9		
	CFA	7	-106 (-518 to 306)	345 (-51 to 741)	383 (52 to 714)	433 (206 to 660)	274 (-149 to 697)		
Asahi Kasei Pharma	Sham	7	-87 (-524 to 350), P = 0.04	-68 (-493 to 357), P = 0.1	-78 (-507 to 351), P = 0.04	140 (-356 to 636), P = 0.2	-42 (-605 to 521), P = 0.8		
	CFA	6	-881 (-1355 to -407)	-632 (-1093 to -171)	-864 (-1329 to -399)	-439 (-976 to 98)	54 (-556 to 664)		
University of Manchester	Naive	6	119 (-179 to 417), P = 0.2	399 (11 to 787), P = 0.9	479 (38 to 920), P = 0.6	481 (5 to 957), P = 0.5	187 (-411 to 785), P = 0.7		
	Sham	6	580 (282 to 878), P = 0.4	838 (450 to 1226), P = 0.1	726 (285 to 1167), P = 0.2	857 (381 to 1333), P = 0.1	716 (116 to 1316), P = 0.1		
	CFA	6	385 (87 to 683)	379 (-7 to 765)	324 (-117 to 765)	243 (-233 to 719)	-11 (-609 to 587)		
Eli Lilly and Company1 (United Kingdom)	Study 1	Naive	8	-302 (-661 to 57), P = 0.02	-361 (-161 to -106), P = 0.04	-7 (-521 to 507), P = 0.3	69 (-433 to 571), P = 0.3	-367 (-1012 to 278), P = 0.8	
		Sham	7	-499 (-862 to -136), P = 0.1	-232 (-518 to 54), P = 0.01	-357 (-902 to 188), P = 0.9	-579 (-1112 to -46), P = 0.5	-220 (-904 to 464), P = 0.6	
		CFA	7	-958 (-1321 to -595)	-798 (-1068 to -528)	-401 (-944 to 142)	-308 (-841 to 225)	-506 (-1188 to 176)	
	Study 2	Sham	8	15 (-248 to 278), P = 0.2	49 (-243 to 341), P = 0.08	-50 (-358 to 258), P = 0.5	105 (-179 to 389), P = 0.9	-33 (-335 to 269), P = 0.9	
		CFA	10	-259 (-494 to -24)	-322 (-583 to -61)	-216 (-490 to 58)	69 (-184 to 322)	-19 (-288 to 250)	
	Study 3	Sham	12	266 (-108 to 640), P = 0.03	404 (102 to 706), P = 0.02	312 (-23 to 647), P = 0.1	214 (-162 to 590), P = 0.5	195 (-205 to 595), P = 0.4	
		CFA	13	-351 (-698 to -4)	-135 (-425 to 155)	-96 (-419 to 227)	30 (-333 to 393)	-51 (-435 to 333)	
	Imperial College London	Study 1	Naive	4	660 (107 to 1213), P = 0.008	229 (-283 to 741), P = 0.9	316 (-111 to 743), P = 0.5	-117 (-701 to 467), P = 0.5	-258 (-766 to 250), P = 0.6
			Sham	3	-162 (-815 to 491), P = 0.1	-114 (-720 to 492), P = 0.5	-246 (-752 to 260), P = 0.07	-184 (-874 to 506), P = 0.4	-140 (-973 to 693), P = 0.5
Study 2		CFA	3	-985 (-1614 to -356)	190 (-392 to 772)	552 (66 to 1038)	230 (-432 to 892)	-564 (-1366 to 238)	
		Naive	4	-674 (-1186 to -162), P = 0.8	-193 (-640 to 254), P = 0.05	-10 (-622 to 602), P = 0.09	272 (-445 to 989), P = 0.4	-543 (-1237 to 151), P = 0.9	
		Sham	3	63 (-531 to 657), P = 0.08	317 (-200 to 834), P = 0.009	49 (-661 to 759), P = 0.09	70 (-761 to 901), P = 0.6	118 (-686 to 922), P = 0.2	
CFA	4	-752 (-1266 to -238)	-952 (-1401 to -503)	-891 (-1504 to -278)	-241 (-960 to 478)	-598 (-1294 to 98)			
Eli Lilly and Company (USA)	Sham	10	48 (-164 to 260), P = 0.0001	46 (-109 to 201), P = 0.005	226 (8 to 444), P = 0.3	12 (-186 to 210), P = 0.4	—		
	CFA	10	-742 (-954 to -530)	-335 (-490 to -180)	37 (-181 to 255)	-127 (-325 to 71)	—		
% of naive vs CFA comparisons reporting significant suppression of burrowing		—	57	42	14	14	0		
% of sham vs CFA comparisons reporting significant suppression of burrowing		—	50	50	30	10	10		

Data shown as mean (95% confidence interval) and analysed using a restricted maximum likelihood (REML)-based mixed model for repeated measures (MMRM) approach with fixed effects (group (naive, sham, complete Freund adjuvant [CFA]), time of assessment (days 1, 2, 3, 7, 10), group-by-time and baseline burrowing-by-time interactions). P values given for naive and sham groups refer to comparison to the CFA group. Significance level has been set at P < 0.05, bold values indicate significant differences to the CFA treated groups. CFA, complete Freund adjuvant.




Figure 4. Burrowing behaviour in individual studies—mean change from the baseline amount burrowed. (A, C, E) Total population ($n = 249$): burrowing behaviour in individual studies in naive (A), sham (C), and complete Freund adjuvant (CFA) (E) groups. Gray area represents the mean with 95% confidence interval (CI). (B, D, F) Selected population ($n = 200$): burrowing behaviour in individual studies in naive (B), sham (D), and CFA (F) groups. Gray area represents the mean with 95% CI.

chance to detect a true effect. We have calculated sample size recommendations for a range of mean differences and SD (Supplemental digital content 5, <http://links.lww.com/PAIN/A306>) to provide guidance for future studies. Increasing the sample size of underpowered studies would result in a more accurate estimated effect size and could reduce variability within studies. Group allocation, time of assessment, laboratory ID, group-by-time interaction, and baseline burrowing-by-time interaction all significantly contributed to the heterogeneity across studies. Adjustment for laboratory ID showed a change from baseline similar to data without adjustment, demonstrating that

suppression of burrowing is robust across laboratories, despite the variability in the effect size.

We also observed effects of local protocol variations on burrowing behaviour. No statistical analyses were performed on these observations, as the study was not designed to formally detect such effects. Confounding factors such as uneven distribution of animals across groups and variables reported only by one centre should be considered when interpreting the data. Observations made were reported both for transparency and to identify variability factors in burrowing behaviour meriting future study. Although strain differences have been reported for other




Figure 5. Time course of change from baseline in the amount burrowed. (A) Total population (n = 249). (B) Selected population (n = 200). Data shown as mean with 95% confidence interval (CI) and analysed using a restricted maximum likelihood (REML)-based mixed model for repeated measures (MMRM) approach with fixed effects [group (naive, sham, complete Freund adjuvant [CFA]), time of assessment (days 1, 2, 3, 7, 10), laboratory ID, group-by-time, and baseline burrowing-by-time interactions]. *Significant difference between the naive and CFA groups. #Significant difference between the sham and CFA group. */# P < 0.05.

outcome measures,^{8,42,63} in this study, no strain differences between Sprague-Dawley and Wistar rats were observed for suppressed burrowing. Animals with a lower body weight developed an increased burrowing deficit, whereas burrowing in sham or naive groups was unaffected. As all animals received the same volume and concentration of CFA, it may be that smaller animals received relatively more CFA per paw mass, resulting in a more severe inflammatory response leading to a larger burrowing deficit. The suppression of burrowing observed with a smaller amount of provided substrate is most likely due to the reduced amount of substrate available. In naive animals, we observed an increased burrowing deficit with male experimenters as compared with female experimenters or mixed experimenter teams. Because a male experimenter was only reported by one centre and only 4 rats were in this group, this result may be due to chance or a centre-specific effect. A study in mice showed that the experimenter’s sex affects pain outcome measures in mice.⁶⁷ Further studies are required to verify whether burrowing behaviour in rats is also affected by the experimenter’s sex. Additional studies would be required to assess the impact of the substrate size and number of training sessions on burrowing, specifically whether a small sized substrate is superior to larger sized substrate and whether an increased number of training sessions reduces burrowing behaviour.

To assess the effect of excluding “poor” burrowers, we analysed the TP (all animals allocated to treatment groups) and

an SP (all allocated animals that burrowed above 500 g at baseline), an approach adapted from the intention-to-treat and per-protocol analyses used in clinical trials.²² In both populations, the pattern of suppressed burrowing was comparable, which suggests that suppressed burrowing is a robust measure. Therefore, we recommend not to exclude “poor” burrowers. Excluding animals would increase variability, resulting in a less accurate effect size estimates. Exclusions could also result in attrition bias, an issue particularly important for studies using relatively small sample sizes.^{1,13,24}

Adding nonevoked ethologically relevant outcome measures to assess the global impact of pain previously has been suggested as a potential means to improve translation.^{49,56,68,75} Development and validation of these measures is of key relevance, particularly as spontaneous pain, functioning, and quality of life are primary outcome measures in clinical trials. Suppression of burrowing reflects the global impact of purportedly pain-induced reduction of general “well-being”.^{27–29,74} Although suppressed burrowing is not a pain-specific test, treatment with known analgesics has been shown to attenuate decreased burrowing behaviour in various pain models, suggesting a pain-specific component.^{2,9,35,61,62} A limitation of our study was the lack of an independent validation of suppressed burrowing as indicative of a pain-specific outcome measure; it will be crucial to address the interdependence of this connection in future studies. Correlation with other nonevoked

Table 6
Type 3 tests of fixed effects.

Fixed effect	Total population				Selected population		
	Nominator, df	Denominator, df	F	Pr > F	Denominator, df	F	Pr > F
Experimental group (naive, sham, CFA)	2	239	13.44	<0.0001*	190	16.85	<0.0001*
Time of assessment (days 1, 2, 3, 7, 10)	4	238	5.75	0.0002*	189	2.49	0.0444*
Laboratory ID (8 participating centres)	7	239	4.74	<0.0001*	189	7.68	<0.0001*
Group-by-time interaction	8	340	3.59	0.0005*	270	4.24	<0.0001*
Baseline burrowing-by-time interaction	5	249	14.50	<0.0001*	193	10.34	<0.0001*

Data were analysed using a restricted maximum likelihood (REML)-based mixed model for repeated measures (MMRM).

* Significant difference between variables within fixed effect.

CFA, complete Freund adjuvant; df, degrees of freedom; Pr, probability.




Figure 6. Burrowing performance 24 hours after complete Freund adjuvant (CFA) injections in subgroups factoring in protocol variations (total population). (A) Burrowing dependent on strain. (B) Burrowing dependent on the weight of animals at the start of study. (C) Burrowing dependent on the substrate size. Data shown as single values (diamonds) and mean (square) with 95% confidence interval (whiskers).

pain-related outcomes would be essential, particularly as a lack of correlation has been shown between burrowing performance and evoked mechanosensory thresholds in a neuropathic pain rat model,³⁵ which suggests that suppressed burrowing may reflect a pain component not directly linked to sensory gain. Furthermore, pharmacological validation studies should be conducted, preferably informed by robust meta-analyses of clinical

trials to guide both drug and dose selection.^{17,53} First, it should be demonstrated to what extent clinically efficacious drugs reverse suppression of burrowing. In the CFA model, ibuprofen has been shown to reverse suppressed burrowing, whereas gabapentin, which has a large body of evidence supporting efficacy in neuropathic, but not inflammatory, pain, is appropriately ineffective,^{2,20,62} suggesting good pharmacologic sensitivity of suppressed burrowing.

Second, compounds such as neurokinin 1 antagonists and cannabinoid 2 receptor agonists that have been shown to be efficacious in animal pain models measuring evoked endpoints but have failed in clinical trials^{48,65,75} should be tested to assess the degree of pain specificity of burrowing.

We chose the CFA model, as we expected a high likelihood of ongoing spontaneous pain.^{19,46,47} To validate suppressed burrowing as a pain-relevant outcome measure for neuropathic pain conditions, burrowing behaviour should be measured in a range of models. An important aspect of modelling neuropathic pain is gender generalizability. Animal studies use mainly male rodents, whereas clinical trials enrol both sexes.⁷ Importantly, sex differences in behavioural responses have been shown in rodents.^{15,42,66} As the primary interest of our study was to evaluate burrowing across centres and not establish the model's predictive validity as a pain outcome measure, only male rats were used. However, future validation studies should include female rats.

A multicentre approach for preclinical studies is very novel. Similar to clinical multicentre trials, the study design should be of a high standard, and results should be reported transparently. In this study, all participating centres followed a basic protocol that was previously reviewed and agreed upon by all parties; however, minor changes were permitted to pragmatically accommodate for local variations in laboratory practice and procedures. No detailed specifications were given regarding the scope of these changes, inevitably resulting in some degree of uncontrolled heterogeneity between studies. An external review of the protocol could have identified this issue before study start. Variations were also reported concerning bias reduction procedures. Although guidance on Good Laboratory Practice was given, there was variability between centres as to the extent to which such practice was followed, most notably as a result of constraints imposed by established local procedures. Future preclinical multicentre studies should not only provide Good Laboratory Practice training and validation but also establish an independent central monitor, similar to phase III clinical trials, to ensure protocol compliance and bias reduction.⁶ It should be noted that despite following bias reduction procedures, because of CFA-induced paw oedema, allocation concealment and blinding could not be maintained in all studies, potentially resulting in an overestimation of the effect size.^{12,13,64} As it was not possible to control for all model-specific factors, it is crucial to report data as transparently as possible to clearly highlight study limitations related to internal validity issues. In clinical trials, the near-universal implementation of the CONSORT reporting guidelines has noticeably improved reporting rigour and transparency.⁷⁰ Although ARRIVE guidelines^{32,43} and other recommendations^{34,55,69} provide a similar framework for preclinical studies, they are not yet as well established as CONSORT^{4,32}; however, a similar positive impact on preclinical studies is expected as these guidelines achieve broader acceptance and implementation. In this study, we reported according to ARRIVE guidelines and presented the data as transparently as possible. Recommendations for future studies, based on our practical experience, are summarised in **Table 7**. An audio abstract of this study is available in the supplemental material (Supplemental digital content 7, Audio, <http://links.lww.com/PAIN/A310>).

In conclusion, our approach demonstrates how implementation of a multicentre study design to evaluate novel preclinical outcome measures can yield robust data and can help accelerate the validation of outcome measures, pain models,

Table 7**Recommendations for future multicentre animal studies.****Recommendation**

Study design

Appoint study manager

Protocol development and refinement: each participating centre has to agree on protocol

External review might be beneficial to identify problems

Required training (inclusive Good Laboratory Practice) should be identified and provided at this stage

Plan for statistical analysis should be agreed upon in the protocol—changes to the analysis should be reported and explained

Independent protocol registration might be beneficial to ensure compliance during experimental phase

Experimental phase

Centralised administration: centralised coordination and monitoring, Web-based data entry portal, randomisation process

Enables easier monitoring and quality assurance

Proactive site monitoring and audit through central administration could be beneficial to identify problems during the experimental phase

Reporting

Transparent reporting such as ARRIVE guidelines—it may be that some adaptation of these guidelines is required to assist reporting of preclinical multicentre studies

and pharmacological interventions. This hopefully may help inform the design and conduct of similar future multicentre studies.

Conflict of interest statement

The following authors are/were employees of the following companies at the time this study was undertaken: R. Wodarski, M. Ligocki, D. Li, and J. D. Kennedy: employees of Eli Lilly and Company; C. Ultenius and C. Stenfors: employees of Astra Zeneca; L. A. Bryden and A. Pekcec: employees of Boehringer Ingelheim; T. Christoph, A. Robens, and K. Rutten: employees of Grünenthal; K. Uto, S. Koyama, and K. Yamamoto: employees of Asahi Kasei; A. Lindsten and M. Segerdahl: employees of Lundbeck. Imperial College London: A. S.C. Rice also received research funding from Pfizer and Astellas. Heidelberg University: R.-D. Treede also received research funding from AbbVie, Astellas, and Boehringer Ingelheim. The other authors have no conflicts of interest to declare.

Joint first authors: R. Wodarski, A. Delaney, C. Ultenius, and R. Morland; Joint senior authors: K. Rutten and A. Rice.

Acknowledgements

This European project has received support from the Innovative Medicines Initiative (IMI) Joint Undertaking (under grant agreement number 115007), resources of which are composed of financial contributions from the European Union's Seventh Framework Programme (FP7/20072013) and European Federation of Pharmaceutical Industries and Associations (EFPIA) companies in-kind contribution (see <http://www.imieuropain.org/> for details). Eli Lilly and Company United Kingdom: We would like to thank Dr Gary Gilmour for his assistance in bringing forward this publication. Karolinska Institute: A. Delaney also received funding from Ulla & Gustaf af Ugglas Foundation; EU Project FP7-Health-2013-Innovation-1602919-2. Boehringer Ingelheim: We would like to thank Stacey Gould for technical assistance.

Appendix A. Supplemental Digital Content

Supplemental Digital Content associated with this article can be found online at <http://links.lww.com/PAIN/A302>, <http://links.lww.com/PAIN/A303>, <http://links.lww.com/PAIN/A304>, <http://links.lww.com/PAIN/A305>, <http://links.lww.com/PAIN/A306>, <http://links.lww.com/PAIN/A307>.

Supplemental media

A supplemental video accompanying this article can be found online at <http://links.lww.com/PAIN/A310>.

Article history:

Received 11 January 2016

Received in revised form 19 May 2016

Accepted 20 May 2016

References

- Abdel-Sattar M, Krauth D, Anglemeyer A, Bero L. The relationship between risk of bias criteria, research outcomes, and study sponsorship in a cohort of preclinical thiazolidinedione animal studies: a meta-analysis. *Evidence Based Preclin Med* 2014;1:11–20.
- Andrews N, Legg E, Lisak D, Issop Y, Richardson D, Harper S, Pheby T, Huang W, Burgess G, Machin I, Rice AS. Spontaneous burrowing behaviour in the rat is reduced by peripheral nerve injury or inflammation associated pain. *Eur J Pain* 2012;16:485–95.
- Andrews NA, Latremoliere A, Basbaum AI, Mogil JS, Porreca F, Rice ASC, Woolf CJ, Currie GL, Dworkin RH, Eisenach JC, Evans S, Gewandter JS, Gover TD, Handwerker H, Huang W, Iyengar S, Jensen MP, Kennedy JD, Lee N, Levine J, Lidster K, Machin I, McDermott MP, McMahon SB, Price TJ, Ross SE, Scherrer G, Seal RB, Sena ES, Silva E, Stone L, Svensson CI, Turk DC, Whiteside G. Ensuring transparency and minimization of methodologic bias in preclinical pain research: PPRECISE considerations. *PAIN* 2016;157:901–9.
- Baker D, Lidster K, Sottomayor A, Amor S. Two years later: journals are not yet enforcing the ARRIVE guidelines on reporting standards for preclinical animal studies. *PLoS Biol* 2014;12:e1001756.
- Barnett SA. *The rat: A study in behavior*. Chicago: University of Chicago Press, 1975.
- Bath PM, Macleod MR, Green AR. Emulating multicentre clinical stroke trials: a new paradigm for studying novel interventions in experimental models of stroke. *Int J Stroke* 2009;4:471–9.
- Beery AK, Zucker I. Sex bias in neuroscience and biomedical research. *Neurosci Biobehav Rev* 2011;35:565–72.
- Benoliel R, Eliav E, Tal M. Strain-dependent modification of neuropathic pain behaviour in the rat hindpaw by a priming painful trigeminal nerve injury. *PAIN* 2002;97:203–12.
- Bryden LA, Nicholson JR, Doods H, Pekcec A. Deficits in spontaneous burrowing behavior in the rat bilateral monosodium iodoacetate model of osteoarthritis: an objective measure of pain-related behavior and analgesic efficacy. *Osteoarthr Cartil* 2015;23:1605–12.
- Caspani O, Reitz MC, Ceci A, Kremer A, Treede RD. Tramadol reduces anxiety-related and depression-associated behaviors presumably induced by pain in the chronic constriction injury model of neuropathic pain in rats. *Pharmacol Biochem Behav* 2014;124:290–6.
- Ciaramitaro P, Mondelli M, Logullo F, Grimaldi S, Battiston B, Sard A, Scarinzi C, Migliaretti G, Faccani G, Cocito D. Traumatic peripheral nerve injuries: epidemiological findings, neuropathic pain and quality of life in 158 patients. *J Peripher Nerv Syst* 2010;15:120–7.
- Crossley NA, Sena E, Goehler J, Horn J, van der Worp B, Bath PM, Macleod M, Dirnagl U. Empirical evidence of bias in the design of experimental stroke studies: a metaepidemiologic approach. *Stroke* 2008;39:929–34.
- Currie GL, Delaney A, Bennett MI, Dickenson AH, Egan KJ, Vesterinen HM, Sena ES, Macleod MR, Colvin LA, Fallon MT. Animal models of bone cancer pain: systematic review and meta-analyses. *PAIN* 2013;154:917–26.
- Deacon RM. Burrowing in rodents: a sensitive method for detecting behavioral dysfunction. *Nat Protoc* 2006;1:118–21.
- DeLeo JA, Rutkowski MD. Gender differences in rat neuropathic pain sensitivity is dependent on strain. *Neurosci Lett* 2000;282:197–9.
- Dermanovic Dobrota V, Hrabac P, Skegro D, Smiljanic R, Dobrota S, Prkacin I, Brkljacic N, Peros K, Tomic M, Lukinovic-Skudar V, Basic Kes V. The impact of neuropathic pain and other comorbidities on the quality of life in patients with diabetes. *Health Qual Life Outcomes* 2014;12:171.
- Finnerup NB, Attal N, Haroutounian S, McNicol E, Baron R, Dworkin RH, Gilron I, Haanpaa M, Hansson P, Jensen TS, Kamerman PR, Lund K, Moore A, Raja SN, Rice AS, Rowbotham M, Sena E, Siddall P, Smith BH, Wallace M. Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. *Lancet Neurol* 2015;14:162–73.
- Freitas KC, Hillhouse TM, Leitt MD, Negus SS. Effects of acute and sustained pain manipulations on performance in a visual-signal detection task of attention in rats. *Drug Dev Res* 2015;76:194–203.
- Gould HJ III. Complete Freund's Adjuvant-induced hyperalgesia: a human perception. *PAIN* 2000;85:301–3.
- Gould SA, Doods H, Lamla T, Pekcec A. Pharmacological characterization of intraplantar Complete Freund's Adjuvant-induced burrowing deficits. *Behav Brain Res* 2016;301:142–51.
- Grace PM, Strand KA, Maier SF, Watkins LR. Suppression of voluntary wheel running in rats is dependent on the site of inflammation: evidence for voluntary running as a measure of hind paw-evoked pain. *J Pain* 2014;15:121–8.
- Gupta SK. Intention-to-treat concept: a review. *Perspect Clin Res* 2011;2:109–12.
- Hasnie FS, Breuer J, Parker S, Wallace V, Blackbeard J, Lever I, Kinchington PR, Dickenson AH, Pheby T, Rice AS. Further characterization of a rat model of varicella zoster virus-associated pain: relationship between mechanical hypersensitivity and anxiety-related behavior, and the influence of analgesic drugs. *Neuroscience* 2007;144:1495–508.
- Holman C, Piper SK, Grittner U, Diamantaras AA, Siegerink B, Kimmelman J, Dimagl U. Where have all the rodents gone? The effects of attrition on preclinical research on stroke and cancer. *PLoS Biol* 2016;14:e1002331.
- Huang W, Calvo M, Karu K, Olausen HR, Bathgate G, Okuse K, Bennett DL, Rice AS. A clinically relevant rodent model of the HIV antiretroviral drug stavudine induced painful peripheral neuropathy. *PAIN* 2013;154:560–75.
- International Association for the Study of Pain. IASP guidelines for the use of animals in research. 2014. Available at: <http://www.iasp-pain.org/Education/Content.aspx?ItemNumber=1217>.
- Jirkof P. Burrowing and nest building behavior as indicators of well-being in mice. *J Neurosci Methods* 2014;234:139–46.
- Jirkof P, Cesarovic N, Rettich A, Nicholls F, Seifert B, Arras M. Burrowing behavior as an indicator of post-laparotomy pain in mice. *Front Behav Neurosci* 2010;4.
- Jirkof P, Leucht K, Cesarovic N, Caj M, Nicholls F, Rogler G, Arras M, Hausmann M. Burrowing is a sensitive behavioural assay for monitoring general wellbeing during dextran sulfate sodium colitis in laboratory mice. *Lab Anim* 2013;47:274–83.
- Kandasamy R, Calsbeek JJ, Morgan MM. Home cage wheel running is an objective and clinically relevant method to assess inflammatory pain in male and female rats. *J Neurosci Methods* 2016;263:115–22.
- Kerins CA, Carlson DS, Hinton RJ, Hutchins B, Grogan DM, Marr K, Kramer PR, Spears RD, Bellinger LL. Specificity of molar pattern analysis as an animal model of determining temporomandibular joint inflammation/pain. *Int J Oral Maxillofac Surg* 2005;34:425–31.
- Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving bioscience research reporting: the ARRIVE guidelines for reporting animal research. *PLoS Biol* 2010;8:e1000412.
- Kramer PR, Kerins CA, Schneiderman E, Bellinger LL. Measuring persistent temporomandibular joint nociception in rats and two mice strains. *Physiol Behav* 2010;99:669–78.
- Landis SC, Amara SG, Asadullah K, Austin CP, Blumenstein R, Bradley EW, Crystal RG, Darnell RB, Ferrante RJ, Fillit H, Finkelstein R, Fisher M, Gendelman HE, Golub RM, Goudreau JL, Gross RA, Gubitza AK, Hesterlee SE, Howells DW, Huguenard J, Kelner K, Koroshetz W, Krainc D, Lasic SE, Levine MS, Macleod MR, McCall JM, Moxley RT III, Narasimhan K, Noble LJ, Perrin S, Porter JD, Steward O, Unger E, Utz U, Silberberg SD. A call for transparent reporting to optimize the predictive value of preclinical research. *Nature* 2012;490:187–91.
- Lau W, Dykstra C, Thevarkunnel S, Silenieks LB, de Lannoy IA, Lee DK, Higgins GA. A back translation of pregabalin and carbamazepine against evoked and non-evoked endpoints in the rat spared nerve injury model of neuropathic pain. *Neuropharmacology* 2013;73:204–15.
- Lemons LL, Wiley RG. Neuropeptide Y receptor-expressing dorsal horn neurons: role in nociceptive reflex and operant responses to aversive cold after CFA inflammation. *Neuroscience* 2012;216:158–66.
- Lewejohann L, Reinhard C, Schrewe A, Brandewiede J, Haemisch A, Gortz N, Schachner M, Sachser N. Environmental bias? Effects of housing conditions, laboratory environment and experimenter on behavioral tests. *Genes Brain Behav* 2006;5:64–72.

- [38] Macleod MR, Fisher M, O'Collins V, Sena ES, Dirnagl U, Bath PM, Buchan A, van der Worp HB, Traystman R, Minematsu K, Donnan GA, Howells DW. Good laboratory practice: preventing introduction of bias at the bench. *Stroke* 2009;40:e50–2.
- [39] Maier C, Baron R, Tolle TR, Binder A, Birbaumer N, Birklein F, Gierthmuhlen J, Flor H, Geber C, Hugel V, Krumova EK, Landwehrmeyer GB, Magerl W, Maihofner C, Richter H, Rolke R, Scherrens A, Schwarz A, Sommer C, Tronnier V, Uceyler N, Valet M, Wasner G, Treede RD. Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): somatosensory abnormalities in 1236 patients with different neuropathic pain syndromes. *PAIN* 2010;150:439–50.
- [40] Mandillo S, Tucci V, Holter SM, Meziane H, Al Banchaabouchi M, Kallnik M, Lad HV, Nolan PM, Ouagazzal AM, Coghill EL, Gale K, Golinzi E, Jacquot S, Krezel W, Parker A, Riet F, Schneider I, Marazziti D, Auwerx J, Brown SDM, Chambon P, Rosenthal N, Tocchini-Valentini G, Wurst W. Reliability, robustness, and reproducibility in mouse behavioral phenotyping: a cross-laboratory study. *Physiol Genomics* 2008;34:243–55.
- [41] Miletic T, Kovacevic-Jovanovic V, Stanojevic S, Vujic V, Kosec D, Mitic K, Dimitrijevic M. Strain differences and the role for HSP47 and HSP70 in adjuvant arthritis in rats. *Scand J Immunol* 2006;64:623–32.
- [42] Najj-Esfahani H, Vaseghi G, Safaeian L, Pilehvarian AA, Abed A, Rafieian-Kopaei M. Gender differences in a mouse model of chemotherapy-induced neuropathic pain. *Lab Anim* 2016;50:15–20.
- [43] National Centre for the Replacement, Refinement and reduction of animals in research. ARRIVE guidelines. Available at: www.nc3rs.org.uk/arrive-guidelines.
- [44] National Institute of Neurological Disorders and Stroke. Improving the quality of NINDS-supported preclinical and clinical research through rigorous study design and transparent reporting. 2015 Available at: http://www.ninds.nih.gov/funding/transparency_in_reporting_guidance.pdf. Accessed October 22, 2015.
- [45] Negus SS, Bilsky EJ, Do Carmo GP, Stevenson GW. Rationale and methods for assessment of pain-depressed behavior in preclinical assays of pain and analgesia. *Methods Mol Biol* 2010;617:79–91.
- [46] Okun A, DeFelice M, Eyde N, Ren J, Mercado R, King T, Porreca F. Transient inflammation-induced ongoing pain is driven by TRPV1 sensitive afferents. *Mol Pain* 2011;7:4.
- [47] Parent AJ, Beaudet N, Beaudry H, Bergeron J, Berube P, Drolet G, Sarret P, Gendron L. Increased anxiety-like behaviors in rats experiencing chronic inflammatory pain. *Behav Brain Res* 2012;229:160–7.
- [48] Percie du Sert N, Rice AS. Improving the translation of analgesic drugs to the clinic: animal models of neuropathic pain. *Br J Pharmacol* 2014;171:2951–63.
- [49] Pereira A, Chappell A, Dethy J, Hoeck H, Arendt-Nielsen L, Verfaillie S, Boulanger B, Jullion A, Johnson M, McNearney T. A Proof-of-Concept (POC) Study Including Experimental Pain Models (EPMs) to assess the effects of a CB2 agonist (LY2828360) in the treatment of patients with Osteoarthritic (OA) knee pain. *Clin Pharmacol Ther* 2013;93:S56–7.
- [50] Peters SM, Pothuizen HHJ, Spruijt BM. Ethological concepts enhance the translational value of animal models. *Eur J Pharmacol* 2015;759:42–50.
- [51] Pfau DB, Krumova EK, Treede RD, Baron R, Toelle T, Birklein F, Eich W, Geber C, Gerhardt A, Weiss T, Magerl W, Maier C. Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): reference data for the trunk and application in patients with chronic postherpetic neuralgia. *PAIN* 2014;155:1002–15.
- [52] Phillips TJ, Brown M, Ramirez JD, Perkins J, Woldeamanuel YW, Williams AC, Orenco C, Bennett DL, Bodi I, Cox S, Maier C, Krumova EK, Rice AS. Sensory, psychological, and metabolic dysfunction in HIV-associated peripheral neuropathy: a cross-sectional deep profiling study. *PAIN* 2014;155:1846–60.
- [53] Phillips TJ, Cherry CL, Cox S, Marshall SJ, Rice AS. Pharmacological treatment of painful HIV-associated sensory neuropathy: a systematic review and meta-analysis of randomised controlled trials. *PLoS One* 2010;5:e14433.
- [54] Prinz F, Schlange T, Asadullah K. Believe it or not: how much can we rely on published data on potential drug targets? *Nat Rev Drug Discov* 2011;10:712–U81.
- [55] Raising the standards. *Nature* 2009;459:1033–4.
- [56] Rakel B, Vance C, Zimmerman MB, Petsas-Blodgett N, Amendola A, Sluka KA. Mechanical hyperalgesia and reduced quality of life occur in people with mild knee osteoarthritis pain. *Clin J Pain* 2015;31:315–22.
- [57] Rice AS, Cimino-Brown D, Eisenach JC, Kontinen VK, Lacroix-Fralish ML, Machin I, Preclinical Pain C, Mogil JS, Stohr T. Animal models and the prediction of efficacy in clinical trials of analgesic drugs: a critical appraisal and call for uniform reporting standards. *PAIN* 2008;139:243–7.
- [58] Rice ASC, Morland R, Huang W, Currie GL, Sena ES, Macleod MR. Transparency in the reporting of in vivo pre-clinical pain research: the relevance and implications of the ARRIVE (Animal Research: reporting in Vivo Experiments) guidelines. *Scand J Pain* 2013;4:58–62.
- [59] Richter SH, Garner JP, Auer C, Kunert J, Wurbel H. Systematic variation improves reproducibility of animal experiments. *Nat Methods* 2010;7:167–8.
- [60] Roeska K, Doods H, Arndt K, Treede RD, Ceci A. Anxiety-like behaviour in rats with mononeuropathy is reduced by the analgesic drugs morphine and gabapentin. *PAIN* 2008;139:349–57.
- [61] Rutten K, Robens A, Read SJ, Christoph T. Pharmacological validation of a refined burrowing paradigm for prediction of analgesic efficacy in a rat model of sub-chronic knee joint inflammation. *Eur J Pain* 2014;18:213–22.
- [62] Rutten K, Schiene K, Robens A, Leipelt A, Pasqualon T, Read SJ, Christoph T. Burrowing as a non-reflex behavioural readout for analgesic action in a rat model of sub-chronic knee joint inflammation. *Eur J Pain* 2014;18:204–12.
- [63] Schaap MW, van Oostrom H, Doornenbal A, van 't Klooster J, Baars AM, Arndt SS, Hellebrekers LJ. Nociception and conditioned fear in rats: strains matter. *PLoS One* 2013;8:e83339.
- [64] Serethy M, Currie GL, Sena ES, Ramnarine S, Grant R, MacLeod MR, Colvin LA, Fallon M. Incidence, prevalence, and predictors of chemotherapy-induced peripheral neuropathy: a systematic review and meta-analysis. *PAIN* 2014;155:2461–70.
- [65] Sindrup SH, Graf A, Sfikas N. The NK1-receptor antagonist TKA731 in painful diabetic neuropathy: a randomised, controlled trial. *Eur J Pain* 2006;10:567–71.
- [66] Sorge RE, Mapplebeck JC, Rosen S, Beggs S, Taves S, Alexander JK, Martin LJ, Austin JS, Sotocinal SG, Chen D, Yang M, Shi XQ, Huang H, Pillion NJ, Bilan PJ, Tu Y, Klip A, Ji RR, Zhang J, Salter MW, Mogil JS. Different immune cells mediate mechanical pain hypersensitivity in male and female mice. *Nat Neurosci* 2015;18:1081–3.
- [67] Sorge RE, Martin LJ, Isbester KA, Sotocinal SG, Rosen S, Tuttle AH, Wieskopf JS, Acland EL, Dokova A, Kadoura B, Leger P, Mapplebeck JC, McPhail M, Delaney A, Wigerblad G, Schumann AP, Quinn T, Frasneli J, Svensson CI, Sternberg WF, Mogil JS. Olfactory exposure to males, including men, causes stress and related analgesia in rodents. *Nat Methods* 2014;11:629–32.
- [68] Spruijt BM, Peters SM, de Heer RC, Pothuizen HHJ, van der Harst JE. Reproducibility and relevance of future behavioral sciences should benefit from a cross fertilization of past recommendations and today's technology: "Back to the future". *J Neurosci Methods* 2014;234:2–12.
- [69] Thut PD, Hermansteyne TO, Flake NM, Gold MS. An operant conditioning model to assess changes in feeding behavior associated with temporomandibular joint inflammation in the rat. *J Orofac Pain* 2007;21:7–18.
- [70] Turner L, Shamseer L, Altman DG, Weeks L, Peters J, Kober T, Dias S, Schulz KF, Plint AC, Moher D. Consolidated standards of reporting trials (CONSORT) and the completeness of reporting of randomised controlled trials (RCTs) published in medical journals. *Cochrane Database Syst Rev* 2012;11:MR000030.
- [71] Wallace VC, Blackbeard J, Pheby T, Segerdahl AR, Davies M, Hasnie F, Hall S, McMahon SB, Rice AS. Pharmacological, behavioural and mechanistic analysis of HIV-1 gp120 induced painful neuropathy. *PAIN* 2007;133:47–63.
- [72] Wallace VC, Blackbeard J, Segerdahl AR, Hasnie F, Pheby T, McMahon SB, Rice AS. Characterization of rodent models of HIV-gp120 and anti-retroviral-associated neuropathic pain. *Brain* 2007;130:2688–702.
- [73] Wallace VC, Segerdahl AR, Blackbeard J, Pheby T, Rice AS. Anxiety-like behaviour is attenuated by gabapentin, morphine and diazepam in a rodent model of HIV anti-retroviral-associated neuropathic pain. *Neurosci Lett* 2008;448:153–6.
- [74] Whittaker AL, Lymn KA, Nicholson A, Howarth GS. The assessment of general well-being using spontaneous burrowing behaviour in a short-term model of chemotherapy-induced mucositis in the rat. *Lab Anim* 2015;49:30–9.
- [75] Woolf CJ. Overcoming obstacles to developing new analgesics. *Nat Med* 2010;16:1241–7.
- [76] Zimmermann M. Ethical guidelines for investigations of experimental pain in conscious animals. *PAIN* 1983;16:109–10.