

# Rolling of grains to prevent winter kill damage

## Problem

Extreme frost in cold winters can lead to soil movements that cause tears in plant roots and the hypocotyls of the grain. This makes growth more difficult in spring and makes the grain very susceptible to a lack of moisture up until the 3-leaves growth stage. This results in crops with gaps.

## Solution

Rolling the grains in spring reconnects soil crumbs with the soil and supports the soil water capillarity and water availability in the topsoil.

## Outcome

The pressure of the roller stimulates not only the formation of roots and improves access to water, but stimulates also the tiller of the grain increasing its stability and thus reducing the risk of falling over.

## Applicability box

### Theme

Crop-specific measure

### Geographical coverage

In regions of wheat and other winter cereal cultivation

### Application time

After the last frost in spring when grains start to shoot.

### Required time

One pass with a roller.

### Period of impact

Current crop

### Equipment

Corrugated roller (cambridge roller) or a smooth roller

### Best in

Wheat and other winter grains.

## Practical recommendation

- Rolling requires dry soil in order to prevent the soil from sticking to the roller and ripping out the plants (especially when using smooth rollers).
- The corrugated roller must not damage the plants too much. Take special care when using sharp-edged rollers.
- Every mechanical interference puts the plants under stress. For this reason, it is advised to go at a maximum rolling speed of 5 km/h.

## Notes

- Even in densely sown or strongly fertilized spelt crops, the rolling is recommended. It weakens the main shoot and increases growth on side shoots, which improves the stability of the plants at the expense of longitudinal growth. Spelt is often rolled later (up until the 1-node stage) and more aggressively than winter wheat.



**Figure 1:** Rolling winter wheat (with corrugated roller in the photo) is supporting seed-soil contact and stimulates tillering. (Photos: [www.gut-derenburg.de](http://www.gut-derenburg.de))



Wheat crop about 4 weeks after rolling.

### Practical testing

If this method seems to be suitable for your farm, we recommend that you test it as follows:

1. Make a part of the grain into a testing area and mark it with poles.
2. Apply the new method on the testing area and cultivate the rest of the field as usual.

### Evaluation and sharing of the results

- **Visual evaluation:** Evaluate the tillering, the crop density and the stability of the crop in various stages of development in both areas. Photographs help documenting the results for a later analysis.
- **Quantitative evaluation:** Compare the yields in the testing area with the yields in the rest of the field.

Share your experience with other farmers, advisors and scientists! Use the comment section on the [Farmknowledge platform!](#) If you have any questions concerning the method, please contact the author of the practice abstract.

### Further information

#### Video

- The video "[Wintergetreide walzen](#)" (in German with Dutch subtitles) demonstrates the technique. In the commentary section, there are additional explanations on how to use this method.

#### Links

- Check the [Farmknowledge tool database](#) for more practical recommendations on grain crops.

### About this practice abstract and OK-Net Arable

#### Publishers:

Research Institute of Organic Agriculture (FiBL), Switzerland  
Ackerstrasse 113, Postfach 219, CH-5070 Frick  
Phone +41 62 865 72 72, [info.suisse@fibl.org](mailto:info.suisse@fibl.org), [www.fibl.org](http://www.fibl.org)

IFOAM EU, Rue du Commerce 124, BE-1000 Brussels  
Phone +32 2 280 12 23, [info@ifoam-eu.org](mailto:info@ifoam-eu.org), [www.ifoam-eu.org](http://www.ifoam-eu.org)

**Authors:** Hansueli Dierauer, Franziska Siegrist und Gilles Weidmann (FiBL)

**Contact:** [hansueli.dierauer@fibl.org](mailto:hansueli.dierauer@fibl.org)

**Translation:** Andreas Basler

**Language editing:** Simon Moakes

**Permalink:** [Orgprints.org/31034](https://orgprints.org/31034)

**OK-Net Arable:** This practice abstract was elaborated in the Organic Knowledge Network Arable project. The project is running from March 2015 to February 2018. OK-Net Arable promotes exchange of knowledge

among farmers, farm advisers and scientists with the aim to increase productivity and quality in organic arable cropping all over Europe.

**Project website:** [www.ok-net-arable.eu](http://www.ok-net-arable.eu)

**Project partners:** IFOAM EU Group (project coordinator), BE; Organic Research Centre, UK; Bioland Beratung GmbH, DE; Aarhus University (ICROFS), DK; Associazione Italiana, per l'Agricoltura Biologica (AIAB), IT; European Forum for Agricultural and Rural Advisory Services (EUFRAS); Centro Internazionale di Alti Studi Agronomici Mediterranei - Istituto Agronomico Mediterraneo Di Bari (IAMB), IT; FiBL Projekte GmbH, DE; FiBL Österreich, AT; FiBL Schweiz, CH; Ökológiai Mezőgazdasági Kutatóintézet (ÖMKI), HU; Con Marche Bio, IT; Estonian Organic Farming Foundation, EE; BioForum Vlaanderen, BE; Institut Technique de l'Agriculture Biologique, FR; SEGES, DK; Bioselena, Bulgaria

© 2017

