

The Open Access Israeli Journal of Aquaculture – Bamidgeh

As from **January 2010** The Israeli Journal of Aquaculture - Bamidgeh (IJA) will be published exclusively as **an on-line Open Access (OA)** quarterly accessible by all AquacultureHub (<http://www.aquaculturehub.org>) members and registered individuals and institutions. Please visit our website (<http://siamb.org.il>) for free registration form, further information and instructions.

This transformation from a subscription printed version to an on-line OA journal, aims at supporting the concept that scientific peer-reviewed publications should be made available to all, including those with limited resources. The OA IJA does not enforce author or subscription fees and will endeavor to obtain alternative sources of income to support this policy for as long as possible.

Editor-in-Chief

Dan Mires

Editorial Board

Rina Chakrabarti	Aqua Research Lab, Dept. of Zoology, University of Delhi, India
Angelo Colorni	National Center for Mariculture, IOLR Eilat, Israel
Daniel Golani	The Hebrew University of Jerusalem Jerusalem, Israel
Hillel Gordin	Kibbutz Yotveta, Arava, Israel
Sheenan Harpaz	Agricultural Research Organization Beit Dagan,
Gideon Hulata	Agricultural Research Organization Beit Dagan,
George Wm. Kissil	National Center for Mariculture, IOLR, Eilat, Israel
Ingrid Lupatsch	Swansea University, Singleton Park, Swansea, UK
Spencer Malecha	Dept. of Human Nutrition, Food & Animal Sciences, CTAHR, University of Hawaii
Constantinos Mylonas	Hellenic Center for Marine Research, Crete, Greece
Amos Tandler	National Center for Mariculture, IOLR Eilat, Israel
Emilio Tibaldi	Udine University Udine, Italy
Jaap van Rijn	Faculty of Agriculture, The Hebrew University of Jerusalem, Israel
Zvi Yaron	Dept. of Zoology, Tel Aviv University, Tel Aviv, Israel

Published under auspices of
The Society of Israeli Aquaculture and Marine Biotechnology (SIAMB),
 University of Hawai'i at Mānoa Library

&

University of Hawai'i at Mānoa
Aquaculture Program
 in association with
AquacultureHub

<http://www.aquaculturehub.org>



UNIVERSITY
 of HAWAII[®]
 MĀNOA
 LIBRARY



AquacultureHub.org

AquacultureHub
 educate · learn · share · engage

ISSN 0792 - 156X

© Israeli Journal of Aquaculture - BAMIGDEH.

PUBLISHER:

Israeli Journal of Aquaculture - BAMIGDEH -
 Kibbutz Ein Hamifratz, Mobile Post 25210,
 ISRAEL


Phone: + 972 52 3965809

<http://siamb.org.il>

Copy Editor **Ellen Rosenberg**



The IJA appears exclusively as a peer-reviewed on-line open-access journal at <http://www.siamb.org.il>. To read papers free of charge, please register online at [registration form](#).
Sale of IJA papers is strictly forbidden.



Dietary Histidine Requirement for Juvenile Large Yellow Croaker, *Pseudosciaena crocea* R.

Yan Li ^{1, 2}, Zhenyan Cheng ¹, Kangsen Mai ¹, Qinghui Ai ^{1*},
Jianhua Shi ²

¹The key laboratory of Mariculture, Ministry of Education, Ocean University of China, Qingdao 266003, P. R. China

²Shanghai Fisheries Research Institute, Shanghai 200433, P. R. China

(Received 16.01.13, Accepted 08.08.13)

Key words: large yellow croaker; histidine; growth; requirement

Abstract

Dietary histidine requirements for large yellow croaker (initial average weight, 6.0 ± 0.10 g) were quantified by feeding isonitrogenous (crude protein 44%) and isocaloric (20 KJ/g) amino acid test diets with graded levels of histidine [0.45% (D1), 0.66% (D2), 0.78% (D3), 0.98% (D4), 1.24% (D5) and 1.40% (D6) of dry diet]. Each diet was randomly assigned to triplicate floating sea cages (1.0 × 1.0 × 1.5 m), 60 fish/cage. At the end of the 51 day experiment, the final weight (FW) and weight gain (WG) of large yellow croaker showed a positive correlation to increasing dietary histidine content (up to 0.78%), and thereafter declined. The growth of fish fed the D6 diet was significantly lower than fish maintained on the D2 and D3 diets; however, there were no significant differences in the growth of large yellow croaker among all the dietary treatments except D6. Fish fed the D3 diet had the highest FW and WG. The shift in feed efficiency (FE) values of fish fed D1 to D5 diets increased as dietary histidine content increased, and significantly decreased in fish fed the D6 diet. Based on the second-degree polynomial regression analyses of the growth data, optimum histidine requirement for juvenile large yellow croaker was 8.7 g/kg dry diet, (18.8 g/kg–20.8 g/kg of dietary protein within 95% confidence interval).

* Corresponding author. Qinghui Ai; e-mail: qhai@ouc.edu.cn

Introduction

Histidine (HIS) is an essential amino acid (EAA) for fish growth and affects health if present at sub-optimal or super-optimal levels in the diet (Khan and Abidi, 2009). HIS is a precursor of histamine, carnosine and anserine, which have been found to act as antioxidants in the lens of eyes (Glover and Wood, 2008). Previous studies indicated that Atlantic salmon (*Salmo salar* L.) fed a diet with lower HIS content had significantly higher mean cataract scores than individuals in the treatment with a higher level of HIS (Breck et al. 2005). When fed higher than optimal levels of HIS, the growth of fish such as Indian major carp *C. catla* (Ravi and Devaraj, 1991), *Labeo rohita* (Abidi and Khan, 2004), and African catfish *Clarias gariepinus* (Khan and Abidi, 2009), could be inhibited. The reason for the decline in weight gain has not been established but may be due to fish intolerance to amino acids above the optimum dietary level (Millamena et al., 1999; Khan and Abidi, 2009). Determining the optimal HIS requirement for fish is imperative when formulating high efficiency feed.

Large yellow croaker is an economically important food fish in China, and has been widely cultured in recent years. Trash fish is the main food for culturing this fish in sea cages. However, trash fish cannot meet the nutritional requirements of fish, is difficult to store, and pollutes aquaculture environments. The formulation of a nutritionally adequate and cost-effective feed is most important for the successful culture of large yellow croaker (Zhang et al. 2008). One of the prerequisites for developing high efficiency diets for fish requires extensive knowledge of its nutritional requirements, especially the essential amino acids (EAA) requirements (Mai et al. 2006). So far only two EAA have been tested for large yellow croaker, methionine (Mai et al. 2006) and lysine (Zhang et al. 2008). The purpose of the present study was to quantify the dietary histidine requirement for large yellow croaker.

Materials and Methods

Experimental diets: The basal diets contained 44% protein and were comprised of fish meal, corn gluten meal, gelatin and brewer's yeast as the intact protein source (Table 1).

Table 1. Formulation and composition of the test diets used for the histidine requirement for juvenile large yellow croaker (g/kg dry matter).

Ingredients	Diet no/Supplementation level					
	D1(0.0)	D2(2.0)	D3(4.0)	D4(6.0)	D5(8.0)	D6(10.0)
Fish meal ¹	140.0	140.0	140.0	140.0	140.0	140.0
Corn gluten meal ¹	110.0	110.0	110.0	110.0	110.0	110.0
Gelatin ¹	60.0	60.0	60.0	60.0	60.0	60.0
brewer's yeast ¹	30.0	30.0	30.0	30.0	30.0	30.0
Amino acid mixture ²	193.0	193.0	193.0	193.0	193.0	193.0
Fish oil	50.0	50.0	50.0	50.0	50.0	50.0
Soybean oil	40.0	40.0	40.0	40.0	40.0	40.0
Mineral mixture ³	20.0	20.0	20.0	20.0	20.0	20.0
Vitamin mixture ⁴	20.0	20.0	20.0	20.0	20.0	20.0
Attractant ⁵	3.0	3.0	3.0	3.0	3.0	3.0
Mold inhibitor ⁶	1.0	1.0	1.0	1.0	1.0	1.0
Antioxidant ⁷	0.5	0.5	0.5	0.5	0.5	0.5
Lecithin	30.0	30.0	30.0	30.0	30.0	30.0
Dextrin	260.0	260.0	260.0	260.0	260.0	260.0
Microcrystalline	32.5	32.5	32.5	32.5	32.5	32.5
Histine	0.0	2.0	4.0	6.0	8.0	10.0
Glutamic acid	10.0	8.0	6.0	4.0	2.0	0.0
Composition analysis (g/kg, on a dry weight basis)						
Crude protein	441.0	437.0	440.0	438.0	443.0	440.0
Crude lipid	136.0	138.0	131.0	135.0	140.0	136.0
Total energy (KJ/g)	202.0	206.0	204.0	206.0	206.0	204.0
Histidine	4.5	6.6	7.8	9.8	12.4	14.0

¹Fish meal (white fish meal): obtained from Hangzhou Wensli Biology Science and Technology Corporation (Zhejiang, China), crude protein 675 g/kg dry matter, crude lipid 78 g/kg dry matter; Corn gluten meal: obtained from commercial market, crude protein 655 g/kg dry matter, crude lipid 49 g/kg dry matter; Beer yeast, crude protein 571 g/kg dry matter, crude lipid 35 g/kg dry matter; Gelatin, crude protein 953 g/kg dry matter, crude lipid 20 g/kg dry matter.

²Amino acid mixture (g/kg diet): arginine, 2.23; isoleucine, 0.95; leucine, 1.66; lysine, 2.33; methionine, 0.88; phenylalanine, 0.84; threonine, 1.12; aspartic acid, 2.33; serine, 0.85; alanine, 1.26; glycine, 0.28; cysteine, 0.08; tyrosine, 0.61; glutamic acid, 2.91; valine, 0.95.

³Mineral premix (mg or g/kg diet), NaF, 2 mg; KI, 0.8 mg; CoCl₂·6H₂O (1%), 50 mg; CuSO₄·5H₂O, 10 mg; FeSO₄·H₂O, 80 mg; ZnSO₄·H₂O, 50 mg; MnSO₄·H₂O, 60 mg; MgSO₄·7H₂O, 1,200 mg; Ca (H₂PO₄)₂·H₂O, 3,000 mg; NaCl, 100 mg; Zoelite, 15.448 g.

⁴Vitamin premix (mg or g/kg diet), thiamin, 25 mg; riboflavin, 45 mg; pyridoxine-HCl, 20 mg; vitamin B12, 0.1 mg; vitamin K3, 10 mg; inositol, 800 mg; pantothenic acid, 60 mg; niacin acid, 200 mg; folic acid, 20 mg; biotin, 1.20 mg; retinol acetate, 32 mg; cholecalciferol, 5 mg; alpha-tocopherol, 120 mg; ascorbic acid, 2000 mg; choline chloride, 2000 mg, ethoxyquin, 150 mg, wheat middling, 14.52 g.

⁵Attractant, glycine and betaine.

⁶Mold inhibitor, p-Aminobenzoic acid.

⁷Antioxidant: Ethoxyquin.

The feeds were supplemented with crystalline amino acids premix to simulate the whole body amino acid pattern of large yellow croakers, except for HIS.

Table 2. Amino acid composition of the experimental diets (% dry matter).

AA ¹	FM ¹	CGM ¹	BY ¹	GL ¹	Total ²	Addition ³	43%Whole body ⁴
Arg	0.59	0.19	0.08	0.45	1.31	2.23	3.54
His	0.19	0.11	0.03	0.08	0.42	Varied	0.88
Ile	0.41	0.25	0.07	0.08	0.81	0.95	1.76
Leu	0.67	0.95	0.11	0.16	1.88	1.66	3.54
Lys	0.73	0.10	0.12	0.21	1.15	2.33	3.48
Met	0.27	0.14	0.02	0.02	0.45	0.88	1.33
Phe	0.37	0.37	0.07	0.11	0.92	0.84	1.76
Val	0.50	0.29	0.09	0.14	1.02	0.95	1.97
Thr	0.40	0.20	0.07	0.10	0.78	1.12	1.90
Asp	0.90	0.36	0.15	0.32	1.74	2.33	4.07
Ser	0.42	0.30	0.08	0.18	0.98	0.85	1.83
Gly	0.60	0.16	0.07	1.34	2.18	0.28	2.46
Ala	0.55	0.52	0.11	0.51	1.68	1.26	2.94
Cys	0.06	0.05	0.01	0.00	0.12	0.08	0.20
Tyr	0.32	0.30	0.05	0.02	0.70	0.61	1.31
Glu	1.23	1.29	0.17	0.56	3.26	2.91	6.17
Total addition ⁵						19.30	

¹AA, amino acid; FM, fish meal; CGM, corn gluten meal; BY, brewer's yeast; GL, gelatin.

²Total, the sum of each kind of amino acids in FM, CGM, BY and GL.

³Addition = the content of each kind of amino acids in 43% Whole body protein – the total of each kind of amino acids in all the protein ingredients.

⁴43% Whole body protein, the amino acids content of large yellow croaker in 43% whole body protein.

⁵Total addition, the sum of each kind of amino acids addition.

The basal diet contained 4.5 g/kg HIS and served as control diet. Crystalline L-HIS was added to the basal diet at six graded levels from 0 g/kg to 10 g/kg, and the HIS content in the diets were 4.5 g/kg, 6.6 g/kg, 7.8 g/kg, 9.8 g/kg, 12.4 g/kg and 14.0 g/kg which were analyzed by an amino acids analyzer (Biochrom Ltd®, England). A combination of fish oil, soybean oil and lecithin was used as a source of lipid. Vitamin and mineral premixes were prepared using the method described in Mai et al., (2006). By adjusting dextrin and microcrystalline, cellulose diets were isonitrogenous (crude protein 44%) and isocaloric (20 KJ/g). The diets were marked as D1, D2, D3, D4, D5, and D6, respectively.

The ingredients were ground into fine powder through 320 µm mesh. All protein ingredients, apart from the amino acids were mixed until homogenous. Amino acids were individually weighed and mixed. The attractant, mold inhibitor, antioxidant, mineral, and vitamin mixture, were mixed separately with microcrystalline cellulose, and then mixed with the above protein mixture. The oils and lecithin were blended and added to the mixture. Water was added to produce stiff dough. The pH of diets was adjusted to 7.0-8.0 by gradually adding 6 mol/l NaOH solution to neutralize the acidity of amino acids (Nose et al. 1974). The homogenous dough was passed through a pelletizer (F-26 (II), South China University of Technology) with two different diameter die (1.5mm and 2.5mm). The moist pellets were dried in an oven at 45 °C for 12 h. The dry pellets were crushed and sieved to obtain suitable pellet sizes (1.5 × 2.0 and 2.5 × 3.0 mm), then sealed in bags and stored at -15 °C until used.

Experimental procedure: Fish were obtained from a commercial farm in Ningbo, China. Prior to the feeding trial, the fish were reared in floating sea cages (3.0 × 3.0 × 3.0 m), and fed the control diet (Diet 1) for 2 weeks for acclimation to experimental conditions.

Juvenile large yellow croaker (initial average weight 6.0 ± 0.10 g) were taken from the above acclimated stock and randomly assigned to 18 floating sea cages ($1.0 \times 1.0 \times 1.5$ m) 60 fish/cage. Each diet was randomly assigned to triplicate cages. Fish were fed test diets to satiation twice daily at 05:00 and 17:30. The duration of the feeding trial was 51 days. During the experimental period, the water temperature ranged between 26-32°C, salinity from 26 g/l-30 g/l, and concentration of dissolved oxygen was approximately 7mg/l.

Measurement and analysis: At the end of the experiment, fish were not fed for 24 h prior to harvest. Before weighing and counting, eugenol (1:10000) was used to anesthetize the fish (Shanghai Reagent Corp, China). Five fish from each pooled replicate were sampled and stored frozen (-20°C) for proximate analysis of whole body composition. Proximate analyses on feeds, diets and fish were performed according to the standard methods of AOAC (1995). The samples of whole fish and diets were dried at 105°C to a constant weight to determine % moisture. The dried fish from each replicate were crushed and mixed thoroughly, then sealed in a plastic bag and stored frozen at -20°C until analysis for protein, lipid, and ash content. Crude protein was determined by measuring nitrogen ($\text{N} \times 6.25$) using the Kjeldahl method after acid digestion. Crude lipid was measured by ether extraction using the Soxhlet method. Ash content was placed in a muffle furnace and incinerated at 600°C for 12 h. The feed ingredients or experimental diets were freeze-dried, and then hydrolyzed with 6 N HCl at 110°C for 22 h and analyzed by amino acids analyzer (Biochrom Ltd®, England).

Calculations and statistical analysis: The following variables were calculated: Weight gain (WG) = (final weight – initial weight) / initial weight $\times 100$. Feed efficiency (FE) = wet weight gain in g / dry diet fed in g. All data were subjected to analysis of variance (ANOVA) and regression analysis where appropriate, using SPSS 13.0 for windows. Differences between the means were tested by Tukey's multiple range tests. The level of significance was reported at $P < 0.05$.

Results

Growth performance: Dietary HIS significantly affected the growth of large yellow croaker ($P < 0.05$). Final weight (FW), and weight gain (WG) (%) of fish ranged from 11.8g (96.3%) to 13.5g (125.0%) as presented in Table 3.

Table 3. Effects of dietary histidine on final weight (FW), weight gain (WG), feed efficiency (FE) and survival of juvenile large yellow croaker (*Pseudosciaena crocea*)¹.

Diet no. (HIS content g/kg dry diet)	FW ²	WG (%) ³	FE ⁴	Survival (%)	
D1(4.5)	12.4ab	106.7ab	0.43ab	100.0	¹ Values are means of three replicate groups (n = 3). Means with different letter in the same column differ significantly ($P < 0.05$). ² FW: final weight ³ WG: weight gain ⁴ FE: feed efficiency ⁵ S.E.M.: standard error of means ⁶ ANOVA: one-way analysis of variance
D2(6.6)	13.2a	120.2a	0.41ab	100.0	
D3(7.8)	13.5a	125.0a	0.45ab	100.0	
D4(9.8)	12.9ab	115.5ab	0.46a	99.4	
D5(12.4)	12.9ab	115.5ab	0.49a	100.0	
D6(14.0)	11.8b	96.3b	0.32b	100.0	
Pooled S.E.M. ⁵	0.20	2.82	0.02	0.10	
ANOVA ⁶					
F-value	4.648	4.481	3.876	1.000	
P-value	0.014	0.016	0.025	0.458	

Initially FW and WG showed increasing trends with increasing dietary HIS content up to 0.78%, but thereafter declined. The growth of fish was significantly lower in fish fed D6 with the highest HIS content (14.0g/kg dry diet) than in fish fed the D2 (6.6 g/kg) and D3 (7.8 g/kg) dry diet; however, there were no significant differences in the growth of the fish between all the dietary treatments except D6. Fish fed D3 with 0.78% HIS content had the highest FW and WG. The shift in feed efficiency (FE) values of fish fed D1 to D5 increased with increasing dietary HIS content, and significantly decreased when fish were fed with D6. No significant differences in survival were found between the different dietary treatments.

For WG (y) in relation to dietary HIS levels (x) to second-degree polynomial regression analysis, the equation was $y = -5.864x^2 + 63.907x - 24.454$. The optimum HIS was the 8.7 g/kg diet (Fig.1), corresponding to 19.7 g/kg of dietary protein for juvenile large yellow croaker.




Fig.1. Effects of dietary histidine on weight gain of juvenile large yellow croaker (*Pseudosciaena crocea*) fed experimental diets. Each point represents the mean of three groups of fish within a treatment with all the survival fish per group. Requirements derived with the second-degree polynomial regression analyses for weight gain is 8.7 g/kg diet (19.8 g/kg dietary protein).

Whole body composition: The body composition of juvenile large yellow croaker was not significantly affected by dietary HIS content. The range of body moisture of large yellow croaker was 788.2 g/kg - 801.6 g/kg; crude protein was 140.0 g/kg - 146.6 g/kg; crude lipid was 22.7 g/kg - 31.4 g/kg; ash was 40.0 g/kg - 41.2 g/kg (Table 4).

Table 4. Effect of dietary histidine on body composition of juvenile large yellow croaker (*Pseudosciaena crocea*) fed experimental diets ¹.

Diet no (histidine Content g/kg dry diet)	Whole - Body composition (g/kg fresh weight)			
	Moisture	Crude protein	Crude lipid	Ash
D1 (4.5)	788.2	144.0	25.3	40.6
D2 (6.6)	800.7	141.3	25.4	40.0
D3 (7.8)	801.6	140.0	22.8	40.4
D4 (9.8)	790.7	146.6	31.4	40.3
D5 (12.4)	801.5	143.0	22.7	41.2
D6 (14.0)	796.7	143.3	26.4	41.4
Pooled S.E.M. ²	2.06	1.07	1.09	0.32
ANOVA ³				
F-value	1.573	0.688	1.937	0.330
P-value	0.241	0.642	0.168	0.885

¹Values are means of three replicate groups (n = 3)

²S.E.M.: standard error of means

³ANOVA: one-way analysis of variance.

Discussion

The data available on the dietary HIS requirements for fish varied between 0.9% and 2.5% of the dietary protein (Abidi and Khan, 2004; Khan and Abidi, 2009). In this study, the optimum dietary HIS requirement for juvenile large yellow croaker was estimated to be 1.97% of dietary protein which falls in the above range. This is probably due to the different requirements of the tested fish. The results were higher than the requirements reported for African catfish fry, *Clarias gariepinus* 1.0-1.05% of dietary protein (Khan and Abidi, 2009), Turbot, *Psetta maxima* 1.5% of dietary protein (Kaushik, 1998), Rainbow trout, *Oncorhynchus mykiss* 1.6% of dietary protein (Ogino, 1980), gilthead sea bream, *Sparus auratus* 1.7% of dietary protein (Kaushik, 1998), European sea bass, *Dicentrarchus labrax* 1.6% of dietary protein (Kaushik, 1998), lower than Japanese eel, *A. japonica* 2.1% of dietary protein (Nose, 1979).

Many previous studies which estimated amino acid requirements for fish also used the second order polynomial regression model (Tibaldi and Tulli, 1999; Ahmed and Khan, 2004; Mai et al., 2006; Khan and Abidi, 2007). Before estimating the amino acid requirements, the significance value (P) and the coefficient of the estimation (R²) of

different models were compared. The second order polynomial regression model better describes the relationship between dietary HIS levels and WG.

An attempt was made with the basal diet of this experiment to lower the content of HIS. However, while still trying to maintain 42%-44% dietary proteins, many ingredients in the diet did not decrease the HIS content of the basal diet. Thus the values of HIS in diets (0.45% dry diet) were higher than expected. The growth of fish fed the lowest dietary HIS content was not inhibited. However excessive HIS did inhibit the growth of fish in this study. This phenomenon was also found in African catfish *Clarias gariepinus* (Khan and Abidi, 2009), Indian major carp, *C. catla* (Ravi and Devaraj, 1991), and rohu, *Labeo rohita* (Murthy and Varghese, 1995). This may have occurred because excessive HIS disrupted the balance of dietary amino acids, leading to toxicity in the tissues, (Mertz, 1972) or extensive necrosis in the epithelial cells of the hepatopancreas (Recodo, 1991) thereby influencing fish growth.

In conclusion, the optimum HIS requirement for juvenile large yellow croaker was found to be 8.7 g/kg dry diet, corresponding to 18.8 g/kg–20.8 g/kg of dietary protein within 95% confidence interval.

Acknowledgement

This study was supported by the National Key Technologies R&D Program for the 15th Five-year Plan of China (Grant No.: 2004BA526B-06) and Program for New Century Excellent Talents in University (NCET-07-0776). We thank W.B. Zhang, X.J. Wang, H.M. Ma, J.K. Shentu and Q.Y. Duan for their assistance in the study.

Reference

- Abidi S.F. and M.A. Khan**, 2004. Dietary histidine requirements of fingerling Indian major carp, *Labeo rohita* (Hamilton). *Isr. J. Aquacult. - Bamidgeh*, 56(3), 2004, pp.200-208.
- Ahmed I. and M.A. Khan**, 2004. Dietary lysine requirement of fingerling Indian major carp, *Cirrhinus mrigala* (Hamilton). *Aquaculture*, 235, 2004, 499-511.
- Ahmed I. and M.A. Khan**, 2005. Dietary histidine requirement of fingerling Indian major carp, *Cirrhinus mrigala* (Hamilton). *Aquaculture Nutrition*, 11, 2005, 359-366.
- Association of Official Analytical Chemists (AOAC)**, 1995. Official Methods of Analysis of Official Analytical Chemists International, 16th edn. Association of Official Analytical Chemists. Arlington, VA.
- Borlongan I.G. and R.M. Coloso**, 1993. Requirements of juvenile milkfish (*Chanos chanos* Forsskal) for essential amino acids. *J. Nutrition*, 123, 1993, pp.125-132.
- Breck O., Bjerkas E., Sanderson J., Waagbo R. and P. Campbell**, 2005. Dietary histidine acts lens protein turnover and synthesis of N-actylhistidine in Atlantic salmon (*Salmo salar* L.) undergoing parr-smolt transformation. *Aquacult. Nutr.*, 11, 2005, pp.321-332.
- Glover C.N. and C.M. Wood**, 2008. Histidine absorption across apical surface of freshwater rainbow trout intestine: mechanistic characterization and the influence of copper. *J. Membrane Biol.*, 217:87-95.
- Kaushik S.J.**, 1998. Whole body amino acid composition of European seabass (*Dicentrarchus labrax*), gilthead seabream (*Sparus aurata*) and turbot (*Psetta maxima*) with an estimation of their IAA requirement profiles. *Aquat. Living Resour.*, 11(5):355-358.
- Khan M.A. and S.F. Abidi**, 2007. Total aromatic amino acid requirement of Indian major carp *Labeo rohita* (Hamilton) fry. *Aquaculture*, 267:111-118.
- Khan M.A. and S.F. Abidi**, 2009. Optimum histidine requirement of fry African catfish, *Clarias gariepinus* (Burchell). *Aquac. Res.*, 40:1000-1010.
- Mai K.S., Wan J.L., Ai Q.H., Xu W., Liufu Z.G., Zhang L., Zhang C.X. and H.T. Li**, 2006. Dietary methionine requirement of large yellow croaker, *Pseudosciaena crocea* R. *Aquaculture*, 253:564-572.
- Mertz E.T.**, 1972. The protein and amino acid needs. In: Halver, J.E. Ed., *Fish Nutrition*. Academic Press, New York. 105-143pp.

- Millamena O.M., Teruel M.B., Kanazawa A. and S. Teshima,** 1999. Quantitative dietary requirements of postlarval tiger shrimp, *Penaeus monodon*, for histidine, isoleucine, leucine, phenylalanine and tryptophan. *Aquaculture*, 179:169-179.
- Murthy H.Y. and T.J. Varghese,** 1995. Arginine and histidine requirements of the Indian major carp, *Labeo rohita* (Hamilton). *Aquacult. Nutr.*, 1:235-239.
- Nose T.,** 1979. Summary report on the requirements of essential amino acids for carp. In: Halver, J.E., Tiews, K. (Eds.), *Finfish Nutrition and Fishfeed Technology*, vol. I. Heenemann, Berlin, Federal Republic of Germany. 145-156pp.
- Nose T., Arai S., Lee D. L. and Y. Hashimoto,** 1974. A note on amino acids essential for growth of young carp. *Bull. Jpn. Soc. Sci. Fish.*, 40:903-908.
- Ogino C.,** 1980. Requirements of carp and rainbow trout for essential amino acids. *Bull. Jpn. Soc. Sci. Fish.*, 46:171-174.
- Quam D.D., Morris J.G. and Q.R. Rogers,** 1987. Histidine requirement of kittens for growth, haematopoiesis and prevention of cataracts. *Brit. J. Nutr.*, 58:521-532.
- Recodo A.G.,** 1991. Histophysiological effects of various histidine levels in the diet on the hepatopancreas of *Penaeus monodon*. Masters thesis. Institute of Biology, College of Science, U.P. Diliman, Quezon City, Philippines. 104 pp.
- Ravi J. and K.V. Devaraj,** 1991. Quantitative essential amino acid requirements for growth of *Catla catla* (Hamilton). *Aquaculture*, 96:281-289.
- Robinson E.H., Wilson R.P. and W.E. Poe,** 1981. Arginine requirement and apparent absence of a lysine-arginine antagonist in fingerling channel catfish. *J. Nutr.* 111:46-52.
- Santiago C.B. and R.T. Lovell,** 1988. Amino acid requirements for growth of Nile tilapia. *J. Nutr.* 118:1540-1546.
- Tibaldi E. and F.Tulli,** 1999. Dietary threonine requirement of juvenile european sea bass *Dicentrarchus labrax*. *Aquaculture*, 175:155-166.
- Unprasert N.G.,** 1994. An evaluation of the use of 'Ideal protein' concept to estimate essential amino acid requirements of Clarias hybrid (*Clarias macrocephalus* × *Clarias gariepinus*). PhD dissertation, Mississippi State University, Mississippi, USA.
- Walton M.J., Adron J.W., Coloso R.M. and C.B. Cowey,** 1986. Dietary requirements of rainbow trout for tryptophan, lysine and arginine as determined by growth and biochemical measurements. *Fish Physiol. Biochem.* 2:161-169.
- Wilson R.P., Allen O.W., Robinson E.H. and W.E. Poe,** 1978. Tryptophan and threonine requirements of fingerling channel catfish. *J. Nutr.* 108:1595-1599.
- Zeitoun I.H, Ullrey D.E and W.T. Magee,** 1976. Quantifying nutrient requirements of fish. *J. Fish Res. Board Can.*, 33:167-172.
- Zhang C.X., Ai Q.H., Mai K.S., Tan B.P., Li H.T. and L. Zhang,** 2008. Dietary lysine requirement of large yellow croaker, *Pseudosciaena crocea* R., *Aquaculture*, 283:123-127.
- Zhang C.X., Mai K.S., Ai Q.H., Zhang W.B., Duan Q.Y., Tan B.P., Ma H.M, Xu W., Liufu Z.G. and X.J. Wang,** 2006. Dietary phosphorus requirement of juvenile Japanese seabass, *Lateolabrax japonicas*. *Aquaculture*, 255:201-209.