

The Open Access Israeli Journal of Aquaculture – Bamidgeh

As from **January 2010** The Israeli Journal of Aquaculture - Bamidgeh (IJA) will be published exclusively as **an on-line Open Access (OA)** quarterly accessible by all AquacultureHub (<http://www.aquaculturehub.org>) members and registered individuals and institutions. Please visit our website (<http://siamb.org.il>) for free registration form, further information and instructions.

This transformation from a subscription printed version to an on-line OA journal, aims at supporting the concept that scientific peer-reviewed publications should be made available to all, including those with limited resources. The OA IJA does not enforce author or subscription fees and will endeavor to obtain alternative sources of income to support this policy for as long as possible.

Editor-in-Chief

Dan Mires

Editorial Board

Rina Chakrabarti	Aqua Research Lab, Dept. of Zoology, University of Delhi, India
Angelo Colorni	National Center for Mariculture, IOLR Eilat, Israel
Daniel Golani	The Hebrew University of Jerusalem Jerusalem, Israel
Hillel Gordin	Kibbutz Yotveta, Arava, Israel
Sheenan Harpaz	Agricultural Research Organization Beit Dagan,
Gideon Hulata	Agricultural Research Organization Beit Dagan,
George Wm. Kissil	National Center for Mariculture, IOLR, Eilat, Israel
Ingrid Lupatsch	Swansea University, Singleton Park, Swansea, UK
Spencer Malecha	Dept. of Human Nutrition, Food & Animal Sciences, CTAHR, University of Hawaii
Constantinos Mylonas	Hellenic Center for Marine Research, Crete, Greece
Amos Tandler	National Center for Mariculture, IOLR Eilat, Israel
Emilio Tibaldi	Udine University Udine, Italy
Jaap van Rijn	Faculty of Agriculture, The Hebrew University of Jerusalem, Israel
Zvi Yaron	Dept. of Zoology, Tel Aviv University, Tel Aviv, Israel

Published under auspices of
The Society of Israeli Aquaculture and Marine Biotechnology (SIAMB),
 University of Hawai'i at Mānoa Library

&

University of Hawai'i at Mānoa
Aquaculture Program
 in association with
AquacultureHub

<http://www.aquaculturehub.org>



UNIVERSITY
 of HAWAII[®]
 MĀNOA
 LIBRARY



AquacultureHub.org

AquacultureHub
 educate • learn • share • engage

ISSN 0792 - 156X

© Israeli Journal of Aquaculture - BAMIGDEH.

PUBLISHER:

Israeli Journal of Aquaculture - BAMIGDEH -
 Kibbutz Ein Hamifratz, Mobile Post 25210,
 ISRAEL

Phone: + 972 52 3965809

<http://siamb.org.il>

Copy Editor **Ellen Rosenberg**



The IJA appears exclusively as a peer-reviewed on-line open-access journal at <http://www.siamb.org.il>. To read papers free of charge, please register online at [registration form](#).
Sale of IJA papers is strictly forbidden.



Use of Herbal Extracts for Controlling Reproduction in Tilapia Culture: Trends and Prospects - a Review

Ndakalimwe Naftal Gabriel^{1,3}, Jun Qiang^{1,2}, Mathew D. Kpundeh¹,
Pao Xu^{1,2*}

¹Wuxi Fisheries College, Nanjing Agricultural University, Wuxi 214081, China

²China Key Laboratory of Freshwater Fisheries and Germplasm Resources Utilization, Ministry of Agriculture, Freshwater Fisheries Research Center, Chinese Academy of Fishery Sciences, Wuxi 214081, Jiangsu, China

³Directorate of Aquaculture, Ministry of Fisheries and Marine Resources, Namibia

(Received 4.1.2015, Accepted 28.3.2015)

Keywords: in vivo administration, monosex methods, phytoestrogens, reproductive physiology, reproduction control, tilapia culture

Abstract

The use of synthetic chemicals in the production of food for human consumption has been condemned by many nations, due to their potential health and environmental hazards. In tilapia farming in particular, synthetic sex reversal hormones have been commonly used to produce all male tilapia progenies. Recently, several herbal extracts have been reported to possess estrogenic properties, some of which are capable of inducing antifertility, abortifacient, and sex inversion in animals. Hence, herbal extracts could be used as safe alternative agents to control precocious tilapia maturity and prolific breeding in production systems. Dietary *Basella alba*, *Quillaja saponaria*, *Trigonella foenum-graecum*, *Glycine max* and *Tribulus terrestris* extracts have been reported to shift tilapia sex ratio in favor of males. Moreover, *Moringa oleifera*, *Carica papaya*, *Aloe vera*, *Azadirachta indica*, and *Hibiscus rosasinesis* extracts demonstrated a direct effect on gonad morphology and delayed maturation in some tilapia species. However, there are limitations, which make it hard for this novel development to progress from experimental trials to widespread adoption by farmers because of lack of adequate knowledge on phytoestrogen extraction methods, their extract concentrations, and identification methods. Hence, there is a need for more research to standardize every aspect concerning the use of phytoestrogens in tilapia culture. The aim of this paper is to provide an overview of the available studies on the use of herbal extracts as potential alternatives to control tilapia reproduction in aquaculture, while also discussing limitations in the existing knowledge and finding a way forward.

Corresponding author: E-mail: xup@ffrc.cn

Introduction

About half a decade ago, 14.3 % of the world’s human population (6.7 billion) was declared undernourished, and the global population is expected to reach 9.3 billion by 2050 (UN, 2010). Agriculture, livestock and fisheries (inland and marine) which are the most important food producing sectors, are close to reaching their maximum capacity in terms of sustainable production (FAO, 2006). Thus the current food shortage is expected to be a long-term problem, unless major strides are taken towards meeting future demands for food. Aquaculture is considered one of the fastest growing animal food producing sectors globally, with a continuous annual growth rate of 8%, increasing production from 16.8 million tons in 1990 to 78.88 million tons in 2010 (FAO, 2012). Apart from being one of the few suitable options for feeding the growing global population with cheap and beneficial animal proteins, aquaculture is also regarded as a way of preserving wild fish stock, providing employment, and is an essential component of integrated rural development (Srinath et al., 2000).

In freshwater aquaculture, tilapia species are widely acknowledged as one of the most important internationally traded fish (FAO/GLOBEFISH, 2013), with a significant increase in production from 383,654 metric tons (mt) in 1990 to 3,500,000 mt in 2011 (Fitzsimmons, 2012) (Fig. 3). In addition to new intensive farming technologies, tilapia possess an impressive range of qualities that have made them very suitable for aquaculture in the 21st century. Their attributes include: their ability to reproduce easily in captivity; rapid growth, reaching their marketable size (in about six months); tolerance of a wide range of environmental conditions; resistance to stress and disease; occupying a low trophic level, with versatile feeding habits; acceptance of artificial feed immediately after yolk-sac absorption; adaptation to a variety of culture systems; and marketability: nutritious and palatable. (Teichert-Codding et al., 1992; Altun et al., 2006; Fitzsimmons et al., 2011; Ghosa and Chakraborty, 2014). Among tilapia aquaculture species, Nile tilapia (*Oreochromis niloticus*) has long been responsible for the significant increase in global production from freshwater aquaculture (FAO, 2002) (Fig.1),




Fig. 1. Percent share of global tilapia aquaculture production by species (Data source; FAO, 2002).

This has been reported to account for a production rate above 3 million mt per year (Fitzsimmons et al., 2012) (Fig.3). On a global scale, China has long been the largest tilapia producer (Fitzsimmons et al., 2012; Tveteras, 2013), with production estimated at 1.3 million mt in 2011, about 40% of the global production (Tveteras, 2013) (Fig.2).




Fig. 2. Countries that significantly contributed to the world tilapia production from 2011- 2013 (adapted from Tveteras, 2013).

Regardless of the widely reported global tilapia aquaculture production and progress over the years, there are several challenges commonly associated with their reproductive ability that inhibit their full aquaculture potential. The most common setback in tilapia aquaculture is their precocious maturity and frequent breeding behavior (Mires, 1995).




Fig. 3: Tilapia global aquaculture production from source; FAO, 2002).

Thus various techniques to control unwanted tilapia reproduction have been developed. Previous methods developed to control undesirable reproduction in tilapia production systems have led to the farming of all male progenies that grow faster and avoid the stunting of cultured populations (Guerrero, 1975; Mires, 1987; Jiménez-Badillo, 2006). An underlying reason for superior growth under these conditions is that they expend more energy on growth rather

than on reproductive behavior (Phelps and Popma, 2000). Methods of producing all male tilapia include manual separation of sexes, hybridization, hormonal sex reversal, and genetic manipulation (Mires, 1977; Mair and Little, 1991), of which, the latter two methods are widely applied in commercial fish farming (Phelps and Poma, 2000; Wassermann and Afonso, 2002). To date, existing mono-sex production methods have suffered from limitations, rendering them ineffective, laborious, unsustainable, and to a large extent inaccessible to fish farmers, especially, rural small scale fish farmers (Hulata et al., 2004). It is important therefore that alternative methods of controlling precocious maturation and prolific breeding in tilapia culture and other aquaculture species be developed, to ensure a more cost effective and sustainable aquaculture industry.

Several studies have highlighted the benefits of medicinal herbs as potentially suitable alternatives to chemicals and drugs (Eckstein and Spira, 1965; Citarasu, 2010; Ganzera et al., 2001; Francis et al., 2002; Dabrowski et al., 2005; Dongmeza et al., 2006; Green and Kelly, 2009; Bai et al., 2012; Felicitta et al., 2013; Hu et al., 2014; Ghosal and Chakraborty, 2014; Gabriel et al., 2015). In addition to anti-stress, growth promotion, appetite stimulation, tonic and immune-stimulation, and antimicrobial properties (Citarasu, 2010; Chakraborty and Hancz, 2011; Ghosal and Chakraborty, 2014), medicinal herbs have also been reported to possess antifertility and abortifacient properties when orally administered to animals (Obaroh and Chionye-Nzeth, 2011). Herbal compounds (phytoestrogens) such as isoflavonoids, flavonoids, lignans, and coumestans among others, are believed to mimic, or function as sex hormones, and are able to block biosynthesis and estrogen by acting as aromatase inhibitors and antagonists to nuclear estrogen receptors in gonad germ cells (Das et al., 2012), and may therefore potentially induce sex reversal, or to delay maturity in fish. The use of medicinal herbs at all levels of aquaculture may not only improve production, but could also enhance the safety and quality of aquatic products, thus increasing the use of these products across the globe.

The aim and scope of this article is: (1) to provide an overview of the available studies on the use of herbal extracts as potential alternatives in the control of tilapia reproduction in aquaculture, (2) to review the reproductive physiology of tilapia and factors affecting it, (3) to briefly discuss early-developed methods of producing monosex tilapia population in aquaculture, and (4) to increase knowledge and prospective use of herbal extracts as an alternative way to produce monosex tilapia in aquaculture.

Tilapia reproductive behaviors and physiology

The ability of all living organisms to reproduce (sexually or asexually) is a pre-requisite for their existence. Understanding reproductive behavior and physiology of animals may be important in the development of sustainable food production, conservation of biodiversity, habitat protection, and establishing restoration initiatives. Many studies have dealt with reproductive behavior and physiology of teleost fish. Besides eco-morphological characteristics, tilapia genera (*Oreochromis*, *Sarotherodon* and *Tilapia*) are

classified mainly based on their reproductive behavior (Trewavas, 1982). While some fish change sex during their lifetime, tilapia species are described as gonochoristics, where individuals sexually differentiate into males (pair testes) or females (pair ovaries) and remain the same sex throughout their life (Yamazaki, 1983; Nakumura et al., 1998). As part of their reproductive behavior, all tilapia species exhibit a high degree of parental care, seen in their nest building and substrate spawning nature (Pullin and McConnell, 1982; Trewavas, 1982). There are both differences and similarities in brooding of eggs and fry in tilapia species. *Oreochromis* and *Sarotherodon* species are both mouth brooders, meaning that eggs are fertilized in the nest, and then stored in their parents' mouth for incubation. The eggs are held in the mouth and the fry are held in the mouth for several days after hatching (Nandlal and Pickering, 2004). *Oreochromis* species possess a maternal mouth brooding nature, whereas, *Sarotherodon* species exercise either paternal or bi-parental mouth brooding behavior (Trewavas 1982; Nandlal and Pickering, 2004).

The easy and rapid propagation of tilapias in various tropical and sub-tropical environments make them ideal aquaculture species, however, this judgment is challenged by their reproductive efficiency combined with precocious maturation (as early as 3 months) (Pullin and McConnell, 1982; Phelps and Popma, 2000). In teleost fish and other vertebrates, reproduction is affected by many external factors such as temperature, photoperiod (Taranger et al., 2010; Bairwa et al., 2013), social patterns (Tubert et al., 2012; Maruska and Fernald, 2013) and internal factors, involving neuroendocrine system, (Schulz et al., 2010; Zohar et al., 2010; Bairwa et al., 2013).

The interaction between environmental factors, the neuroendocrine system, and reproduction in fish has not yet been fully explored, however, a considerable number of related studies have been published. Similar to other vertebrates, reproduction in fish depends on the neuroendocrine system, which initiates and controls gametogenesis (formation of female oocyte and male spermatozoon) and steroidogenesis (formation of steroids), through the activation of hypothalamus-pituitary-gonad axis (HPG) or the gonadotropic axis (Levavi-Sivan and Yaron, 1993; Kah et al., 1993; Van der Kraak et al., 1998; Fink, 2000; Tena-Sempere and Huhtaniemi, 2003; Kajimura et al., 2004; Marchesan et al., 2005; Pinilla et al., 2012). The initiation and completion of reproductive activity in teleost fish is controlled by photoperiod, which ensures an appropriate reproduction season favorable for offspring survival (Bromage et al., 2001). In addition, water temperature is believed to modulate HPG and affect the rate of gametogenesis or allow or inhibit gametogenesis to proceed beyond certain stages and/or completion (Mañafios et al., 1997; Prat et al., 1999). Furthermore, water temperature can affect the onset of reproduction or puberty indirectly in fish through its effect on somatic growth and energy storing (Taranger et al., 2010).

In tilapia species, the highest reproductive activity is associated with increasing photoperiod and warmer temperature, while low spawning rates are associated with lower temperatures and shorter photoperiod (Bairwa et al., 2013). One study concluded that a photoperiod of 12h light/dark cycle in Nile tilapia aquaculture ensured maximum fecundity, seed production, and spawning frequency (El-Sayed and Kawanna, 2007), although a minimum temperature range of 20-23^o is reported to be suitable for breeding in most tilapia species (Bairwa et al., 2013). Under subtropical and temperate conditions, where temperature or photoperiods are more variable, a well-defined breeding season needs to be determined for most tilapia species (Bromage et al., 2001; Bairwa et al., 2013).

Methods to control tilapia reproduction in production systems

Wild spawns in culture ponds of tilapia have led to precocious maturity and uncontrolled reproduction in farmed tilapia. This has necessitated the development of various methods to mitigate these behaviors.

The main, method used to control reproduction in tilapia is the culture of all-male tilapia, attained through manual separation of sexes, hybridization, hormone induced sex reversal, and genetic manipulation (androgenesis, gynogenesis, polyploidy and transgenesis) (Eckstein et al., 1965; Guerrero, 1975; Mires, 1977; Lovshin et al., 1990; Mair and Little, 1991; Fortes, 2005). Other methods include intermittent harvest, high stocking density culture, biological control, sterilization, culture in cages (Mair and Little,

1991) and using organic toxicants and/ or other chemicals (Fortes, 2005). Among the methods listed above, hormonal sex reversal has been widely adopted in aquaculture across the globe, and has become a reliable all-male tilapia production technique responsible for the global success of tilapia production over the years (Pandian and Sheela, 1995). Recently, this method has been combined with genetic manipulation (Acosta and Gupta, 2010). However, some of these methods have limitations when implemented outside experimental studies or developmental trials.

Manual separation of sexes

In addition to tilapia species being gonochoristic, they also possess sexual dimorphic characteristics, which make it easy to sort them into males and females. Manual separation of sexes as a method of obtaining all male/monosex tilapia populations is strictly based on separating males from females by visual inspection of external urogenital pores, often with the aid of dye applied (Fortes, 2005; Fuentes-Silva et al., 2013). The genital papilla of male is simple and smaller with two openings; the urogenital opening, where the milt and urine are excreted and the anus, for the discharge of fecal waste, whereas the female has a flatter and larger papilla with three openings; the anus, the urethra (for excretion of urine) and the oviduct, where the eggs pass through. In addition to sexual dimorphism, secondary sex characteristics can also be used to help differentiate sexes in tilapia, for instance, looking at the dorsal and anal fins which are pointed in males but rounded in the females (Chervinski and Rothbard, 1982).

With these methods, sex separation is carried out before fish reach sexual maturity (Mair and Little, 1991), when they are large fingerlings (50- 80g), however the reliability of sexing depends on the skill of the workers, the species to be sorted, and its size (Fortes, 2005). Although this method is feasible, it is tedious, and difficult even for skilled workers to achieve 90 percent accuracy in sexing. Therefore, breeding and reproduction are rarely completely controlled (Mair and Little, 1991; Penman and McAndrew, 2000). Furthermore, this method may impact on economic returns of the fish farmer since in addition to the high cost of skilled laborers, about 40-50 percent of the female fingerlings are normally discarded (Fortes, 2005). Therefore this method is generally applied at subsistence level farming where fish populations are normally small; it may not be useful in commercial farming.

Hybridization

Hybridization, which is mating of genetically different individuals or groups, may involve crosses within a species (also known as line crossing or strain crossing) or crosses between species (Bartley et al., 2001). The rationale of this technique is to produce a hybrid or strain of superior quality than the parent species (Essa and Haroun, 1998). In aquaculture, hybridization is not only used to manipulate sex ratios or produce sterile fish, but also to increase growth rate, improve flesh quality, increase disease resistance, improve environmental tolerance, and improve a variety of other traits to make fish production more profitable (Bartley et al., 2001). Hybridization between tilapia species has attracted a lot of research (Table 1), which has subsequently led to the adoption of some tilapia hybrids (i.e. *Oreochromis aureus* x *O. mossambicus*; and *O. aureus* x *O. niloticus*) in commercial farms (Chapman, 1992).

Despite the fact that hybridization is associated with the production of a high number of male progeny, this development is surrounded by numerous constraints, which make it unsustainable. Some of these constraints include: limited fecundity of parent fish which restrict fry production, difficulty in producing sufficient number of hybrid fry due to spawning incompatibility between parent species (Mires, 1977; Varadaraj and Pandian, 1989), and inasmuch as not all crosses produce 100% males, the hybrids may still require manual separation of sexes or hormone augmentation. In addition, widespread adoption of this method would result in the introgression of tilapia species with deleterious implications for the conservation of tilapia genetic resources (Mair and Little, 1991). More research is needed to either perfect this method or replace it with easily manageable and environmental friendly techniques.

Table 1. Hybridization of some tilapia species and proportion of male progeny produced

Cosses (♂ x ♀)	Males %	References
<i>O. aureus</i> x <i>O. niloticus</i> (Ugadan strain)	96-100	Pruginin, 1967;
<i>O. aureus</i> x <i>O. vulcani</i>	98-100	Pruginin, 1975;
<i>O.hornorum</i> x <i>O. spilurus</i>	100	Hulata et al. 1983
<i>O. macrochir</i> x <i>O. niloticus</i>	100	Wohlfarth et al., 1994
<i>O.urolepsis hormorum</i> x <i>O. nigra</i>	98-100	Wohlfarth et al., 1994
<i>O.urolepsis hormorum</i> x <i>O. vulcani</i>	98-100	Majumdar et al. 1983
<i>O. macrochir</i> x <i>O. mossambicus</i>	100	Majumdar et al., 1983
<i>O.hornorum</i> x <i>O. niloticus</i>	100	Wohlfarth et al., 1994
<i>O.urolepsis hormorum</i> x <i>O. niloticus</i>	100	Wohlfarth et al., 1990
<i>O. aureus</i> x <i>O. niloticus</i> (Stirling strain)	100	Marengoni et al., 1998
<i>O.aureus</i> x <i>O. mossambicus</i>	100	Beadmore et al., 2001
<i>O.hornorum</i> x <i>O. mossambicus</i>	100	Hickling, 1960

Hormonal sex reversal

Constraints associated with manual separation of sexes and hybridization techniques have subsequently led to another technique called hormonal sex reversal. Hormonal sex reversal is the most efficient and commonly used method for mass production of all male tilapia in both small and large scale tilapia production (Pandian and Sheela, 1995; Phelps and Popma, 2000); and the success of global tilapia production is due to this technique.

Tilapia larvae are believed to be sexually undifferentiated up until 2 weeks after hatching, and at this time, larvae produce equal proportions of sex hormones; androgen (male), and estrogen (female) (Fuentes-Silva et al., 2013). Therefore, intervention or augmentation by exogenous steroid hormones such as androgen (male) or estrogen (female) during gonadal development or before sexual differentiation would influence the larvae to become either male or female depending on the hormone applied (Fortes, 2005; Fuentes-Silva et al., 2013). Two synthetic androgen hormones namely methyltestosterone (MT) and ethynyltestosterone have been widely used for masculinizing genotypic female tilapia (Mair and Little, 1991; Phelps and Popma, 2000; Forbes, 2005). Although, there is a wide range of hormonal administration methods, the commonly used treatment is oral administration for varying periods of 18-60 days in tanks or aquaria, with the dosage ranging from 10-60-mg/kg diet which is widely acknowledged as being effective for sex reversal (Guerrero, 1979).

Sex reversal success is achieved when uniform age fry or larvae eat only the hormone treated feed during the period of treatment, thus making it difficult to apply this technique in earthen ponds (Mires, 1995; Phelps and Popma, 2000). While many studies concluded that this technique offers a practical and economic approach for the control of tilapia reproduction (Guerrero, 1979), widespread use of large quantities of sex reversal hormones in hatcheries may pose a health risk to workers, consumers, and the environment (Phelps and Popma, 2000). In addition, hormones may be difficult to obtain in some countries, and hatchery facilities and skilled laborers are required (Forbes, 2005). Thus, this technology may create adverse effects in developing countries such as Sub-Saharan Africa, where aquaculture is still in the infancy stage, where there is poor infrastructure, poor or no protective equipment, and no effective guidelines on the use of hormones. More studies on the safe use of hormones in fish are crucial, for exploring affordable, environmentally friendly, and appropriate technology.

Genetic manipulation

The problems involved in the direct application of hormones to produce monosex (all-male tilapia) have eventually led to another alternative strategy in the production of an all-male tilapia population. Given the fact that tilapia species exhibit a predominantly monofactorial genotypic system similar to humans, male heterogamety (XY) and female homogamety (XX) (Penman et al., 1987; Mair et al., 1990; Mair and Little, 1991), a model was proposed for the production of monosex male progeny by genetic

manipulation of sex determination in *O. niloticus* (Mair and Little, 1991; Herrera and Cruz, 2001). This technique is based on the production of numbers of "supermale" of the novel genotype "YY" which should yield all male progeny when crossed with normal females (Mair and Little, 1991). YY male technology was first shown in medaka, *Oryzias latipes* using the technology of hormonal sex reversal and selective breeding (Yamamoto, 1958). YY males were also produced in guppy, *Poecilia reticulata* (Yamamoto, 1963) in goldfish, *Carassius auratus* (Yamamoto, 1975), and then in tilapia, *O. niloticus* and *mossambicus* (Varadaraj and Pandian, 1989).

The procedures to produce YY supermale tilapia to yield all male tilapia (XY) have been well documented. The technology involves a series of stages of feminization and progeny testing (Herrera and Cruz, 2001). By adopting the Tuan et al. (1999) model to produce YY supermales, sex reversed females can be produced by oral application of a synthetic hormone, diethylstilbestrol (DES). The sex reversed females (XY) can be identified from DES-treated females by progeny testing with sex-reversed males (XX) of the same strain. Three confirmed sex-reversed females (XY) are then crossed with three normal males (XY) to generate YY supermales. The YY supermale is crossed with a normal female, which will eventually give all male progeny, called genetically generated male tilapia (GMT) (Figure 4). GMT and the YY supermales produced using this technology are not considered genetically modified organisms (Fortes, 2005). Compared to the hormonal sex reversal method, YY supermale technology is believed to be more feasible on a commercial scale, and is environmentally friendly since use of hormones on broodstock is limited and no hormonal residues are detected in consumed fish (Mair and Little, 1991). The genetic integrity of species or strain is also not affected, thus the fish produced for culture maintain a normal genotype (Trombka and Avtalion, 1993). Although GMT technology has been proven to be better than other tilapia monosex producing techniques (Mair et al., 1995), dissemination of information about its current application is limited especially in poor communities worldwide. In addition, the technique can be complex, time consuming, tedious, and still require sex hormones at its initial stage; it is only suitable for homogametic species (XX/XY) (Mair and Little, 1991).




Fig.4: Schematic diagram for producing super male (YY) and all male tilapia (XY) (Tave, 1993).

Novel techniques to control tilapia reproduction

As discussed above, early methods of producing a monosex (all-male) tilapia population involved technical limitations that make these methods inappropriate for small aquaculture farms. The main concerns regarding these methods (especially synthetic sex hormone application) in tilapia production include; potential health risks caused by improper implementation of this system by farm workers, detrimental impacts on the environment, and social constraints (Mair and Little, 1991). To date, there is no substantial proof of any environmental damage or harm to humans caused by the synthetic hormone used for sex inversion, however the quantity of MT used in this practice is large compared to the actual dose required for sex reversal (Mlalila et al.,

2015), and this may lead to accumulation of this chemical in tilapia production systems.

This is in accordance with the previous study that reported that oral administration of MT to fish resulted in detectable levels of steroids in recirculating systems (Hulak et al., 2008). If this contaminated water is released into natural waters, this may have far-reaching effects of possible sex reversal in some natural populations, thus altering the sex dynamics of the stock, producing more males which ultimately may lead to recruitment failure (Megbowon, 2011). Therefore, the tilapia-farming sector is currently faced with a major challenge of finding sex control alternative methods, which are non-hazardous, cost effective, consumer and environmentally friendly. Studies have reported that water temperature and herbal extracts (phytochemicals) can influence hormone biosynthesis and the gonadal sex differentiation process (Baroiller et al., 2009; Bairwa et al., 2013; Fuentes-Silva et al., 2013), thus, they could provide alternative means of producing monosex fish populations in aquaculture, particularly in tilapia culture.

Temperature-induced monosex tilapia production

Tilapia is a thermo-sensitive species, its male to female ratio increases with temperature and/or ovarian differentiation induced by low temperatures (Fuentes-Silva et al., 2013). How does this happen? Studies have indicated that temperature is influential at a critical stage of sex differentiation in larval fish relatively similar to the hormone sensitive period. Inhibition of an enzyme called aromatase which catalyzes the conversion of androgen to estrogen during sex differentiation occurs, at high temperatures, thereby shifting larval or fry sex ratio to male (Brodie et al., 1999; Baroiller and D'Cotta, 2001). Masculinization of tilapia was possible at temperatures above 32°C (Baroiller et al., 1995). It has been suggested that this technique could be more effective when applied at least 10 days after fertilization. Sex determination in tilapia is governed by complex genetic interaction of exogenous factors including temperature (Baroiller et al. 2009). This method is novel therefore little of the mode of action is understood. Studies on temperature induced sex determination in tilapia are limited, thus, the mechanisms involved are yet to be established, given that, physiological, genetic or ecological studies cannot totally be depended upon, to better understand the dynamics of environmental sensitivity (Baroiller et al., 2009).

The use of temperature to produce monosex tilapia populations may be environmentally friendly and does not pose a health hazard to humans, however, at present, this method lacks sufficient control and complete sex reversal that is required to ensure its commercial application has yet to be established (Fuentes-Silva et al., 2013). Even if this method becomes effective in controlling tilapia reproduction, it is still not cost effective, and may be hard to apply to small-scale fish farming in poor rural areas. To the best of our knowledge there is no report to date of the commercial application of this technique in tilapia culture. Therefore, more economically viable tilapia biology and environmental research is needed.

Potential of Herbal extracts in producing monosex tilapia

The idea of using herbal extracts to produce monosex tilapia populations in aquaculture is novel, and operates on the principles of synthetic sex reversal hormones in fish culture. Several herbal extracts contain phytochemicals (phytoestrogens) which are structurally and/ or functionally similar to the steroid hormones i.e. 17- β estradiol (E2) in animals. They are capable of producing estrogenic effects in animals (Fowler, 1983; Lehtinen and Tana, 2001). The impaired effects of phytoestrogen were first discovered in the fertility of sheep and cattle that grazed on clover (*Trifolium subterraneum*) (Bennetts et al. 1946). These findings of could have set the pace for research on phytoestrogens, which subsequently led to the discovery of phytoestrogenic properties in several human foodstuffs including rice, soybeans, wheat, grain, potatoes, etc., and the isolation of two potential estrogenic substances from *Trifolium subterraneum* (Bradburry and White, 1954).

Furthermore, the reported potential of phytochemical/phytoestrogens to control reproduction in tilapia, have wide ranging consequences on various physiological processes in animals such as anti-stress, growth promotion, appetite, stimulation, tonic and immune-stimulation, and antimicrobial properties (Citarasu, 2010; Chakraborty and Hanz, 2011; Chakraborty et al., 2012). Nowadays, herbal extracts are preferred to synthetic drugs because they are cost effective, environmentally friendly, and less likely

to produce disease resistance due to their diversity (Logambal et al., 2000; Olusola et al., 2013). In humans, many women prefer phytoestrogens as alternatives to hormone replacement therapy (HRT) and estrogen replacement therapy (ERT), as they do not pose a risk of breast, endometrial cancer, or irregular bleeding (Brzezinski and Debi, 1999; Wade et al., 1999; Wagner et al., 2001).

Scientific databases such as Science Direct reported that the level of research and adoption of phytoestrogens in humans is more advanced than research in this area in aquaculture. Although the shift from synthetic drugs to herbal extracts has been increasing in aquaculture, more attention has been directed to the study of herbal extracts as growth, digestive and immune-stimulating enhancers in fish. Little research has been carried out on phytoestrogens as reproductive inhibitors in tilapia culture. Research which has paved the way to further studies on use of phytoestrogens in tilapia culture has investigated herbal extracts from; *Quillaja saponaria* (Angeles et al., 2015) and *Trigonella-foenum-graecum* (Francis et al., 2001; 2002; 2005; Stadlander et al., 2008), *Azadirachta indica* (Jegade and Fagbenro, 2008), *Hibiscus rosa-sinensis* (Jegade, 2010), *Carica papaya* (Jegade, 2011; Abdelhak et al., 2013; Ampofo-Yeboah, 2013); *Aloe vera* (Jegade, 2011) *Moringa oleifera* (Ampofo-Yeboah, 2013), *Basella alba* (Ghosal and Chakraborty, 2014), Soybean (El-Sayed et al., 2012), and *Tribulus terrestris* (Omitoyin et al., 2013) (Table 2).

A study was conducted showing swollen spermatid nuclei, increased interstitial cells, and focal necrosis in testes, and hydropic degeneration, ruptured follicles, granulomatous inflammation in the interstitium, and necrotic ovaries when Neem leaves, *Azadirachta indica*, were incorporated in *Tilapia zilli* basal diet at 2.0 g/kg (Jegade and Fagbenro, 2008), (Table 3). Endocrine disrupting compounds (EDCs) including phytoestrogens may impair animal reproduction either by affecting gonad differentiation, by either directly affecting the gonad, or delaying maturation. Neem leaves may be an effective reproductive inhibitor in *Tilapia zilli*. Similar gonadal histological changes, due to incorporation of *Hibiscus rosa-sinensis* leaves at 3.0 g/kg in tilapia diet were reported (Jegade, 2010). *Aloe vera* latex was incorporated into a Nile tilapia diet at 2.0 mL/kg (Jegade, 2011) (Table 3). Similar findings were noted when pawpaw (*Carica papaya*) seeds were included in Nile tilapia basal diet at 120g/kg diets (Abdelhak et al., 2013) (Table 3). During these studies significant gonadal histological changes were reported at high doses, and a study confirmed that the effects (i.e. sterility) of papaya seed at high dose were permanent, while medium and low dose may have reversible effects (Abdelhak et al., 2013). These findings support the early studies, which reported that fertility rates in cheetahs in captivity were reduced when fed a feline diet of a soybean product. This effect was reversed when the soybean product was removed from the diet (Setchell et al., 1987).

Furthermore, Saponin extracts from *Quillaja saponaria* (QS) (Francis et al., 2002), fenugreek (*Trigonella foenum-graecum*) and soapbark tree (*Quillaja saponaria*) (Stadlander et al., 2008), and *Tribulus terrestris* (Omitoyin et al., 2013), reportedly shifted the normal 50:50 male to female sex ratio of Nile tilapia larvae in favor of males when incorporated in their diet, and a high percentage of males was recorded with high concentrations (Table 3). These masculinization effects of saponin extracts on tilapia larvae may be explained by the fact that saponin is able to elevate testosterone production (Ganzera et al., 2001), and as a result, plants that contain saponin compounds, particularly *Tribulus terrestris* have been used to treat impotence in humans (Adaikan et al., 2000; Gauthaman et al., 2002).

Table 2. Phytoestrogenic compounds and in vitro (human cells) status of some herbs.

Family/ sub-family	Scientific / common name	Phytoestrogenic compounds	Phytoestrogenic	References
			activity in vitro	
Liliaceae	<i>Aloe vera</i>	Flavonoids, Saponins, Anthraquinones	not confirmed	Patel et al. 2012
		Phenol, Steroids, Alkaloids		Kumar et al. 2012
	<i>Aloe arborescens</i>	not identified	confirmed	Matsuda et al. 2001
	<i>Aloe forex</i>	not identified	confirmed	Matsuda et al. 2001
	<i>Aloe dracaena loureiroi</i>	Chromanone	confirmed	Ichikawa et al.1997
Cannabaceae	<i>Humulus lupulus</i>	Flavone	confirmed	Hesse et al. 1981; Zava et al. 1998
				Milligan et al.1999
Moringaceae	<i>Moringa oleifera</i>	Flavonoids, phenol	not confirmed	Dillard and German, 2000
				Siddhuraja and Becker, 2003
Agavaceae	<i>Yucca sp.</i>	not identified	confirmed	Zava et al. 1998
Agavaceae	<i>Serenoa repens</i>	not identified	confirmed	Di Silverio et al. 1992
Cariceae	<i>Carica papaya</i>	Alkaloids, Flavonoids, Anthraquinone	not confirmed	Bamisaye et al. 2013
		Saponin, Phenol		
Poaceae	<i>Avena sativa</i> (oat)	Lignan	confirmed	Mazur, 1998
Annonaceae	<i>Annaxagorea luzonensis</i>	Prenylflavonoids	confirmed	Kitaoka et al. 1998
Euphorbiaeae	<i>Jatropha curcas</i>	Alkaloids, Flavonoids, Saponin	not confirmed	Linda et al. 2014
		Phenol		
Meliaceae	<i>Azadirachta indica</i> (Neem)	Flavonoids, Phenol, Saponin, Alkaloids	not confirmed	Linda et al. 2014
Papavenaceae	<i>Sanguinaria Canadensis</i> (bloodroot)	Flavonoids, Phenol, Saponin, Alkaloids	confirmed	Zava et al. 1998
Rubiaceae	<i>Coffea arabica</i> (Coffee)	Lignan	confirmed	Kitts, 1987
	<i>Uncaria tomentosa</i>	Not identified	confirmed	Salazar and Jayme, 1998
Fabaceae	<i>Glycine max</i> (soy bean)	Lignan	confirmed	Rafi et al. 2000
Linaceae	<i>Linum usitatissimum</i> (Flax)	not identified	confirmed	Adlercreutz et al. 1992
Apiaceae	<i>Angelica sinensis</i> (dong quai)	not identified	confirmed	Dixon-Shanies and Shaikh, 1999
Verbenaceae	<i>Verbena officinalis</i>	not identified	confirmed	Dixon-Shanies and Shaikh, 1999
	<i>Vitex agnus-castus</i>	not identified	confirmed	Liu et al. 2001
Lamiaceae	<i>Leonurus cardiac</i>	not identified	confirmed	Zava et al. 1998
Basellaceae	<i>Basella alba</i>	Phenol, Flavonoids, Saponin	confirmed	Thirupathi and Rao 2014
Malvaceae	<i>Hibiscus macranthus</i>	Flavonoids, Tannin, Sterol	confirmed in bull leyding cells	Moundipa et al. 2005
	<i>Hibiscus rosasinesis</i>	Flavonoids, Tannin, Sterol	not confirmed	Soni et al. 2011
	<i>Hibiscus sabdariffa</i> Linn	Flavonoids, Tannin, Sterol	not confirmed	Kumar et al. 2012
Moraceae	<i>Maclura pomifera</i>	Isoflavone	confirmed	Maier et al. 1995
	<i>Morus microphyla</i>	not identified	confirmed	Maier et al. 1995
Rosaceae	<i>Pranus africanum</i>	not identified	confirmed	Maier et al., 1995
Zingiberaceae	<i>Curcuma longa</i>	not identified	confirmed	Zava et al. 1998
Polygonaceae	<i>Fallopia multiflora</i>	anthraquinone	confirmed	Matsuda et al. 2001
	<i>Rheum officinale</i>	anthraquinone	confirmed	Matsuda et al. 2001

Contrary to what is expected of phytoestrogens in tilapia culture, soybean meal was reported to sharply reduce the percentage of males when added to Nile tilapia diet and tilapia farmers were cautioned to avoid the use of soybean as a source of protein during sex reversal treatment (El-Sayed et al., 2012) (Table 3).

The effects of phytoestrogens on animals depend on their ratio to endogenous estrogen, aromatase activity, animal species, reproduction status, length of exposure, and method of administration (Bennetau-Pelissero et al., 2001; Green and Kelly, 2009; Monteiro et al., 2000), and eventually, they may either exert the same effect as estrogen or block the effect of the estrogen (Andersen et al. 2003; Trant et al., 2001; Tsai et al., 2000). These facts therefore explain the estrogenic effects of soybean meal in Nile tilapia. We can see that phytoestrogens could be an alternative method to potentially control tilapia reproduction in production systems.

Table 3. In vivo herbal extracts inhibitory studies in tilapia reproduction

Scientific name	Extracts	Delivery	Concentrations	Exposure (days)	Study	Effective doses	Tilapi spp.	References
<i>Moringa oleifera</i>	Seed powder	Oral	0.5, 1.0, 5.0, 10, 15g/kg	60	Gonad histology	5.0 g/kg	<i>O. mossambicus</i> fingerlings	Ampofo-Yebo, 2013
	Seed powder	Oral	15 g/kg	90	Sex ratio	15 g/kg	<i>O. mossambicus</i> larvae	Ampofo-Yebo, 2013
<i>Canca papya</i>	Seed powder	Oral	60, 90, 120 g/kg	60	Gonad histology	120 g/kg	<i>O. niloticus</i> fingerlings	Abdelhak et al. 2013
	Seed powder	Oral	0.5, 1.0, 5.0, 10, 15g/kg	60	Gonad histology	5.0 g/kg	<i>O. mossambicus</i> fingerlings	Ampofo-Yebo, 2013
	Seed powder	Oral	15 g/kg	90	Sex ratio	15 g/kg	<i>O. mossambicus</i> fingerlings	Ampofo-Yebo, 2013
<i>Hibiscus rosasinesis</i>	Leaves	Oral	1.0, 2.0, 3.0, 4.0 g/kg	60	Gonad histology	3.0 g/kg	<i>O. niloticus</i> fingerlings	Jegede, 2010
<i>Azadirachta indica</i>	Leaves	Oral	0.5, 1.0, 1.5, 2.0 g/kg	60	Gonad histology	2.0 g/kg	<i>T. zillii</i> fingerlings	Jegede & Fagbenro, 2008
<i>Aloe vera</i>	Latex	Oral	0.5, 1.0, 1.5, 2.0 mg/kg	60	Gonad histology	2.0 mg/kg	<i>O. niloticus</i> fingerlings	Jegede, 2011
<i>Basella alba</i>	Aqueous leaves	Immersion	0.05, 0.1, 0.15, g/l	30	Sex ratio	0.1 g/l	<i>O. niloticus</i> larvae	Ghosal & Charabarty, 2014
<i>Ouillaja saponins</i>	Saponin	Oral	50, 150, 300, 500, 700 mg/kg	60	Sex ratio	700 mg/kg	<i>O. niloticus</i> larvae	Francis et al. 2002
	Methanol saponin	Oral	40, 60, 80%, (150 & 1000ppm)	70	Sex ratio	80%, 150 ppm	<i>O. niloticus</i> larvae	Francis et al. 2002
<i>T. foenum-fraecum</i>	Methanol saponin	Oral	40, 60, 80%, (150 & 1000ppm)	70	Sex ratio	80%, 150 ppm	<i>O. niloticus</i> larvae	Statlander et al., 2008
<i>Glycine max (soybean)</i>	Genistein/Daidzein	Oral	10, 20, 30 mg/kg	28	Sex ratio	30 mg/kg	<i>O. niloticus</i> larvae	El Sayed et al. 2012
<i>Tribulus terrestris Prododioscin</i>	Saponin	Oral	1.0, 1.5, 2.0, 2.5 g/kg	42	Sex ratio	2.5 g/kg	<i>O. niloticus</i> larvae	Omitoyin et al, 20013

Phytoestrogens and their modes of actions

As discussed above, phytoestrogens are plant-derivatives that are structurally and /or functionally similar to mammalian estrogen 17β -estradiol (E2), therefore capable of producing estrogenic actions in animals (Price and Fenwick, 1985; Knight and Eden, 1996; Lehtinen and Tana, 2001; Ososki and Kennelly, 2003). The common groups of phytoestrogens include isoflavonoids (i.e. flavonols & isoflavans), coumestans (i.e. coumestrol), and lignan (Lehtinen and Tana, 2001). Among these groups, Isoflavonoids (e.g. genistein and daidzen) are the most well-known of the phytoestrogen and are by monocarboxylic derivatives of flavonoids with a carbon skeleton of 15 atoms similar to that of the coumestans (coumestrole) (Kaldas and Hughes, 1989) (Figure 5) whereas, the lignans have a skeleton based on 18 atoms (Adlercreutz, 1995). In addition to these groups of phytoestrogens, other classes of phytoestrogens include anthraquinones (Matsuda et al., 2001), Chalcones (Rafi et al., 2000), flavones (Milligan et al., 1999), prenylflavonoids, (Kitaoka et al., 1998) and saponins (Chan et al., 2002) (Table 2).




Fig 5. Structural formulas for some of the phytoestrogenic derivatives

The effects of phytoestrogens are partly based on the stability of their natural structures and low molecular weight, which allow them to pass through cell membranes and interact with enzymes and ERs to subsequently cause an estrogenic response (Adlercreutz, 1998), thus enhancing the success of masculinization in the sex inversion process of tilapia larvae. These effects may either be estrogenic or anti-estrogenic (Lehtinen and Tana, 2001; Ososki and Kennelly, 2003). Estrogenic potentials are able to mimic endogenous

estrogens and cause estrogenic effects, whereas the anti-estrogenic potentials may block or alter estrogen receptors (ER) and prevent estrogenic activity, thus causing inverse estrogenic effects (Brzezinski and Debi, 1999); Ososki and Kennelly, 2003; Matozzo et

al., 2008). Phytoestrogens can be classified as selective estrogen receptor modulators (SERMs) (Brzezinski and Debi, 1999) with non-steroidal chemicals having similar structure to E2 and an affinity for estrogen receptors, and function either as agonists or antagonists (Gruber et al., 2002). Phytoestrogenic responses depend on several factors including methods of administration, dosage, metabolism of the target organism, intake of other chemical substances (Kelly et al., 1995; Xu et al., 1995), target tissues, number and types of ERs, and the presence or absence of endogenous estrogen (Glazier and Bowman, 2001).

There is insufficient knowledge on the mechanism of phytoestrogens in inducing estrus in animals. However, they are believed to exert estrogenic effects on the central nervous system, induce estrus, and stimulate growth of the genital tract in female animals (Ososki and Kennelly, 2003). In humans and rats, phytoestrogens are understood to bind to two ERs namely, ER β and ER α , generally found in the brain, bladder, lungs, ovary, prostate, testis, uterus, spleen, and thymus tissues (Mosselman et al., 1996; Kuiper et al., 1997). In addition, a third ER called ER γ has been recently reported in different tissues of Atlantic croaker fish (Hawkins et al., 2000).

It has been observed that not all plants that show estrogenic activity can induce estrus; there are plant substances which are not phytoestrogens yet induce estrus (Ososki and Kennelly, 2003). This further illustrates the ambiguity in the mechanism of phytochemical inducement of estrus. Therefore, there is a need for more extensive research of phytoestrogens in a holistic manner.

Phytoestrogen screening and extraction techniques

There are a large number of techniques used to verify, quantify, and extract phytoestrogens from herbs and foodstuffs. Some of the quantification methods of phytoestrogens include bioassays, cyclodextrin-modified micellar electrokinetic chromatography (CD-MEKC), micellar electrokinetic capillary chromatography (MECC), nonaqueous capillary electrophoresis (NACE), gas chromatography coupled with a mass spectrometer (GC-MS), and high pressure liquid chromatography (HPLC) (Liang et al., 2009). As far as phytochemical assessments are concerned, these techniques have their own limitations. For instance, GC-MS as a quantification technique is reported to involve extensive purification procedures before analysis, thus making it more labor intensive and time consuming with much sample preparation (Wang et al., 2002), whereas, MEKC, MECC and NACE are associated with low reproducibility (Liang et al., 2009); bioassay techniques on the other hand have different level of sensitivity (Diel et al., 1999). Despite their limitations, these techniques are still useful, and are reported to be more effective when combined (Ososki and Kennelly, 2003; Ebuehi and Okorie, 2009).

Of these techniques, HPLC is the most commonly used method, since it involves limited sample preparation (Wang et al., 2002), and its usage is highly adaptable. Thus, it is most frequently combined with other techniques to improve phytochemical screening effectiveness. The combination of thin layer chromatography (TLC), infrared spectroscopy (IRS) and HPLC has been reported to be very effective to analyze and quantify flavonoids in *Jatropha curcas* leaves (Ebuehi and Okorie, 2009). In an attempt to quantify three phytoestrogens (Triterpene acid, Oleanolic acid, and Ursolic acid) in pawpaw and moringa seed powder respectively through ultra-high performance liquid chromatography combined with electrospray ionization mass spectrometer detection (UHPLC-ESI-MS/MS) (Ampofo-Yeboah, 2013), only Oleanolic acid in moringa could be quantified. Phytochemical screening is complicated, thus, it is always difficult to compare phytoestrogen content levels between herbs/plants and foodstuffs (Mazur, 1998).

In addition to the different phytoestrogen quantification techniques, there are also different kinds of phytoestrogen extraction methods such as solvents, solid phase extraction, and supercritical extraction (Asimi and Sahu, 2013). Among the three methods, solvents with different polarities such as petrol, ether, toluene, acetone, ethanol, ethyl acetate and water is the common phytochemical extraction method (Asimi and Sahu, 2013; Susmitha et al., 2013). A phytochemical screening study on Neem (*Azadirachta indica*) was able to extract flavonoids, alkaloids, steroids, saponin, and tanin in ethanol, methanol, acetone solution, respectively, whereas, water and ether were unable to extract steroids, saponins, and tanins (Susmitha et al., 2013). Meanwhile, Bamisaye et al. (2013) was able to extract all five phytoestrogens (flavonoids, alkaloids,

steroids, saponin, and tannin) extracted by Susmitha et al. (2013) from aqueous extracts of leaves and roots of *Carica papaya*.

Similar to phytoestrogens quantification techniques, combinations of solvents have been effective in extracting phytochemicals. For instance, a mixture of distilled water, ethanol, and acetone solution was reported to extract tannins, saponins, and flavonoids from *Aloe vera* leaves using GC-MS techniques (Arunkumar and Muthuselvam, 2009). Nevertheless, methanol/water/HCl (70:29: 1, v/v/v) mixture is reported to be the best of several solvent combinations used to extract phytochemicals (Asimi and Sahu, 2013). Therefore, the types of extraction solvents/techniques and phytochemical quantification/verification techniques remain very important aspects to consider in phytoestrogenic studies.

Administration of herbal extracts in aquaculture

In aquaculture, herbal extracts may be administered orally, by injection, and/or by immersion (Table 3). Injection and immersion may be the most effective methods of administering herbal extracts to fish, however, they may not be practical in aquaculture at all stages of production as they are expensive, labor intensive, and stressful to fish. Although oral administration is considered less effective, because absorption by fish is slow (Harikrishnan et al., 2009), this method allows a large number of fish to be treated with reduced stress and at lower cost, and labor input (Sakai, 1999), thus making it a potentially more suitable option in aquaculture. Herbal extracts incorporated in fish diets at different doses (Bulfon et al., 2013) depend on the extracts, size of the animals, farming system, and purpose of administration. There is a lack of standardization in herbal extract administration in aquaculture therefore more studies are needed to try and standardize methodology of extraction procedures, extract dosage, method of administration, and quantifying effects of herbal extracts in different aquaculture fish species.

Limitations in the existing knowledge and the path forward

Precocious maturity and prolific breeding of tilapia species in aquaculture presents numerous production challenges such as stunted populations, feed utilization, health and welfare. Consequently, the use of synthetic sex reversal hormones as a popular approach to mitigate this problem is more production oriented, while overlooking the impact of these hormones on the environment, and on humans, and animals. The idea that medicinal herbal extracts possess the ability to control tilapia reproduction in intensive production systems has been proposed [Francis et al. (2002); Jegede & Fagbenro 2008); Stadlander et al. (2008); Jegede (2010; 2011); El- Sayed et al. (2012); Abdelhak et al. (2013); Ampofo-Yeboah (2013); Omitoyin et al. (2013)] however, this technology has yet to progress from experimental trials to widespread adoption by farmers. Compared to synthetic sex hormones, medicinal herbal extracts are easily accessible especially by small scale fish farmers, simple to apply, and may be safe for both the environment and humans as they tend to be more biodegradable (Logambal et al. 2000; Dabrowski et al., 2005; Olusola et al., 2013; Reverter et al., 2014).

The general application of herbal extracts in aquaculture is limited. The gap in existing knowledge limits the successful implementation of herbal extracts especially phytoestrogens. The problems include phytoestrogen identification methods, extraction methods, types of extracts (organic, aqueous, methanol, ethanol among others), harmful and beneficial doses for each aquaculture fish species, mode of application, mode of action, effects of different classes of phytoestrogens or herbal extracts in general on fish reproductive systems and other physiological processes such as energy metabolism, growth hormones and protein turnover. Saponin compounds found in most medicinal plants are believed to be promising tilapia masculinizing agents but reduce feed intake in fish (Dongmeza et al., 2006). Fish fed diets containing methanol extracted moringa leaf have been reported to show a better feed intake and growth performance than those fed with diets containing raw moringa leaf meal extract (Afuang et al., 2003). As stipulated above, methods of extraction and types of extracts are among the factors that determine the beneficial properties and efficacy of herbal extracts on the health of cultured fish. Several bioactive compounds are considered to be toxic (Dongmeza et al. 2006), and at present there is lack of knowledge on the toxicity of some herbal extracts in fish, consumers, and the environment. Furthermore, the effects of interaction of phytoestrogens and exogenous parameters including dietary inclusion levels are still

unknown. Therefore, the need to research more and standardize every aspect of the use of herbal extracts in aquaculture species is overwhelming. There is a need to explore and implement the use of herbs in a sustainable manner, as adopting them in aquaculture may double the pressure already exerted by agricultural sectors and humans. In conclusion, this review reveals that there are numerous medicinal plants with potential to control early maturity and prolific breeding in tilapia aquaculture. However, the lack of sufficient knowledge on herbal extracts limits the cost effective use of phytoestrogens in aquaculture. Therefore, more research is required to further validate the use of herbal extracts with their allied phytoestrogenic activity, extraction methods, and extract concentration.

Acknowledgements

This work was jointly supported by Nanjing Government-Nanjing Agricultural University (NJG-NAU), Wuxi fisheries college of NAU, Department of Biotechnology. This article is part of an MSc thesis partly financially supported by NJG-NAU international scholarship program (No. 2013NJ0906) and Namibian Government scholarship & training program (NGSTP) awarded to Ndakalimwe N Gabriel.

References

- Abdelhak M.E., Madkour F.F., Ibrahim M.A., Sharaf M.S., Sharaf M.M. and D.A, Mohammed,** 2013. Effects of pawpaw, *Carica papaya* seeds meal on the productive performance and histological characters of gonads in Nile tilapia, *Oreochromis niloticus*. *IJAR*, 3 (12): 34-37.
- Acosta O.B. and V.M. Gupta,** 2010. The genetic improvement of farmed tilapias project: impact and lessons learned. pp. 149-171. In: S.S. De Silva and F.B. Davy (eds.). *Success Stories in Asian Aquaculture*. Springer, Netherland. 218 pp.
- Adaikan P.G., Gauthaman K., Prasad R.N. and S.C. Ng,** 2000. Proerectile pharmacological effects of *Tribulus terrestris* extract on the rabbit corpus cavernosum. *Ann. Acad. Med. Singap.*, 29: 22-26.
- Adlercreutz H,** 1998. Epidemiology of phytoestrogens. *Baillieres Clin. Endocrinol. Metab.*, 12: 605-623.
- Adlercreutz H,** 1995. Phytoestrogens: epidemiology and a possible role in cancer protection. *Environ. Health Perspect.*, 103: 103-112.
- Adlercreutz H., Mousavi Y., Clark J., Höckerstedt K., Hamalainen E., Wahala K., et al.,** 1992. Dietary phytoestrogens and cancer: in vitro and in vivo studies. *J. Steroid Biochem.*, 41: 331-337.
- Afuang W., Siddhuraju P. and K. Becker,** 2003. Comparative nutritional evaluation of raw, methanol extracted residues and methanol extracts of moringa (*Moringa oleifera* lam.) leaves on growth performance and feed utilization in Nile tilapia, *Oreochromis niloticus*. *Aquac. Res.*, 34: 1147-1159.
- Altun T., Tekelioglu N. and S. Danaba,** 2006. Tilapia culture and its problem in Turkey. *J. Fish Aquat. Sci*, 23: 473-478.
- Angeles I.P. and Y-H Chien,** 2015. Dietary effect of *Quillaja saponaria* and /or *Yucca schidigera* extract on growth and survival of common carp *Cyprinus carpio* and their antioxidant capacity and metabolic response upon low dissolved oxygen. *The Israeli Journal of Aquaculture-Bamidgeh*, IJA_64.2014.1165.
- Ampofo-Yeboah A,** 2013. Effect of phytochemical feed additives on gonadal development in Mozambique Tilapia, *Oreochromis mossambicus*. *PhD thesis*, 254 pp. Stellenbosch University, South Africa, <http://scholar.sun.ac.za>.
- Andersen L., Bjerregaard P. and B. Korsgaard,** 2003. Vitellogenin induction and brain aro- matase activity in adult male and female zebra fish exposed to endocrine disrupters. *Fish Physiol. Biochem.*, 28: 319-321.
- Arunkumar S. and M. Muthuselva,** 2009 Analysis of phytochemical constituents and antimicrobial activities of *Aloe vera* against clinical pathogens. *W. J. Agri Sci.*, 5: 572-576. **Asimi O.A. and N.P Sahu,** 2013. Herbs/spices as feed additives in aquaculture. *Sci. J. Pure Appl. Sci.*, 2: 284-292.

- Bai D., Wu X., Zhu G., Guo Y., Yang G., Ning B et al.,** 2012. Astragalus polysaccharides enhance cellular immune response and disease resistance in yellow catfish. *The Israel Journal of Aquaculture-Bamidgeh*, [IJA_65.2013.688](#).
- Bairwa K.M., Saharan N., Rawat D.K., Jakhar K.J. and A. Bera,** 2013. Photoperiod, melatonin and its importance in fish reproduction. *Cent. Eur. J. Exp. Biol.*, 3: 7-15.
- Bamisaye F.A, Ajani E.O. and J.B. Minari,** 2013. Prospects of ethnobotanical uses of pawpaw (*Carica papaya*). *J. Med. Plants Stud.*, 1: 171-177.
- Baroiller J.F. and H. D’Cotta,** 2001. Environment and sex determination in farmed fish. *Comp. Biochem. Phys. C*, 130: 399-409.
- Baroiller J.F., Chourrou D., Fostier A. and B. Jalabert,** 1995. Temperature and sex chromosomes govern sex ratios of the mouthbrooding cichlidfish *Oreochromis niloticus*. *J. Exp. Zool.*, 273:216-223.
- Baroiller J.F., D’Cotta H., Bezault E., Wessels S. and G. Hoerstgen-Schwark,** 2009. Tilapia sex determination: Where temperature and genetics meet. *Comp. Biochem. Phys. A*, 153: 8-30.
- Bartley M.D., Rana K. and J.A. Immink.,** 2001. The use of inter-specific hybrids in aquaculture and Fisheries. *Rev. Fish Biol. Fisher.*, 10: 325-337.
- Beardmore J.A., Mair G.C. and R.I. Lewis,** 2001. Monosex male production in finfish as exemplified by tilapia: applications, problems, and prospects. *Aquaculture*, 197: 283-301.
- Bennetau-Pelissero C, Breton B, Bennetau B, Corraze G, LeMenn E, Davail-Cuisset B et al.,** 2001. Effect of Genistein enriched diets on the endocrine process of gametogenesis and on reproduction efficiency of the rainbow trout, *Onchrohynchus mykiss*. *Gen. Comp. Endocr.*, 121: 173-187.
- Bennetts H.W., Underwood E.J. and F.L. Shier,** 1946. A specific breeding problem of sheep on subterranean clover pastures in Western Australia. *Aust. Vet. J.*, 22: 2-12.
- Bradbury R.B. and D.E. White,** 1954. Estrogens and related substances in plants. *Vitamins and Hormones*, 12: 207-233.
- Brodie A., Lu Q. and B. Long,** 1999. Aromatase and its inhibitors. *J. Steroid Biochem. Mol. Biol.*, 69: 205-210.
- Bromage N., Porter M. and C. Randall,** 2001. The environmental regulation of maturation in farmed finfish with special reference to the role of photoperiod and melatonin. *Aquaculture*, 197: 63-98.
- Brzezinski A. and A. Debi,** 1999. Phytoestrogens: the ‘natural’ selective estrogen receptor modulators? *Eur. J Obstet. Gyn R B.*, 85: 47-51.
- Bulfon C., Volpatti D. and M. Galeotti,** 2013. Current research on the use of plant derived products in farmed fish. *Aquac. Res.*, doi: 10.1111/are.12238.
- Chakraborty BS., Molnar T. and C. Hancz,** 2012. Effects of methyltestosterone, tamoxifen, ganistein, and *Basella alba* extract on masculinization of guppy (*Poecilia reticulata*). *J. Appl. Pharm. Sci.*, 2(12), 48-52.
- Chakraborty S.B. and C. Hancz,** 2011. Application of phytochemicals as immunostimulant, antipathogenic and antistress agents in finfish culture. *Reviews in Aquaculture*, 3: 103-119.
- Chan R.Y., Chen W.F., Dong A., Guo D. and M.S. Wong,** 2002. Estrogen-like activity of ginsenoside Rg1 derived from *Panax notoginseng*. *J. Clin. Endocr. Metab.*, 87: 3691-3695.
- Chapman F.A,** 1992. Culture of hybrid tilapia: A reference profile. University of Florida, IFAS extension, *CIR1051*, <http://edis.ifas.ufl.edu>.
- Chervinski J. and S. Rothbard,** 1982. An aid in manually sexing tilapia, brief technical note. *Aquaculture*, 26: 389.
- Citarasu T,** 2010. Herbal biomedicines: a new opportunity for aquaculture industry. *Aquacult. Int.*, 18: 403-414.
- Dabrowski K., Gustavo R. and G.A.M. Abiado,** 2005. Use of phytochemicals as an environmentally friendly method to produce sex reversed Nile tilapia. In: Burright J, Flemming C, Egna, H. (eds.) *22nd annual technical report*, pp. 287-330. Aquaculture CRSP, Corvallis, Oregon.
- Das R., Rather M.A., Basavaraja N., Sharma R. and U. K. Udit,** 2012. Effect of nonsteroidal aromatase inhibitor on sex reversal of *Oreochromis mossambicus* (Peter, 1852). *The Israel Journal of Aquaculture-Bemidgeh*, 64-69, [IJA_64.2012.703](#).

- Di Silverio F., D'Eramo G., Lubrano C., Flammia G.P. and A. Sciarra**, 1992. Evidence that *Serenoa repens* extract displays an antiestrogenic activity in prostatic tissue of benign prostatic hypertrophy patients. *Eur. Urol.*, 21: 309–314.
- Diel P., Smolnikar K. and H. Michna**, 1999. In vitro test systems for the evaluation of the estrogenic activity of natural products. *Planta Medica*, 65: 197–203.
- Dillard C.J. and J.B German**, 2000. Phytochemicals: nutraceuticals and human health: a review. *J. Sci. Food Agr.*, 80: 1744–1756.
- Dixon-Shanies D. and N. Shaikh**, 1999. Growth inhibition of human breast cancer cells by herbs and phytoestrogens. *Oncology Report*, 6: 1383–1387.
- Dongmeza E., Siddhuraju P., Francis G. and K. Becker**, 2006. Effects of dehydrated methanol extracts of *Moringa (Moringa oleifera Lam.)* leaves and three of its fractions on growth and feed nutrient assimilation in Nile tilapia (*Oreochromis niloticus L.*). *Aquaculture*, 261: 407-422.
- Ebuehi O.A. and N.A. Okorie**, 2009. Phytochemical screening and quantification of flavonoids from leaf extract of *Jatropha curcas* Linn. *Niger Q. J. Hosp. Med.*, 19: 200-205.
- Eckstein B. and M. Spira**, 1965. Effect of sex hormone on gonadal differentiation in a cichlid, *Tilapia aurea*. *The Biology Bulletin (Woods Hole, Mass)*, 129: 482–489.
- El-Sayed A.F.M. and M. Kawanna**, 2007. Effects of photoperiod on growth and spawning efficiency of Nile tilapia (*Oreochromis niloticus L.*) broodstock in a recycling system. *Aqua. Res.*, 38: 1242-1247.
- El-Sayed M.A.B., Abdel-Aziz H.E.S. and M.H. Abdel-Ghani**, 2012. Effects of phytoestrogens on sex reversal of Nile tilapia (*Oreochromis niloticus*) larvae fed diets treated with 17 α -methyltestosterone. *Aquaculture*, 360: 58-63.
- Essa A.M. and R.M. Haroun**, 1998. Cross-breeding experiments on some important fishes of family cichlidae (genus *Oreochromis*) and evaluation of their hybrids. *Egypt. J. Aquat. Biol. Fish.*, 2: 43-61.
- FAO, 2002.** *The State of World Fisheries and Agriculture*. FAO, Rome, p150.
- FAO, 2006.** *State of World Aquaculture 2006*:FAO Fisheries Technical Paper No. 500. FAO, Rome, 134p.
- FAO, 2012.** *The State of World Fisheries and Aquaculture 2012*. FAO, Rome, p209.
- FAO/GLOBEFISH**, 2013. *GLOBEFISH highlights, a quarterly update on world seafood market 3/2013*. FAO, Rome, p. 61.
- Felicitta J., Manju R. A., Ronald J., Sakthika T., Nagarajan R. and G. Chelladurai**, 2013. The effect of different concentrations garlic (*Allium sativum*) and onion (*Allium cepa*) on growth, survival, and hematology of juvenile tilapia (*Oreochromis mossambicus*). *Isr. J. Aquacult.- Bamidgeh*, [IJA 65. 2013. 822](#).
- Fink G**, 2000. Neuroendocrine regulation of pituitary function: general principle. In: Conn PM, Freeman ME, Totowa NJ (eds.) *Neuroendocrinology in physiology & Medicine*, pp. 107-134. Humana Press, Totowa, New Jersey.
- Fitzsimmons K, Martinez R. and P. Ramotar**, 2012. Global production and market situation in 2012-tilapia continue to climb the chart. *AQUA 2012 meeting abstract*. World aquaculture society, Baton Rouge, Louisiana.
- Fitzsimmons K., Garcia M.R. and G.P. Alanis**, 2011. Why tilapia is becoming the most important food fish on the planet. In: K. Fitzsimmons, L. Liping (eds.). *Proceedings of the 9th International symposium on tilapia aquaculture*, pp.8-16. Aquafish CRSP, Shanghai, China.
- Fortes D.R**, 2005. *Review of techniques and practice in controlling tilapia population and identification of methods that may have practical applications in Nauru including a national tilapia plan*. Agdex Pacific Islands 492/679, New Caledonia, France.
- Fowler M.E**, 1983. Plant poisoning in free-living wild animals: a review. *J Wildl Dis*, 19: 34- 43.
- Francis G., Levavi-Sivan B., Avitan A. and K. Becker**, 2002. Effects of long term feeding *Quillaja saponins* on sex ratio, muscle and serum cholesterol and LH levels in Nile tilapia (*Oreochromis niloticus L.*). *Comp. Biochem. Phys. C*, 133: 593-603.
- Francis G., Makkar H.P.S. And K. Becker**, 2001. Effects of *Quillaja saponins* on growth, metabolism, egg production, and muscle cholesterol in individually reared Nile tilapia (*Oreochromis niloticus*). *Comp. Biochem. Phys. C*, 129:105–114.

- Francis G., Makkar H.P.S. and K. Becker**, 2005. *Quillaja* saponins – a natural growth promoter for fish. *Anim. Feed Sci. Tech.*, 121: 147–157.
- Fuentes-Silva C., Soto-Zaraźua M.G., Torres-Pacheco I. and A. Flores-Rangel**, 2013. Male tilapia production techniques: A mini review. *Afr. J. Biotechnol.*, 12: 5496-5502.
- Gabriel N.N., Qiang J., He J., Ma Y.X., Kpundeh M.D. and P. Xu**, 2015. Dietary Aloe vera supplementation on growth performance, some haemato-biochemical parameters and disease resistance against *Streptococcus iniae* in tilapia (GIFT). *Fish and Shellfish Immunol.*, doi: 10.1016/j.fsi.2015.03.002.
- Ganзера M., Bedir E. and I.A. Khan**, 2001. Determination of steroidal saponins in *Tribulus terrestris* by reversed-phase high-performance liquid chromatography and evaporative light scattering detection. *J. Pharm. Sci.*, 90: 1752–1758.
- Gauthaman K., Adaikan P.G. and R.N.V Prasad**, 2002. Aphrodisiac properties of *Tribulus terrestris* extract (Protodioscin) in normal and castrated rats. *Life Sciences*, 71: 1385–1396.
- Ghosal I. and B.S. Chakraborty**, 2014. Effects of the aqueous leaf extract of *Basella alba* on sex reversal Nile tilapia, *Oreochromis niloticus* L. *IOSR J. Pharm. Biol. Sci* 9: 162-164.
- Glazier M.G. and M.A. Bowman**, 2001. A review of the evidence for the use of phytoestrogens as a replacement for traditional estrogen replacement therapy. *Archives of Internal Medicine*, 161: 1161– 1172.
- Green C.C. and A.M. Kelly**, 2009. Effects of the estrogen mimic genistein as a dietary component on sex differentiation and ethoxyresorufin-O-deethylase (EROD) activity in channel catfish *Ictalurus punctatus*. *Fish Physiol. Biochem.*, 35: 377–384.
- Gruber C.J., Tschugguel W., Schneeberger C. and J.C. Huber**, 2002. Production and actions of estrogens. *New Engl. J. Med.* 346: 340–352.
- Guerrero D.R.**, 1979. Use of hormonal steroids for artificial sex reversal of tilapia. *Proc. Indian Natl. Sci. Acad.*, 45: 512-514.
- Guerrero R.D.**, 1975. Use of androgens for the production of all-male *Tilapia aurea* (Steindachner). *Transaction of the American Fisheries Society*, 104: 342–348.
- Harikrishnan R., Balasundaram C., Kim M.C., Kim J.S., Han Y.J. and M.S. Heo**, 2009. Innate immune response and disease resistance in *Carassius auratus* by triherbal solvent extracts. *Fish Shellfish Immunol.*, 27:508–515.
- Hawkins M.B., Thornton J.W., Crews D., Ski K., Dotte A. and P. Thomas**, 2000. Identification of a third distinct estrogen receptor and reclassification of estrogen receptors in teleosts. *Proc. Natl. Acad. Sci. USA*, 97: 10751–10756.
- Herrera A.A., Cruz R.R. and FGBPGM-GIFTP**, 2001. Development biology of the supermale YY tilapia (*Oreochromis niloticus*) Histogenesis of the reproductive system. *Science Diliman*, 13 (1): 33-40.
- Hesse R., Hoffmann B., Karg H. and H. Vogt**, 1981. Demonstration of phytoestrogens in feed plants and hops by means of receptor test. *Zentralbl Veterinarmed A*, 28: 422–454.
- Hickling C.F.**, 1960. The Malacca Tilapia Hybrid. *J. Genet*, 57: 1-10.
- Hu H., Mai K., Zhang Y., Ai Q., Xu W., Zhang W., et al**, 2014. Effects of dietary xylan on growth performance, digestive enzyme activity and intestinal morphology of juvenile turbot (*Scophthalmus maximus* L). *Isr. J. Aquacult.- Bamidgeh*, [IJA 67. 2015. 1115.](#)
- Hulak M., Paroulek M., Simek P., Kocour M., Gela D., Rodina M et al.**, 2008. Water polluted by 17 α -methyltestosterone provides successful male sex inversion of common carp (*Cyprinus carpio* L.) from gynogenetic offspring. *J. Appl. Ichthyol.*, 24: 707-710.
- Hulata G., Cnaani A., Slossman T. and G.A.E. Graham**, 2004. Fertility problems in the second generation of a four-species tilapia cross. *The Israeli Journal of Aquaculture – Bamidgeh* [56\(3\), 159-165.](#)
- Ichikawa K., Kitaoka M., Taki M., Takaish I.S., Iijima Y., Boriboon M., et al**, 1997. Retrodihydrochalcones and homoisoflavones isolated from Thai medicinal plant *Dracaena loureiri* and their estrogen agonist activity. *Planta Medica*, 63: 540–543.
- Hulata G., Wohlfarth G. and S. Rothbard**, 1983. Progeny-testing selection of tilapia broodstocks producing all-male hybrid progenies-preliminary results. *Aquaculture*, 33: 263-268.

- Jegade T**, 2010. Control of reproduction in *Oreochromis niloticus* (Linnaeus 1758) using *Hibiscus rosa-sinensis* (Linn.) leaf meal as reproduction inhibitor. *J. Agr. Sci.*, 2: 149-154.
- Jegade T**, 2011. Effects of *Aloe vera* (Liliaceae) on the gonad development in Nile tilapia (*Oreochromis niloticus*) (Linnaeus 1758). In: Fitzsimmons, K., & Liping, L (eds), *Proceedings of the 9th International symposiums on tilapia aquaculture*, 222-227 pp. Aquafish CRSP, Shanghai, China.
- Jegade T. and O. Fagbenro**. 2008. Histology of gonads in *Tilapia zillii* (Gervais) fed Neem (*Azadirachta indica*) leaf meal diets. *8th International symposium on tilapia aquaculture*. Cairo, Egypt.
- Jiménez-Badillo L**, 2006. Age-growth models for tilapia (*Oreochromis aureus*) (perciformers, Cichlidae) of the infiernillo reservoir, Mexico and reproductive behavior: Review. *Int. J. Trop. Biol.*, 54: 577-588.
- Kah O, Anglade I, Leprêtre E, Dubourg P. and D. De Monbrison**, 1993. The reproductive brain in fish. *Fish Physiol. Biochem.*, 11:85-98.
- Kajimura S., Kawaguchi N., Kaneko T., Kawazoe I., Hirano T., Visitacion N., et al.**, 2004. Identification of the growth hormone receptor in an advanced teleost, the tilapia, *Oreochromis mossambicus* with special reference to its distinct expression pattern in the ovary. *J. Endocrinol.*, 181: 65-76.
- Kaldas RS, Hughes CL Jr.**, 1989. Reproductive and general metabolic effects of phytoestrogens in mammals. *Reprod. Toxicol.*, 3: 81-89.
- Kelly G.E., Joannou G.E., Reeder A.Y., Nelson C. M.A. Waring**, 1995. The variable metabolic response to dietary isoflavones in humans. *Proc. Soc. Exp. Biol. Med.*, 208: 40-43.
- Kitaoka M., Kadokawa H., Sugano M., Ichikawa K., Taki M., Takaishi S et al.**, 1998. Prenylflavonoids: a new class of non-steroidal phytoestrogen (Part 1). Isolation of 8-isopentenylnaringenin and an initial study on its structure-activity relationship. *Planta Medica*, 64: 511- 515.
- Kitts D.D**, 1987. Studies on the estrogenic activity of a coffee extract. *J. Toxicol. Environ. Health*, 20: 37-49.
- Knight D.C. and J.A. Eden**, 1996. A review of the clinical effects of phytoestrogens. *Ob. Gyn.*, 87: 897-904.
- Kuiper G.G.J.M., Carlsson B.O., Grandien K., Enmark E., Haggblad J., Nilsson, S et al.**, 1997. Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptor alpha and beta. *Endocrinology*, 138: 863-870.
- Kumar K.H.N., Chandana E., Preethi S.D. and B.J. Chauhan**, 2012a. In vitro antimicrobial activity and phytochemical screening of *Aloe vera* Linn. *Int. J. Curr. Pharmaceut. Res.*, 4: 45-47.
- Kumar M., Garg R. and R. Garg**, 2012b. Phytochemical properties and antioxidant activity of *Hibiscus Sabdariffa* Linn. *Int. J. Chem. Pharmaceut. Sci.*, 1:1236-1240.
- Lehtinen K.J. and J. Tana**, 2001. Review of endocrine disrupting natural compounds and endocrine effects of pulp and paper mill and municipal sewage effluents. *The Finnish environment: Environmental protection*, 447:1-73.
- Levavi-Sivan B. and Z. Yaron**, 1993. Intracellular mediation of GnRH action on GTH release in tilapia. *Fish Physiol. Biochem.*, 11(1-6): 51-59.
- Liang Z., Jiang Z., Fong D.W. and Z. Zhao**, 2009. Determination of Oleanolic Acid and Ursolic Acid in *Oldenlandia diffusa* and its Substitute Using High Performance Liquid Chromatography. *J. Food Drug Anal.*, 17:69-77.
- Linda J.H.A. and E.O.S. Okon**, 2014. Comparative study of the phytochemical properties of *Jatropha curcas* and *Azadirachta indica* plant extracts. *J. poisonous med. plants res.*, 2:21-24. **Liu A., Yang Z., Zhu M. and J. Huo**, 2001. Estrogenicity of black cohosh (*Cimicifuga racemosa*) and its effect on estrogen receptor level in human breast cancer MCF-7 cells. *J. Hyg. Res.*, 30: 77-80.
- Logambal S.M., Venkatalakshmi S. and R.D. Michael**, 2000. Immunostimulatory effect of leaf extract of *Ocimum sanctum* Linn. In *Oreochromis mossambicus* (peters). *Hydrobiologia*, 430:113-120.

- Lovshin L.L., Da Silva A.B., Carneiro-Sobrinho A. and F.R. Melo**, 1990. Effects of *Oreochromis niloticus* females on the growth and yield of male hybrids (*O. niloticus* female × *O. hornorum* male) cultured in earthen ponds. *Aquaculture*, 88: 55–60.
- Mair G.C., Abucay S.J., Beardmore A.J. and D.O.F. Skibinski**, 1995. Growth performance trials of genetically male tilapia (GMT) derived from YY-males in *Oreochromis niloticus* L.: on station comparisons with mixed sex and sex reversed male populations. *Aquaculture*, 137: 313–323.
- Mair G.C., Beardmore J.A. and D.O.F. Skibinski**, 1990. Experimental evidence for environmental sex determination in *Oreochromis* species. In: Hirano RI, Hanyu, I (eds.) *2nd Asian Fisheries Forum*, pp. 555–558. Asian Fisheries Society, Manila, Philippines.
- Mair G.C. And C.D. Little**, 1991. Population control in farmed tilapia. *NAGA, ICLARMQ* 17: 8–13.
- Majumdar K.C. and B.J. McAndrew**, 1983. Sex ratio from interspecific crosses within the tilapias. *Int. symp. on Tilapia in aquaculture*, pp.261–269. Nazareth, Israel.
- Mañanós E., Zanuy S. and M. Carrillo**, 1997. Photoperiodic manipulations of the reproductive cycle of sea bass (*Dicentrarchus labrax*) and their effects on gonadal development, and plasma 17 β -estradiol and vitellogenin levels. *Fish Physiol. Biochem.*, 16: 211–222.
- Marchesan M., Spoto M., Verginella L. and E.A. Ferrero**, 2005. Behavioural effects of artificial light on fish species of commercial interest Ferrero. *Fish. Res.*, 73: 171–185.
- Marengoni N.G. and Y. Onoue**, 1998. Ultraviolet-induced androgenesis in Nile tilapia *Oreochromis niloticus* (L.), and hybrid Nile × blue tilapia, *O. aureus* (Steindachner). *Aquac. Res.*, 29:359–366.
- Maruska P.K. D.R. Fernald**, 2013. Social regulation of male reproduction plasticity in an African cichlidae fish. *Soc. Intergr. Comp. Biol.*, 53: 938–950.
- Matozzo V., Gagne F., Marin G.M., Ricciardi F. and C. Blaise**, 2008. Vitellogenin as a biomarker of exposure to estrogen compounds in aquatic Invertebrates: A review. *Environ. Int.*, 34: 531–545.
- Matsuda H., Shimoda H., Morikawa T. and M. Yoshikawa**, 2001. Phytoestrogens from the roots of *Polygonum cuspidatum* (Polygonaceae): structure-requirement of hydroxyanth- raquinones for estrogenic activity. *Bioorg. Med Chem. Lett.*, 11: 1839–1842.
- Mazur W**, 1998. Phytoestrogen content in foods. *Baillieres Clin. Endocrinol. Metab.*, 12: 729–742.
- Megbowon I**, 2011. Tilapia Production in Nigeria. *Fisheries Society of Nigeria Quarterly Publication*, 4: 18–22.
- Milligan SR., Kalita J.C., Heyerick A., Rong H., Cooman L. and D. Keukeleire**, 1999. Identification of a potent phytoestrogen in hops (*Humulus lupulus* L.) and beer. *J. Clin. Endocr. Metab.*, 83: 2249–2252.
- Mires D.** 1977. Theoretical and practical aspects of the production of all male tilapia hybrids. *The Israeli Journal of Aquaculture –Bamidgeh*, 29: 94–101.
- Mires D., 1987.** A Technical evaluation of tilapia cultures. *Proceedings of the International Symposium on Tilapia in Aquaculture* 8–13 May 1983. Nazareth, Israel. Tel Aviv University, Israel.
- Mires D., 1995.** The tilapias in Nash C.E., and Novotny A.J., World aquatic sciences (C8), *Production of aquatic animals: Fishes*. Elsevier Science, New York, pp133–152.
- Mlalila M., Mahika C., Kalombo L., Swai H. and A. Hilonga**, 2015. Human food safety and environmental hazards associated with the use of methyltestosterone and other steroids in production of all-male tilapia. *Environ. Sci. Pollut. Res.*, Doi 10.1007/s11356-015-4133-3.
- Monteiro P.R.R., Reis-Henriques, M.A. and J. Coimbra**, 2000. Polycyclic aromatic inhibit in vitro ovarian steroidogenesis in the flounder (*Platichthys flesus* L.). *Aquat Toxicol.*, 48: 549–559.
- Mosselman S., Polman J. and R. Dijkema**, 1996. ER β : identification and characterization of a novel human estrogen receptor. *FEBS Letters*, 392: 49–53.

- Moundipa F.P., Beboy E.S.N., Zelefact F., Ngouela S., Tsamo E., and W.B. Schill,** 2005. Effects of *Basella alba* and *Hibiscus macranthus* extracts on testosterone production of adult rat and bull leydig cells. *Asian J. Androl.*, 7: 411-417.
- Nakamura M., Kobayashi T., Chang X. and Y. Nagahama,** 1998. Gonadal sex differentiation in teleost fish. *J. Exp. Zool.*, 281:362-372.
- Nandlal S. and T. Pickering,** 2004. *Tilapia fish farming in Pacific Island countries: Tilapia hatchery operation*. Noumea, New Caledonia PMid: 15255439.
- Obaroh I.O. and C.G. Achionye-Nzeth,** 2011. Effects of crude extract of *Azadirachta indica* leaves at controlling prolific breeding in *Oreochromis niloticus* (Linnaeus, 1978). *Asian J. Agri. Res.*, 5: 277-282.
- Olusola S.E., Emikpe B.O. and F.E. Olaifa,** 2013. The potentials of medical plants extracts as bio-antimicrobial in aquaculture. *Int. J. Med. Arom. Plants.*, 3: 404-412.
- Omitoyin B.O., Ajani E.K. and H.O. Sadiq,** 2013. Preliminary investigation of *Tribulus terrestris* (Linn, 1753) extracts as natural sex reversal agent in *Oreochromis niloticus* (Linn. 1758) larvae. *Int. J. Aquacult.*, 3:133-137.
- Ososki A.L. and E.J. Kenelly,** 2003. Phytoestrogens: A review of the present state of research. *Phytother Res.*, 17: 845-869.
- Pandian T.J. and S.G. Sheela,** 1995. Hormonal induction of sex reversal in fish. *Aquaculture*, 138:1-22.
- Patel K.D., Patel K. and S.P. Dhanabal,** 2012. Phytochemical standardization of *Aloe vera* extract by HPTLC techniques. *J. acute dis.*, doi: 10.1016/S2221-6189 (13) 60054-2
- Penman D.J. and B.J. McAndrew,** 2000. Genetics for the management and improvement of cultured Tilapia. In: Bereridge MCM, McAndrew BJ (eds.) *Tilapia: Biology and exploitation*, Kluwer Academic publishers, Dordrecht, Boston, London. pp.227-266.
- Penman D.J., M.S. Shah, A.J. Beardmore and F.O.D. Skibinski,** 1987. Sex ratios of gynogenetic and triploid tilapia. *Proceedings on the World Symposium on Selection, hybridization and genetic engineering in Aquaculture*, 78: 237-276.
- Phelps P.R. and J.T. Popma,** 2000. Sex reversal of tilapia. In: Costa-pierce AB, Rakocy EK (eds) *Tilapia aquaculture in the Americas*, 2:39-59. The World Aquaculture Society: Baton Rouge, Louisiana.
- Pinilla L., Aguilar E., Dieguez C., Millar P.R. and M. Tena-Sempere,** 2012. Kiss peptins and reproduction: physiological roles and regulatory mechanisms. *Physiol. Rev.*, 92:1235-1316.
- Prat F., Zanuy S., Bromage N. and M. Carrillo,** 1999. Effects of constant short and long photoperiod regimes on the spawning performance and sex steroid levels of female and male sea bass. *J. Fish Biol.*, 54: 125-137.
- Price K.R. and G.R. Fenwick,** 1985. Naturally occurring oestrogens in foods – a review. *Food Addit. Contam.*, 2: 73-106.
- Pruginin Y,** 1967. The culture of carp and tilapia hybrid in Uganda. *FAO fisheries Report*, 44: 223-229.
- Pruginin Y., Rothbard S., Wohlfarth G., Halevy A., Moav R. and G. Hulata,** 1975. All-male broods of *Tilapia nilotica* x *T. aurea* hybrids. *Aquaculture*, 6: 11-21.
- Pullin R.S.V. and R.H. Lowe-McConnell (eds.),** 1982. The biology and culture of tilapias. *International Conference on the Biology and Culture of Tilapias*, 7: 432. ICLARM, Bellagio, Italy.
- Rafi M.M., Rosen R.T., Vassail A., Ho C.T., Zhang H., Ghai G., et al.,** 2000. Modulation of bcl-2 and cytotoxicity by licochalcone-A, a novel estrogenic flavonoid. *Anticancer Res.*, 20: 2653-2658.
- Reverter M., Bontemps N., Lecchini D., Banaigs B. and P. Sasal,** 2014. Use of plant extracts in fish aquaculture as an alternative to chemotherapy: current status and future perspectives. *Aquaculture*, 433: 50-60.
- Sakai M,** 1999. Current research status of fish immunostimulant. *Aquaculture*, 172: 63-92.
- Salazar E.L. and V. Jayme,** 1998. Depletion of specific binding sites for estrogen receptor by *Uncaria tomentosa*. *Proc. West. Pharmacol. Soc.*, 41: 123-124.
- Schulz WR., de Franca R.L., Lareyre J.J., LeGac F., Chiarini-Garcia H., Nobrega H.R., et al.,** 2010. Spermatogenesis in fish. *Gen. comp. Endocr.*, 165: 390-411.

- Setchell K.D.R., Gosselin S.J., Welsh M.B., Johnston J.O., Balistreri W.F., Kramer L.W., et al.**, 1987. Dietary estrogens – a probable cause of infertility and liver disease in captive cheetahs. *Gastroenterology*, 93: 225–233.
- Siddhuraju P. and K. Becker**, 2003. Antioxidant properties of various solvent extracts of total phenolic constituents from three different agro-climatic origins of drumstick tree (*Moringa oleifera* Lam.). *J. Agr. Food Chem.*, 15: 2144–2155.
- Soni D., Gupta A., Solanki R., Jana K.G.**, 2011. Pharmacognostical phytochemical and physiochemical findings over the root extract of *Hibiscus rosa-sinensis* (Malvaceae). *J. Nat. Prod. Plant Res.*, 1: 73-79.
- Srinath K., Sridhar M., Kartha P.N.R. and A.N. Mohanan**, 2000. Group farming for sustainable aquaculture. *Ocean and coastal management*, 43: 557-571.
- Stadtlander T, Focken U, Levavi-Sivan B, Dweik H, Qutob M, Abu-Lafi S et al.**, 2008. Treatment with saponins from *Trigonella foenum-graecum* and *Quillaja saponaria* influences sex ratio in Nile tilapia (*Oreochromis niloticus*) larvae. In: Fitzsimmons K, Liping L (eds.) proceedings of the 8th international symposiums on tilapia aquaculture. Cairo, Egypt. 355-364 pp.
- Susmitha S., Vidyamol K.K., Ranganayaki P. and R. Vijayaragavan**, 2013. Phytochemical extraction and antimicrobial properties of *Azadirachta indica* (Neem). *Global J. Pharmacol.*, 7: 316-320.
- Taranger L.G., Carrillo M., Schulz., W.R., Fontaine P., Zanuy S., Felip A., et al.**, 2010. Control puberty in farmed fish. *Gen. comp Endocr.*, 165: 483-515.
- Tave D**, 1993. *Genetics for fish hatchery managers*, 2nd ed. Van-Nostrand Reinhold, New York.
- Teichert-Coddington D.R., Green B.W. and R.P. Phelps**, 1992. Influence of site and season on water quality and tilapia production in Panama and Honduras. *Aquaculture*, 105: 297-314.
- Tena-Sempere M. and I. Huhtaniemi**, 2003. Gonadotropins and gonadotropin receptors. In: B.C.J.M. Fauser (ed.) *Reproductive Medicine: Molecular, cellular and genetic fundamentals*. Parthenon, New York. pp. 225-244.
- Thirupathi B. and G.S. Rao**, 2014. Comparative phytochemical quantification in green and purple varieties of *Basella alba* for preventive management of nutritional deficiency and stress related diseases. *World J. pharm. Pharm. Sci.*, 3:1061-1071.
- Trant J.M., Gavasso S., Ackers J., Chung B.C. and A.R. Place**, 2001. Developmental expression of cytochrome P450 aromatase genes (CYP19a and CYP19b) in zebrafish fry (*Danio rerio*). *J. Exp Zool.*, 290: 475–483.
- Trewavas E**, 1982. Tilapias: taxonomy and speciation. In: R.S.V. Pullin, R.H Lowe-McConnell (eds.) *proceedings of the 7th ICLARM Conference, Biology and Culture of Tilapia*. Manila, Philippines. pp. 3-13.
- Trombka D. and R. Avtalion**, 1993. Sex determination in tilapia-a review. *Bamidgeh*, 45: 26-37.
- Tsai C.L., Wang L.H., Chang C.F. and C.C. Kao**, 2000. Effects of gonadal steroids on brain serotonergic and aromatase activity during the critical period of sexual differentiation in tilapia, *Oreochromis mossambicus*. *J. Neuroendocrinol.*, 49:894-898.
- Tuan P.A, Mair G.C., Little D.C. J.A. Beardmore**, 1999. Sex determination and the feasibility of genetically male tilapia production in the Thai-Chitralada strain of *Oreochromis niloticus* (L.). *Aquaculture*, 173: 257-269.
- Tubert C, Nostro L.F., Villafane V. and C. Pandolfi**, 2012. Aggressive behavior and reproductive physiology in females of the social cichlid fish *Cichlasoma dimerus*. *Physiol. Behav.*, 106: 193-200.
- Tveteras R**, 2013. Global fish production and trends in 2012-2013. *Vietfish international*, 10: 40-44.
- UN**, 2010. *World Population Prospects: The 2010 Revision*. Department of Economic and Social Affairs, Population Division, New York.
- Van der Kraak G., Chang P.J. and M.D. Janz**, 1998. *Reproduction: The physiology of fishes*. CRC Press, Boca Ratón, Florida.
- Varadaraj K. and T.J. Pandian**, 1989. First Report on production of supermale tilapia by integrating endocrine sex reversal with gynogenetic technique. *Curr. Sci.*, 58: 434-441.

- Wade C., Kronenberg F., Kelly A. and P.A. Murphy,** 1999. Hormone-modulating herbs: implications for women's health. *J. Am. Med. Womens Assoc.*, 54: 181-183.
- Wagner J.D., Anthony M.S. and J.M. Cline.** 2001. Soy phytoestrogens: research on benefits and risks. *Clin. Obst. Gyn.*, 44: 843- 852.
- Wang C.C., Prasain J.K. and S. Barnes,** 2002. Review of the methods used in the determination of phytoestrogens. *J. chromatogr B.*, 777: 3-28.
- Wassermann J.G. and B.O.L. Afonso,** 2002. Validation of aceto-carminic technique for evaluating phenotypic sex in Nile tilapia (*Oreochromis niloticus*) fry. *Crecial Rural, Santa Maria*, 32:133-139.
- Wohlfarth G.W., Hulata G. and A. Halevy,** 1990. Growth, survival and sex ratio of some tilapia species and their interspecific hybrids. *European Aquaculture society Special Publication*, 11: 87-101.
- Worhlfarth G.W,** 1994. The unexploited potential of tilapia hybrids in Aquaculture. *Aquacult. Fish. Manage.*, 25: 781-788.
- Xu X., Harris K.S., Wang H.J., Murphy P.A. and S. Hendrich,** 1995. Bioavailability of soybean isoflavones depends upon gut microflora in women. *J. Nutr.*, 125: 2307-2315.
- Yamamoto T,** 1958. Artificial induction of functional sex reversal in genotypic males of *O. latipes*. *J. Exp Zool.*, 137: 227-264.
- Yamamoto T,** 1963. Induction of reversal in sex differentiation of YY zygotes in medaka, *Oryzias latipes*. *Genetics*, 48: 293-306.
- Yamamoto, T,** 1975. A YY male goldfish from mating estrone-induced XY female and normal male. *J. Hered*, 662: 2-4.
- Yamazaki F,** 1983. Sex control and manipulation in fish. *Aquaculture*, 33: 329-354.
- Zava D.T., Dollbaum C.M. and M. Blen,** 1998. Estrogen and progestin bioactivity of foods, herbs, and spices. *Proceedings of the Society for Experimental Biology and Medicine*, 217:369-378.
- Zohar Y., Muñoz-Cueto J.A., Elizur A. and O. Kah,** 2010. Neuro-endocrinology of reproduction in teleost fish. *Gen. comp. Endocr.*, 165: 438-455.