

The Open Access Israeli Journal of Aquaculture – Bamidgen

As from **January 2010** The Israeli Journal of Aquaculture - Bamidgen (IJA) will be published exclusively as **an on-line Open Access (OA)** quarterly accessible by all AquacultureHub (<http://www.aquaculturehub.org>) members and registered individuals and institutions. Please visit our website (<http://siamb.org.il>) for free registration form, further information and instructions.

This transformation from a subscription printed version to an on-line OA journal, aims at supporting the concept that scientific peer-reviewed publications should be made available to all, including those with limited resources. The OA IJA does not enforce author or subscription fees and will endeavor to obtain alternative sources of income to support this policy for as long as possible.

Editor-in-Chief

Dan Mires

Editorial Board

Rina Chakrabarti	Aqua Research Lab, Dept. of Zoology, University of Delhi, India
Angelo Colorni	National Center for Mariculture, IOLR Eilat, Israel
Daniel Golani	The Hebrew University of Jerusalem Jerusalem, Israel
Hillel Gordin	Kibbutz Yotveta, Arava, Israel
Sheenan Harpaz	Agricultural Research Organization Beit Dagan,
Gideon Hulata	Agricultural Research Organization Beit Dagan,
George Wm. Kissil	National Center for Mariculture, IOLR, Eilat, Israel
Ingrid Lupatsch	Swansea University, Singleton Park, Swansea, UK
Spencer Malecha	Dept. of Human Nutrition, Food & Animal Sciences, CTAHR, University of Hawaii
Constantinos Mylonas	Hellenic Center for Marine Research, Crete, Greece
Amos Tandler	National Center for Mariculture, IOLR Eilat, Israel
Emilio Tibaldi	Udine University Udine, Italy
Jaap van Rijn	Faculty of Agriculture, The Hebrew University of Jerusalem, Israel
Zvi Yaron	Dept. of Zoology, Tel Aviv University, Tel Aviv, Israel

Published under auspices of
The Society of Israeli Aquaculture and Marine Biotechnology (SIAMB),
University of Hawai'i at Mānoa Library

&

University of Hawai'i at Mānoa
Aquaculture Program
in association with
AquacultureHub

<http://www.aquaculturehub.org>



UNIVERSITY
of HAWAII[®]
MĀNOA
LIBRARY



AquacultureHub.org

AquacultureHub
educate · learn · share · engage

ISSN 0792 - 156X

© Israeli Journal of Aquaculture - BAMIGDEH.

PUBLISHER:

Israeli Journal of Aquaculture - BAMIGDEH -
Kibbutz Ein Hamifratz, Mobile Post 25210,
ISRAEL


Phone: + 972 52 3965809

<http://siamb.org.il>

Copy Editor **Ellen Rosenberg**



The IJA appears exclusively as a peer-reviewed on-line open-access journal at <http://www.siamb.org.il/>. To read papers free of charge, please register online at [registration form](#).
Sale of IJA papers is strictly forbidden.



Growth, Survival and Fatty Acid Composition of Freshwater Crayfish (*Astacus leptodactylus*) Juveniles Fed Enriched *Daphnia magna* as an Alternative to *Artemia*

Seval Bahadir Koca^{*1}, Nalan Ozgur Yigit¹, Esra Uzunmehmetoglu¹, Zekiye Guclu², Gurkan Diken¹ and Hasan Eralp¹

¹Suleyman Demirel University, Egirdir Fisheries Faculty, Isparta, Turkey

²Agriculture and Rural Development Support Institution, Burdur Provincial Coordinator, Burdur, Turkey

(Received 20.2.2015, Accepted 12.4.2015)

Key words: enrichment; *Daphnia magna*; anchovy oil emulsion; *Astacus leptodactylus*; growth; survival

Abstract

This experiment was conducted to investigate the effects *Daphnia magna* enriched with lipid emulsions as an alternative to *Artemia*, on growth, survival rate, and fatty acid composition of freshwater crayfish (*Astacus leptodactylus* Esch. 1823). The four treatment groups were (a) unenriched *Artemia* (UEA), (b) unenriched *D. magna* (UED), (c) *D. magna* enriched with redpepper emulsion (DER), and (d) *D. magna* enriched with anchovy oil emulsion (DEA). All tests were carried out in triplicate for 30 days. The crayfish (mean weight 0.12g) were fed ad libitum once daily. At the end of experiment, the highest eicosapentaenoic acid, 20:5n-3 (EPA) level was found in the DEA group (5.77%). The highest DHA (docosahexaenoic acid, 22:6n-3) level was found in the DER group (2.73%) which was statistically similar to the DEA group. In addition, high n-3 HUFA (high unsaturated fatty acid) levels were detected in enriched *D. magna* groups with emulsions. However, high EPA levels in enriched *D. magna* groups with emulsions were not reflected in crayfish tissues, but DHA level was reflected in crayfish tissues fed with anchovy oil emulsion. The crayfish fed with *D. magna* showed similar growth to that of the *Artemia* fed groups. The growth of the enriched *D. magna* groups did not differ.

* Corresponding author. Seval Bahadir Koca: e-mail: sevalkoca1@hotmail.com

Introduction

Astacus leptodactylus is a native crayfish species in Turkey (Köksal, 1988). There is no crayfish aquaculture in Turkey and all production is obtained from wild harvests (Harlioglu et al, 2012). Turkey was the largest supplier of *A. leptodactylus* to Western Europe from 1970 until 1986 but harvests were severely reduced in most wild populations due to infection by crayfish plague, (*Aphanomyces astaci*) after 1985 (Köksal, 1988; Ackefors, 2000). The harvest in 2012 was only 492 tons (Anonymous, 2014). To replenish native stocks of *A. leptodactylus*, exogenous production of juvenile crayfish is needed. However, the nutritional requirements of this species in culture are unknown.

Artemia nauplii are a main food source for larval forms of crustaceans (Immanuel et al., 2007). However, these are expensive and scarce (Das et al., 2007). Crayfish in nature feed mainly on rotifers and benthonic crustaceans (cladocerans and copepods) and, later on larvae of aquatic insects (Hessen, 1989). Cladocerans can possibly substitute for *Artemia* in cultured freshwater crustaceans (Alam, 1995; Das et al. 2007). Commercial emulsions are expensive, therefore, both *Artemia* and commercial emulsions are not practical in freshwater crayfish culture.

In shrimp fed cladoceran enriched with highly unsaturated fatty acids, (HUFA), growth and survival rates increased (Das, et al., 2007). HUFA, eicosapentaenoic acid (EPA), and docosahexaenoic acid (DHA), have been recognized as very important nutrients for the growth of crustaceans (Alam, 1995; Das et al., 2007). However, there is no literature on crayfish fed with enriched cladocerans.

In the present study, the effects of enriched *Daphnia magna* with lipid emulsions as an alternative to *Artemia* on growth, survival rate, and fatty acid profile in *A. leptodactylus*, were investigated.

Materials and Methods

Fatty acid profile of anchovy oil and ingredient of anchovy oil emulsion. The ingredients of anchovy oil emulsion, and the fatty acid profile of anchovy oil for emulsion, are given in Tables 1 & 2.

Table 1. Ingredients of anchovy oil emulsion

<i>Ingredients</i>	<i>Amount</i>
Water (ml)	100
Anchovy oil (ml)	50
Egg yolk (ml)	20
Vitamin mix(g)	0.50

Vitamin mix (Rovimix BCK for 100g): 500mg vitamin B1; 1000mg vitamin B2; 500mg vitamin B6; 3mg vitamin B12; 10000mg vitamin C; 5000mg niacin; 300mg vitamin K3; 10mg D-Biotin; 1500mg Cal.D. Pantothenate; folic acid 150mq.

Table 2. Fatty acid profile of anchovy oil as percentage of total lipid

Fatty acid	Anchovy oil
12:0	0.06±0.00
14:0	6.03±0.00
15:0	0.13±0.00
16:0	15.69±0.18
17:0	0.35±0.01
18:0	2.91±0.03
20:0	0.26±0.01
23:0	0.11±0.01
24:0	1.02±0.02
14:1	0.25±0.01
16:1	9.11±0.40
17:1	0.63±0.01
20:1	0.29±0.01
18:1 n9	13.00±0.15
18:1n7	3.04±0.06
22:1n9	0.17±0.02
18:2 n6	1.91±0.01
20:3n6	0.11±0.00
20:4 n6	0.54±0.01
18:3 n3	1.22±0.01
20:5 n3 (EPA)	11.29±0.09
22:6 n3 (DHA)	18.43±0.24
20:2	1.07±0.00
22:2	0.17±0.02
SFA	26.53±0.23
MUFA	26.48±0.29
PUFA	34.72±0.31
HUFA	31.60±0.30

Table3. Fatty acid profile of redpepper and anchovy oil emulsions as percentage of total lipids.

Fatty Acid	Redpepper emulsion	Anchovy oil emulsion	a
12:0	0.11±0.01	0.06±0.00	*
14:0	5.57±0.46	5.66±0.16	ns
14:1	0.18±0.01	0.25±0.01	*
15:0		0.12±0.00	-
16:0	34.90±1.77	15.61±0.08	**
16:1	1.87±0.03	8.90±0.28	**
17:0	0.07±0.01	0.33±0.01	**
17:1	0.10±0.01	0.59±0.00	**
18:0	0.88±0.05	2.87±0.03	**
18:1 n9	2.45±0.26	17.46±0.19	**
18:1n7		0.18±0.01	-
18:2 n6	1.48±0.01	2.78±0.03	**
18:3 n3	0.40±0.02	1.20±0.01	**
20:0	0.17±0.02	0.30±0.02	*
20:1	0.21±0.01	0.31±0.02	*
20:2	2.24±0.11	1.06±0.00	**
20:3n6	0.09±0.03	0.10±0.01	ns
20:4 n6	0.54±0.09	0.28±0.02	ns
20:5 n3 (EPA)	2.12±0.13	10.82±0.10	**
22:1n9	0.55±0.14	0.20±0.06	ns
22:2	0.32±0.02	0.28±0.03	ns
24:0	0.41±0.04	0.96±0.00	**
22:6 n3 (DHA)	27.59±0.92	18.10±0.12	**
SFA	42.10±1.19	25.92±0.28	**
MUFA	5.35±0.17	27.87±0.14	**
PUFA	34.76±1.26	34.61±0.02	ns
HUFA	32.88±1.24	30.64±0.03	ns

Enrichment and culture of *D. magna*. Two concrete ponds (20 tons capacity) were prepared for mass culture of *D. magna* on a rotation basis. Inorganic fertilizers containing nitrogen and phosphate were used for culture of *D. magna*. The water quality parameters for the rearing of *D. magna* were temperature, 22-24°C; pH, 7.0-7.5; and dissolved oxygen, 5.2-6.7 mg/L throughout the culture period. Each separate *D. magna* group was kept in 10 L capacity plastic buckets containing 4 L of water. The enrichment was conducted in emulsions containing anchovy oil (1.25g/L) and redpepper (0.75 g/L) (Akuamaks, Ankara/TURKEY) in two parts for a time period of 24 h with mild aeration. Fatty acid profiles of emulsions are given in Table 3.

Level of statistical significance (a): ns = $p > 0.05$, * = $p < 0.05$, ** = $p < 0.01$

***Artemia* culture.** 0.5-1.0g of encapsulated *Artemia salina* cysts (INVE Aquaculture, Izmir/TURKEY) were hatched and harvested according to Sorgeloos et al., 1986.

Rearing conditions of crayfish. The crayfish were obtained from broodstock captured from Egirdir Lake, Turkey. A 30 day feeding trial was conducted with crayfish (initial weight 0.12g). The crayfish were stocked into 12 aquaria (40 x 70 cm base area) at a density of 35 individuals /m². Pipes (2cm diameter and 4 cm length) were placed in each aquarium to provide shelter. The physicochemical water conditions were: temperature 18.3±0.07°C, dissolved oxygen 6.2±0.09 mg/L, ammonium 0.01±0.07 mg/L, pH 7.5-8.5, calcium 64.8 mg/L, magnesium 11.2 mg/L, hardness 77.1 mg/L, photoperiod 12/12 which were all kept constant throughout the experiment. Feed remains and feces were siphoned daily and 20% of water was exchanged. Four treatments UEA, UED, DER, and DEA, were applied in triplicate. The crayfish were fed ad libitum once daily (Calvo et al.,

2013). They were measured to the nearest 1 mm from the tip of the rostrum to the end of telson, and weighed to the nearest 0.1 mg after removing the excess water.

Growth parameters. The growth parameters were calculated at the end of the feeding experiment:

Weight gain (g) = final weight - initial weight

Specific Growth Rate (SGR) % = (ln final weight - ln initial weight) / days x100

Survival rate (%) = (Final crayfish number/Initial crayfish number) x100

Fatty acid analyses. The live food, crayfish tissue, and emulsions were stored until further analysis at -80°C. Fatty acid analyses of live food, crayfish tissue, and emulsions, were analyzed by GC Clarus 500 with autosampler (Perkin Elmer, USA) equipped with a flame ionization detector and a fused silica capillary SGE (scientific glass engineering) column (30 mX 0.32 mm ID X 0.25 µm BP20 0.25 UM, USA). The oven temperature was 140°C, held for 5 min, raised to 200°C at rate 4°C/min and held at 220°C at rate 1°C/min, while the injector and the detector temperature were set at 220°C and 280°C, respectively. The sample size was 1 µl and the carrier gas was controlled at 16 ps. The split used was 1:50. Fatty acids were identified by comparing the retention times of FAME with the standard 37 component FAME mixture. Two replicate *gas chromatography* (GC) analyses were performed and the results expressed in GC area % as a mean value and ± standard deviation.

Statistical analyses. The results were examined with a one-way analysis of variance (ANOVA) using the SPSS 13.0 computer program (SPSS Inc., Chicago, USA). Mean comparisons were tested using Duncan's test ($p < 0.05$). Fatty acid profile results of emulsions are presented as mean±SE and subjected to independent-samples t-test for determining significant differences between treatment means.

Results

Fatty acid profile of emulsions. The highest EPA content (10.82%) was found in anchovy oil emulsions while the highest DHA content was found in redpepper emulsions ($P < 0.01$). There was no significant difference in PUFA and HUFA contents between the different emulsions. The highest SFA content (42.10 %) was found in redpepper emulsions while the highest MUFA content was found in anchovy oil emulsions ($P < 0.01$).

Growth and survival. Final weight, carapace length, total length, weight gain, and SGR showed no significant differences among the groups ($P > 0.05$). However, enriched groups showed a trend towards increased final weight, weight gain, and SGR. UED and DER groups showed higher survival rate ($P < 0.05$) than UEA and DEA groups (Table 4).

Table 4 Survival and growth performance of *A. leptodactylus*

	UEA	UED	DEA	DER
Initial weight (g)	0.12±0.02	0.12±0.01	0.12±0.02	0.12±0.02
Final weight (g)	0.28±0.03	0.29±0.02	0.30±0.03	0.30±0.02
Weight gain (g)	0.16±0.01	0.17±0.02	0.18±0.01	0.18±0.01
Final carapax (mm)	11.26±0.36	11.42±0.29	11.74±0.33	11.56±0.29
Final total length (mm)	21.96±0.77	22.52±0.57	22.97±0.62	22.47±0.59
Survival rate (%)	60.00±1.92 ^b	90.00±3.85 ^a	70.00±1.92 ^b	88.88±2.94 ^a
SGR (%)	2.88±0.45	2.96±0.29	3.11±0.45	3.03±0.51

All the values are given in mean± SE. Values with different superscript letters were significantly different ($P < 0.05$) from others in the same row.

Fatty acid profile in tissues of *D. magna* and *Artemia*. A total of sixteen acids were detected and the fatty acid profiles in tissues of *D. magna* and *Artemia* were recorded (Table 5). The enrichment process affected the fatty acid profile of *D. magna*. The amount of EPA and DHA increased in enriched *D. magna* groups. DEA group showed the highest level (5.77 %) of EPA ($P < 0.05$). This value was considerably higher than in the other groups. The highest level of DHA (2.73 %) was found in DER group ($P < 0.05$). Contents of n-3 HUFA were higher in enriched *D. magna* tissues than in unenriched *D. magna* and *Artemia* ($P < 0.05$). The highest n-6 HUFA content was found in the *Artemia* tissues ($P < 0.05$). The n-6/n-3 ratio ranged from 0.29 to 0.96.

Table 5. Fatty acid profile in tissues of *D. magna* and *Artemia*

Fatty Acid	UEA	UED	DEA	DER
<i>Saturated fatty acids</i>				
14:0	0.69±0.02 ^b	3.16±0.03 ^a	3.51±0.27 ^a	3.13±0.14 ^a
15:0	-	0.19±0.02	0.29±0.12	0.20±0.01
16:0	7.21±0.22 ^d	19.55±0.04 ^a	16.40±0.17 ^c	18.12±0.66 ^b
17:0	0.63±0.02 ^a	0.62±0.02 ^a	0.35±0.01 ^b	0.23±0.01 ^c
18:0	4.48±0.05 ^a	2.24±0.07 ^c	3.25±0.01 ^b	1.94±0.43 ^c
20:0	0.51±0.05 ^a	0.09±0.02 ^b	0.17±0.01 ^b	0.19±0.04 ^b
Total SFA	13.52±0.25 ^c	25.85±0.08 ^a	23.95±0.21 ^b	23.80±0.39 ^b
<i>Monounsaturated fatty acids</i>				
18:1n-7	-	3.82±0.42 ^c	8.35±0.55 ^a	6.25±0.15 ^b
18:1n-9	12.40±0.10 ^{ab}	16.42±0.38 ^a	12.31±0.32 ^{ab}	9.36±2.56 ^b
20:1	-	0.18±0.02	0.34±0.03	0.61±0.33
Total MUFA	12.40±0.10 ^b	20.41±0.05 ^a	21.01±0.84 ^a	16.22±3.01 ^{ab}
<i>Polyunsaturated fatty acids</i>				
18:2n-6	4.99±0.03 ^c	10.76±0.23 ^a	4.81±0.06 ^c	6.14±1.94 ^b
20:3n-6	1.22±0.06 ^a	0.07±0.00 ^c	0.34±0.00 ^b	0.15±0.09 ^{bc}
20:4n-6	1.70±0.21 ^a	0.27±0.03 ^b	0.33±0.02 ^b	0.33±0.14 ^b
18:3n-3	24.71±0.77 ^a	10.53±0.10 ^b	3.21±0.20 ^c	3.83±1.23 ^c
20:5n-3 (EPA)	1.84±0.13 ^b	0.17±0.06 ^c	5.77±0.77 ^a	3.25±0.10 ^b
22:6n-3 (DHA)	0.35±0.10 ^c	0.89±0.12 ^{cb}	1.99±0.52 ^{ab}	2.73±0.18 ^a
20:2	0.36±0.03 ^d	0.93±0.05 ^c	1.72±0.03 ^b	1.99±0.12 ^a
Total PUFA	35.15±1.17 ^a	23.61±0.08 ^b	18.15±0.03 ^c	18.41±1.50 ^c
Total n-6	7.91±0.13 ^b	11.10±0.20 ^a	5.48±0.05 ^c	6.61±0.66 ^{bc}
Total n-3	26.89±1.27 ^a	11.58±0.07 ^b	10.96±0.05 ^b	9.81±0.95 ^b
n-6/n-3	0.29±0.02 ^d	0.96±0.02 ^a	0.50±0.01 ^c	0.67±0.00 ^b
Total n-6 HUFA	2.92±0.15 ^a	0.34±0.03 ^b	0.67±0.02 ^b	0.48±0.23 ^b
Total n-3 HUFA	2.18±0.03 ^c	1.05±0.17 ^d	7.76±0.25 ^a	5.98±0.28 ^b

All the values are given in mean± SE. Values with different superscript letters were significantly different (P<0.05) from the others in the same row.

Fatty acid composition of A. leptodactylus. Fatty acid profile in tissues of crayfish is given in Table 6. The highest DHA content in tissue was found in crayfish fed DEA (9.07%). This was similar to the DER group. The high DHA content in enriched *D. magna* groups was reflected in crayfish tissues. The highest EPA level (5.77%) was found in the *D. magna* enriched with anchovy oil emulsion, EPA levels in crayfish tissues were similar in all groups (P>0.05). n-3 HUFA contents were similar in all groups (P<0.05).

Table 6. Fatty acid composition in tissues of *A. leptodactylus*

Fatty Acid	UEA	UED	DEA	DER
<i>Saturated fatty acids</i>				
14:0	2.36±0.23 ^b	5.13±0.57 ^a	5.37±0.07 ^a	4.96±0.74 ^a
15:0	0.29±0.08 ^a	0.10±0.00 ^b	0.07±0.01 ^b	0.08±0.00 ^b
16:0	14.68±0.89 ^b	17.25±0.02 ^{ab}	18.23±0.04 ^a	17.72±0.58 ^a
18:0	4.16±0.22	4.38±0.24	3.84±0.04	4.46±0.04
20:0	0.26±0.03	0.28±0.06	0.21±0.02	0.34±0.13
Total SFA	21.74±0.80 ^b	27.14±0.63 ^a	27.71±0.03 ^a	27.56±0.25 ^a
<i>Monounsaturated fatty acids</i>				
15:1	0.25±0.06 ^d	0.58±0.01 ^c	0.78±0.02 ^b	1.36±0.05 ^a
16:1	3.66±0.18 ^b	4.43±0.08 ^a	4.54±0.10 ^a	3.84±0.14 ^b
18:1n-9	22.70±1.07 ^{ab}	24.66±0.29 ^a	23.55±0.13 ^{ab}	22.37±0.02 ^b
22:1n-9	0.05±0.01 ^c	-	0.26±0.02 ^a	0.18±0.01 ^b
Total MUFA	26.65±0.95 ^b	29.66±0.20 ^a	29.12±0.03 ^a	27.75±0.13 ^{ab}
<i>Polyunsaturated fatty acids</i>				
18:2n-6	6.88±0.60	7.69±0.21	7.83±0.05	6.51±0.28
20:3n-6	1.05±0.03 ^a	0.22±0.01 ^b	0.18±0.00 ^b	0.22±0.00 ^b
20:4n-6	0.79±0.08 ^a	0.31±0.02 ^b	0.35±0.05 ^b	0.29±0.02 ^b
18:3n-3	1.14±0.11 ^c	1.84±0.01 ^a	1.67±0.05 ^{ab}	1.58±0.03 ^b
20:5n-3 (EPA)	6.16±0.17	5.35±0.45	5.72±0.55	6.01±1.07
22:6n-3 (DHA)	6.67±0.44 ^c	7.54±0.40 ^{bc}	9.07±0.08 ^a	7.99±0.01 ^{ab}
Total PUFA	22.67±0.33 ^{ab}	22.93±0.23 ^{ab}	24.81±0.67 ^a	22.59±0.76 ^b
Total n-6	8.72±0.70 ^a	8.21±0.18 ^{ab}	8.36±0.10 ^{ab}	7.02±0.30 ^b
Total n-3	13.96±0.38	14.73±0.05	16.46±0.58	15.58±1.06
n-6/n-3	0.63±0.07 ^a	0.56±0.01 ^{ab}	0.51±0.01 ^{ab}	0.45±0.05 ^b
Total n-6 HUFA	1.85±0.11 ^a	0.53±0.03 ^b	0.53±0.05 ^b	0.51±0.06 ^b
Total n-3 HUFA	12.82±0.27	12.89±0.06	14.79±0.62	14.00±1.08

All the values are given in mean± SE values with different superscript letters were significantly different (P<0.05) from the others in the same row

Discussion

To our knowledge there are no published studies on the effects of *Daphnia magna* enriched with lipid emulsions used as an alternative to *Artemia*, on growth and fatty acid composition of crustaceans.

In this study, *D. magna* and *Artemia* fed groups showed similar growth performance. The enriched *D. magna* groups did not show differences in final weight, weight gain, and SGR. Growth rate and survival increased in shrimp fed enriched cladoceran with HUFA (Alam, 1995; Das et al., 2007). These positive effects were also observed on growth and survival rate in crustacea fed enriched *Artemia* (Millamena et al., 1988; Abelin 1991; Romdhane et al., 1995; Citarasu et al., 1998; Immanuel et al., 2001; Immanunuel et al. 2004; Chakraborty et al. 2010). High levels of n-3 HUFA in diets had no growth promoting effects in shrimps (Rees et al. 1994). In the present study, n-3 HUFA content increased in enriched *D. magna* tissues (DEA and DER groups) compared with unenriched groups (UEA and UED groups). The growth and survival rates did not increase with increasing n-3 HUFA levels. The optimum HUFA levels in *A. leptodactylus* still need investigation.

Growth performance of shrimp increased with increasing EPA and DHA rates in enriched *Artemia* (Immanuel et al., 2001; 2004; Rees et al., 1994). Growth and survival rates of *Macrobrachium rosenbergii* larvae fed enriched *Moina micrura* increased with increasing amount of EPA and DHA in the diets (Das et al. 2007). Increasing the level of both EPA and DHA above a certain inclusion level resulted in a decrease in growth of *Penaeus monodon* (Glencross & Smith, 2001). In the present study, the highest EPA content was found in *D. magna* enriched with anchovy oil while the highest DHA content was found in *D. magna* enriched with redpepper. These enrichments did not result in enhanced growth.

Increasing levels of EPA and DHA affected the n-6:n-3 balance and adversely affected growth (Glencross & Smith 2001), however, n-6/n-3 in the diet ratios showed a decline in enriched *Moina micrura* and improved growth of *M. rosenbergii* post larvae fed the lowest n-6/n-3 ratio (Das et al. 2007). In the present study, n-6/n-3 ratio was lower in the enriched groups of *D. magna* than in the unenriched ones. The lowest n-6/n-3 ratio was determined in the *Artemia* fed group. This may be because the C18:3 n3 level in *Artemia* was much higher than in the *D. magna* groups.

EPA and DHA levels of *Moina micrura* increased with enrichment (Das et al. 2007). Similarly, in the current study, EPA and DHA contents showed significant increase with enrichment of *D. magna*. Although EPA and DHA levels of *D. magna* enriched with emulsions were the highest we believe that these were not sufficient to enhance growth. DHA and EPA contents in tissues of wild and captive adult *A. leptodactylus* ranged between 7.77%-23.10% and 6.02%-13.51% respectively (Harlioglu et al., 2012). In the current study, DHA levels were higher or equal while EPA levels were lower and therefore need to be enhanced.

Post larvae fed higher dietary n-3 fatty acids showed higher n-3 fatty acids in tissues (Das et al. 2007). In contrast, the highest n-3 fatty acids were seen in the *Artemia* diet in the current study, but n-3 fatty acids in crayfish tissues were similar in all groups.

Crayfish groups fed with *D. magna* showed similar growth to the *Artemia* group however growth of crayfish groups fed enriched *D. magna* with emulsions was not affected. DHA levels were high in crayfish tissues of groups fed anchovy oil emulsion yet we believe that HUFA levels of emulsions were not sufficient in the enriched groups and that the HUFA requirements of *A. leptodactylus* still need to be determined.

References

- Abelin P**, 1991. Development and evolution of unconventional forms of *Artemia* sp. as food for penaeid shrimp. PhD Thesis, University of Ghent, Belgium.
- Ackefors H**, 2000. Freshwater crayfish farming technology in the 1990s: a European and global perspective. *Fish. Fish.*, 1(4): 337–359.
- Alam M.J., Ang K.J. and M. Begum**, 1995. Use of egg custard augmented with cod liver oil and *Moina micrura* on production of freshwater prawn post larvae. *Aquac. Int.*, 3: 249–259.

- Calvo, N.S., Stumpf, L. Sacristán, H.J. and L.S. López Greco,** 2013. Energetic reserves and digestive enzyme activities in juveniles of the red claw crayfish *Cherax quadricarinatus* nearby the point-of-no-return. *Aquaculture*, 416–417: 85–91.
- Chakraborty, K., Chakraborty, D.R., Radhakrishnan, E. and K.K.Vijayan,** 2010. Fatty acid profiles of spiny lobster (*Panulirus homarus*) phyllosoma fed enriched *Artemia*. *Aquac Res.*, 41: 393-403.
- Citarasu, T., Immanuel, G. and M.P. Marian,** 1998. Effect of feeding *Artemia* enriched with stresstol and cod-liver oil on growth and stress resistance in the Indian white shrimp *Penaeus indicus* post larvae. *Asian Fish. Sci.*, 12: 1-7.
- Das, S.K., Tiwari, V.K, Venkateswarlu, G., Reddy, A.K., Parhi, J., Sharma P. and J.K., Chettri,** 2007. Growth, survival and fatty acid composition of *Macrobrachium rosenbergii* (De Man, 1879) post larvae fed HUFA-enriched *Moina micrura*. *Aquaculture*. 269: 464–475.
- Glencross, B.D. and D.M. Smith,** 2001. Optimising the dietary levels of eicosapentaenoic and docosahexaenoic essential fatty acids for the prawn, *Penaeus monodon*. *Aquac Nutr.*, 7: 101–112.
- Harlioglu, A.G., Aydin, S. and O. Yilmaz,** 2012. Fatty acid, cholesterol and fat-soluble vitamin composition of wild and captive freshwater crayfish (*Astacus leptodactylus*). *Int. J. Food Sci. Technol.*, 18: 93–100
- Hessen, D.O.,** 1989. Crayfish food and nutrition. In. Skurdal J, Westman K and Bergan P I (Eds.), *Crayfish Culture in Europe*. Report from the workshop on crayfish culture, 16–19 Nov 1987, Trondheim, Norway, pp. 164–174.
- Immanuel, G., Palavesam, A. and M. Petermarian,** 2001. Effects of feeding lipid enriched *Artemia* nauplii on survival, growth, fatty acids and stress resistance of postlarvae *Penaeus indicus*. *Asian Fish. Sci.*, 14: 377-388.
- Immanuel, G., Palavesam, A., Sivaram, V., Michael, B.M., M.P. Marian,** 2004. Feeding trashfish *Odonus niger* lipid enriched *Artemia* nauplii on growth, stress resistance and HUFA requirements of *Penaeus monodon* postlarvae. *Aquaculture*, 237: 301–313.
- Immanuel, G., Citarasu, T., Sivaram, V., Shankar, S. and A. Palavesam,** 2007. Bioencapsulation strategy and highly unsaturated fatty acids (HUFA) enrichment in *Artemia franciscana* nauplii by using marine trash fish *Odonus niger* liver oil. *Afr J Biotechnol.*, 6: 2043-2053.
- Köksal, G.,** 1988. *Astacus leptodactylus* in Europe. In. Holdich DM and Lowery RS (Eds) *Freshwater Crayfish. Biology, Management and Exploitation*. London. Timber Press, Oregon, 365–400.
- Lavens, P. and P. Sorgeloos,** 2000. The history, present status and prospect of the availability of the *Artemia* cysts for aquaculture. *Aquaculture*, 181: 397-403.
- Millamena, D.M, Bombeo, R.F., Jumalon, N.A. and K.L. Simpson,** 1988. Effect of various diets on the nutritional value of *Artemia* species as food for the tiger prawn *Penaeus monodon*. *Mar. Biol.* 98: 217-221.
- Rees, J.F. and Cure, K.,** 1994. Piyatiratitivorakul, S., Sorgeloos, P., Menasveta, P., Highly unsaturated fatty acid requirements of *Penaeus monodon* post larvae. An experimental approach based on artemia enrichment. *Aquaculture*, 122: 193–207.
- Romdhane, M.S., Devresse, B., Leger, P.H. and P. Sorgeloos,** 1995. Effects of feeding (w-3) HUFA-enriched *Artemia* during a progressively increasing period on the larviculture of freshwater prawns. *Aquacult. Int.* 3: 236-242.
- Sorgeloos, P., Lavens, P., Leger, P., Tackaert, W. and D. Versickele,** 1986. Manual for the Culture and Use of Brine Shrimp, *Artemia* in Aquaculture. 319pp
- Anonymous,** 2014. Su Ürünleri İstatistikleri. Türkiye İstatistik Kurumu, Yayın No. 4349, Ankara