

# The Open Access Israeli Journal of Aquaculture – Bamidgen

As from **January 2010** The Israeli Journal of Aquaculture - Bamidgen (IJA) will be published exclusively as an **on-line Open Access (OA)** quarterly accessible by all AquacultureHub (<http://www.aquaculturehub.org>) members and registered individuals and institutions. Please visit our website (<http://siamb.org.il>) for free registration form, further information and instructions.

This transformation from a subscription printed version to an on-line OA journal, aims at supporting the concept that scientific peer-reviewed publications should be made available to all, including those with limited resources. The OA IJA does not enforce author or subscription fees and will endeavor to obtain alternative sources of income to support this policy for as long as possible.

## Editor-in-Chief

Dan Mires

## Editorial Board

**Rina Chakrabarti** Aqua Research Lab, Dept. of Zoology, University of Delhi, India

**Angelo Colorni** National Center for Mariculture, IOLR, Eilat, Israel

**Daniel Golani** The Hebrew University of Jerusalem, Israel

**Hillel Gordin** Kibbutz Yotveta, Arava, Israel

**Sheenan Harpaz** Agricultural Research Organization, Beit Dagan, Israel

**Gideon Hulata** Agricultural Research Organization Beit Dagan, Israel

**George Wm. Kissil** National Center for Mariculture, IOLR, Eilat, Israel

**Ingrid Lupatsch** Swansea University, Singleton Park, Swansea, UK

**Spencer Malecha** Dept. of Human Nutrition, Food & Animal Sciences, CTAHR, University of Hawaii

**Constantinos Mylonas** Hellenic Center for Marine Research, Crete, Greece

**Amos Tandler** National Center for Mariculture, IOLR, Eilat, Israel

**Emilio Tibaldi** Udine University, Udine, Italy

**Jaap van Rijn** Faculty of Agriculture, The Hebrew University of Jerusalem, Israel

**Zvi Yaron** Dept. of Zoology, Tel Aviv University, Israel

**Copy Editor** Miriam Klein Sofer

Published under auspices of  
**The Society of Israeli Aquaculture and Marine Biotechnology (SIAMB)**  
 &  
**University of Hawai'i at Mānoa**  
 &  
**AquacultureHub**  
<http://www.aquaculturehub.org>



UNIVERSITY  
of HAWAII  
MĀNOA  
LIBRARY



**AquacultureHub**  
educate • learn • share • engage

ISSN 0792 - 156X

© Israeli Journal of Aquaculture - BAMIGDEH.

PUBLISHER:

Israeli Journal of Aquaculture - BAMIGDEH -  
Kibbutz Ein Hamifratz, Mobile Post 25210,  
ISRAEL


Phone: + 972 52 3965809

<http://siamb.org.il>



The IJA appears exclusively as a peer-reviewed on-line open-access journal at <http://www.siamb.org.il>. To read papers free of charge, please register online at [registration form](#).

Sale of IJA papers is strictly forbidden.



## Effect of Partial Substitution of Fish Meal with Sunflower Meal on Feed Utilization, Intestinal Digestive Enzyme, Hematological Indexes, Intestinal, and Liver Morphology on Juvenile Turbot (*Scophthal musmaximus* L.)

Huihui Zhou<sup>#</sup>, Chaoqun Li<sup>#</sup>, Fuyun Bian, Mingsan Man, Kangsen Mai, Wei Xu, Gen He\*

Key Laboratory of Aquaculture Nutrition and Feed, Ministry of Agriculture, and the Key Laboratory of Mariculture, Ministry of Education, Ocean University of China, Qingdao 266003, P. R. China

<sup>#</sup> These authors contributed equally to this study and share first authorship

**Keywords:** turbot; sunflower meal; replacement of fish meal; growth performance; hematological antioxidant defense system; intestinal and liver morphology

### Abstract

A 70-day feeding trial was conducted to evaluate effects of partial substitution of fish meal (FM) by sunflower meal (SFM) on juvenile turbot (*Scophthal musmaximus* L.). Five isonitrogenous and isoenergetic diets were formulated with 0%, 15%, 25%, 35%, and 45% replacement of FM protein with protein from SFM. Triplicate groups of juvenile turbot (30 fish per group), were hand-fed twice daily to apparent satiation. Final body weight (FBW), specific growth rate (SGR), and weight gain rate (WGR), were not significantly influenced by type of plant protein at the 15% level ( $P>0.05$ ), while higher levels showed significant reduction of FBW, SGR, WGR. Feed efficiency ratio (FER) and feed intake (FI) were significantly influenced when FM protein was replaced up to 45% ( $P<0.05$ ). Body composition parameters were not affected by SFM substitution but body crude lipid was lowest and ash was highest at 45% ( $P<0.05$ ). Trypsin and diastase values did not vary with experimental diets but lipase activity was significantly reduced ( $P<0.05$ ). Catalase (CAT) values were significantly lower than the control ( $P<0.05$ ) when substitution level reached or exceeded 35%; no significant differences were observed in total antioxidant capacity (T-AOC) and malondialdehyde (MDA) values ( $P>0.05$ ). In the SFM diet groups, all superoxide dismutase (SOD) values were significantly higher than the control ( $P<0.05$ ); villi length and enterocytes were significantly reduced ( $P<0.05$ ), but there was no significant difference ( $P>0.05$ ) in microvilli height between diets; parenchyma structure of liver was severely damaged; smaller hepatocyte areas and areas with high levels of hepatocyte vacuolization and disorganization were present. All results indicated that SFM protein can partially replace FM protein in juvenile turbot diets without adverse effects.

## Introduction

Turbot, a marine, benthic, carnivorous fish, is widely cultured for its high economic value in Europe and East Asia, especially the northern area of China (Miao et al., 2016). However, dietary protein requirement for turbot is high (Danielssen and Hjertnes, 1993; Regost et al., 1999). Aquaculture of turbot is still heavily reliant upon high-quality fish meal.

Fish meal has been widely used in the aquafeed industry as a high-quality feed ingredient because of its well-balanced amino acid composition, good palatability, and low carbohydrate content. Along with decreasing natural fishery resources and increasing demands for fishery products, FM resources have been greatly depleted, resulting in rising prices (Shiu et al., 2015) and supply instability. Therefore, the search for new protein sources, which are cheaper and more abundant, has become imperative. Studies have been undertaken to examine the replacement of FM in diets for turbot by plant proteins such as soybean meal (Yigit et al., 2010, Zhang et al., 2016), corn gluten meal (Regost et al., 1999), rapeseed meal (Burel et al., 2000) and sunflower meal (SFM) (Nogales Merida et al., 2010).

A low-cost, locally available, and highly palatable plant protein source, SFM has produced good results as an alternative plant protein to fish meal. Based on previous studies, FM can be replaced by up to 30% SFM in sea bream juvenile diets (Nogales Merida et al., 2010), 33% in Atlantic salmon (Gill et al., 2006) and 22% in rainbow trout (*Onchorynchus mykiss*) (Martínez, 1984) without any adverse effects on growth performance of fish. Although SFM has high fibre and lignin content (and a low level of lysine), it is rich in methionine with low levels of anti-nutrition factors (Olvera-Novoa et al., 2002, Nogales Merida et al., 2010). Consequently SFM has been identified as a promising ingredient for fish feed, but little is known about the use of SFM in turbot diets. This study investigated the potential of SFM for use as an alternative plant protein source to FM in a carnivorous fish, the juvenile turbot (*Scophthal musmaximus* L.), by studying the effects of substituting FM with SFM on growth, feed utilisation, intestinal digestive enzymes, hematological antioxidant indices, and intestinal and liver morphology. This experiment also provides a reference for further application of plant protein sources in aquaculture and provides more comprehensive information about the effects on flatfish fed diets containing graded levels of SFM.

## Materials and Methods

**Feeding ingredients and diet formulations.** The experimental diets were based on the nutritional requirement of juvenile turbot using FM and SFM as primary protein sources, fish oil and soy lecithin as main lipid sources, and wheat meal as carbohydrate source. SFM, the plant protein resource was supplied by BUNGE. Five isonitrogenous (approximately 50% crude protein) and isoenergetic (approximately 20.0 KJ/g diet of gross energy) diets were formulated to replace 0% (control), 15% (SFM15), 25% (SFM25), 35% (SFM35) and 45% (SFM45) of the FM proteins by SFM proteins. Crystal amino acids (lysine and methionine) were supplemented to meet the essential amino acid (EAA) requirements of juvenile turbot based on the amino acid composition of FM diet. Y<sub>2</sub>O<sub>3</sub> (0.1%) was supplemented as the indicator for apparent digestibility determination according to Glencross (2007).

Diets and experimental ingredients were determined prior to diet formulation. All ingredients were ground into fine powder through 178 µm mesh and gradually mixed. The premix was blended with lipid source ingredients, then water which contained crystal amino acid and choline chloride was added. Pellets (3 mm × 4 mm) were made automatically using a pellet-making machine acquired from South China University of Technology and dried for 12h in a ventilated oven at 55°C. All diets were stored at -20°C until use. Composition of all diets is shown in Table 1.

**Table 1.** Formulas and proximate composition of the experimental diets (% dry matter)

Ingredients	Treatments				
	FM	SFM15	SFM25	SFM35	SFM45
Fish meal <sup>a</sup>	60.00	51.00	45.00	39.00	33.00
Sunflower meal <sup>b</sup>	0.00	12.90	21.50	30.10	38.70
Wheat meal <sup>a</sup>	27.00	20.18	15.82	11.29	6.92
Wheat gluten meal <sup>a</sup>	1.50	2.60	3.30	4.00	4.70
Fish oil	4.00	4.70	5.10	5.60	6.00
Soy lecithin	2.00	2.00	2.00	2.00	2.00
Ca(H <sub>2</sub> PO <sub>4</sub> ) <sub>2</sub>	0.00	0.80	1.30	1.80	2.30
Amino acid	0.00	0.32	0.48	0.72	0.88
Choline chloride	0.30	0.30	0.30	0.30	0.30
Taurine	1.00	1.00	1.00	1.00	1.00
Vitamin premix <sup>c</sup>	1.00	1.00	1.00	1.00	1.00
Mineral premix <sup>d</sup>	2.00	2.00	2.00	2.00	2.00
Calcium propionic acid	0.05	0.05	0.05	0.05	0.05
Ethoxyquinoline	0.05	0.05	0.05	0.05	0.05
Attractant <sup>e</sup>	0.50	0.50	0.50	0.50	0.50
Sodium alginate	0.50	0.50	0.50	0.50	0.50
Yttrium oxide	0.10	0.10	0.10	0.10	0.10
Total	100.00	100.00	100.00	100.00	100.00
Energy (KJ/g)	20.04	20.05	20.04	20.05	20.04
Crude protein	50.32	50.34	50.31	50.32	50.29
Crude lipid	11.10	11.32	11.41	11.59	11.67

Note: FM, diet fish meal; SFM15, replacement of 15% fish meal protein by sunflower meal protein; SFM25, replacement of 25% fish meal protein by sunflower meal protein; SFM35, replacement of 35% fish meal protein by sunflower meal protein; SFM45, replacement of 45% fish meal protein by sunflower meal protein.

<sup>a</sup> Supplied by Great seven Bio-Tech(Qingdao, China); fish meal, crude protein, 73.77%, crude lipid, 7.58%; wheat meal, crude protein, 17.82%, crude lipid, 2.24%; wheat gluten meal, crude protein, 83.31%, crude lipid, 1.75%.

<sup>b</sup> Supplied by BUNGE; sunflower meal, crude protein, 51.47%, crude lipid, 2.67%.

<sup>c</sup> Vitamin premix (mg kg<sup>-1</sup> diet): retinal palmitate, 32; cholecalciferol, 5; DL- $\alpha$ -tocopherol acetate, 240; menadione, 10; thiamin-HCl, 25; riboflavin, 45; pyridoxine-HCl, 20; cyanocobalamin, 10; D-calcium pantothenate, 60; amine nicotinic acid, 200; folic acid, 20; biotin, 60; mesoinositol, 800; ascorbyl polyphosphate (contained 35% ascorbic acid), 2000; microcrystalline cellulose, 16473.

<sup>d</sup> Mineral premix (mg kg<sup>-1</sup> diet): MgSO<sub>4</sub> · 7H<sub>2</sub>O, 1200; CuSO<sub>4</sub> · 5H<sub>2</sub>O, 10; FeSO<sub>4</sub> · H<sub>2</sub>O, 80; ZnSO<sub>4</sub> · H<sub>2</sub>O, 50; MnSO<sub>4</sub> · H<sub>2</sub>O, 45; CoCl<sub>2</sub> · 6H<sub>2</sub>O (1%), 50; Na<sub>2</sub>SeO<sub>3</sub> (1%), 20; calcium iodine, 60; zoelite, 8485.

<sup>e</sup> Attractant: betaine: dimethyl-propiothetin: glycine: alanine: 5-phosphate inosine = 4:2:2:1:1.

**Experimental procedure.** Experiments were conducted in Yi Haifeng Aquatic Product CO. Ltd (Qingdao, China). Juvenile turbot were obtained from a fish rearing farm (Yantai, China) and acclimated to the system for 2 weeks during which they were fed with commercial diets until onset of experiment. After being fasted for 24 h, healthy juvenile turbot (average initial weight 6.77 ± 0.02g) were selected and randomly assigned to 15 experimental fiber glass tanks (30 fish/tank). Seawater was continuously pumped and passed through sand filters into each tank at approximately 3 L/min. Each diet was randomly assigned to three replicate fiber glass tanks. The diets were hand-fed to the fish twice daily at 7:00 and 19:00 to apparent satiation for 70 days. Remaining feed and feces were cleaned immediately after feeding to ensure high water quality. During the experimental period, the water temperature ranged from 19°C-22°C, and salinity from 30‰-33‰, pH ranged from 7.5-8.0, and dissolved oxygen was approximately 7mg/L.

**Sample collection.** After approximately five-weeks of the feeding trial, fecal samples were collected from each tank by siphoning 3-4 hours after feeding. Collected fecal samples were stored at -20°C. At the end of the feeding trial, all the fish were starved for 24h. Total number and body weight of the fish in each tank were determined. Six fish from each tank were randomly sampled and stored at -20°C for whole body composition analysis. The entire intestine and liver were sampled and placed in Bouin's fixative solution and then transferred into 70% ethanol after 24 h for histological evaluation. After being fasted for 24 h, blood samples were taken from the caudal vasculature of six fish from each tank using disposable sterile syringes of 1.0 ml. The blood samples were then centrifuged at 4000g for 10min at 4°C to obtain serum samples which were stored

at -80°C prior to analysis. Intestines were collected and frozen in liquid nitrogen at -80°C for digestive enzyme analysis.

*Chemical analyses:*

*Body composition.* Moisture, crude protein, crude lipid and ash were analyzed for fish samples as well as for diets and experimental ingredients according to standard method (AOAC, 1995). Moisture was measured by drying the samples to constant weight at 105°C. Crude protein ( $N \times 6.25$ ) was determined by using the Kjeldahl method (Kjeltec TM 8400, FOSS, Sweden) after acid digestion. Crude lipid was extracted with diethyl ether using Soxhlet method (Buchi 36680, Switzerland). Ash was measured after combustion in a muffle furnace at 550°C for 16 h.

*Digestibility determination.*  $Y_2O_3$  (0.1%) was supplemented as the indicator for apparent digestibility determination of dry matter and crude protein (Glencross et al., 2007).  $Y_2O_3$  (0.1%) content in feces and diets were determined by Inductively Coupled Plasma-atomic Emission Spectrophotometer (ICP-OES, VISTA-MPX) after acid digestion with perchloric acid.

*Intestinal digestive enzyme and hematological antioxidant indexes determination.* Protease activity was detected using Folin-Phenol method. Lipase and amylase activity was detected using kits from Nanjing Jiancheng Engineering Institute Co., Ltd., China. Catalase (CAT), superoxide dismutase (SOD), and malondialdehyde (MDA) were detected using visible Spectrometry, WST-1, and thiobarbituric acid (TBA) methods (kits were obtained from Nanjing Jiancheng Engineering Institute Co., Ltd., China) respectively. Total antioxidant capacity (T-AOC) was also detected using kits from Nanjing Jiancheng Engineering Institute Co., Ltd., China.

*Intestinal and liver histology.* Samples of distal intestine and liver tissue were fixed and transferred in 70% ethyl alcohol. Approximately 1cm length segments of fixed tissues were cut, routinely dehydrated in an alcohol series, and embedded in paraffin. Then approximately 7µm sections of the tissues were cut, mounted on albumin coated slides, and then stained with matoxylin and eosin (H&E). Morphological structure of these tissues was observed using an imaging microscope (Olympus, DP72, Nikon, Japan).

Intestinal Images were analyzed to determine the ratio (R) between the villi height (VH) or enterocytes height (EH) or microvilli height (MH) and the lumen diameter (LD) of the gut [ $R1 = VH / LD$ ,  $R2 = EH / LD$ ,  $R3 = MH / LD$ , arbitrary units (AU)]. A high R value indicates high villi height or enterocyte height or microvilli height. Liver images were analyzed for hepatocyte area (HA). All data was able to reflect the possible alterations in intestine and liver tissue structure.

*Calculation methods.* The following variables were calculated:

$$\text{Survival rate (SR, \%)} = (\text{final fish number} / \text{initial fish number}) \times 100\%$$

$$\text{Weight gain rate (WGR, \%)} = (\text{final body weight} - \text{initial body weight}) / \text{initial body weight} \times 100\%$$

$$\text{Specific growth rate (SGR, \%)} = (\ln \text{ final body weight} - \ln \text{ initial body weight}) / \text{days} \times 100\%$$

$$\text{Feed intake (FI, \%)} = \text{dry total feed intake} / [(\text{final total body weight} + \text{initial total body weight}) / 2] / \text{days} \times 100\%$$

$$\text{Feed efficiency ratio (FER)} = \text{wet total weight gain (g)} / \text{dry total feed intake (g)}$$

$$\text{Apparent digestibility coefficients (ADC, \%)} = (1 - Y_2O_3 \text{ in the diet} / Y_2O_3 \text{ in feces} \times \text{nutrient in feces} / \text{nutrient in diets}) \times 100\%$$

*Statistical analysis.* All statistical evaluations were analyzed using SPSS 19.0 software. Data were obtained using one-way analysis of variance (ANOVA) followed by Tukey's test. Differences were regarded as significant when  $P < 0.05$ . Data were expressed as means  $\pm$  standard error.

## Results

*Growth performance and feed utilization.* No significant differences ( $P > 0.05$ ) were found in SR among diets (Table 2). There were no obvious differences ( $P > 0.05$ ) in FBW, SGR, and WGR of fish fed 15% SFM (SFM15) replacement of fish meal protein compared with fish fed the FM diet; higher levels of SFM replacement of FM protein significantly reduced ( $P < 0.05$ ) FBW, SGR, and WGR. No significant differences ( $P > 0.05$ ) were found in FER but there was a significant reduction at the 45% level ( $P < 0.05$ ); no significant differences

( $P > 0.05$ ) were found in FI but a significant increase ( $P < 0.05$ ) was found at 45% (see Table 2).

**Table 2.** Growth parameters and feed utilization of juvenile turbot fed the experimental diets

Treatments	Final body weight (g) /FBW	Weight gain rate (%) /WGR	Specific growth rate (%/d) /SGR	Survival rate (%) /SR	Feed efficiency ratio (g/g) /FER	Feed intake (%/d) /FI
FM	59.38±1.76 <sup>a</sup>	776.16±25.96 <sup>a</sup>	3.1±0.04 <sup>a</sup>	100±0.00	1.36±0.02 <sup>a</sup>	1.81±0.03 <sup>a</sup>
SFM15	51.19±2.11 <sup>ab</sup>	656.31±32.05 <sup>ab</sup>	2.89±0.06 <sup>ab</sup>	100±0.00	1.32±0.02 <sup>a</sup>	1.79±0.01 <sup>ab</sup>
SFM25	46.87±2.24 <sup>b</sup>	593.92±34.68 <sup>b</sup>	2.76±0.07 <sup>b</sup>	98.89±1.11	1.24±0.04 <sup>a</sup>	1.86±0.03 <sup>a</sup>
SFM35	45.64±3.10 <sup>bc</sup>	572.5±45.94 <sup>bc</sup>	2.71±0.10 <sup>b</sup>	98.89±1.11	1.19±0.05 <sup>ab</sup>	1.79±0.05 <sup>a</sup>
SFM45	36.5±1.19 <sup>c</sup>	438.63±17.32 <sup>c</sup>	2.4±0.04 <sup>c</sup>	97.78±2.22	1.05±0.06 <sup>b</sup>	1.98±0.03 <sup>b</sup>

Note: Values show mean ± standard error, n = 3; values in the same column with different superscript letters indicate significant difference ( $P < 0.05$ ).

**Body composition.** No significant differences were found in moisture and crude protein contents of fish body among treatments ( $P > 0.05$ ). Clear differences in whole body crude lipid and ash contents were observed in juvenile turbot fed 45% SFM (SFM45) replacement of fish meal protein. They also had the lowest body crude lipid and highest ash content ( $P < 0.05$ ) (see Table 3).

**Table 3.** Whole body composition of juvenile turbot fed the experimental diets

Treatments	Moisture	Crude protein	Crude lipid	Ash
FM	76.56±0.20	16.38±0.13	3.82±0.09 <sup>a</sup>	3.56±0.05 <sup>a</sup>
SFM15	77.54±0.52	15.68±0.45	3.36±0.24 <sup>ab</sup>	3.57±0.06 <sup>a</sup>
SFM25	77.27±0.10	16.16±0.19	3.3±0.18 <sup>ab</sup>	3.73±0.11 <sup>ab</sup>
SFM35	77.5±0.20	16.11±0.20	3.21±0.15 <sup>ab</sup>	3.75±0.08 <sup>ab</sup>
SFM45	77.5±0.17	16.18±0.18	2.87±0.27 <sup>b</sup>	4.03±0.16 <sup>b</sup>

Note: Values show mean ± standard error, n = 3; values in the same column with different superscript letters indicate significant difference ( $P < 0.05$ ).

**Apparent digestibility coefficients.** When FM protein was replaced with up to 35% of SFM protein, apparent digestibility coefficients (ADC) of dry matter decreased significantly with increasing replacement levels of fish meal protein in diets, while ADC of crude protein of SFM45 was significantly lower ( $P < 0.05$ ) than in an FM diet (Table 4).

**Table 4.** Apparent digestibility coefficients (% ADC) for dry matter and crude protein of the experimental diets

Treatments	Dry matter	Crude protein
FM	54.22±1.90 <sup>a</sup>	85.55±1.07 <sup>ab</sup>
SFM15	54.78±0.63 <sup>a</sup>	89.28±0.19 <sup>a</sup>
SFM25	52.54±0.38 <sup>a</sup>	88.23±0.66 <sup>a</sup>
SFM35	38.68±2.80 <sup>b</sup>	86.2±1.21 <sup>ab</sup>
SFM45	36.09±0.89 <sup>b</sup>	83.64±0.81 <sup>b</sup>

Note: Values show mean ± standard error, n = 3; values in the same column with different superscript letters indicate significant difference ( $P < 0.05$ ).

*Intestinal digestive enzyme activity and hematological antioxidant indexes.* Trypsin and diastase values showed no variation among experimental diets ( $P > 0.05$ ). There was a significant ( $P < 0.05$ ) reduction in lipase activity in the fish meal replacement groups compared with the control (see Table 5).

**Table 5.** Activity of intestinal digestive enzyme of juvenile turbot fed the experimental diets

Treatments	trypsin (10U/mg)	diastase AMS(U/mg)	lipase LPS(U/mg)
FM	11.81±0.89	0.58±0.11	70.10±2.45 <sup>a</sup>
SFM15	11.82±0.23	0.93±0.02	54.85±2.05 <sup>bc</sup>
SFM25	12.12±1.06	0.92±0.09	50.95±1.09 <sup>c</sup>
SFM35	12.03±0.12	0.65±0.14	51.21±0.97 <sup>c</sup>
SFM45	12.22±0.23	0.69±0.15	59.77±0.78 <sup>b</sup>

Note: Values show mean ± standard error, n = 3; values in the same column with different superscript letters indicate significant difference ( $P < 0.05$ ).

T-AOC and CAT values were affected by treatment used: CAT values were significantly lower than the control when substitution level reached, or exceeded, 35%, while no significant differences were obtained in total antioxidant capacity (T-AOC) and malondialdehyde (MDA) values compared to the control ( $P > 0.05$ ). All values of superoxide dismutase (SOD) in SFM treatments were significantly higher ( $P < 0.05$ ) than the control. (see Table 6).

**Table 6.** Antioxidant indexes of juvenile turbot fed the experimental diets

Treatments	T-AOC(U/ml)	CAT(U/ml)	SOD(U/ml)	MDA(U/ml)
FM	8.36±0.41 <sup>ab</sup>	2.00±0.02 <sup>ab</sup>	15.17±0.66 <sup>a</sup>	8.57±0.22
SFM15	9.74±0.26 <sup>a</sup>	2.33±0.67 <sup>a</sup>	34.16±0.54 <sup>d</sup>	7.29±0.49
SFM25	8.14±0.50 <sup>ab</sup>	1.67±0.12 <sup>bc</sup>	26.15±0.26 <sup>c</sup>	8.90±0.33
SFM35	8.12±0.20 <sup>ab</sup>	1.48±0.14 <sup>c</sup>	24.21±0.26 <sup>b</sup>	8.90±0.41
SFM45	7.52±0.58 <sup>b</sup>	1.27±0.09 <sup>c</sup>	22.74±0.34 <sup>b</sup>	7.48±0.34

Note: Values show mean ± standard error, n = 3; values in the same column with different superscript letters indicate significant difference ( $P < 0.05$ ).

*Intestinal and liver histology.* Intestinal and liver samples of fish from different treatments were compared to those of the control. SFM replacement diets significantly ( $P < 0.05$ ) reduced the length of villi and enterocytes, but microvilli height showed no significant differences ( $P > 0.05$ ) among diets (Table 7, Fig. 1). The parenchyma structure of liver from fish fed diets containing graded levels of SFM was significantly damaged. Smaller hepatocyte areas, and areas with high levels of hepatocyte vacuolisation and disorganisation, were present in some samples from fish fed SFM diets (Fig. 2).

**Table 7.** Distal intestinal and liver histology indexes of juvenile turbot fed the experimental diets

Treatments	$R_1^a$ ( $10^{-1}$ )	$R_2^b$ ( $10^{-2}$ )	$R_3^c$ ( $10^{-3}$ )	Hepatocyte area ( $\mu\text{m}^2$ )
FM	3.48±0.04 <sup>a</sup>	1.97±0.08 <sup>a</sup>	1.79±0.06	377.7±22.14 <sup>a</sup>
SFM15	2.88±0.07 <sup>b</sup>	1.62±0.03 <sup>b</sup>	1.8±0.08	315.89±9.89 <sup>b</sup>
SFM25	2.77±0.06 <sup>b</sup>	1.62±0.05 <sup>b</sup>	1.8±0.09	311.07±9.66 <sup>b</sup>
SFM35	2.47±0.06 <sup>bc</sup>	1.52±0.06 <sup>b</sup>	1.57±0.05	326.28±10.41 <sup>b</sup>
SFM45	2.64±0.07 <sup>c</sup>	1.39±0.03 <sup>b</sup>	1.54±0.08	313.95±4.66 <sup>b</sup>


Note: Values show mean ± standard error, n = 3; values in the same column with different superscript letters indicate significant difference ( $P < 0.05$ ).

<sup>a</sup>  $R_1$  = villi height/ lumen diameter

<sup>b</sup>  $R_2$  = enterocytes height/ lumen diameter


<sup>c</sup>  $R_3$  = microvilli height/ lumen diameter





**Fig.1** Transverse section photomicrographs of turbot's distal-intestine. Villi height (VH) was analyzed at a magnification of  $\times 40$ ; (B) enterocytes height (EH) and microvilli height (MH) were analyzed at higher magnification of  $\times 400$ .





**Fig. 2** Transverse section photomicrographs of turbot's liver (magnification  $\times 200$ )

### Discussion

The present study indicated that FM could be replaced by SFM at the level of 15% without any adverse effects on growth and feed utilisation of juvenile turbot. However, FBW, SGR, WGR and FER were significantly reduced with increasing levels of FM substitution. The acceptable replacement rate by SFM in this study was lower than that by soybean meal in the diet of turbot (Zhang *et al.*, 2016). SFM substitution at 14% had no adverse effects on the growth performance on tilapia (Furuya *et al.* (2000). SFM substitution level of FM up to 33% had no adverse effects on growth performance in Atlantic salmon (Gill *et al.* 2006).

Many authors attributed lower growth with increasing levels of SFM substitution to essential amino acid (EAA) deficiency; improved growth was observed in rainbow trout fed diets containing SFM supplemented with Met, Lys, and Leu than diets without EAA supplementation (Sanz *et al.*, 1994). EAA could be the first limiting factor for replacement of FM by high levels of SFM (Olvera-Novoa *et al.* 2002). In the present study, despite the supplemented lysine and methionine based on the EAA composition of an FM diet, other EAA such as arginine, associated with fish growth (Fournier *et al.*, 2002), could cause

reduction in growth. Some authors regard the suppression of feed intake (FI) as a reason for unsatisfactory growth performance with plant protein replacement in diets for fish (Chen *et al.*, 2011; Peng *et al.*, 2013). Together with rising substitution levels of FM, FI fluctuation was observed in this study. It is possible that higher fibre content in SFM causes reduced growth of fish. Similar results were seen in sea bream when the fibre content in diets was increased (Sanchez-Lozano *et al.* 2007).

High fibre and lignin contents in SFM could affect the digestibility of protein or dry matter in carnivorous fish (Olvera-Novoa *et al.*, 2002, Stickney *et al.*, 1996). In the present study, no adverse effects on apparent digestibility coefficients (ADC) of dry matter were obtained up to 35% of FM protein replacement. Significant ADC reduction of crude protein was observed only at 45% level. Results confirmed that relatively low FM substitution levels by SFM did not affect the ADC of protein or dry matter in turbot.

In general, plant protein such as soybean meal containing anti-nutrition factors (ANFs) could affect digestive enzyme activity (Yu *et al.*, 2013); however, in this study, values of trypsin and diastase content showed no variation among treatments but there was a significant reduction in lipase activity with increasing substitution levels of FM. The adverse influence on lipase activity could be due to patho-morphological changes in the distal intestine and the presence of ANFs in SFM. Reduction in lipase activity indicated that high substitution levels of FM impaired digestion and absorption of lipids.

Animals have a powerful antioxidant defence system, which can reduce the physiological damage to the body by removing oxygen free radicals. SOD, CAT, T-AOC, and MDA play extremely important roles in this antioxidant defence system. MDA is a final product of lipid peroxidation which can reflect the degree of endogenous oxidative damage (Ding *et al.*, 2015). Hematological antioxidant parameters showed the lowest MDA level and highest T-AOC, SOD, and CAT activities in SFM15. This indicated that antioxidant defence mechanisms in the body had been developed to reduce oxidative stress and protect biological systems from free radical toxicity (Nordberg and Arner, 2001) when the substitution level reached 15%. An increasing trend in MDA and decreasing trend in T-AOC, SOD, and CAT were observed when the substitution level reached, or exceeded 25%. This indicated that higher SFM levels (above 15%) in diets could cause severe oxidative stress and free radical toxicity which could not be alleviated by the antioxidant defence system of the fish body, consistent with results seen in other fish fed diets containing plant proteins (Lin and Li, 2011; Peng *et al.* 2013). The anti-nutrition factors (ANFs) in SFM were considered to have been the principal cause of observed damage.

The parenchyma structure of the liver from fish fed SFM diet was significantly damaged. This differed from findings of Nogales Merida *et al.* (2010). Areas with high levels of hepatocyte vacuolisation and disorganisation were present and the volume of hepatocytes showed significant shrinkage in some samples from fish fed SFM diets. As the main site for toxin accumulation in the animal body, both the structure and function of the liver would be damaged by higher levels of ANFs from SFM. Further research is needed to establish the effects of prolonged feeding with SFM.

This study indicated that SFM substitution treatments significantly reduced the height of both villi and enterocytes. These could be signs of damage to the structure of the distal intestine at high substitution levels of FM. However the microvilli height in the present study showed no significant differences among treatments, which suggests that intestinal function was not significantly impaired by higher substitution levels of FM. This mirrored the result of Nogales Merida *et al.* (2010).

In conclusion, this study investigated the potential for SFM to be used as an alternative plant protein to FM in juvenile turbot (*Scophthal musmaximus* L.) diets. It was found that FM could be replaced by SFM at the 15% level with no significant adverse effects on growth and feed utilisation in turbot. Low growth associated with increased SFM substitution levels was considered to be due to the adverse effects on ADC of dry matter, intestinal digestive enzyme activity, the hematological antioxidant defence system, and the histology of intestine and liver.

### Acknowledgements

The study was supported by Special Fund for Agro-scientific Research in the Public Interest (201303053) to G.H., 973 program (2014CB138602) to K.M., and Shandong Provincial Natural Science Foundation (JQ201206) to G.H.

### References

- Association of Official Analytical Chemists (AOAC)**, 1995. Official Methods of Analysis of Official Analytical Chemists International 16<sup>th</sup> ed. Association of Official Analytical Chemists, Arlington, VA, USA
- Burel, C., Boujard, T., Kaushik, S.J., Boeuf, G., Van der Geyten, S., Mol, K.A., Kuhn, E.R., Quinsac, A., Krouti, M., Ribaillier, D.**, 2000. Potential of plant-protein sources as fish meal substitutes in diets for turbot (*Psetta maxima*): growth, nutrient utilisation and thyroid status. *Aquaculture*. 188, 363-382.
- Chen, W., Ai, Q., Mai, K., Xu, W., Liufu, Z., Zhang, W., Cai, Y.**, 2011. Effects of dietary soybean saponins on feed intake, growth performance, digestibility and intestinal structure in juvenile Japanese flounder (*Paralichthys olivaceus*). *Aquaculture*. 318, 95-100.
- Danielssen, D.S., Hjertnes, T.**, 1993. Effect of dietary protein levels in diets for turbot (*Scophthalmus maximus* L.) to market size. [INRA Colloquia; Fish nutrition in practice]. 61, 89-96.
- Ding, Z.L., Zhang, Y.X., Ye, J.Y., Du, Z.Y., Kong, Y.Q.**, 2015. An evaluation of replacing fish meal with fermented soybean meal in the diet of *Macrobrachium nipponense*: Growth, nonspecific immunity, and resistance to *Aeromonas hydrophila*. *Fish Shellfish Immunol*. 44, 295-301.
- Fournier, V., Guillou-Coustans, M.F., Metailler, R., Vachot, C., Guedes, M.J., Tulli, F., Oliva-Teles, A., Tibaldi, E., Kaushik, S.J.**, 2002. Protein and arginine requirements for maintenance and nitrogen gain in four teleosts. *British J Nutr*. 87, 459-469.
- Furuya, V.R.B., Furuya, W.M., Hayshi, C., Soares, C.M.**, 2000. Niveles de inclusión de harina de girasol en la alimentación de la tilapia del Nilo (*Oreochromis niloticus*) en etapa juvenil. *Zootecnia Trop*. 18 (1), 1-10.
- Gill, N., Higgs, D.A., Skura, B.J., Rowshandeli, M., Dosanjh, B.S., Mann, J., Gannam, A.L.**, 2006. Nutritive value of partially dehulled and extruded sunflower meal for post-smolt Atlantic salmon (*Salmo salar* L.) in sea water. *Aquacult Res*. 37, 1348-1359.
- Glencross, B., Hawkins, W., Veitch, C., Dods, K., McCafferty, P., Hauler, R.**, 2007. The influence of dehulling efficiency on the digestible value of lupin (*Lupinus angustifolius*) kernel meal when fed to rainbow trout (*Oncorhynchus mykiss*). *Aquacult Nutr*. 13, 462-470.
- Lin, S., Luo, L.**, 2011. Effects of different levels of soybean meal inclusion in replacement for fish meal on growth, digestive enzymes and transaminase activities in practical diets for juvenile tilapia, *Oreochromis niloticus* x *O. aureus*. *Anim Feed Sci Technol*. 168, 80-87.
- Martínez, C.A.**, 1984. Advances in the substitution of fish meal and soybean meal by sunflower meal in diets of rainbow trout (*Salmo Gairdneri* L.). *An. Inst. Cienc. Mar Limnol. Univ. Nac. Auton. Méx.* 13 (2), 345-350.
- Miao S., Nie Q., Miao H., Zhang W.\*, Mai K.**, 2016. Effects of dietary carbohydrates with different molecular complexity on growth performance, feed utilization and metabolic responses of juvenile turbot *Scophthalmus maximus*. *Isr. J. Aquacult.-Bamidgeh*, IJA\_68.2016.1307, 9 pgs.
- Nogales Merida, S., Tomas-Vidal, A., Martinez-Llorens, S., Jover Cerda, M.**, 2010. Sunflower meal as a partial substitute in juvenile sharpnose sea bream (*Diplodus puntazzo*) diets: Amino acid retention, gut and liver histology. *Aquaculture*. 298, 275-281.
- Nordberg, J., Arner, E.S.J.**, 2001. Reactive oxygen species, antioxidants, and the mammalian thioredoxin system. *Free Rad Biol Med.*, 31, 1287-1312.

- Olvera-Novoa, M.A., Olivera-Castillo, L., Martinez-Palacios, C.A.,** 2002. Sunflower seed meal as a protein source in diets for *Tilapia rendalli* (Boulanger, 1896) fingerlings. *Aquacult Res.*, 33, 223-229.
- Peng, M., Xu, W., Ai, Q., Mai, K., Liufu, Z., Zhang, K.,** 2013. Effects of nucleotide supplementation on growth, immune responses and intestinal morphology in juvenile turbot fed diets with graded levels of soybean meal (*Scophthalmus maximus* L.). *Aquaculture*. 392, 51-58.
- Regost, C., Arzel, J., Kaushik, S.J.,** 1999. Partial or total replacement of fish meal by corn gluten meal in diet for turbot (*Psetta maxima*). *Aquaculture*. 180, 99-117.
- Sanchez Lozano, N.B., Tomas Vidal, A., Martinez-Llorens, S., Nogales Merida, S., Espert Blanco, J., Monino Lopez, A., Pla Torres, M., Jover Cerda, M.,** 2007. Growth and economic profit of gilthead sea bream (*Sparus aurata*, L.) fed sunflower meal. *Aquaculture*. 272, 528-534.
- Sanz, A., Morales, A.E., Delahiguera, M., Cardenete, G.,** 1994. Sunflower meal compared with soybean-meal as partial substitutes for fish-meal in rainbow-trout (*Oncorhynchus mykiss*) diets - protein and energy- utilization. *Aquaculture*. 128, 287-300.
- Shiu, Y.L., Hsieh, S.L., Guei, W.C., Tsai, Y.T., Chiu, C.H., Liu, C.H.,** 2015. Using *Bacillus subtilis* E20-fermented soybean meal as replacement for fish meal in the diet of orange-spotted grouper (*Epinephelus coioides*, Hamilton). *Aquaculture Research*. 46, 1403-1416.
- Stickney, R.R., Hardy, R.W., Koch, K., Harrold, R., Seawright, D., Masee, K.C.,** 1996. The effects of substituting selected oilseed protein concentrates for fish meal in rainbow trout *Oncorhynchus mykiss* diets. *J World Aquacult Soc.*, 27, 57-63.
- Wang, X.-F., Li, X.-Q., Leng, X.-J., Shan, L.-L., Zhao, J.-X., Wang, Y.-T.,** 2014. Effects of dietary cottonseed meal level on the growth, hematological indices, liver and gonad histology of juvenile common carp (*Cyprinus carpio*). *Aquaculture*. 428, 79-87.
- Yigit, M., Ergun, S., Turker, A., Harmantepe, B., Erteken, A.,** 2010. Evaluation of soybean meal as a protein source and its effect on growth and nitrogen utilization of black sea turbot (*Psetta maeotica*) juveniles. *J Marine Sci Technol-Taiwan*. 18, 682-688.
- Yu, D.H., Gong, S.Y., Yuan, Y.C., Lin, Y.C.,** 2013. Effects of replacing fish meal with soybean meal on growth, body composition and digestive enzyme activities of juvenile Chinese sucker, *Myxocyprinus asiaticus*. *Aquacult Nutr.*, 19, 84-90
- Zhang B., He G.\*, Wang X., Pi X.\*, Mai K., Zhou H.,** 2016. Effects of replacing fish meal with soybean meal or fermented and phytase-treated soybean meal respectively on growth performance, feed utilization, and apparent digestibility coefficients in juvenile turbot (*Scophthalmus maximus* L.). [Isr. J Aquacult.-Bamidgeh](#), IJA\_68.2016.1283, 8 pgs.