



Sveriges lantbruksuniversitet
Swedish University of Agricultural Sciences

This is an author produced version of a paper published in *Journal of Soils and Sediments*.

This paper has been peer-reviewed but may not include the final publisher proof-corrections or pagination.

Citation for the published paper:

Gunnar Börjesson, Holger Kirchmann, Thomas Kätterer. (2014) Four Swedish long-term field experiments with sewage sludge reveal a limited effect on soil microbes and on metal uptake by crops. *Journal of Soils and Sediments*. Volume: 14, Number: 1, pp 164-177.

<http://dx.doi.org/10.1007/s11368-013-0800-5>.

Access to the published version may require journal subscription.

Published with permission from: Springer.

Standard set statement from the publisher:

“The final publication is available at Springer via <http://dx.doi.org/10.1007/s11368-013-0800-5>”

Epsilon Open Archive <http://epsilon.slu.se>

1 SOILS, SEC # • RESEARCH ARTICLE

2

3 **Four Swedish long-term field experiments with sewage sludge reveal a limited effect on**
4 **soil microbes and on metal uptake by crops**

5

6 **G. Börjesson • H. Kirchmann • T. Kätterer**

7

8 **Affiliations-addresses:**

9 Gunnar Börjesson (✉) • Holger Kirchmann

10 Department of Soil and Environment, Swedish University of Agricultural Sciences,

11 P.O. Box 7014, 75007 Uppsala, Sweden

12 e-mail: gunnar.borjesson@slu.se

13

14 Thomas Kätterer

15 Department of Ecology, Swedish University of Agricultural Sciences, P.O. Box 7044, 75007

16 Uppsala, Sweden

17

18

19 (✉) **Corresponding author:**

20 Telephone: +46 18 67 27 53

21 Fax: +46 18 67 31 56

22 e-mail: gunnar.borjesson@slu.se

23

24 **Abstract**

25 Purpose: To study the effect of sewage sludge amendment on crop yield and on microbial
26 biomass and community structure in Swedish agricultural soils.

27 Materials and methods: Topsoil samples (0-0.20 m depth) from four sites where sewage
28 sludge had been repeatedly applied during 14-53 years were analysed for total C, total N, pH
29 and PLFAs (phospholipid fatty acids). Heavy metals were analysed in both soil and plant
30 samples, and crop yields were recorded.

31 Results and discussion: At all four sites, sewage sludge application increased crop yield and
32 soil organic carbon. Sludge addition also resulted in elevated concentrations of some heavy
33 metals (mainly Cu and Zn) in soils, but high concentrations of metals (Ni and Zn) in plant
34 materials were almost exclusively found in the oldest experiment, started in 1956. PLFA
35 analysis showed that the microbial community structure was strongly affected by changes in
36 soil pH. At those sites where sewage sludge had caused low pH, Gram-positive bacteria were
37 more abundant. However, differences in community structure were larger between sites than
38 between the treatments.

39 Conclusions: At all four sites long-term sewage sludge application increased the soil organic
40 carbon and nitrogen content, microbial biomass and crop yield. Long-term sewage sludge
41 application led to a decrease in soil pH. Concentrations of some metals had increased
42 significantly with sewage sludge application at all sites, but the amounts of metals added to
43 soil with sewage sludge were found not to be toxic for microbes at any site.

44

45

46 **Keywords** Heavy metals • Long-term field experiments • Mycorrhiza • Phospholipid fatty
47 acids • Microbial community structure

48

49 **1 Introduction**

50 Soil organic matter is a key attribute of soil fertility. The pool of soil organic C can be
51 increased either indirectly by applying mineral fertilisers, through higher yields, or directly by
52 adding organic amendments such as sewage sludge (Kätterer et al. 2013). Sewage sludge
53 provides organic matter and essential plant nutrients, but may also contain unwanted metals,
54 xenobiotic substances and pathogens (Singh and Agrawal 2008).

55 In Sweden, the political goal is that all kinds of waste are recycled and that "by 2015 at
56 least 60 per cent of phosphorus compounds present in wastewater will be recovered for use on
57 productive land" (Environmental Objectives Portal 2012). The production of sewage sludge
58 from 402 wastewater treatment plants in Sweden was 203 500 Mg dw in 2010 and 25% of this
59 was used in agriculture (Statistics Sweden 2012). This is less than in the Scandinavian
60 neighbour countries Norway and Denmark (57% and 54% in 2009; Eurostat, 2012), but more
61 than in Finland (3%; Lindfors 2012). The ReVAQ project in Sweden has led to a certification
62 system for sludge which is run by the Swedish Water and Wastewater Association (Kärman
63 et al. 2007). The aim is to reduce heavy metals and other contaminants in wastewater, a
64 process which allows for continuous improvement of the incoming wastewater, as a way of
65 improving the quality of the sludge applied on agricultural land. Today there are over 37
66 wastewater treatment plants producing sludge certified by ReVAQ, corresponding to over
67 50% of Swedish sludge from wastewater (Revaq 2012), but the use of sewage sludge as a
68 fertiliser in agriculture is still the subject of much debate, mostly concerning the possibilities
69 for accumulation of heavy metals, especially cadmium (*e.g.* Linderholm et al. 2012).

70 The main objective of this study was to investigate the effect of continuous application of
71 sewage sludge on soil microbial biomass and community structure. Long-term experiments
72 are needed to be able to detect effects on the microbial biomass (Diacono and Montemurro
73 2010). Amendment with organic materials such as sewage sludge generally increases the
74 microbial biomass, but high doses may cause negative effects on enzymatic activities in soil
75 (Singh and Agrawal 2008), and inconclusive results presented in the literature makes it
76 uncertain if sludge addition will affect microbial community structure, and whether these
77 potential changes are negative for soil health and fertility (MacDonald et al. 2011).

78 Cadmium (Cd) was also of special interest due to its high toxicity, long body retention
79 time and high mobility in the environment (Alloway and Jackson 1991). Its concentration has
80 increased in soils as a result of industrial emissions with subsequent atmospheric deposition,
81 but emissions have decreased since the mid-1970s, due to cleaning at point sources (Pacyna et

82 al. 2009). The average concentrations of Cd in sewage sludge in Sweden are now below the
83 Swedish threshold value of 2 mg kg⁻¹ DM (Suppl. Table A). Due to less atmospheric
84 deposition and use of P fertiliser with a low Cd content, the Cd content in Swedish wheat
85 grains has decreased from the peak concentrations recorded during the 1970s (Kirchmann et
86 al. 2009). Therefore, we hypothesized that the effect of heavy metals on soil microorganisms,
87 as a result of repeated sewage sludge amendment, was not significant.

88

89 **2 Methods**

90 2.1 Sites

91 Soil samples were taken for analysis from four sites where sewage sludge has been repeatedly
92 applied in long-term field experiments situated in different parts of Sweden; Ultuna, Lanna,
93 Petersborg and Igelösa (Table 1). In these four experiments, at least one sewage sludge
94 treatment is included in the experimental design. At Ultuna and Lanna, the sewage sludge
95 treatments are being compared with other organic and mineral fertilisers, while at Igelösa and
96 Petersborg comparisons are being made between different levels of sewage sludge and
97 different levels of mineral fertiliser. All sites are designed as block experiments, with four full
98 replicates per treatment.

99 In the Ultuna experiment, all organic fertilisers, including sewage sludge, are applied
100 every second year, corresponding to 4 Mg C ha⁻¹. The Lanna experiment has a similar design,
101 with 8 Mg ash-free dry matter ha⁻¹ applied every second year. At Lanna, an additional sewage
102 sludge treatment is included in which metal salts (Cd 0.098, Cu 3.036 and Ni 6.250 kg ha⁻¹)
103 are added together with the sludge. The experiments at Petersborg and Igelösa have an
104 identical design, with three levels of sewage sludge (0, 4 or 12 Mg dry matter ha⁻¹ every
105 fourth year) compared with three levels of NPK fertiliser (0 N, ½ normal N and normal N).
106 The plots are 6 m × 20 m at Igelösa and Petersborg, 8 (or 6) m × 14 m at Lanna . These three
107 sites have normal conventional management, including mould-board ploughing to a depth of
108 22-24 cm. At Ultuna, the plots are 2 m × 2 m and the soil is manually managed within frames,
109 down to a depth that should be comparable to the other sites.

110 The sewage sludge applied to plots is obtained from nearby wastewater plants,
111 Kungsängsverket in Uppsala (Ultuna), Ryaverket in Gothenburg (Lanna), Sjölund in Malmö
112 (Petersborg) and Källby in Lund (Igelösa). All of these in principle use the same method for P
113 removal, which is precipitation with FeCl₃. Prior to 1976, AlSO₄ was used for precipitation in
114 Uppsala. The concentration of heavy metals has decreased since the 1970's in all these

115 wastewaters, exemplified by a ten-fold decrease of Cd and Pb in the Uppsala sludge
116 (Börjesson et al. 2012). Recent mean values for sewage sludge in Sweden are given in Suppl.
117 Table A.

118 In the Ultuna experiment, different spring-sown crops have been grown. At Lanna, only
119 cereals, mainly oats and barley, have been used. Petersborg and Igelösa have a 4-year crop
120 rotation, with a few exceptions consisting of winter wheat, sugar beet, spring barley and
121 oilseed rape.

122

123 2.2 Soil sampling

124 Topsoil samples (0-20 cm depth) from the four sites were taken from plots that had received
125 sewage sludge, and for comparison, from unfertilised and N-fertilised plots. Ultuna was
126 sampled in September 2009 and Lanna in (autumn) 2010, just before the biennial application
127 of sludge. Igelösa and Petersborg were sampled in June 2011, two years after sludge
128 spreading. Samples from Igelösa were taken from all plots with different levels of sewage
129 sludge, but without mineral N fertilisation, while at Petersborg sampling also included the
130 highest level of N fertilisation combined with the highest level of sewage sludge and the
131 control without sewage sludge.

132

133 2.3 Analyses

134 Grain, dried green mass, and soil samples were analysed after digestion in 2M HNO₃ with an
135 inductively coupled plasma-mass spectrometer (Elan 6100 ICP-MS; Perkin Elmer, Waltham
136 MA 02451, USA) for eight metals (Al, Cd, Cu, Fe, Mn, Ni, Pb and Zn) in topsoil and nine
137 metals (Cd, Co, Cr, Cu, Mo, Ni, Pb, Se and Zn) in crop material. Concentrations of total soil
138 C and total N were determined by dry combustion (LECO CNS Analyzer; LECO
139 Corporation, St Joseph MI 49085, USA) and pH was measured in distilled water.

140 Phospholipid fatty acids (PLFA) were analysed according to the method described by
141 Börjesson et al. (2012), whereby 1 g freeze-dried sample was fractionated with 5 ml
142 chloroform, 20 ml acetic acid and 5 ml methanol. The methanol phase was used for PLFA
143 analysis in all samples. The chloroform phase was also saved for determination of the neutral
144 lipid fatty acid (NLFA) 16:1 ω 5 in samples from Petersborg and Igelösa. For principal
145 component analysis (PCA), 25 PLFAs that were present at all sites were converted to
146 percentage of total moles for evaluation of microbial community structure.

147 Statistical analyses were done with the use of JMP software ver. 6.0.3 (SAS Institute Inc.,
148 Cary NC 27513, USA); including analysis of variance, comparison of treatment effects
149 (Tukey-Kramer HSD), and principal components obtained from covariance matrices of PLFA
150 data.

151

152 **3 Results**

153 3.1 Yields and soil pH

154 At all four sites, sewage sludge had exerted a positive effect on crop yields (4% per year or
155 more compared with unfertilised plots) and on soil organic matter levels (Fig. 1, Table 2 and
156 3). It should be noted that normal fertilisation differs between the sites. It is 80 kg N ha⁻¹ year⁻¹
157 at Ultuna and Lanna, while the dosage at Petersborg and Igelösa is according to fertiliser
158 recommendations in the area, i.e. 140 kg N ha⁻¹ for wheat and slightly lower for other grains.

159 Soil pH had dropped in the sewage sludge treatment at the Ultuna site from 6.5 at the
160 start of the experiment to 4.9 at the time of sampling in 2009, while pH had remained between
161 6.1 and 6.7 in the other treatments. At Lanna, the pH had also dropped in the plots treated
162 with sewage sludge, although so far only from 6.6 in 1996 to 6.1 in 2010. The other two sites
163 were limed in 1998, and in 2010 the pH values ranged between 6.9 and 7.3 at Petersborg and
164 between 7.4 and 7.7 at Igelösa.

165

166 3.2 Metals in soil

167 The concentrations of metals were generally much higher in soil samples from Ultuna than
168 soils from the other sites, not only in the sewage sludge-treated soil but also in the normally
169 fertilised and unfertilised soils, which can be regarded as background (Table 4), due to the
170 geological history in this region (Eriksson et al. 2010). At Ultuna, the concentrations of Cd,
171 Ni, Pb and Zn in the sewage sludge-treated soil were higher than background levels,
172 seemingly due to high inputs during the early years of the experiment. For example, the
173 concentration of Cd in sewage sludge amounted to 9.1 mg kg⁻¹ in 1972, but then decreased
174 rapidly to around 2 mg kg⁻¹, and in 2009 it was down at 0.65 mg kg⁻¹. No increase in heavy
175 metal concentrations has been observed since the 1970s, according to historical data (Table
176 4). An exception is Cu, soil concentrations of which have increased steadily because Cu levels
177 in sludge have remained high. High concentrations of Fe in the Ultuna sludge-treated soil are
178 an effect of addition of FePO₄, produced by P removal through precipitation with FeCl₃
179 during wastewater treatment.

180 At Lanna, metal concentrations in soils were generally low (Table 5), but those of Cu and
181 Zn had increased in all sewage sludge-treated plots, while Cd and Ni concentrations were only
182 significantly higher when these metals were added as soluble salts together with the sludge
183 (Treatment G). For the other metals, including Pb, no significant differences were observed.

184 Data from the experiments at Igelösa and Petersborg are reported in detail by Andersson
185 (2012). However, it should be noted that average Cu concentrations were also higher at these
186 sites through sewage sludge addition (Table 6). For the other metals, there has been a trend
187 for a slight increase, although barely significant.

188 Sewage sludge amendment decreased soil pH and the correlations between pH values and
189 metal concentrations were strongly negative (Suppl. Table B), most obviously for Zn ($r=-$
190 0.858).

191 192 3.3 Metals in plants

193 Analysis of metals in samples of silage maize harvested in 2009 in the Ultuna experiment
194 showed that only Ni and Zn levels were significantly higher in plant material from the sewage
195 sludge-treated plots, compared to the control plots (Table 7). Cadmium levels in crop samples
196 from both the unfertilised and the sewage sludge treated-plots were considerably higher than
197 in the corresponding treatments at the other sites. In order to compare silage maize with
198 previous metal uptake by crops in this treatment, data from fodder rape, which was also
199 harvested as green biomass, were used (Table 7). With the exception of Cu and Zn, which had
200 similar uptake, metal concentrations were considerably lower in 2009 than in 1974.

201 Analyses of metals in wheat grain at Lanna (Table 8) showed that there was significantly
202 more Ni in sewage sludge plots when spiked with Ni in salt solution. Similarly, there were
203 elevated levels of Cd in the treatment with sludge spiked with metals, but for Cr, Co, Cu, Pb
204 and Zn no significant differences were found. Molybdenum was found to be higher in crops
205 fertilised with calcium nitrate (CaNO_3). However, Mo in grain was correlated with soil pH
206 ($r=0.94$; $p=0.0061$), similarly to what was found in silage maize from Ultuna.

207 An interesting observation is the effect of N fertilisation on Cd uptake in wheat grain at
208 Igelösa (Fig. 2). Accumulation of Cd was low in that soil and plant uptake was not affected by
209 sewage sludge application, but concentrations increased in crops as an effect of NPK
210 fertilisation, with N was applied as ammonium nitrate (N27).

211 212 3.4 PLFAs

213 The concentrations of total PLFAs were highest in the sewage sludge-treated soils at all sites
214 (Tables 9-11). Correlations between soil organic matter and total PLFA content were highly
215 positive at all sites, and PLFA:soil C ratios were similar independent of site or experimental
216 treatment (Fig. 1).

217 Data from the Ultuna experiment showed that almost all the PLFAs identified had
218 increased in the sewage sludge treatment. The variation within treatments was rather high,
219 with CV (coefficient of variation = standard deviation divided by mean value) up to 25% in
220 the sewage sludge treatment. There was a shift in the sludge plots towards more branched
221 PLFAs (*e.g.* i16:0, i17:0, br18:0, 10Me17:0), indicating relatively more Gram-positive
222 bacteria compared with other bacterial groups (Table 9). There was also a strong effect on
223 cy17:0 and cy19:0.

224 In samples from Lanna, the highest total values were obtained from the two sewage
225 sludge treatments, and the addition of extra heavy metals as soluble salts did not seem to
226 affect the microbes at all (Table 10). The sludge-treated plots also had the highest
227 concentrations of most individual PLFAs. In contrast to the Ultuna site, the effect of sludge
228 treatments on mono-unsaturated PLFAs, *e.g.* 16:1 ω 7 and 18:1 ω 7, was not negative in Lanna.

229 In samples from Petersborg and Igelösa, sludge amendment caused a 31% increase in
230 total PLFAs at Petersborg and a 33% increase at Igelösa, comparing the highest application
231 rate with the unfertilised control (Table 11). Almost all individual PLFAs increased in equal
232 proportions due to sludge amendment on these two sites, *i.e.* no inhibitory effect could be
233 observed. If we calculate the PLFAs as concentrations (nmol PLFA g soil⁻¹), some of the
234 PLFAs had more than doubled, *e.g.* i14:0 at Igelösa and 18:2 at Petersborg. At Petersborg,
235 fertilisation with mineral N caused a further increase in microbial biomass (Table 11),
236 although a lower proportion of carbon was incorporated into microbes according to the PLFA
237 analysis (Fig. 1).

238 Analysis of microbial community structures with PCA of individual PLFAs showed that
239 differences between sites were more pronounced than differences between treatments, with
240 samples from the same sites clustered together (Fig. 3). However, some of the samples from
241 Ultuna were extremely separated. Comparisons (Suppl. Table B) showed that PC (principal
242 component) 1 was dominated by the presence of PLFA 16:0, which is ubiquitous in most
243 organisms and thus of limited use for explaining differences. PC 2 had strong positive
244 correlations with most of the mono-unsaturated PLFAs, while the eukaryotic biomarkers
245 PLFAs 18:2+18:3 were negatively correlated with some branched PLFAs (i16:0, i17:0, br18:0

246 and 10me17:0), while other branched PLFAs (i14:0, a15:0, and 10Me18) had a positive
247 correlation. The PLFA cy19:0, regarded as an indicator of stressed production of mono-
248 unsaturated PLFAs (Kaur et al. 2005), was also negatively correlated with PC 2.

249 The PCA were also interpreted in relation to sites and treatments by comparing Tables 9-
250 11. Some PLFAs appeared to be site-specific, *e.g.* at Ultuna i14:0, a15:0 and 16:1 ω 5 had low
251 values, while 10me16:0 was more common than at the other sites. Petersborg had high values
252 of 14:0, while Lanna had high values of i15:0 and 17:1, but low values of 18:2+3 and 18:1 ω 9.
253 The treatments at Ultuna differed greatly, with values for the sewage sludge treatment being
254 particularly extreme, with high ratios of the PLFAs 16:0, i17:0 and cy19:0, but low 18:1 ω 7,
255 18:1 ω 5 and 10me18:0.

256 A finding that can be regarded as more general was a positive correlation between pH and
257 the mono-unsaturated PLFAs at all sites (*e.g.* 18:1 ω 7 with $r=0.855$ for $N=60$; Suppl. Table B).
258 This was also obvious in the PCA (Fig. 4), where these PLFAs clustered together in the left
259 part of the panel. There was also a negative correlation between the mono-unsaturated PLFAs
260 and metals in the soil samples (Table 4; *cf.* Suppl. Table B).

261

262 3.5 Neutral lipid fatty acids (NLFAs)

263 NLFAs, among which 16:1 ω 5 is regarded as a biomarker for arbuscular mycorrhiza (*e.g.*
264 Bååth 2003), were only found in samples from Igelösa and Petersborg (Fig. 5). Mineral
265 fertilisation with NPK had a strong negative effect on the NLFA 16:1 ω 5 at Petersborg, which
266 declined in concentration from 3.3 to 1.3 and 0.6 nmol NLFA g DM soil⁻¹ in the fully
267 fertilised treatments, while only a small but non-significant decrease could be observed in the
268 plots with the highest sewage sludge dose. At Igelösa, no significant change was observed,
269 but in contrast a trend for an increase in the NLFA 16:1 ω 5 at the lower dose of sewage
270 sludge, as also seen at Petersborg.

271

272 4 Discussion

273 4.1 Metals in soil and crops

274 Amending soil with sewage sludge increased soil organic matter and microbial biomass at all
275 four sites investigated. In addition, lower soil pH values in sludge-treated soil, for example at
276 Ultuna (pH 4.9 after 53 years), did not cause lower microbial biomass. However, our results
277 contradict earlier reports from the 1980s and 1990s about negative effects of sewage sludge
278 fertilisation on microbial biomass and activity on soils in Britain (Brookes and McGrath

279 1984), but also in the Ultuna experiment (*e.g.* Witter et al. 1993). In the latter report, negative
280 effects were attributed to metal toxicity. In order to test this hypothesis, we can first compare
281 estimates of microbial biomass in the sludge-treated plots at Ultuna: Using the factor 5.85 mg
282 ATP g⁻¹ biomass C according to Tate and Jenkinson (1982) and 0.172 mmol PLFA g⁻¹
283 biomass C according to Joergensen and Emmerling (2006), there has been an increase from
284 231 µg biomass C g⁻¹ DM soil in 1990 (ATP data from Witter et al. 1993) to 517 µg biomass
285 C g⁻¹ DM soil in 2009. For the other treatments in the Ultuna experiment, the two methods
286 have given similar biomass values (within 10% difference).

287 Over the last 20 years, emissions of heavy metals in Europe have declined and for most
288 metals, deposition in 2005 was only around 20% of that in 1980 (Pacyna et al. 2009). A
289 similar decline has taken place in metal concentrations in sewage from Swedish wastewater
290 plants (*e.g.* Börjesson et al. 2012). In the Ultuna experiment, most of the metal enrichment in
291 sewage sludge-treated soil originated from applications made before 1974. Since then,
292 application of sludge to soil has only mainly affected the Cu content in soil. Copper from
293 water pipes has dissolved to a high extent into Uppsala's drinking water, causing Cu
294 enrichment in sewage sludge. However, it should be mentioned that since 2008 these levels
295 have also decreased radically due to improved treatment techniques (Uppsala Vatten 2012). A
296 conclusion regarding the sludge treatment at Ultuna is that lower inputs of heavy metals
297 during recent decades have allowed the microbial biomass in the soil to increase and utilise a
298 large proportion of the organic matter (*cf.* Börjesson et al. 2012). Chaperon and Sauvé (2007)
299 investigated the toxicities of Ag, Cu, Hg and Zn to the enzymes dehydrogenase and urease in
300 an agricultural soil, and calculated EC₅₀-values (concentrations giving 50% of the enzymatic
301 activity in the control) to be 268 mg (4.221 mmol) Cu kg soil⁻¹ for dehydrogenase and 491 mg
302 Cu kg soil⁻¹ for urease. For Zn, the corresponding values were 926 mg Zn kg soil⁻¹ for
303 dehydrogenase and 1178 mg Zn kg soil⁻¹ for urease. These values are lower than what could
304 be found in the sewage sludge-treated soil at Ultuna (Table 4), but other enzymes may be
305 more sensitive, such as N₂-fixation: McGrath et al. (1995) estimated that free-living
306 cyanobacteria were reduced by 50 % at metal concentrations 114 mg Zn, 33 mg Cu, 17 mg
307 Ni, 2.9 mg Cd, 80 mg Cr and 40 Pb mg kg soil⁻¹. Since these values are lower than the Ultuna
308 values, an inhibitory effect could be expected, but we must also take the effect of organic
309 matter into account, and thus the threshold value, the exact mechanism and the identification
310 of the most inhibitory metal need further investigation.

311 Crop yields have increased at all four sites through sewage sludge amendment and even
312 the much higher metal concentrations at Ultuna than at the other sites have not caused any
313 decrease in crop yields. Furthermore, metal concentrations in crops treated with sewage
314 sludge have not been elevated. This was very obvious for Cu, for which high concentrations
315 in the soil and an increase over recent years did not result in elevated levels in plants (for data
316 on Igelösa and Petersborg; see Andersson 2012).

317 A metal of special interest is Cd, since high contents in food can cause human kidney
318 damage (*e.g.* Åkesson et al. 2005). Strict regulations limiting the use of sewage sludge on
319 agricultural soils are in place in Sweden and other countries. The sludge-treated soil at Ultuna
320 has higher Cd concentrations ($0.73 \text{ mg Cd kg}^{-1}$) than are currently permissible due to
321 additions of metal-rich sewage sludge before 1975. At the other sites, concentrations are
322 below the threshold limit of $0.4 \text{ mg Cd kg}^{-1}$ soil, despite the fact that Cd has been added as a
323 soluble salt together with sludge on eight occasions at Lanna, and large amounts of sewage
324 sludge have been added during 30 years at Petersborg and Igelösa. In an earlier investigation
325 of the Ultuna site, Bergkvist et al. (2003) analysed total Cd in HNO_3 extracts in archived soil
326 samples from the sewage sludge treatment and found an increase from around $0.15 \text{ mg Cd kg}^{-1}$
327 DM soil in original samples from 1956 up to $0.84 \text{ mg Cd kg}^{-1} \text{ DM}$ in 1997. However, they
328 found no elevated concentrations below the topsoil (0.1-0.2 m). Bergkvist (2003, p. 33) found
329 that the net transport of Cd through the soil profile was very low and suggested that there was
330 crop-driven recirculation, with Cd leaching from the topsoil to the subsoil. Bergkvist et al.
331 (2003) also found large variations in Cd concentrations in grain and straw fractions of a barley
332 crop from the Ultuna site, measured in samples from five different years (1978-1997), but the
333 concentration never exceeded $0.05 \text{ mg Cd kg}^{-1} \text{ DM}$ in grain samples from sewage sludge-
334 amended plots. It can be added that the permissible limit for Cd in flour for some Swedish
335 mills is $0.08 \text{ mg Cd kg}^{-1}$, while EU legislation cites $0.2 \text{ mg Cd kg}^{-1}$, limiting the use of the
336 grain as food. All cereal samples from the sites in our study had Cd concentrations below both
337 these limits (Table 8, Fig. 2). This means that they were also below what Kirkham (2006)
338 regarded as the normal Cd concentration in plants, namely $0.1 \text{ mg Cd kg}^{-1}$. The general trend
339 for Cd in Swedish wheat grain has been a decline from 0.08 to $0.04 \text{ mg Cd kg dw}^{-1}$ since the
340 peak concentrations in the 1970s (Kirchmann et al. 2009). The Cd contents in European
341 topsoils are lowest ($<0.1 \text{ ppm}$) in Scandinavia and in Portugal (Pan et al., 2010).

342 An increase in Cd uptake by crops fertilised with CaNO_3 was observed at Igelösa. This is
343 a well-known phenomenon which has been observed in several studies in Sweden. For

344 example in a study using two cultivars and three sites, Wångstrand et al. (2007) showed that
345 Cd concentrations in wheat grain were significantly correlated with N fertilisation rate. The
346 most likely cause is a decrease in available sorption sites due to sorption of Ca, which
347 increases the activity of Cd in the soil solution and makes it available for plant uptake (Gray
348 et al. 2002). Cadmium accumulation also varies between species and varieties of the same
349 species, although it does not seem to be bound to a specific region of the chromosome (Ci et
350 al. 2012). It may also be influenced by soil type, where high pH values and high contents of
351 clay and organic matter can reduce the solubility of Cd (Giller et al. 1998; Mitchell et al.
352 1999; Gibbs et al. 2006; Eriksson 2009; Gao et al. 2010; Larsson Jönsson and Asp 2011).
353 However, in the Igelösa samples in our study, the Cd uptake in plants was positively
354 correlated with soil pH ($r=0.39$, $p=0.018$), which adds support to the hypothesis that an anion
355 (in this case NH_4^+) from the fertiliser applied replaces Cd at active soil particle sites and
356 makes Cd more plant-available, as suggested by Lorenz et al. (1994).

357 A dependence of plant uptake of Mo on soil pH was observed in samples from all sites,
358 due to increased solubility of Mo with increasing pH (Tyler and Olsson 2001).

359 A review by Smith (2009) concluded that Zn was the most crucial metal for soil
360 microbial processes due to the increase in labile forms, particularly under acidic conditions.
361 This can also be manifested as higher crop uptake of Zn (Smith 2009), as found in maize
362 silage in our study (Table 7). Villar and Garcia (2008) showed that bioleaching, *i.e.*
363 solubilisation of metal ions with the aid of *Thiobacillus* spp., needed a pH of 2-3 to initiate
364 release of Cr and Cu, while solubilisation of Ni and Zn started already at pH 6-6.5. At a site in
365 New Zealand that had received over 1000 Mg ha⁻¹ of wet sewage sludge, corresponding to
366 >150 Mg dry weight, Speir et al. (2003) observed a strong relationship between plant uptake
367 of Zn and Zn in soil solution, but not for other metals, and attributed this to lowered pH. This
368 supports our findings of increased Ni and Zn in silage maize from sewage sludge-treated plots
369 at Ultuna, where soil pH had decreased to 4.88 after more than 50 years of application.

370

371 4.2 Microbial composition (PLFAs and NLFAs)

372 The microbial community analysis showed that there was a larger difference between soil
373 types than between treatments. This has also been reported by others, *e.g.* Bünemann et al.
374 (2008) in a study with C and P additions. Among samples from the Ultuna experiment, there
375 was a large variation between treatments. This effect is also evident in the PCA plot (Fig. 3),
376 where the separation of Ultuna samples from the other samples is partly due to different

377 lengths of experimental period. However, it is most likely that these differences were caused
378 by the specific soil tillage applied at this site. The Ultuna soil is manually managed; all tillage
379 is done by hand so that soil compaction through machinery is avoided. As a result, large
380 differences in bulk density could develop between treatments due to different soil C contents
381 (Kätterer et al. 2011).

382 As reported earlier (Börjesson et al. 2012), the microbial community structure, measured
383 as PLFAs, was significantly affected by differences in soil pH in the Ultuna experiment. Soil
384 pH values in the sewage sludge-treated plots at Ultuna and Lanna have decreased over the
385 years, while acidification has been prevented through liming at Petersborg and Igelösa. At
386 Ultuna, the decrease in mono-unsaturated PLFAs was inversely proportional to the increase in
387 branched PLFAs, indicating a shift from Gram-negative to Gram-positive bacteria. High
388 ratios of cy17:0 and cy19:0, indicating stress among Gram-negative bacteria, were most
389 obvious in the sewage sludge treatment at Ultuna. The effect of pH on Gram-negative bacteria
390 is well documented (*e.g.* Aciego Petri and Brookes 2009; Fernández-Calviño et al. 2010), but
391 Gram-negative bacteria are also known to increase after nitrogen fertilisation, both through
392 mineral fertilisers and manure (Peacock et al. 2001). Petersen et al. (2003) also found
393 increased concentrations compared with the control of cy17:0 in Danish soils amended with
394 high levels of sewage sludge, but the assumed stress effect on Gram-negative bacteria
395 gradually decreased during the growing season.

396 Metal contamination in soils has previously been reported by Abaye et al. (2005), who
397 investigated agricultural soils in England fertilised with sewage sludge from 1942 to 1961.
398 They found that Gram-negative bacteria were much more common than Gram-positive.
399 Likewise, Sandaa et al. (2001) found higher amounts of bacteria belonging to the α -
400 subdivision in soils amended with contaminated or heavy metal-spiked sewage sludge in
401 Germany, and Piotrowska-Seget et al. (2005) found that metal-tolerant bacteria isolated from
402 soils were mainly Gram-negative, some of them with plasmid-associated tolerance to Zn and
403 Cd. This may also point towards the profound effect of pH caused by sewage sludge at
404 Ultuna, since Gram-negative bacteria were much less abundant there, and therefore the effect
405 of metals can be ruled out. Gram-positive bacteria are also assumed to consume older organic
406 material (*e.g.* Waldrop and Firestone 2004), which fits with our observation of a decline in
407 soil organic carbon during the last 15 years in the sewage sludge plots at Ultuna (Börjesson et
408 al. 2012). Furthermore, in a long-term experiment in Germany, where metal-amended sewage
409 sludge had been applied on five occasions between 1980 and 1989, the PLFA profile of the

410 most metal-contaminated soils was very similar to that in the Ultuna sludge samples, with a
411 decrease in 16:1 ω 5, 18:1 ω 7, 10Me18:0 and an increase in 14:0, i15:0, 16:0 and cy19:0
412 relative to the mineral fertilised treatment (Witter et al. 2000). Those authors also performed a
413 test for metal tolerance and found that the study soils had developed a tolerance for Zn. A
414 tolerance to Zn has presumably also developed at Ultuna, since Zn values have been at
415 constantly high levels for a long time.

416 Some PLFAs seem to be site-specific: At Lanna, very low ratios of both the PLFAs 18:2
417 and 18:1 ω 9 (Table 10) indicate that fungi are less abundant there compared with the other
418 three sites. Some PLFAs were obviously treatment-specific, but the effects were not
419 consistent between the different sites, thus providing further proof that the sites, and their
420 chemical and physical soil properties, were more important than the treatments for
421 determining the microbial community structure.

422 The lack of significant change in the amounts of the PLFA 18:2 indicates that there was
423 no obvious effect on fungi in the sludge-amended plots in our study, which is in accordance
424 with some previous reports (e.g. Marschner et al. 2003, Suhadolc et al. 2010). However it
425 contradicts some others in which metal inputs decreased bacterial-fungal ratios of biomarker
426 PLFAs (Khan and Scullion 2000), but negative effects on fungal gene fragments in Cu-
427 containing sewage sludge have also been reported (Macdonald et al. 2011).

428 The lack of response in NLFA content in the soil samples from sewage sludge-treated
429 plots at Petersborg and Igelösa contradicts previously reported negative effects of sewage
430 sludge on the mycorrhizal fungus *Glomus mossae* (Jacquot-Plumey et al. 2003), but their
431 result were obtained with metal-spiked sludge mixed with quartz. This further exemplifies the
432 necessity to perform studies in long-term field experiments and the significance of adsorption
433 or covalent binding of metals with organic matter (Alloway and Jackson 1991). At normal
434 doses, P added with sewage sludge, mainly as insoluble FePO₄, obviously did not affect the
435 mycorrhiza in this study, although this could have been expected (Jansa et al. 2006).

436

437 **5 Conclusions**

- 438 • At all four sites investigated, long-term sewage sludge application increased soil fertility in
439 terms of soil organic matter (total C and N), microbial biomass and crop yield.
- 440 • Long-term sewage sludge application led to a decrease in soil pH values.
- 441 • Concentrations of some metals, mainly Cu and Zn, had increased significantly with sewage
442 sludge application at all sites.

- 443 • The amounts of metals added to soil with sewage sludge were found not to be toxic for
444 microbes at any site. Earlier observations of lower microbial biomass at the Ultuna site
445 were no longer detectable.

446

447 **Acknowledgements** We are grateful to Per-Göran Andersson at the Swedish Rural Economy
448 and Agricultural Society Malmöhus for access to the sites and data on metal analyses and
449 yields at Petersborg and Igelösa. Lena Ek and Inger Juremalm carried out metal analyses at
450 the Department of Soil and Environment and Elisabet Börjesson, Department of
451 Microbiology, SLU, assisted with GC analysis of PLFAs. The work was conducted with
452 support from SLF (The Swedish Farmers' Foundation for Agricultural Research) under
453 contract H1033139, and from the 'Agricultural landscape' section within the program for
454 Environmental Monitoring and Assessment at the Swedish University of Agricultural
455 Sciences.

456

457 **References**

- 458 Abaye DA, Lawlor K, Hirsch PR, Brookes PC (2005) Changes in the microbial community of
459 an arable soil caused by long-term metal contamination. *Eur J Soil Sci* 56:93-102. doi:
460 [10.1111/j.1365-2389.2004.00648.x](https://doi.org/10.1111/j.1365-2389.2004.00648.x)
- 461 Aciego Petri JC, Brookes PC (2009) Substrate inputs and pH as factors controlling microbial
462 biomass, activity and community structure in an arable soil. *Soil Biol Biochem* 41:1396-
463 1405. doi: [10.1016/j.soilbio.2009.03.017](https://doi.org/10.1016/j.soilbio.2009.03.017)
- 464 Åkesson A, Lundh T, Vahter M, Bjellerup P, Lidfeldt J, Nerbrand C, Samsioe G, Strömberg
465 U, Skerfving S (2005) Tubular and glomerular kidney effects in Swedish women with
466 low environmental cadmium exposure. *Environ Health Perspect* 113:1627-1631. doi:
467 [10.1289/ehp.8033](https://doi.org/10.1289/ehp.8033)
- 468 Alloway BJ, Jackson AP (1991) The behaviour of heavy metals in sewage sludge-amended
469 soils. *The Science of the Total Environment* 100, 151-176. doi: [10.1016/0048-
470 9697\(91\)90377-Q](https://doi.org/10.1016/0048-9697(91)90377-Q)
- 471 Andersson A, Nilsson KO (1975) Influence on the levels of heavy metals in soil and plant
472 from sewage sludge as fertilizer (In Swedish: Effekter på tungmetallhalterna i mark och
473 växt vid tillförel av rötslam som växtnäringskälla och jordförbättringsmedel). *Rapporter
474 från Avdelningen för växtnäringslära Nr 96, Lantbrukshögskolan, Uppsala. ISBN 91-
475 7088-373-4*

- 476 Andersson P-G (2012) Slamspridning på åkermark. Fältförsök med kommunalt avloppsslam
477 från Malmö och Lund under åren 1981-2011. (In Swedish) Hushållningssällskapets
478 rapportserie 16. Swedish Rural Economy and Agricultural Societies, Malmö, Sweden.
479 ISBN 91-88668-74-6
- 480 Bååth E (2003) The use of neutral lipid fatty acids to indicate the physiological conditions of
481 soil fungi. *Microb Ecol* 45:373-383. doi: [10.1007/s00248-003-2002-y](https://doi.org/10.1007/s00248-003-2002-y)
- 482 Bergkvist P (2003) Long-term fate of sewage-sludge derived cadmium in arable soils. PhD
483 diss., Agraria 410, Swedish University of Agricultural Sciences, Uppsala, Sweden
- 484 Bergkvist P, Jarvis N, Berggren D, Carlgren K (2003) Long-term effects of sewage sludge
485 applications on soil properties, cadmium availability and distribution in arable soil. *Agric*
486 *Ecosys Environ* 97:167-179. doi: [10.1016/S0167-8809\(03\)00121-X](https://doi.org/10.1016/S0167-8809(03)00121-X)
- 487 Börjesson G, Menichetti L, Kirchmann H, Kätterer T (2012) Soil microbial community
488 structure affected by 53 years of nitrogen fertilisation and different organic amendments.
489 *Biol Fertil Soils* 48:245–257. doi: [10.1007/s00374-011-0623-8](https://doi.org/10.1007/s00374-011-0623-8)
- 490 Brookes PC, McGrath SP (1984) Effects of metal toxicity on the size of the soil microbial
491 biomass. *J Soil Sci* 35:341-346. doi: [10.1111/j.1365-2389.1984.tb00288.x](https://doi.org/10.1111/j.1365-2389.1984.tb00288.x)
- 492 Bünemann EK, Smernik RJ, Marschner P, McNeill AM (2008) Microbial synthesis of organic
493 and condensed forms of phosphorus in acid and calcareous soils. *Soil Biol*
494 *Biochem* 40:932-946. doi: [10.1016/j.soilbio.2007.11.012](https://doi.org/10.1016/j.soilbio.2007.11.012)
- 495 Chaperon S, Sauvé S (2007) Toxicity interaction of metals (Ag, Cu, Hg, Zn) to urease and
496 dehydrogenase activities in soils. *Soil Biol Biochem* 39:2329–2338. doi:
497 [10.1016/j.soilbio.2007.04.004](https://doi.org/10.1016/j.soilbio.2007.04.004)
- 498 Ci D, Jiang D, Li S, Wollenweber B, Dai T, Cao W (2012) Identification of quantitative trait
499 loci for cadmium tolerance and accumulation in wheat. *Acta Physiol Plant* 34:191-202.
500 doi: [10.1007/s11738-011-0818-5](https://doi.org/10.1007/s11738-011-0818-5)
- 501 Diacono M, Montemurro F (2010) Long-term effects of organic contaminants on soil fertility.
502 A review. *Agron Sustain Dev* 30:401-422. doi: [10.1051/agro/2009040](https://doi.org/10.1051/agro/2009040)
- 503 Environmental Objectives Portal (2012) Sweden's environmental objectives - An introduction.
504 [http://www.miljomal.se/Environmental-Objectives-Portal/Undre-meny/About-the-
505 Environmental-Objectives/15-A-Good-Built-Environment/Interim-targets/Waste/](http://www.miljomal.se/Environmental-Objectives-Portal/Undre-meny/About-the-Environmental-Objectives/15-A-Good-Built-Environment/Interim-targets/Waste/).
506 Accessed 2013-08-14

- 507 Eriksson J (2009) Strategi för att minska kadmiumbelastningen i kedjan mark-livsmedel-
508 människa (in Swedish). Report MAT21 nr 1/2009, 59 pp. ISBN 978-91-86197-26-1
509 <http://www-mat21.slu.se/publikation/pdf/Cd%20rapport2009.pdf>. Accessed 2012-05-25
- 510 Eriksson J, Mattsson L, Söderström M (2010) Current status of Swedish arable soils and
511 cereal crops. Data from the period 2001-2007 (In Swedish, with English summary).
512 Report 6349. Swedish Environmental Protection Agency, Stockholm, Sweden. URL:
513 <http://www.naturvardsverket.se/Documents/publikationer/978-91-620-6349-8.pdf>.
514 Accessed 2013-10-11
- 515 Eurostat (2012) Theme: Environment and energy – sewage sludge from urban wastewater.
516 <http://epp.eurostat.ec.europa.eu>. Publish date : 18-Apr-2012. Accessed 2012-05-25
- 517 Fernández-Calviño D, Martín AP, Arias-Estévez M, Bååth E, Díaz-Raviña M (2010)
518 Microbial community structure of vineyard soil with different pH and copper content.
519 *Appl Soil Ecol* 46:276-282. doi: [10.1016/j.apsoil.2010.08.001](https://doi.org/10.1016/j.apsoil.2010.08.001)
- 520 Gao XP, Brown KR, Racz GJ, Grant CA (2010) Concentration of cadmium in durum wheat
521 as affected by time, source and placement of nitrogen fertilization under reduced and
522 conventional-tillage management. *Plant Soil* 337:341-354. doi: [10.1007/s11104-010-
523 0531-y](https://doi.org/10.1007/s11104-010-0531-y)
- 524 Gibbs PA, Chambers BJ, Chaudri AM, McGrath SP, Carlton-Smith CH, Bacon JR, Campbell
525 CD, Aitken MN (2006) Initial results from a long-term, multi-site field study of the
526 effects on soil fertility and microbial activity of sludge cakes containing heavy metals.
527 *Soil Use Manage* 22:11-21. doi: [10.1111/j.1475-2743.2006.00003.x](https://doi.org/10.1111/j.1475-2743.2006.00003.x)
- 528 Giller KE, Witter E, McGrath SP (1998) Toxicity of heavy metals to microorganisms and
529 microbial processes in agricultural soils: A review. *Soil Biol Biochem* 30:1389-1414. doi:
530 [10.1016/S0038-0717\(97\)00270-8](https://doi.org/10.1016/S0038-0717(97)00270-8)
- 531 Gray CW, Moot DJ, McLaren RG, Reddecliffe T (2002) Effect of nitrogen fertiliser
532 applications on cadmium concentrations in durum wheat (*Triticum turgidum*) grain. *New
533 Zeal J Crop Hort* 30, 291-299. doi: [10.1080/01140671.2002.9514226](https://doi.org/10.1080/01140671.2002.9514226)
- 534 Jacquot-Plumey E, Caussanel JP, Gianinazzi S, Van Tuinen D, Gianinazzi-Pearson V (2003)
535 Heavy metals in sewage sludges contribute to their adverse effects on the arbuscular
536 mycorrhizal fungus *Glomus mosseae*. *Folia Geobot* 38, 167-176. doi:
537 [10.1007/BF02803149](https://doi.org/10.1007/BF02803149)

- 538 Jansa J, Wiemken A, Frossard E (2006) The effects of agricultural practices on arbuscular
539 mycorrhizal fungi. Geological Society, London, Special Publications 266:89-115. doi:
540 [10.1144/GSL.SP.2006.266.01.08](https://doi.org/10.1144/GSL.SP.2006.266.01.08)
- 541 Joergensen RG, Emmerling C (2006) Methods for evaluating human impact on soil
542 microorganisms based on their activity, biomass, and diversity in agricultural soils. J
543 Plant Nutr Soil Sci 169:295-309. doi: [10.1002/jpln.200521941](https://doi.org/10.1002/jpln.200521941)
- 544 Kätterer T, Bolinder MA, Andrén O, Kirchmann H, Menichetti L (2011) Roots contribute
545 more to refractory soil organic matter than above-ground crop residues, as revealed by a
546 long-term field experiment. Agric Ecosys Environ 141:184-192. doi:
547 [10.1016/j.agee.2011.02.029](https://doi.org/10.1016/j.agee.2011.02.029)
- 548 Kätterer T, Bolinder MA, Berglund K, Kirchmann H (2013) Strategies for carbon
549 sequestration in agricultural soils in northern Europe. Acta Agr Scand Section A 62:181-
550 198. doi: [10.1080/09064702.2013.779316](https://doi.org/10.1080/09064702.2013.779316)
- 551 Kärman E, Malmqvist P-A, Rydhagen B, Svensson G (2007) Evaluation of the ReVAQ
552 project (In Swedish: Utvärdering av ReVAQ-projektet). Report 2007-02, Svenskt Vatten
553 Utveckling. Available at: http://vav.griffel.net/filer/Rapport_2007-02.pdf. Accessed
554 2013-08-14
- 555 Kaur A, Chaudhary A, Kaur A, Choudhary R, Kaushik R (2005) Phospholipid fatty acid - A
556 bioindicator of environment monitoring and assessment in soil ecosystem. Curr Sci India
557 89:1103-1112
- 558 Khan M, Scullion J (2000) Effect of soil on microbial responses to metal contamination.
559 Environ Poll 110:115-125. doi: [10.1016/S0269-7491\(99\)00288-2](https://doi.org/10.1016/S0269-7491(99)00288-2)
- 560 Kirchmann H, Mattsson L, Eriksson J (2009) Trace element concentration in wheat grain:
561 results from the Swedish long-term soil fertility experiments and national monitoring
562 program. Environ Geochem Health 31:561-571. doi: [10.1007/s10653-009-9251-8](https://doi.org/10.1007/s10653-009-9251-8)
- 563 Kirkham MB (2006) Cadmium in plants on polluted soils: Effects of soil factors,
564 hyperaccumulation, and amendments. Geoderma 137, 19-32. doi:
565 [10.1016/j.geoderma.2006.08.024](https://doi.org/10.1016/j.geoderma.2006.08.024)
- 566 Larsson Jönsson EH, Asp H (2011) Influence of nitrogen supply on cadmium accumulation in
567 potato tubers. J Plant Nutr 34:345-360. doi: [10.1080/01904167.2011.536877](https://doi.org/10.1080/01904167.2011.536877)
- 568 Linderholm, K, Tillman A-M, Mattsson JE (2012) Life cycle assessment of phosphorus
569 alternatives for Swedish agriculture. Resour Conserv Recy 66: 27-39. doi:
570 [10.1016/j.resconrec.2012.04.006](https://doi.org/10.1016/j.resconrec.2012.04.006)

- 571 Lindfors E (2012) Examination of expanded uses for the sewage sludge that is produced in
572 Lotsbroverket (In Swedish: Undersökning av utökade användningsområden för
573 Lotsbroverkets slam). Civil Eng. Exam work, Department of Soil and Environment, Plant
574 nutrition and soil biology, SLU, Uppsala, Sweden. ISSN 1401-5765.
- 575 Lorenz SE, Hamon RE, McGrath SP, Holm PE, Christiansen TH (1994) Applications of
576 fertilizer cations affect cadmium and zinc concentrations in soil solutions and uptake by
577 plants. *Eur J Soil Sci* 45:159-165. doi: [10.1111/j.1365-2389.1994.tb00497.x](https://doi.org/10.1111/j.1365-2389.1994.tb00497.x)
- 578 Macdonald CA, Clark IM, Zhao F-J, Hirsch PR, Singh BK, McGrath SP (2011) Long-term
579 impacts of zinc and copper enriched sewage sludge additions on bacterial, archaeal and
580 fungal communities in arable and grassland soils. *Soil Biol Biochem* 43:932-941. doi:
581 [10.1016/j.soilbio.2011.01.004](https://doi.org/10.1016/j.soilbio.2011.01.004)
- 582 Marschner P, Kandeler E, Marschner B (2003) Structure and function of the soil microbial
583 community in a long-term fertilizer experiment. *Soil Biol Biochem* 35:453-461. doi:
584 [10.1016/S0038-0717\(02\)00297-3](https://doi.org/10.1016/S0038-0717(02)00297-3)
- 585 McGrath SP, Chaudri AM, Giller KE (1995) Long-term effects of metals in sewage sludge on
586 soils, microorganisms and plants. *J Ind Microbiol* 14:94-104. doi: [10.1007/BF01569890#](https://doi.org/10.1007/BF01569890#)
- 587 Mitchell LG, Grant CA, Rac, GJ (1999) Effect of nitrogen application on concentration of
588 cadmium and nutrient ions in soil solution and in durum wheat. *Can J Soil Sci* 80:107-
589 115. doi: [10.4141/S98-085](https://doi.org/10.4141/S98-085)
- 590 Pacyna JM, Pacyna EG, Aas W (2009) Changes of emissions and atmospheric deposition of
591 mercury, lead, and cadmium. *Atmos Environ* 43:117-127. doi:
592 [10.1016/j.atmosenv.2008.09.066](https://doi.org/10.1016/j.atmosenv.2008.09.066)
- 593 Pan JL, Plant JA, Voulvoulis N, Oates CJ, Ihlenfeld C (2010) Cadmium levels in Europe:
594 implications for human health. *Environ Geochem Health* 32:1-12. doi: [10.1007/s10653-](https://doi.org/10.1007/s10653-009-9273-2)
595 [009-9273-2](https://doi.org/10.1007/s10653-009-9273-2)
- 596 Peacock AD, Mullen MD, Ringelberg DB, Tyler DD, Hedrick DB, Gale PM, White DC
597 (2001) Soil microbial community responses to dairy manure or ammonium nitrate
598 applications. *Soil Biol Biochem* 33:1011-1019. doi: [10.1016/S0038-0717\(01\)00004-9](https://doi.org/10.1016/S0038-0717(01)00004-9)
- 599 Petersen SO, Henriksen K, Mortensen GK, Krogh PH, Brandt KK, Sørensen J, Madsen T,
600 Petersen J, Grøn C (2003) Recycling of sewage sludge and household compost to arable
601 land: fate and effects of organic contaminants, and impact on soil fertility. *Soil Till Res*
602 72:139-152. doi: [10.1016/S0167-1987\(03\)00084-9](https://doi.org/10.1016/S0167-1987(03)00084-9)

- 603 Piotrowska-Seget Z, Cycón M, Kozdrój J (2005) Metal-tolerant bacteria occurring in heavily
604 polluted soil and mine spoil. *Appl Soil Ecol* 28:237-246. doi:
605 [10.1016/j.apsoil.2004.08.001](https://doi.org/10.1016/j.apsoil.2004.08.001)
- 606 Revaq (2012) Newsletter (In Swedish: Nyhetsbrev).
607 [http://www.svenskvatten.se/Documents/Kategorier/Avlopp%20och%20milj%c3%b6/RE](http://www.svenskvatten.se/Documents/Kategorier/Avlopp%20och%20milj%c3%b6/REVAQ/REVAQ%20Nyhetsbrev%20nr%202.pdf)
608 [VAQ/REVAQ%20Nyhetsbrev%20nr%202.pdf](http://www.svenskvatten.se/Documents/Kategorier/Avlopp%20och%20milj%c3%b6/REVAQ/REVAQ%20Nyhetsbrev%20nr%202.pdf). Accessed 2012-06-04
- 609 Sandaa R-A, Torsvik V, Enger Ø (2001) Influence of long-term heavy-metal contamination on
610 microbial communities in soil. *Soil Biol Biochem* 33:287-295. doi: [10.1016/S0038-](https://doi.org/10.1016/S0038-0717(00)00139-5)
611 [0717\(00\)00139-5](https://doi.org/10.1016/S0038-0717(00)00139-5)
- 612 Singh RP, Agrawal M (2008) Potential benefits and risks of land application of sewage
613 sludge. *Waste Manage* 28: 347-358 doi: [10.1016/j.wasman.2006.12.010](https://doi.org/10.1016/j.wasman.2006.12.010)
- 614 Smith SR (2009) A critical review of the bioavailability and impacts of heavy metals in
615 municipal solid waste composts compared to sewage sludge. *Environ Int* 35:142-156. doi:
616 [10.1016/j.envint.2008.06.009](https://doi.org/10.1016/j.envint.2008.06.009)
- 617 Speir TW, Van Schaik AP, Percival HJ, Close ME, Pang LP (2003) Heavy metals in soil,
618 plants and groundwater following high-rate sewage sludge application to land. *Water Air*
619 *Soil Poll* 150:319-358. doi: [10.1023/A:1026101419961](https://doi.org/10.1023/A:1026101419961)
- 620 Statistics Sweden, 2012. Discharges to water and sewage sludge production in 2010
621 Municipal wastewater treatment plants, pulp and paper industry and other industry (In
622 Swedish: Utsläpp till vatten och slamproduktion 2010. Kommunala reningsverk,
623 skogsindustri samt övrig industri).
624 [http://www.scb.se/Statistik/MI/MI0106/2010A01/MI0106_2010A01_SM_MI22SM1201.](http://www.scb.se/Statistik/MI/MI0106/2010A01/MI0106_2010A01_SM_MI22SM1201.pdf)
625 [pdf](http://www.scb.se/Statistik/MI/MI0106/2010A01/MI0106_2010A01_SM_MI22SM1201.pdf). Accessed 2013-08-14
- 626 Suhadolc M, Schroll R, Hagn A, Dörfler U, Schloter M, Lobnik F (2010) Single application
627 of sewage sludge - Impact on the quality of an alluvial agricultural soil. *Chemosphere*
628 81:1536-1543. doi: [10.1016/j.chemosphere.2010.08.024](https://doi.org/10.1016/j.chemosphere.2010.08.024)
- 629 Swedling E-O (2011) Personal communication, e-mail 2011-01-12 (Ernst-
630 Olof.Swedling@uppsalavatten.se)
- 631 Tate KR, Jenkinson DS (1982) Adenosine triphosphate measurement in soil: An improved
632 method. *Soil Biol Biochem* 14:331-335. doi: [10.1016/0038-0717\(82\)90002-5](https://doi.org/10.1016/0038-0717(82)90002-5)
- 633 Tyler G, Olsson T (2001) Plant uptake of major and minor mineral elements as influenced by
634 soil acidity and liming. *Plant Soil* 230:307-321. doi: [10.1046/j.1365-2389.2001.t01-1-](https://doi.org/10.1046/j.1365-2389.2001.t01-1-00360.x)
635 [00360.x](https://doi.org/10.1046/j.1365-2389.2001.t01-1-00360.x)

- 636 Uppsala Vatten (2012) Environmental report for Kungsängen wastewater plant (In Swedish).
637 http://www.uppsalavatten.se/Documents/Gemensam/Milj%C3%B6rapporter/2012_Miljorapport_Kungsangsverket.pdf. Accessed 2013-08-14
638
- 639 Villar LD, Garcia O (2002) Solubilization profiles of metal ions from bioleaching of sewage
640 sludge as a function of pH. *Biotechnol Lett* 24:611-614. doi: [10.1023/A:1015010417315](https://doi.org/10.1023/A:1015010417315)
- 641 Waldrop MP, Firestone MK (2004) Altered utilization patterns of young and old soil C by
642 microorganisms caused by temperature shifts and N addition. *Biogeochemistry* 67:235-
643 248. doi: [10.1023/B:BIOG.0000015321.51462.41](https://doi.org/10.1023/B:BIOG.0000015321.51462.41)
- 644 Wångstrand H, Eriksson J, Öborn I (2007) Cadmium concentration in winter wheat as
645 affected by nitrogen fertilization. *Eur J Agron* 26:209–214. doi:
646 [10.1016/j.eja.2006.09.010](https://doi.org/10.1016/j.eja.2006.09.010)
- 647 Witter E, Mårtensson AM, Garcia FV (1993) Size of the soil microbial biomass in a long-term
648 field experiment as affected by different N-fertilizers and organic manures. *Soil Biol
649 Biochem* 25:659-669. doi: [10.1016/0038-0717\(93\)90105-K](https://doi.org/10.1016/0038-0717(93)90105-K)
- 650 Witter E, Gong P, Bååth E, Marstorp H (2000) A study of the structure and metal tolerance of
651 the soil microbial community six years after cessation of sewage sludge applications.
652 *Environ Toxicol Chem* 19:1983-1991. doi: [10.1897/1551-
653 5028\(2000\)019<1983:ASOTSA>2.3.CO;2](https://doi.org/10.1897/1551-5028(2000)019<1983:ASOTSA>2.3.CO;2)
654
655

656 **Figure legends**

657

658 **Fig. 1** Correlation between Total C (% of dry weight soil) and Total PLFAs, given as mean
 659 values for topsoil samples from the four sites. Numbers correspond to the following
 660 treatments: **Petersborg**: 1 = A0 (no sludge, no NPK), 2 = B0 (1 ton sludge, no NPK), 3 = C0
 661 (4 ton sludge, no NPK), 4 = A2 (no sludge, normal NPK), 5 = C2 (4 ton sludge, normal
 662 NPK); **Igelösa**: 6 = A0 (no sludge, no NPK), 7 = B0 (1 ton sludge, no NPK), 8 = C0 (4 ton
 663 sludge, no NPK); **Lanna**: 9 = Cropped, unfertilised, 10 = CaNO₃, 11 = Sewage sludge, 12 =
 664 Sewage sludge + metals; **Ultuna**: 13 = Cropped, unfertilised, 14 = CaNO₃, 15 = sewage
 665 sludge.

666 The fitted line for correlation has the equation: Total PLFAs = 33.01 * Total C (%) – 0.900;
 667 with $r=0.958$ and $p<0.0001$ ($n=15$). Error bars show standard deviation for each treatment
 668 ($n=4$)

669

670 **Fig. 2** Cadmium in soil samples and in wheat grain from Igelösa 2011. Bars = standard
 671 deviation, $n=4$. A= No sewage sludge; B= 4 ton sludge every 4th year; and C=12 ton sludge
 672 every 4th year. 0=No NPK; 1=half of normal NPK; and 2=normal NPK. Different letters
 673 above error bars indicate significant difference (Tukey-Kramer HSD, $\alpha=0.05$)

674

675 **Fig. 3** Principal components of 25 PLFAs in all four experiments (N=60). Data points are
 676 mean values for each treatment, with standard error for $n=4$

677


678 **Fig. 4** Principal component analysis based on correlation matrix of individual PLFAs (mol-
 679 %) and certain variables in topsoil samples from all four sites ($n=15$)

680

681 **Fig. 5** Effects of sewage sludge and mineral N on the mycorrhizal biomarker NLFA 16:1 ω 5
 682 in samples from Petersborg and Igelösa. A = No sewage sludge; B = 4 ton sludge every 4th
 683 year; and C = 12 ton sludge every 4th year. 0 = No NPK; and 2 = normal NPK. Error bars are
 684 standard deviation for $n=4$. Different letters above error bars indicate significant difference
 685 (Tukey-Kramer HSD, $\alpha=0.05$)


686

687



688
689
690
691
692
693

Fig. 1.




694
695

696

697 Fig. 2.

698

699




700
701

702

703 Fig. 3.


704



705
706

707 Fig. 4.

708



709
710

711

712 Fig. 5.

713

Table 1

Characteristics of the four study sites

| Site | Coordinates | Start (Year) | Soil type/Clay content | Amount of sludge applied |
|------------|------------------|-----------------|------------------------|---|
| Ultuna | 59°49'N, 17°39'E | 1956 | Clay loam (36.5% clay) | 4 Mg C ha ⁻¹ every 2nd yr |
| Lanna | 58°21'N, 13°06'E | 1997 | Silty clay (42% clay) | 8 Mg dry matter ha ⁻¹ every 2nd yr |
| Petersborg | 55°32'N, 13°00'E | 1981 | Sandy loam (14% clay) | 4 or 12 Mg dry matter ha ⁻¹ every 4 th yr |
| Igelösa | 55°45'N, 13°18'E | 1981 | Loam (26% clay) | 4 or 12 Mg dry matter ha ⁻¹ every 4 th yr |

Table 2

Properties of topsoil at the four study sites. Ult=Ultuna (measured 2009),
Lan=Lanna (2010), Pet=Petersborg (2011), Ige=Igelösa (2011).

Original = Start years: Ultuna 1956, Lanna 1997, Petersborg and Igelösa 1981

| Treatment | Total C (%) | | | | pH | | | |
|--------------------------|-------------|------|------|--------------------|------|------|------------------|------------------|
| | Ult | Lan | Pet | Ige | Ult | Lan | Pet ^b | Ige ^b |
| <i>Original at start</i> | 1.50 | 1.98 | 1.2 | 1.9 | 6.5 | 6.65 | 6.8 | 7.0 |
| Unfert. cropped | 1.12 | 1.87 | 0.94 | 1.88 | 6.23 | 6.45 | 6.80 | 7.11 |
| Normal fertilised | 1.37 | 1.92 | 1.14 | (2.0) ^a | 6.68 | 6.55 | 6.62 | n.m. |
| Sewage sludge | 2.66 | 2.23 | 1.30 | 2.21 | 4.88 | 6.08 | 6.51 | 7.05 |

^a = measured 2006

^b It should be noted that all experiments at Petersborg and Igelösa were limed in 1998

n.m. = not measured

Table 3

Crop yields in the long-term field experiments, given as mean values (\pm standard deviation) relative to the fully NPK fertilised treatment as a reference at each site, $n=4$.

For Petersborg and Igelösa, treatments were: A = No sewage sludge, B = 4 Mg sludge every 4th year, C = 12 Mg sludge every 4th year; 0 = No NPK, 1 = half of normal NPK, 2 = normal NPK. Treatments with yields in italics are set to 100

| Ultuna 1956-2009 | | Lanna 1997-2010 | | Petersborg 1982-2011 | | Igelösa 1982-2011 | |
|--------------------------|--------------|--------------------------|--------------|-------------------------|--------------|----------------------|--|
| B. Cropped, unfertilised | 56.5 (30.7) | I. Cropped, unfertilised | 36.6 (7.7) | A0. | 49.7 (17.5) | 56.1 (19.5) | |
| C. Calcium nitrate | <i>100</i> | B. Calcium nitrate | <i>100.0</i> | A1. | 85.6 (8.7) | 80.4 (10.6) | |
| O. Sewage sludge | 139.3 (67.8) | F. Sewage sludge | 105.7 (25.6) | A2. | <i>100.0</i> | <i>100.0</i> | |
| | | G. Sewage sludge+metals | 102.9 (26.3) | B0. | 58.0 (23.6) | 62.0 (20.4) | |
| | | | | B1. | 90.1 (14.5) | 87.2 (10.3) | |
| | | | | B2. | 103.7 (11.0) | 104.2 (6.0) | |
| | | | | C0. | 64.6 (20.6) | 70.8 (18.5) | |
| | | | | C1. | 91.3 (12.1) | 90.9 (10.7) | |
| | | | | C2. | 106.7 (6.8) | 104.7 (11.6) | |

Table 4

Mean metal concentrations (mg kg⁻¹ DM) in soil samples from the Ultuna experiment (\pm standard deviation) on earlier analysis occasions and in those taken in 2009. Different letters within rows indicate significant difference (Tukey-Kramer HSD, $\alpha=0.05$)

| | Sewage sludge | | | Calcium nitrate 2009 | Unfertilised 2009 |
|--------------------------|---------------|-------------|--------------------------|--------------------------|--------------------------|
| | 1968* | 1974* | 2009 | | |
| Al (g kg ⁻¹) | n.m. | n.m. | 20.7 (0.6) ^a | 20.2 (0.2) ^a | 20.0 (0.39) ^a |
| Cd | 0.74 | 0.72 (0.04) | 0.73 (0.02) ^a | 0.24 (0.02) ^b | 0.23 (0.01) ^b |
| Co | 11.6 | 11.6 (0.3) | <i>n.m.</i> | <i>n.m.</i> | <i>n.m.</i> |
| Cu | 69 | 84 (4) | 196 (4.6) ^a | 27.8 (1.95) ^b | 26.7 (0.57) ^b |
| Fe (g kg ⁻¹) | n.m. | n.m. | 34.7 (0.7) ^b | 27.6 (0.4) ^a | 27.4 (0.3) ^a |
| Mn | 413 | 420 (7) | 478 (16) ^a | 464 (6) ^a | 457 (25) ^a |
| Ni | 40.8 | 34.5 (0.4) | 27.2 (0.92) ^a | 22.9 (0.77) ^b | 22.7 (0.79) ^b |
| Pb | 38.5 | 40.4 (0.7) | 41.0 (3.06) ^a | 21.6 (0.70) ^b | 21.3 (0.67) ^b |
| Zn | 272 | 268 (9) | 271 (9.18) ^a | 87.6 (4.22) ^b | 85.9 (3.00) ^b |

* Data from Andersson and Nilsson (1975); extraction with 2M HNO₃ at 100°C (same as for samples from 2009)

n.m. not measured

Table 5

Mean metal concentrations (mg kg^{-1} DM) in soil samples from Lanna 2010 (\pm standard deviation). Different letters within rows indicate significant difference (Tukey-Kramer HSD, $\alpha=0.05$)

| | I. Cropped, unfertilised | B. Calcium nitrate | F. Sewage sludge | G. Sewage sludge + metals |
|---------------------------|-----------------------------|----------------------------|----------------------------|------------------------------|
| Al (g kg^{-1}) | 21.0 (1.8) ^a | 21.8 (2.1) ^a | 21.8 (1.3) ^a | 21.7 (0.7) ^a |
| Cd | 0.121 (0.009) ^b | 0.120 (0.005) ^b | 0.140 (0.015) ^b | 0.287 (0.017) ^a |
| Cu | 9.09 (0.88) ^b | 8.90 (0.79) ^b | 20.8 (1.07) ^a | 25.0 (1.32) ^a |
| Fe (g kg^{-1}) | 25.4 (2.6) ^a | 26.4 (2.4) ^a | 27.7 (1.8) ^a | 26.4 (1.0) ^a |
| Mn | 518 (175) ^a | 483 (159) ^a | 624 (215) ^a | 461 (120) ^a |
| Ni | 10.2 (0.95) ^b | 10.2 (0.44) ^b | 10.4 (0.63) ^b | 17.9 (1.09) ^a |
| Pb | 13.7 (0.81) ^a | 14.1 (0.65) ^a | 14.4 (0.86) ^a | 14.4 (0.45) ^a |
| Zn | 63.2 (6.1) ^b | 65.6 (6.4) ^b | 83.1 (5.6) ^a | 79.3 (4.4) ^a |

Table 6

Mean metal concentrations (mg kg^{-1} DM) in soil samples from Igelösa and Petersborg 2011 (\pm standard deviation for $n = 4$). A = No sewage sludge, B = 4 Mg sludge every 4th year, C = 12 Mg sludge every 4th year; 0 = No NPK, 1 = half of normal NPK, 2 = normal NPK. For each site, different letters within columns indicate significant difference (Tukey-Kramer HSD, $\alpha=0.05$)

| | Cu | Zn | Cd | Pb | Ni |
|-------------------|----------------------------|--------------------------|---------------------------|---------------------------|--------------------------|
| <i>Igelösa</i> | | | | | |
| A0 | 15.3 (6.6) ^a | 47.5 (1.9) ^a | 0.30 (0.03) ^a | 16.3 (0.50) ^a | 12.0 (0.82) ^a |
| B0 | 22.3 (9.2) ^a | 53.8 (1.0) ^b | 0.31 (0.02) ^{ab} | 16.8 (0.50) ^a | 12.0 (0) ^a |
| C0 | 25.8 (1.7) ^a | 58.0 (2.9) ^b | 0.34 (0.019) ^b | 17.3 (0.50) ^a | 12.3 (0.50) ^a |
| <i>Petersborg</i> | | | | | |
| A0 | 9.4 (0.2) ^c | 38.0 (2.9) ^a | 0.24 (0.01) ^a | 13.5 (0.58) ^{ab} | 7.9 (0.39) ^a |
| A2 | 15.3 (12.5) ^{abc} | 38.3 (3.3) ^a | 0.24 (0) ^a | 12.8 (0.50) ^a | 7.7 (0.66) ^a |
| B0 | 14.8 (1.5) ^{abc} | 42.3 (2.8) ^{ab} | 0.27 (0.01) ^a | 14.0 (1.41) ^{ab} | 8.3 (0.92) ^a |
| C0 | 21.0 (1.8) ^a | 45.0 (1.6) ^b | 0.26 (0.01) ^a | 14.5 (0.58) ^b | 8.2 (0.47) ^a |
| C2 | 20.3 (2.6) ^{ab} | 45.3 (2.2) ^b | 0.27 (0.01) ^a | 14.0 (0.82) ^{ab} | 8.3 (0.36) ^a |

Table 7

Mean metal concentrations (mg kg^{-1} DM) in maize silage from the Ultuna experiment 2009 (\pm standard deviation, $n = 4$). Different letters within rows indicate significant difference between treatments (Tukey-Kramer HSD, $\alpha=0.05$). Data in italics from analysis of fodder rape, harvested as green mass at Ultuna 1974 (Andersson and Nilsson 1975)

| | B. Cropped, unfertilised | C. Calcium nitrate | O. Sewage sludge | <i>in fodder rape 1974</i> | |
|----|-----------------------------|----------------------------|----------------------------|----------------------------|----------------|
| Cd | 0.109 (0.017) ^a | 0.059 (0.014) ^b | 0.130 (0.026) ^a | <i>0.303</i> | <i>(0.020)</i> |
| Co | 0.032 (0.010) ^a | 0.019 (0.009) ^a | 0.017 (0.003) ^a | <i>0.094</i> | <i>(0.006)</i> |
| Cr | 0.371 (0.414) ^a | 0.137 (0.083) ^a | 0.146 (0.060) ^a | <i>0.62</i> | <i>(0.18)</i> |
| Cu | 3.56 (0.18) ^a | 3.52 (0.69) ^a | 4.11 (0.36) ^a | <i>5.8</i> | <i>(0.2)</i> |
| Mo | 0.263 (0.083) ^a | 0.392 (0.013) ^a | 0.210 (0.012) ^a | <i>n.m.</i> | |
| Ni | 0.358 (0.094) ^b | 0.237 (0.117) ^b | 0.606 (0.155) ^a | <i>4.43</i> | <i>(1.16)</i> |
| Pb | 0.441 (0.153) ^a | 0.366 (0.134) ^a | 0.451 (0.314) ^a | <i>1.95</i> | <i>(0.50)</i> |
| Se | 0.157 (0.065) ^a | 0.130 (0.040) ^a | 0.139 (0.037) ^a | <i>n.m.</i> | |
| Zn | 13.2 (4.83) ^b | 9.70 (0.78) ^b | 91.0 (18.8) ^a | <i>98.3</i> | <i>(19.4)</i> |

n.m. not measured

Table 8

Mean metal concentrations (mg kg^{-1} DM) in wheat grain from Lanna 2010 (\pm standard deviation). Different letters within rows indicate significant difference (Tukey-Kramer HSD, $\alpha=0.05$)

| | I. Cropped, Unfertilised | B. Calcium nitrate | F. Sewage sludge | G. Sewage sludge + extra metals |
|----|------------------------------|------------------------------|------------------------------|------------------------------------|
| Cd | 0.026 (0.003) ^a | 0.032 (0.004) ^a | 0.026 (0.004) ^a | 0.059 (0.008) ^a |
| Co | 0.0055 (0.0006) ^a | 0.0023 (0.0005) ^a | 0.0035 (0.0006) ^a | 0.0033 (0.0010) ^a |
| Cr | n.d. | n.d. | n.d. | n.d. |
| Cu | 3.59 (0.16) ^a | 3.91 (0.11) ^a | 3.97 (0.12) ^a | 3.93 (0.13) ^a |
| Mo | 0.57 (0.17) ^b | 1.07 (0.38) ^a | 0.30 (0.03) ^b | 0.33 (0.02) ^b |
| Ni | 0.21 (0.06) ^b | 0.12 (0.03) ^b | 0.24 (0.05) ^b | 0.63 (0.11) ^a |
| Pb | 0.0055 (0.0013) ^a | 0.0090 (0.0028) ^a | 0.0063 (0.0010) ^a | 0.0075 (0.0072) ^a |
| Se | 0.032 (0.0319) ^a | 0.072 (0.032) ^a | 0.024 (0.024) ^a | 0.029 (0.039) ^a |
| Zn | 26.8 (1.1) ^a | 21.5 (1.7) ^a | 32.6 (2.7) ^a | 31.6 (2.3) ^a |

n.d. = below detection limit

Table 9

PLFA levels (% of total moles) in topsoil samples from Ultuna 2009, given as mean values for $n = 4$ plots of each treatment. Different letters within rows (between treatments) indicate significant difference (Tukey-Kramer HSD, $\alpha=0.05$)

| | Cropped, unfertilised | Calcium nitrate | Sewage sludge |
|---|--------------------------|--------------------|--------------------|
| i14:0 | 0.29 ^a | 0.20 ^a | 0.11 ^a |
| 14:0 | 1.61 ^a | 1.34 ^a | 1.50 ^a |
| i15:0 | 8.16 ^a | 7.10 ^a | 9.70 ^b |
| a15:0 | 5.23 ^{ab} | 5.61 ^b | 4.30 ^a |
| 15:0 | 0.52 ^a | 0.37 ^a | 0.94 ^a |
| i16:0 | 2.60 ^a | 3.13 ^{ab} | 3.86 ^b |
| 16:1 ω 9 | 1.06 ^a | 1.61 ^a | 0.69 ^a |
| 16:1 ω 7 | 8.67 ^b | 9.31 ^b | 5.02 ^a |
| 16:1 ω 5 | 3.01 ^b | 3.12 ^b | 1.32 ^a |
| 16:0 | 12.88 ^a | 11.62 ^a | 17.43 ^b |
| 17:1 | 4.60 ^b | 4.02 ^{ab} | 3.19 ^a |
| 10me16 | 5.47 ^a | 6.10 ^a | 5.21 ^a |
| i17:0 | 1.88 ^a | 2.14 ^a | 4.77 ^b |
| a17:0 | 3.88 ^a | 3.67 ^a | 2.71 ^a |
| cy17:0 | 2.96 ^a | 3.37 ^a | 5.53 ^b |
| 17:0 | 0.27 ^a | 0.16 ^a | 0.80 ^b |
| br18:0 | 0.98 ^a | 2.27 ^{ab} | 4.19 ^c |
| 10Me17:0 | 1.83 ^a | 1.61 ^a | 2.67 ^a |
| 18:2+18:3 | 9.54 ^a | 6.00 ^a | 5.16 ^a |
| 18:1 ω 9 | 6.62 ^{ab} | 7.89 ^b | 5.36 ^a |
| 18:1 ω 7 | 6.76 ^b | 7.24 ^b | 3.36 ^a |
| 18:1 ω 5 | 1.29 ^a | 1.53 ^a | 0.65 ^a |
| 18:0 | 3.09 ^a | 2.87 ^a | 4.02 ^a |
| 10me18:0 | 3.57 ^b | 4.40 ^b | 1.76 ^a |
| cy19:0 | 3.25 ^a | 3.36 ^a | 5.79 ^b |
| Total PLFAs (nmol g soil ⁻¹) | 46.3 ^a | 41.5 ^a | 89.2 ^b |

Table 10

PLFA levels (% of total moles) in topsoil samples from Lanna 2010, given as mean values for $n = 4$ plots of each treatment. Different letters within rows (between treatments) indicate significant difference (Tukey-Kramer HSD, $\alpha=0.05$)

| | Cropped, unfertilised | Calcium nitrate | Sewage sludge | Sewage sludge + metals |
|---|--------------------------|--------------------|--------------------|---------------------------|
| i14:0 | 0.97 ^{ab} | 1.06 ^b | 0.89 ^{ab} | 0.84 ^a |
| 14:0 | 1.54 ^a | 1.85 ^a | 1.72 ^a | 1.46 ^a |
| i15:0 | 9.64 ^{ab} | 9.41 ^a | 9.97 ^{bc} | 10.36 ^c |
| a15:0 | 8.04 ^a | 8.17 ^a | 8.25 ^a | 8.22 ^a |
| 15:0 | 0.95 ^{ab} | 1.01 ^b | 0.88 ^{ab} | 0.82 ^a |
| i16:0 | 3.18 ^a | 3.17 ^a | 3.08 ^a | 2.98 ^a |
| 16:1 ω 9 | 1.73 ^a | 1.78 ^a | 1.50 ^a | 1.62 ^a |
| 16:1 ω 7 | 7.76 ^a | 7.64 ^a | 7.84 ^a | 8.33 ^a |
| 16:1 ω 5 | 4.25 ^{ab} | 4.44 ^b | 3.55 ^a | 3.87 ^{ab} |
| 16:0 | 11.79 ^a | 11.68 ^a | 11.80 ^a | 11.67 ^a |
| 17:1 | 4.88 ^a | 4.63 ^a | 5.33 ^a | 5.43 ^a |
| 10me16:0 | 4.68 ^a | 4.19 ^a | 4.18 ^a | 4.28 ^a |
| i17:0 | 2.48 ^a | 2.53 ^a | 2.66 ^a | 2.54 ^a |
| a17:0 | 2.54 ^a | 2.63 ^a | 2.71 ^a | 2.57 ^a |
| cy17 | 4.75 ^a | 4.77 ^a | 4.51 ^a | 4.66 ^a |
| 17:0 | 0.42 ^a | 0.15 ^a | 0.55 ^a | 0.54 ^a |
| br18:0 | 1.50 ^a | 1.50 ^a | 1.48 ^a | 1.33 ^a |
| 10me17:0 | 1.89 ^a | 1.80 ^a | 1.92 ^a | 1.80 ^a |
| 18:2 | 1.72 ^a | 1.63 ^a | 1.22 ^a | 1.40 ^a |
| 18:1 ω 9 | 5.78 ^a | 5.96 ^a | 6.31 ^a | 6.05 ^a |
| 18:1 ω 7 | 7.63 ^a | 7.80 ^a | 7.73 ^a | 7.90 ^a |
| 18:1 ω 5 | 1.32 ^a | 1.34 ^a | 1.41 ^a | 1.38 ^a |
| 18:0 | 3.30 ^a | 3.39 ^a | 3.14 ^a | 3.07 ^a |
| 10me18:0 | 3.71 ^{ab} | 3.85 ^b | 3.43 ^{ab} | 3.11 ^a |
| cy19:0 | 3.53 ^a | 3.63 ^a | 3.95 ^a | 3.77 ^a |
| Total PLFAs (nmol g soil ⁻¹) | 49.2 ^{ab} | 46.8 ^{ab} | 72.7 ^{de} | 74.8 ^e |

Table 11

PLFA levels (% of total moles) in topsoil samples from Petersborg and Igelösa 2011, given as mean values for $n = 4$ plots of each treatment. A = No sewage sludge, B = 4 Mg sludge every 4th year, C = 12 Mg sludge every 4th year; 0 = No NPK, 1 = half of normal NPK, 2 = normal NPK. Different letters within rows (between treatments) indicate significant difference (Tukey-Kramer HSD, $\alpha=0.05$)

| PLFA | Petersborg | | | | | Igelösa | | |
|---|----------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|
| | A0 | A2 | B0 | C0 | C2 | A0 | B0 | C0 |
| i14:0 | 1.05 ^b | 1.20 ^{bc} | 1.19 ^{bc} | 1.27 ^c | 1.21 ^{bc} | 0.70 ^a | 1.28 ^c | 1.25 ^c |
| 14:0 | 2.21 ^a | 2.17 ^a | 2.21 ^a | 1.91 ^a | 2.21 ^a | 2.00 ^a | 1.68 ^a | 1.51 ^a |
| i15:0 | 8.11 ^{bc} | 7.66 ^a | 8.09 ^{abc} | 8.25 ^c | 7.81 ^{ab} | 7.82 ^{abc} | 7.84 ^{abc} | 8.01 ^{abc} |
| a15:0 | 7.25 ^{ab} | 6.93 ^a | 7.53 ^{abc} | 7.84 ^{bc} | 7.67 ^{abc} | 7.87 ^{bc} | 7.70 ^{abc} | 8.25 ^c |
| 15:0 | 0.77 ^a | 0.78 ^{ab} | 0.81 ^{abc} | 0.86 ^c | 0.85 ^{bc} | 0.77 ^a | 0.77 ^a | 0.75 ^a |
| i16:0 | 2.93 ^{ab} | 3.04 ^{bc} | 2.88 ^a | 2.93 ^{ab} | 3.04 ^{bc} | 3.07 ^c | 2.96 ^{abc} | 3.02 ^{bc} |
| 16:1 ω 9 | 1.73 ^c | 1.49 ^{ab} | 1.54 ^{abc} | 1.43 ^{ab} | 1.36 ^a | 1.56 ^{bc} | 1.40 ^{ab} | 1.41 ^{ab} |
| 16:1 ω 7 | 7.36 ^a | 7.47 ^{ab} | 7.79 ^{abc} | 7.53 ^{abc} | 7.61 ^{abc} | 8.15 ^{abc} | 8.33 ^c | 8.31 ^{bc} |
| 16:1 ω 5 | 4.14 ^{bc} | 3.96 ^{ab} | 4.03 ^{abc} | 4.01 ^{abc} | 3.69 ^a | 4.30 ^{bc} | 4.37 ^c | 4.26 ^{bc} |
| 16:0 | 11.96 ^d | 11.79 ^{cd} | 11.65 ^{cd} | 11.68 ^{cd} | 11.59 ^{cd} | 11.24 ^{bc} | 10.91 ^{ab} | 10.66 ^a |
| 17:1 | 3.89 ^a | 3.81 ^a | 3.88 ^a | 3.79 ^a | 3.59 ^a | 3.76 ^a | 3.75 ^a | 3.67 ^a |
| 10me16:0 | 5.19 ^a | 4.38 ^a | 4.74 ^a | 4.72 ^a | 4.96 ^a | 5.15 ^a | 4.73 ^a | 4.82 ^a |
| i17:0 | 2.18 ^c | 1.96 ^{ab} | 2.12 ^{bc} | 2.06 ^{abc} | 1.94 ^{ab} | 1.94 ^{ab} | 1.87 ^a | 1.95 ^{ab} |
| a17:0 | 2.72 ^a | 2.85 ^{ab} | 2.86 ^{ab} | 2.94 ^{ab} | 3.15 ^b | 2.80 ^{ab} | 3.02 ^{ab} | 3.15 ^b |
| cy17:0 | 3.71 ^b | 3.56 ^{ab} | 3.54 ^{ab} | 3.52 ^{ab} | 3.56 ^{ab} | 3.43 ^{ab} | 3.38 ^a | 3.41 ^{ab} |
| 17:0 | 0.55 ^{abcd} | 0.56 ^{bcd} | 0.60 ^d | 0.58 ^{cd} | 0.60 ^d | 0.52 ^{abc} | 0.50 ^{ab} | 0.48 ^a |
| br18:0 | 1.96 ^a | 2.05 ^a | 1.80 ^a | 1.77 ^a | 1.88 ^a | 2.40 ^b | 2.44 ^b | 2.43 ^b |
| 10me17:0 | 1.57 ^a | 1.80 ^b | 1.76 ^{ab} | 1.66 ^{ab} | 1.77 ^{ab} | 1.59 ^a | 1.65 ^{ab} | 1.65 ^{ab} |
| 18:2+18:3 | 2.61 ^a | 4.56 ^{ab} | 3.51 ^{ab} | 4.14 ^{ab} | 3.84 ^{ab} | 4.48 ^{ab} | 5.32 ^b | 5.25 ^b |
| 18:1 ω 9 | 6.47 ^a | 7.19 ^{bc} | 6.58 ^{ab} | 6.74 ^{abc} | 7.37 ^c | 6.71 ^{ab} | 6.96 ^{abc} | 7.00 ^{abc} |
| 18:1 ω 7 | 7.95 ^a | 8.09 ^a | 8.33 ^{ab} | 7.90 ^a | 7.90 ^a | 8.98 ^c | 9.07 ^c | 8.84 ^{bc} |
| 18:1 ω 5 | 1.48 ^{cd} | 1.55 ^d | 1.42 ^{cd} | 1.36 ^{bc} | 1.47 ^{cd} | 1.27 ^{ab} | 1.19 ^a | 1.16 ^a |
| 18:0 | 4.96 ^d | 4.03 ^{bc} | 4.33 ^c | 4.01 ^{bc} | 3.78 ^b | 3.25 ^a | 2.95 ^a | 2.89 ^a |
| 10me18:0 | 3.48 ^b | 3.74 ^b | 3.34 ^b | 3.66 ^b | 3.83 ^b | 2.72 ^a | 2.79 ^a | 2.69 ^a |
| cy19:0 | 3.80 ^b | 3.40 ^a | 3.49 ^{ab} | 3.48 ^{ab} | 3.37 ^a | 3.56 ^{ab} | 3.21 ^a | 3.25 ^a |
| Total PLFAs (nmol g soil ⁻¹) | 28.2 ^a | 35.7 ^{ab} | 32.2 ^a | 36.9 ^{ab} | 42.6 ^b | 59.3 ^c | 70.5 ^d | 78.6 ^d |