



**Murdoch**  
UNIVERSITY

**MURDOCH RESEARCH REPOSITORY**

**Authors Version**

**Smyth, W.F. (1991) Sum graphs of small sum number. Colloquia mathematica Societatis János Bolyai, 60 . pp. 669-678.**

<http://researchrepository.murdoch.edu.au/27401/>

Copyright: © 1991 North-Holland Pub. Co.  
It is posted here for your personal use. No further distribution is permitted.

# SUM GRAPHS OF SMALL SUM NUMBER

W. F. Smyth

Department of Computer Science & Systems  
McMaster University

School of Computing Science  
Curtin University of Technology

## ABSTRACT

Given an integer  $r > 0$ , let  $G_r = (V, E)$  denote a graph consisting of a simple finite undirected connected nontrivial graph  $G$  together with  $r$  isolated vertices  $\overline{K}_r$ . Let  $L : V \rightarrow \mathbb{Z}^+$  denote a labelling of the vertices of  $G_r$  with distinct positive integers. Then  $G_r$  is said to be a *sum graph* if there exists a labelling  $L$  such that for every distinct vertex pair  $u$  and  $v$  of  $V$ ,  $(u, v) \in E$  if and only if there exists a vertex  $w \in V$  whose label  $L(w) = L(u) + L(v)$ . For a given subgraph  $G$ , the *sum number*  $\sigma = \sigma(G)$  is defined to be the least number  $r$  for which  $G_r$  is a sum graph; in particular, if  $G_1 = G \cup \overline{K}_1$  is a sum graph, then the subgraph  $G$  is called a *unit graph*. In this paper it is shown that there exist graphs of every order  $n$  and size  $m$  whose sum number is  $O(n)$ . Further, it is shown that for every integer  $m$  satisfying  $\lfloor n^2/4 \rfloor < m \leq \binom{n}{2}$  there exists no unit graph, while for each  $m$  such that  $n - 1 \leq m \leq \lfloor n^2/4 \rfloor$  there exists at least one unit graph. Methods of proof are constructive.

## 1 INTRODUCTION

Sum graphs (defined in the Abstract) were introduced by Harary [Har88,Har89]. Hao [Hao89] showed that a graph of order  $n$  is a sum graph if and only if its size  $m \leq (\binom{n}{2} - \lfloor n/2 \rfloor)/2$ ; for a given graph  $G$ , he also established a lower bound on  $\sigma(G)$  in terms of the degree sequence of  $G$ . Gould and Rödl [GR89] derived complex upper and lower bounds on  $\sigma(G)$ , expressed in terms of the order  $n$  and size  $m$  of  $G$ . In particular, their results show that there exist classes  $\mathcal{G}$  of graphs such that over all  $G \in \mathcal{G}$ ,  $\sigma(G) \in \Omega(n^2)$ . However, they provide no method for the construction of such graphs, and the available constructive results all relate to special graphs  $G$  of small sum number; that is, such that  $\sigma(G) \in O(n)$ . In particular, Ellingham [E89] has shown that for every non-trivial tree  $T$ ,  $\sigma(T) = 1$ ; Bergstrand et al. [BHHJKW89] that for every complete graph  $K_n$ ,  $n \geq 4$ ,  $\sigma(K_n) = 2n - 3$ ; Hartsfield and Smyth [HS89] that for every complete bipartite graph  $K_{p,q}$ ,  $2 \leq p \leq q$ ,  $\sigma(K_{p,q}) = \lceil (3p + q - 3)/2 \rceil$ .

In this paper new results for graphs of small sum number are established; these results make some progress toward resolving an open problem posed by Harary [Har88]: the characterization of unit graphs.

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$

Section 2 uses methods similar to those of Gould and Rödl, but which are however constructive, to show that there exists a connected graph  $G$  of given order  $n$  and size  $m$  ( $n - 1 \leq m < \binom{n}{2} - 1$ ) such that  $\sigma(G) < \sigma(K_n) = 2n - 3$ . In Section 3 unit graphs of order  $n$  are considered. It is shown that for  $m > \lfloor n^2/4 \rfloor$  no unit graph exists, while for every integer  $m$  satisfying  $n - 1 \leq m \leq \lfloor n^2/4 \rfloor$ , there exists a unit graph  $G$  of order  $n$  and size  $m$ .

Generally, in order to simplify notation, and where no ambiguity results, vertices of sum graphs will be referenced by their label under  $L$ .

## 2 CONSTRUCTING GRAPHS OF SMALL SUM NUMBER

In this section we show how to construct connected graphs  $G$  of given order  $n$  and size  $m$  for which  $\sigma(G) \in O(n)$ .

For a sum graph  $G_r = (V, E)$ , denote by  $\{v_1, v_2, \dots, v_r\}$  the labels of the vertices of  $\overline{K}_r$ , where  $v_1 < v_2 < \dots < v_r$ . For each  $v_j, 1 \leq j \leq r$ , let  $\mu_j$  denote the number of edges  $(x, y) \in E$  such that  $x + y = v_j$ . Then  $\mu_j$  is called the *multiplicity* of  $v_j$ . Now consider the special case in which  $G_r = K_n \cup \overline{K}_{2n-3}$ , for some integer  $n \geq 4$ . Then, as shown in [BHHJKW89], a correct labelling of  $G_r$  is achieved by assigning labels to the vertices of  $K_n$  as follows:

$$x_i = 4i - 3, 1 \leq i \leq n.$$

Hence the corresponding labels  $v_j$  of  $\overline{K}_{2n-3}$  are

$$v_j = 4j + 2, 1 \leq j \leq 2n - 3.$$

Let us call this the *standard labelling* of  $K_n \cup \overline{K}_{2n-3}$ . It is then straightforward to establish the following result:

**Lemma 2.1** *For every integer  $n \geq 4$ , and for every positive integer  $j \leq n - 1$ , the standard labelling of  $K_n \cup \overline{K}_{2n-3}$  yields multiplicities*

$$\mu_j = \mu_{2n-j-2} = \lceil j/2 \rceil. \quad \square$$

It is worth noting that in fact the result of Lemma 2.1 holds for *any* correct labelling of  $K_n \cup \overline{K}_{2n-3}$  [AHS90]. Multiplicity patterns for the first few values of  $n$  are shown in Table 2.1.

Multiplicities of $\overline{K}_{2n-3}$ , $4 \leq n \leq 8$	
$n$	$\mu_j, 1 \leq j \leq 2n - 3$
4	{1, 1, 2, 1, 1}
5	{1, 1, 2, 2, 2, 1, 1}
6	{1, 1, 2, 2, 3, 2, 2, 1, 1}
7	{1, 1, 2, 2, 3, 3, 3, 2, 2, 1, 1}
8	{1, 1, 2, 2, 3, 3, 4, 3, 3, 2, 2, 1, 1}

Table 2.1

Observe that if a vertex labelled  $v_j$  is removed from  $\overline{K}_{2n-3}$ , and if at the same time every edge  $(x, y)$  for which  $x + y = v_j$  is removed from  $K_n$ , the resulting graph (a subgraph of  $K_n \cup \overline{K}_{2n-3}$ ) will still be a sum graph. This process (removal of a single vertex  $v_j$  and all corresponding edges) applied to an arbitrary sum graph  $G_r = (V, E)$  is called a *reduction* and written

$$\phi_j : G_r \rightarrow G_{r-1},$$

where  $G_{r-1} = (V - \{v_j\}, E')$  and  $|E'| = |E| - \mu_j$ . We shall show that for every nonnegative integer  $m \leq \binom{n-1}{2}$ , exactly  $m$  edges can be removed from the sum graph  $G_{2n-3}^* = K_n \cup \overline{K}_{2n-3}$  by a sequence of reductions, yielding a graph which consists of a single connected component together with isolated vertices, and which is therefore a sum graph of small sum number.

First consider the case  $m = \binom{n-1}{2}$ , so that the reduced graph contains  $\binom{n}{2} - \binom{n-1}{2} = n - 1$  edges. Suppose that the vertices  $v_1, v_2, \dots, v_{n-2}$  and  $v_{n+1}, v_{n+2}, \dots, v_{2n-3}$  and their corresponding edges are removed, leaving only  $v_{n-1}$  and  $v_n$  from among the original isolated vertices. Since by Lemma 2.1,

$$\mu_{n-1} + \mu_n = \lceil \frac{n-1}{2} \rceil + \lceil \frac{n-2}{2} \rceil = n - 1,$$

the reduced graph  $G_2^*$  certainly contains the prescribed number of edges. Further, let  $E_{n-1}$  and  $E_n$  denote the edges of  $G_2^*$  corresponding to  $v_{n-1}$  and  $v_n$ , respectively; then both  $E_{n-1}$  and  $E_n$  are matchings of  $K_n$ , and in fact  $E_{n-1}$  is a maximum matching while  $|E_n| \geq |E_{n-1}| - 1$ . Since moreover  $E_{n-1} \cap E_n = \emptyset$ , it must be true that  $G_2^* - \{v_{n-1}, v_n\}$  is connected, and we have

**Lemma 2.2** *Suppose that the sum graph  $G_{2n-3}^*$  is repeatedly reduced by the reductions  $\phi_j$  for every integer  $j = 1, 2, \dots, n-2, n+1, n+2, \dots, 2n-3$ . Then the reduced graph  $G_2^*$  consists of a path  $P_n$  on  $n$  vertices together with two isolated vertices  $v_{n-1}$  and  $v_n$ .  $\square$*

It follows from Lemma 2.2 that, provided  $v_{n-1}$  and  $v_n$  are not removed, any repeated reduction of  $G_{2n-3}^*$  will always yield a single connected component together with a subset of the original set  $\overline{K}_{2n-3}$  of isolated vertices. We therefore consider a reduction strategy which leaves  $v_{n-1}$  and  $v_n$  intact.

In general, we seek reductions which transform the sum graph  $G_{2n-3}^*$  into another sum graph  $G_{2n-3-k}^*$  by removing  $k$  vertices and  $m$  edges. Each such transformation can be characterized by a set  $M$  of multiplicities removed from the original set  $\{\mu_1, \mu_2, \dots, \mu_{2n-3}\}$ . For such a set  $M$ ,  $k = |M|$  is called the *vertex remove* and  $m = \sum_{\mu \in M} \mu$  is called the *edge remove*. For example,  $M = \{1\}$  would indicate the removal of  $k = 1$  vertex and  $m = 1$  corresponding edge;  $M = \{1, 1, 1, 2\}$  would indicate the removal

of  $k = 4$  vertices and  $m = 5$  corresponding edges.

In order to uniquely identify the sets  $M$  of removed multiplicities, they are subscripted according to "class" and by an ordinal within each class. For example, the sets corresponding to  $m = 1, 2, \dots, 6$  are written as follows:

$$\begin{aligned} M_{3,1} &= \{1\}; \\ M_{4,1} &= \{1, 1\}, M_{4,2} = \{1, 1, 1\}; \\ M_{5,1} &= \{1, 1, 1, 1\}, M_{5,2} = \{1, 1, 1, 1, 2\}, M_{5,3} = \{1, 1, 1, 1, 2\}. \end{aligned}$$

In general, the  $h^{\text{th}}$  class will contain  $h-2$  sets  $M_{h,1}, M_{h,2}, \dots, M_{h,h-2}$ , where for  $M_{h,h-2}$ ,

$$m_{h,h-2} = \sum_{\mu \in M_{h,h-2}} \mu = \binom{h-1}{2}$$

is the largest number of edges which can be removed from  $K_h$  without necessarily disconnecting it. We state a formal procedure for the iterative computation of the sets in each class based upon the sets in the previous class:

#### Procedure CONSTRUCT-MULTIPLICITY-SETS

**Step 1** {Initialize.}  $M_{3,1} = \{1\}$ .  
**Step 2** {Iterate.} **For**  $h = 4, 5, \dots, n$  **do**  
    **Step 2.1**  $h' \leftarrow \lceil h/2 \rceil - 1$ .  
    **Step 2.2** **If**  $h$  **is even, then**  
        **for**  $i = 1, 2, \dots, h'$  **do**  
             $M_{h,i} \leftarrow M_{h-1,h'+i-1} \cup \{h'\}$ ;  
             $M_{h,h'+i} \leftarrow M_{h,i} \cup \{h'\}$ .  
    **Step 2.3** **If**  $h$  **is odd, then**  
         $M_{h,h'} \leftarrow M_{h-1,h-3} \cup \{h'\}$ ;  
        **for**  $i = 1, 2, \dots, h' - 1$  **do**  
             $M_{h,i} \leftarrow M_{h-1,h'+i-1} \cup \{h' - 1\}$ ;  
             $M_{h,h'+i} \leftarrow M_{h,i} \cup \{h'\}$ .

This procedure generates a sequence  $\mathcal{M}$  of  $\binom{n-1}{2}$  sets

$$M_{3,1}, M_{4,1}, M_{4,2}, M_{5,1}, \dots, M_{n,1}, M_{n,2}, \dots, M_{n,n-2}$$

in strictly increasing order of edge remove from  $m = 1$  ( $M_{3,1}$ ) to  $m = \binom{n-1}{2}$  ( $M_{n,n-2}$ ). It is not difficult to see that every set of  $\mathcal{M}$  is without loss of generality a subset of the multiplicities

$$\{\mu_1, \mu_2, \dots, \mu_{n-2}, \mu_{n+1}, \mu_{n+2}, \dots, \mu_{2n-3}\}.$$

Thus by Lemma 2.2 the sum graph associated with every element of  $\mathcal{M}$  has a connected component of order  $n$  and a set of isolated vertices of cardinality less than  $2n-3$ . Indeed, the exact number of isolated vertices may be computed for each element of  $\mathcal{M}$  as a function of the number of removed edges:

**Theorem 2.1** *For every integer  $n \geq 4$  and every positive integer  $m \leq \binom{n-1}{2}$ , let  $\mathcal{K}_{n \setminus m}$  denote the set of all graphs formed by removing exactly  $m$  edges from  $K_n$ . Then there exists  $G \in \mathcal{K}_{n \setminus m}$  such that*

$$\sigma(G) \leq 2n - 1 - 2i - \lfloor m/i \rfloor,$$

*where  $i$  is the least integer such that  $m \leq \binom{2i+1}{2}$ .*

**Proof** The proof is by induction on  $n$ , based on the multiplicity sets generated by Procedure CONSTRUCT-MULTIPLICITY-SETS. The demonstration is laborious but straightforward, and is omitted.  $\square$

The condition on  $i$  in Theorem 2.1 may be expressed algebraically as

$$i = \lceil (\sqrt{8m+1} - 1)/4 \rceil.$$

Znám [Z91] has discovered an interesting alternative formulation of this result. Let  $i$  be the greatest integer such that  $m \geq 4\binom{i}{2}$ ; let  $j$  be the greatest non-negative integer such that  $m \geq 4\binom{i}{2} + ij$ . Then there exists  $G \in \mathcal{K}_{n \setminus m}$  such that

$$\sigma(G) \leq 2n + 1 - 4i - j.$$

### 3 UNIT GRAPHS

In this section we use the properties of unit graphs to improve upon Theorem 2.1 — to show, in fact, that for every integer  $n \geq 2$  and every integer  $m$  satisfying  $n-1 \leq m \leq \lfloor n^2/4 \rfloor$ , there exists a unit graph of order  $n$  and size  $m$ .

**Lemma 3.1** *Let  $G = (V, E)$  denote a unit graph, and let  $G_1 = G \cup \overline{K}_1$  denote the corresponding sum graph. Suppose that  $G_1$  is correctly labelled. Then the vertex of  $V$  of greatest label  $u$  has degree one.*

**Proof** Suppose  $u$  has degree at least two. Then  $u$  is adjacent to two distinct vertices which we suppose to be labelled  $x_1$  and  $x_2$ . Hence there must exist vertices of  $G_1$  labelled  $v_1 = x_1 + u > u$  and  $v_2 = x_2 + u > u$ . Since neither of these vertices can belong to  $V$ , they must both be isolated, contradicting the assumption that  $G$  is a unit graph. Hence  $u$  must have degree one.  $\square$

**Lemma 3.2** *Let  $d_i$ ,  $1 \leq i \leq n$ , denote the degrees of the vertices of a unit graph  $G$ , where  $d_1 \leq d_2 \leq \dots \leq d_n$ . Then  $d_i \leq i$ .*

**Proof** A consequence of a result of Hao [Hao89] that

$$\sigma(G) > \max_{1 \leq i \leq n} (d_i - i). \quad \square$$

**Lemma 3.3** *There exists a unit graph of order  $n$  containing a clique on  $\nu$  vertices if and only if  $\nu \leq \lfloor n/2 \rfloor + 1$ .*

**Proof** Since  $K_\nu$  contains  $\nu$  vertices of degree  $\nu - 1$ , it is easy to see that if the graph  $G$  of order  $n$  contains  $K_\nu$ , then the vertex of  $(n - \nu + 1)^{th}$  largest degree must have degree at least  $\nu - 1$ . That is, in the symbolism of Lemma 3.2,

$$d_{n-\nu+1} \geq \nu - 1.$$

If moreover  $\nu > \lfloor n/2 \rfloor + 1$ , then

$$d_{n-\nu+1} \geq \nu - 1 > n - \nu + 1,$$

and Lemma 3.2 implies that  $G$  cannot be a unit graph.

To prove the converse, consider the sum graph  $G_r$  generated by vertices labelled consecutively  $1, 2, \dots, n + 1$ . The subgraph  $G \subset G_1$  induced by  $V = \{1, 2, \dots, n\}$  is then in fact a unit graph, and  $G$  contains a clique of order  $\lfloor n/2 \rfloor + 1$ .  $\square$

**Theorem 3.1** *No unit graph of order  $n$  has size  $m > \lfloor n^2/4 \rfloor$ .*

**Proof** The unit graph on vertices  $V = \{1, 2, \dots, n\}$ , together with isolated vertex  $n + 1$ , forms a sum graph of size

$$m = \binom{\lfloor n/2 \rfloor + 1}{2} + \binom{n - \lfloor n/2 \rfloor}{2},$$

which after some manipulation becomes

$$m = \lfloor n^2/4 \rfloor.$$

By Lemmas 3.2 and 3.3, there exists no unit graph of order  $n$  of larger size.  $\square$

Theorem 3.1 may also be deduced from Hao's result, quoted in the Introduction, giving an upper bound on the size of a sum graph.

**Theorem 3.2** *For every integer  $n \geq 2$  and for every integer  $m$  satisfying  $n - 1 \leq m \leq \lfloor n^2/4 \rfloor$ , there exists a unit graph of order  $n$  and size  $m$ .*

**Proof** Consider a sum graph  $G_1 = (V, E)$  generated by  $V = \{1, 2, \dots, n + 1\}$ .  $G_1$  contains a unit graph  $G^{(0)}$  of order  $n$  and size  $m = \lfloor n^2/4 \rfloor$ . We show first that by relabelling some of the vertices of  $V$ , unit graphs  $G^{(i)}$  of order  $n$  and size  $m - i$  can be constructed, for every positive integer  $i \leq \lfloor n/2 \rfloor$ .

Let  $k = \lfloor n/2 \rfloor$  and recall that  $G^{(0)}$  contains  $K_{k+1}$  as a subgraph. Hence separate  $V$  into subsets  $V_1 = \{1, 2, \dots, k + 1\}$  and  $V_2 = \{k + 2, k + 3, \dots, n + 1\}$ . Observe that the edges represented by any vertex label  $v \in V_2$  may be counted in two classes:  $m_1 = m_1(v)$  edges of  $K_{k+1}$  and  $m_2 = m_2(v)$  edges  $(x, y)$  which join  $x \in V_1$  to  $y \in V_2$ .

Suppose that  $k$  is even. Then labels  $v \in V_2$  correspond to these two classes of edge as shown in the following table (the last line occurs only in the case that  $n = 2k + 1$ ):

$v$	$m_1$	$m_2$
$k + 2$	$k/2$	$0$
$k + 3$	$k/2$	$1$
$k + 4$	$\frac{k}{2} - 1$	$2$
$\cdot$	$\cdot$	$\cdot$
$\cdot$	$\cdot$	$\cdot$
$\cdot$	$\cdot$	$\cdot$
$2k$	$1$	$k - 2$
$2k + 1$	$1$	$k - 1$
$2k + 2$	$0$	$k$

Observe that replacing  $V_2$  by another set  $V'_2 = \{v_1, v_2, \dots, v_{n-k+1}\}$  of distinct positive integers, where  $k + 1 < v_1 < v_2 < \dots < v_{n-k}$ , will not affect the total number of edges counted by  $m_2$ , provided  $v_{n-k} - v_1 \leq k + 1$ . But the total number of edges counted by  $m_1$  can be reduced by exactly  $i$  if label  $k + 2i$  of  $V_2$  is replaced by  $2k + 3$ ,  $1 \leq i \leq k/2$ , while all other labels are unchanged. Then this relabelling yields unit graphs  $G^{(i)}$  of order  $n$  and size  $m - i$ , for every positive integer  $i \leq k/2$ . If now label  $k + 2$  of  $V_2$  is replaced by  $2k + 3$ , and if in addition label  $k + 2i + 1$  is replaced by  $2k + 4$ , then the total number of edges counted by  $m_1$  will be reduced by  $i + k/2$ , where  $1 \leq i \leq k/2$ . We see then that unit graphs  $G^{(i)}$  of order  $n$  and size  $m - i$  can be constructed for every positive integer  $i \leq k = \lfloor n/2 \rfloor$ . But since

$$\lfloor n^2/4 \rfloor - \lfloor (n-1)^2/4 \rfloor = \lfloor n/2 \rfloor,$$

it follows that unit graphs of order  $n$  and size  $m$  can be formed, for every integer  $m$  satisfying

$$\lfloor (n-1)^2/4 \rfloor \leq m \leq \lfloor n^2/4 \rfloor.$$

The same conclusion is reached, by an almost identical argument, when  $k$  is odd.

To complete the proof, observe now that by the above result a single vertex and a path of length one can be added to a unit graph of order  $n - 1$  and size

$$\lfloor (n-2)^2/4 \rfloor \leq m \leq \lfloor (n-1)^2/4 \rfloor,$$

thus yielding a unit graph of order  $n$  and size  $m + 1$ . More generally,  $s \geq n - 4$  vertices and a path of length  $s$  can be added to a unit graph of order  $n - s$  and size  $m$  satisfying

$$\lfloor (n-s-1)^2/4 \rfloor \leq m \leq \lfloor (n-s)^2/4 \rfloor,$$

thus yielding a unit graph of order  $n$  and size  $m + s$ . If we consider consecutive values  $s = 1, 2, \dots, n - 4$  and observe that for  $s = n - 3$ ,

$$n - 1 = \lfloor (n-s)^2/4 \rfloor + (n-s),$$

it follows that unit graphs of order  $n$  and every size  $m$  satisfying

$$n - 1 \leq m \leq \lfloor n^2/4 \rfloor$$



can be constructed, as required.  $\square$

It should be noted that Lemma 3.2 and Theorem 3.2 give rise to no converse. That is, there exist graphs whose degree sequence satisfies the conditions of Lemma 3.2 and whose size satisfies the condition of Theorem 3.2, but which are not unit graphs. An example is the star on six vertices with three additional edges joining four of the points of the star:  $V = \{v_1, v_2, \dots, v_6\}$  and  $E = \{v_1v_i, 2 \leq i \leq 6; v_2v_3, v_3v_4, v_4v_5\}$ .

## REFERENCES

- [AHS90] N. Achuthan, N. Hartsfield, W. F. Smyth, *work in progress*.
- [BHHJKW89] D. Bergstrand, F. Harary, K. Hodges, G. Jennings, L. Kuklinski, J. Wiener, **The sum number of a complete graph**, to appear.
- [E89] M. N. Ellingham, **Sum graphs from trees**, submitted for publication.
- [GR89] R. Gould, V. Rödl, **Bounds on the number of isolated vertices in sum graphs**, to appear.
- [Hao89] T. Hao, **On sum graphs**, *Journal of Combinatorial Mathematics & Combinatorial Computing* 6 (1989) 207-212.
- [Har88] F. Harary, **Sum graphs and difference graphs**, presentation at the Nineteenth Southeastern Conference on Combinatorics, Graph Theory & Computing.
- [Har89] F. Harary, **Sum graphs: some open questions**, presentation at the Second International Conference on Graph Theory, Combinatorics, Algorithms & Applications.
- [HS89] N. Hartsfield, W. F. Smyth **The sum number of complete bipartite graphs**, to appear.
- [Z91] S. Zná́m, *private communication*.

## ACKNOWLEDGEMENT

This work was supported in part by Grant No. A8180 of the Natural Sciences & Engineering Research Council of Canada.